

Sorghum and Western Style Breads

Subjects: Food Science & Technology

Submitted by:  Rubina

Rumler

Definition

Due to the extreme weather conditions, caused by the climate change, the usual wheat harvest yield and quality in the Western countries were difficult to maintain in the past few years. The altered wheat quality was primarily characterized by a rising protein content. The application of high protein wheat flours in baking products leads to baking difficulties due to its elastic dough behavior. As these issues will further face the Western cereal cultivation, heat resistant cereals, like sorghum, are attracting increasing interest. A partial substitution of wheat with sorghum might offer one possible solution to address the current challenging situation. To introduce sorghum in the Western cereal and baking industry, the grain and its unique chemical and rheological properties need to be more widely promoted. Until now, several authors have conducted studies in order to emphasize the high potential of sorghum. The aim of the present review is to broaden the current knowledge of the chemical, rheological and baking properties of sorghum in comparison to wheat.

1. Introduction

The wheat grain industry already experienced the effects of the climate change in the past few years. Dry climate with low rainfall was responsible for crop failures of wheat. According to a study, 60% of the current wheat growing areas could be affected by insufficient water supply by the end of this century ^[1]. However, due to the climate change not only wheat harvest losses happened, but also the wheat protein contents increased in the last few years ^[2]. A suitable solution must be found to compensate for the loss of wheat in the grain and baking industry and to deal with the altered quality. A possible way to address these challenges could be the use of food additives or to import wheat from other countries. A more sustainable, customer-friendly and innovative option could be the use of regional grains, preferable gluten-free grains, in wheat products. A partial replacement by gluten-free cereals might offer an option in order to compensate for the changed protein content of wheat. Sorghum, which is gluten-free and heat resistant, presents a promising grain in this context. The use of sorghum in wheat products could be a possible future solution to counteract the high climate-induced protein content of wheat. In addition, end products consisting of sorghum can gain valuable nutrients ^[3].

In order to introduce sorghum into the Western diet, the nutritional and technological challenges have to be considered. Mainly the altered protein composition, due to the absence of gluten results in changed dough rheological parameters and end-product qualities ^{[4][5][6][7]}. However, like other cereals sorghum contains tannins, trypsin inhibitors and phytic acids, which are known to impair protein digestion and mineral absorption ^[8]. Sometimes these values are higher than in other cereals. In addition, sorghum prolamins, called kafirins, are known to have a low digestibility, especially in prepared foods ^[9], which is mainly due to their crosslinking properties upon processing. Sorghum possesses higher crosslinking properties than wheat for example. However, if this protein cross-linking could in turn offer beneficial viscoelastic properties in food is still unknown ^[10].

2. Classification and Morphology of Sorghum

Although sorghum is not well known in the West, it is the fifth most produced cereal worldwide. In Africa sorghum is considered to be the most important cereal. Outside of Africa production takes place mainly in India, USA, Australia, and Argentina, whereby sorghum in the USA and in Australia is mostly used as animal feed. Five races are known under sorghum: bicolor, kafir, caudatum, durra and guinea. All combinations from those mentioned races are called intermediate races. Of all races, Sorghum bicolor Moench is mainly cultivated ^[11]. According to Taylor there are different classification systems of sorghum.

Some of them seem to be complex. However, like wheat and maize sorghum is classified into the grass family Poaceae as can be seen in Figure 1. Sorghum belongs, like maize, to the subfamily Panicoideae and to the tribe Andropogoneae [12]. Due to the botanical relationship sorghum is often compared to maize.




Figure 1. Family tree of sorghum. Modified according to Taylor [12].

The structure of a sorghum kernel is comparable to the structure of a corn kernel, but in general, a sorghum grain is smaller. Awika et al. [13] determined the diameters of different sorghum kernels to range between 1.7-2.7 mm, and the thousand kernel weight of different varieties was between 15.2-38.8 g. The extent of the pericarp thickness and the presence of a testa is variety dependent [13]. Both sorghum and maize, have an endosperm with a flourey and a corneous part [14]. Sorghum kernels appear in color nuances of white, red, brown and black (Figure 2).



Figure 2. Sorghum field in Horsching, Austria (©Rumler, 2020).

However, the color of the pericarp is not always indicative of the appearance of sorghum flour. For example, a sorghum kernel which owns a red pericarp, may have a light endosperm color. Therefore, if special food colors are required, the insertion of sorghum fraction flours may be useful in order to achieve appropriate food appearances [15].

3. Chemical and Nutritional Properties of Sorghum

3.1. Protein

The protein content in sorghum can be comparable to the protein content of wheat [5][16]. However, Tasie and Gebreyes [17] found a wide range of 8.20-16.48% between different sorghum whole grain varieties. Attempts have often been made to identify correlations between nutrients and the sorghum kernels color.

Whether the protein content differs between red and white sorghum is still poorly understood. A research of Galan et al. [18] detected a higher protein content in red whole grain sorghum flour, while Yousif et al. [19] and Srichuwong et al. [20] figured out a higher protein content in whole grain white sorghum flour.

In sorghum protein the predominant fraction is the prolamin fraction, which is called kafirin. According to a study almost half of the sorghum protein belonged to the kafirin fraction (47.7%), followed by the glutelin fraction (24.1%) and the albumin and globulin fraction (18.8%) [21]. Regarding the kafirins, four types of kafirins, occurring in form of protein bodies, can be mentioned: alpha-, beta-, gamma-, and delta-kafirin. Alpha-kafirins are the most common kafirins found in sorghum [10][22]. Detailed amino acid analyses have been carried out by Mohapatra et al. [23]. The most common amino acids found in whole grain sorghum were glutamic acid (44.38 mg/100 g protein), arginine (28.34 mg/100 g protein), aspartic acid (20.50 mg/100 g protein), leucine (16.50 mg/100 g protein) and glycine (16.22 mg/100 g protein) [23]. The limiting acid is lysine, as for most other true cereals [24].

Sorghum proteins have been gaining much attention due to their ability to crosslink, in particular upon cooking. Authors believe that this kafirin cross linking is connected to low protein digestibility of sorghum. In this context a comparison of the digestion of kafirins and zeins was performed by Emmambux and Taylor [9]. The authors analyzed the digestibility of sorghum and maize meal as well as the digestibility of aqueous tertiary butanol extracted kafirin and zein. In order to evaluate the influence of heat on sorghum and maize digestibility, the samples were undertaken a boiling and a pressure cooking treatment. Already in the untreated state, the maize meal and extracted zeins showed a higher digestion rate than the sorghum meal and kafirins. However, after boiling and pressure cooking this difference was even more obvious. Authors detected that sorghum kafirins (obtained from decorticated sorghum kernels) were more polymerized by disulfide bridges during the cooking process than the zeins [9]. Same authors further pointed out, that sorghum proteins were not only able to form complexes between each other, but also had the ability to connect with condensed tannins and tannic acids [25]. This leads to the suggestion that sorghum kernel dehulling or the usage of condensed tannin free sorghum varieties might offer a higher digestibility. However, crosslinking interactions According to Hamaker and Bugusu [10] sorghum proteins are embedded within protein bodies and therefore hard to access, but on the other hand they may show viscoelastic behavior in food. However, several questions remain unanswered at present. Fundamental investigations are still needed in order to determine the properties of kafirins.

3.2. Carbohydrate

Carbohydrate amounts between 67.5–76.4% were found in different whole grain sorghum varieties from Ethiopia [17]. Pontieri et al. [26] found that five varieties of refined sorghum, which were grown in Italy, the USA or Uganda, had an average carbohydrate content of 75.23% in dry matter. Starch is the main component of carbohydrates and is located in the endosperm. Srichuwong et al. [20] reported a starch content of 72.2% and 73.8% in dry matter in white and red whole grain sorghum, with amylose contents of 25.8% and 24.6% respectively. However, there is a wide range of amylose content in sorghum varieties, and even waxy varieties, which have high amylopectin contents occur in the wild [27]. The diameter of starch granules was measured to be between 5–35 µm and they appeared as polygonal and spherical [20], but also more unusual granules shapes, which researchers called “doughnut-shaped”, were occasionally discovered in sorghum kernels [28]. Miafo et al. [29] determined the content of free sugars in sorghum refined flour of 30.35 mg/g sorghum flour. Sucrose made up the largest proportion (12.42 mg/g), followed by maltose (4.25 mg/g) and fructose (4.22 mg/g). The usage of sorghum flour might offer a sugar reduction possibility in bakery products as it has already a unique sweet taste itself. As humans are tending more and more towards balanced diets, sensory trials are required in order to state if sorghum can partially replace household sugar.

Whole sorghum contains a notable amount of dietary fibers, which is mostly located in the outer layers of the sorghum kernel. A study carried out in 2012 showed a total dietary fiber content between 9.13–15.09 g/100 g that has been detected in eight sorghum varieties, whereby the soluble fiber part ranged

between 0.15–0.88 g/100 g [30]. Authors compared the fiber content determined in whole grain sorghum, dehulled sorghum and sorghum bran. It was found that the bran fraction had the highest value with 41.38 g/100 g, followed by 26.34 g/100 g in whole grain sorghum and 11.53 g/100 g in hulled sorghum [31].

3.3. Fat

Whole grain sorghum flour is known for its high fat content (3.5%) especially compared to wheat (2% in whole wheat) [3]. Fat is mostly located in the sorghum germ (Taylor, 2003). The most common fatty acids are linoleic acid (18:2), followed by oleic acid (18:1) and palmitic acid (16:0), with average contents of 43.86%, 37.98% and 13.07% in refined sorghum flour, respectively [26]. The ratio of saturated to unsaturated fatty acids was found to be 0.27–0.37 [32]. Knowledge is lacking if high fat contents in sorghum flour play a crucial role within shelf life. However, [33] were able to extend the storage time of sorghum flour from 15 days to several months by means of heat treatments. As sorghum is mostly harvested in Africa, knowledge about sorghum milling in the Western countries was rare until very recently. It was not clear whether a high fat content in sorghum brings difficulties within a milling process. However, in 2021 authors concluded that sorghum flour can be provided via commercial wheat mills [15].

3.4. Micronutrients

Sorghum is known for having a valuable micronutrient profile. Istianah et al. [5] determined 1.64% ash in dehulled sorghum. However, it seems that micronutrient contents vary to a great extent among different varieties [17][34]. Tasie and Gebreyes (2020) identified ash contents between 1.1–2.3% in dry matter of different whole grain sorghum flours from Ethiopia. Regarding the micronutrients, amounts of phosphorus ranged between 112.5–327.7 mg/100 g, of sodium between 2.2–6.2 mg/100 g, of magnesium between 62.0–207.5 mg/100 g and of calcium between 9.5–67.2 mg/100 g [17]. Varieties with a particularly high mineral content exist, like for example the variety Tabat with phosphorus quantities of 350 mg/100 g, sodium of 14.5 mg/100 g, magnesium of 329 mg/100 g and calcium of 24.5 mg/100 g extracted sorghum flour [35].

3.5. Secondary Plant Products

Although people in the Western hemisphere are increasingly turning to alternative cereals with valuable ingredients, like amaranth or buckwheat, sorghum is still unknown. However, sorghum also offers a lot of promising health-promoting components. A review pointed out that especially sorghum polyphenols, as they have antioxidative effects, can help preventing chronic diseases, like improvement of insulin sensitivity, reduction of fat accumulation, or reduction of mild chronic inflammation [36]. Therefore, many research studies are interested in measuring the polyphenol content of sorghum, regardless of the research question. It can be stated with certainty, that polyphenols vary enormously among the biodiversity of sorghum. However, despite this interest, little research tended to focus on the context between polyphenols and sorghum colors. While researchers found that the polyphenol content of whole grain wheat (92.4 mg/100 dry matter) and whole grain white sorghum (95.2 mg/g dry matter) was similar, whole grain red sorghum had a significantly higher content of 203.5 mg/g dry matter [20]. Higher polyphenol levels in whole grain red sorghum flour than in whole grain white sorghum flour were confirmed by further studies [19][37]. If polyphenols are determined in more detail, flavonoids and phenolic acids are mainly mentioned. Wide ranges of phenolic acids were reviewed in the past. However, in red sorghum, as well as in white sorghum, ferulic acid was found to be the most predominant phenolic acid [38]. Among all flavonoids, special attention has been paid to the tannins, chemically named proanthocyanidins. Tannins are known to reduce sorghum digestibility [8]. There are low tannin and high tannin sorghum varieties, both naturally occurring. Tasie and Gebreyes [17] examined 35 whole grain sorghum flours from Ethiopia and detected tannin contents between 1.3–3337.2 mg/100 g dry matter. They also reported that condensed tannins and tannin acids in sorghum can form complexes with kafirins. Such a complex formation is one reason to lower sorghum protein digestibility. There was no evidence that other phenolic compounds, such as catechin or flavonoids, can bind kafirin [9]. A decreased protein digestion of tannin containing sorghum compared to non-tannin sorghum was also detected by Wedad et

al. [39]. As condensed tannins are not present in wheat [14], it is questionable if using tannin rich sorghum varieties in a wheat blend leads to nutritional advantages. Besides phenolic acids and flavonoids, carotenoids also offer antioxidant properties. Luteolin, zeaxanthin and beta-carotene were found in amounts of 122.3 mg/kg, 25.2 mg/kg and 27.0 mg/kg, respectively, in white sorghum [40].

References

1. Trnka, M.; Feng, S.; Semenov, M.A.; Olesen, J.E.; Kersebaum, K.C.; Rötter, R.P.; Semerádová, D.; Klem, K.; Huang, W.; Ruiz-Ramos, M. Mitigation efforts will not fully alleviate the increase in water scarcity occurrence probability in wheat-producing areas. *Sci. Adv.* 2019, 5, eaau2406.
2. Gagliardi, A.; Carucci, F.; Masci, S.; Flagella, Z.; Gatta, G.; Giuliani, M.M. Effects of genotype, growing season and nitrogen level on gluten protein assembly of durum wheat grown under mediterranean conditions. *Agronomy* 2020, 10, 755.
3. Salim-ur-Rehman Ahmad, M.; Bhatti, I.; Shafique, R.; Mueen-ud-Din, G.; Murtaza, M. Effect of pearling on physico-chemical, rheological characteristics and phytate content of wheat-sorghum flour. *Pak. J. Bot.* 2006, 38, 711–719.
4. Rizk, I.R.; Hemat, E.E.; Gadallah, E.; Abou-Elazm, M.; Bedeir, H. Quality characteristics of sponge cake and biscuit prepared using composite flour. *J. Agric. Soc. Sci.* 2015, 23, 537–547.
5. Istianah, N.; Ernawati, L.; Anal, A.K.; Gunawan, S. Application of modified sorghum flour for improving bread properties and nutritional values. *Int. Food Res. J.* 2018, 25, 166–173.
6. Ognean, C.-F. Technological and sensorial effects of sorghum addition at wheat bread. *Agric. Food* 2015, 3, 209–217.
7. Sibanda, T.; Ncube, T.; Ngoromani, N. Rheological properties and bread making quality of white grain sorghum-wheat flour composites. *Int. J. Food Sci. Nutr. Eng.* 2015, 5, 176–182.
8. Osman, M.A.; Gasseem, M. Effects of domestic processing on trypsin inhibitor, phytic, acid, tannins and in vitro protein digestibility of three Sorghum varieties. *Int. J. Agric. Technol.* 2013, 9, 1187–1198.
9. Emmambux, M.N.; Taylor, J.R.N. Properties of heat-treated sorghum and maize meal and their prolamin proteins. *J. Agric. Food Chem.* 2009, 57, 1045–1050.
10. Hamaker, B.R.; Bugusu, B.A. Workshop on the Proteins of Sorghum and Millets: Enhancing Nutritional and Functional Properties for Africa [CD]; Scientific Research: Pretoria, South Africa, 2003.
11. Cruickshank, A. Sorghum grain, its production and uses: Overview. In *Encyclopedia of Food Grains*, 2nd ed.; Wrigley, C., Corke, H., Seetharaman, K., Faubion, J., Eds.; Academic Press: Oxford, UK, 2015; pp. 153–158.
12. Taylor, J.R.N. Sorghum and millets: Taxonomy, history, distribution, and production. In *Sorghum and Millets: Chemistry, Technology, and Nutritional Attributes*, 2nd ed.; Taylor, J., Duodu, K.G., Eds.; Elsevier: Amsterdam, The Netherlands, 2019; pp. 1–21.
13. Awika, J.M.; McDonough, C.M.; Rooney, L.W. Decorticating sorghum to concentrate healthy phytochemicals. *J. Agric. Food Chem.* 2005, 53, 6230–6234.
14. Taylor, J.R.N. Overview: Importance of sorghum in Africa. In *Proceedings of the Workshop on the Proteins of Sorghum and Millets: Enhancing Nutritional and Functional Properties for Africa*, Pretoria, South Africa, 2–4 April 2003.
15. Rumler, R.; Bender, D.; Speranza, S.; Frauenlob, J.; Gamper, L.; Hoek, J.; Jäger, H.; Schönlechner, R. Chemical and physical characterization of sorghum milling fractions and sorghum whole meal flours obtained via stone or roller milling. *Foods* 2021, 10, 870.
16. Seleem, H.A.; Omran, A.A. Evaluation quality of one layer flat bread supplemented with beans and sorghum baked on hot metal surface. *Food Nutr. Sci.* 2014, 5, 2246–2256.
17. Tasie, M.M.; Gebreyes, B.G. Characterization of nutritional, antinutritional, and mineral contents of thirty-five sorghum varieties grown in Ethiopia. *Int. J. Food Sci.* 2020, 2020, 1–11.
18. Galán, M.G.; Llopart, E.E.; Drago, S.R. Losses of nutrients and anti-nutrients in red and white sorghum cultivars after decorticating in optimised conditions. *Int. J. Food Sci. Nutr.* 2017, 69, 283–290.
19. Yousif, A.; Nhepera, D.; Johnson, S. Influence of sorghum flour addition on flat bread in vitro starch digestibility, antioxidant capacity and consumer acceptability. *Food Chem.* 2012, 134, 880–887.
20. Srichuwong, S.; Curti, D.; Austin, S.; King, R.; Lamothe, L.; Gloria-Hernandez, H. Physicochemical properties and starch digestibility of whole grain sorghums, millet, quinoa and amaranth flours, as affected by starch and non-starch constituents. *Food Chem.* 2017, 233, 1–10.
21. Taylor, J.R.N.; Schussler, L.; van der Walt, W.H. Fractionation of proteins from low-tannin sorghum grain. *J. Agric. Food Chem.* 1984, 32, 149–154.
22. Belton, P.S.; Delgadillo, I.; Halford, N.G.; Shewry, P.R. Kafirin structure and functionality. *J. Cereal Sci.* 2006, 44, 272–286.
23. Mohapatra, D.; Patel, A.S.; Kar, A.; Deshpande, S.S.; Tripathi, M.K. Effect of different processing conditions on proximate composition, anti-oxidants, anti-nutrients and amino acid profile of grain sorghum. *Food Chem.* 2019, 271, 129–135.

24. Gassem, M.A.A.; Osman, M.A. Proximate composition and the content of sugars, amino acids and anti-nutritional factors of three sorghum varieties. *Agric. Res. Cent. King Saud Univ. Res. Bull.* 2003, 125, 5–19.
25. Emmambux, N.M.; Taylor, J.R.N. Sorghum kafirin interaction with various phenolic compounds. *J. Sci. Food Agric.* 2003, 83, 402–407.
26. Pontieri, P.; Di Fiore, R.; Troisi, J.; Bean, S.R.; Roemer, E.; Okot, J.; Alifano, P.; Pignone, D.; Del Giudice, L.; Massardo, D.R. Chemical composition and fatty acid content of white food sorghums grown in different environments. *Maydica* 2011, 56, 1–7.
27. Sang, Y.; Bean, S.; Seib, P.A.; Pedersen, J.; Shi, Y.-C. Structure and functional properties of sorghum starches differing in amylose content. *J. Agric. Food Chem.* 2008, 56, 6680–6685.
28. Singh, H.; Sodhi, N.S.; Singh, N. Characterisation of starches separated from sorghum cultivars grown in India. *Food Chem.* 2010, 119, 95–100.
29. Miafo, A.-P.T.; Koubala, B.B.; Kansci, G.; Muralikrishna, G. Free sugars and non-starch polysaccharides-phenolic acid complexes from bran, spent grain and sorghum seeds. *J. Cereal Sci.* 2019, 87, 124–131.
30. Martinol, H.S.D.; Tomaz, P.A.; Moraes, É.A.; Conceição, L.L.d.; Oliveira, D.d.S.; Queiroz, V.A.V.; Rodrigues, J.A.S.; Pirozi, M.R.; Pinheiro-Sant’Ana, H.M.; Ribeiro, S.M.R. Chemical characterization and size distribution of sorghum genotypes for human consumption. *Rev. Inst. Adolfo Lutz (Impresso)* 2012, 71, 337–344.
31. Moraes, É.A.; Marineli, R.d.S.; Lenquiste, S.A.; Steel, C.J.; Menezes, C.B.d.; Queiroz, V.A.V.; Maróstica Júnior, M.R. Sorghum flour fractions: Correlations among polysaccharides, phenolic compounds, antioxidant activity and glycemic index. *Food Chem.* 2015, 180, 116–123.
32. Osman, R.O.; Abd El-Gelil, F.M.; El-Noamany, H.M.; Dawood, M.G. Oil content and fatty acid composition of some varieties of barley and sorghum grains. *Grasas Y Aceites* 2000, 51, 157–162.
33. Meera, M.S.; Bhashyam, M.K.; Ali, S.Z. Effect of heat treatment of sorghum grains on storage stability of flour. *LWT-Food Sci. Technol.* 2011, 44, 2199–2204.
34. Motlhaodi, T.; Bryngelsson, T.; Chite, S.; Fatih, M.; Ortiz, R.; Geleta, M. Nutritional variation in sorghum [*Sorghum bicolor* (L.) Moench] accessions from southern Africa revealed by protein and mineral composition. *J. Cereal Sci.* 2018, 83, 123–129.
35. Makawi, A.B.; Mustafa, A.I.; Adiamo, O.Q.; Mohamed Ahmed, I.A. Physicochemical, nutritional, functional, rheological, and microbiological properties of sorghum flour fermented with baobab fruit pulp flour as starter. *Food Sci. Nutr.* 2019, 7, 689–699.
36. Girard, A.L.; Awika, J.M. Sorghum polyphenols and other bioactive components as functional and health promoting food ingredients. *J. Cereal Sci.* 2018, 84, 112–124.
37. Beta, T.; Corke, H.; Rooney, L.W.; Taylor, J.R.N. Starch properties as affected by sorghum grain chemistry. *J. Sci. Food Agric.* 2000, 81, 245–251.
38. Serna-Saldivar, S.O.; Espinosa-Ramírez, J. Grain structure and grain chemical composition. In *Sorghum and Millets*; Elsevier: Amsterdam, The Netherlands, 2019; pp. 85–129.
39. Wedad, H.A.; El Tinay, A.H.; Mustafa, A.I.; Babiker, E.E. Effect of fermentation, malt-pretreatment and cooking on antinutritional factors and protein digestibility of sorghum cultivars. *Pak. J. Nutr.* 2008, 7, 335–341.
40. Przybylska-Balcerek, A.; Frankowski, J.; Stuper-Szablewska, K. Bioactive compounds in sorghum. *Eur Food Res. Technol* 2019, 245, 1075–1080.

Keywords

sorghum;western bread;rheology;baking;wheat sorghum blends
