

Applications of Heterocyclic Compounds

Subjects: Chemistry, Medicinal

Contributor: Faiz Ullah, Sami Ullah, Muhammad Farhan Ali Khan, Muhammad Mustaqeem, Rizwan Nasir Paracha, Muhammad Fayyaz ur Rehman, Fariha Kanwal, Syed Shams ul Hassan, Simona Bungau

Fluorescently labelled heterocyclic compounds are useful in bioanalytical applications, including in vivo imaging, high throughput screening, diagnostics, and light-emitting diodes. These compounds have various therapeutic properties, including antifungal, antitumor, antimalarial, anti-inflammatory, and analgesic activities. Different neutral fluorescent markers containing nitrogen heterocycles (quinolones, azafluoranthenes, pyrazoloquinolines, etc.) have several electrochemical, biological, and nonlinear optic applications. Photodynamic therapy (PDT), which destroys tumors and keeps normal tissues safe, works in the presence of molecular oxygen with light and a photosensitizing drugs (dye) to obtain a therapeutic effect.

Keywords: fluorescence ; heterocyclic compounds ; antitumor

1. Introduction

Molecular luminescence approaches include phosphorescence and fluorescence. A photon is absorbed by an analyte molecule, which stimulates a species. The emission spectrum can be used for quantitative and qualitative studies ^{[1][2]}. Because of their potential various functional applications, luminous materials have received significant attention lately ^[3]. They have been widely employed, including in the food, pharmaceutical, optical, and textile sectors ^{[4][5]}. Conventionally, inorganic emitting materials were commonly used; however, organic luminescent materials with brilliant emission have largely replaced them due to their wide range of uses, including emergency lighting, low cost, environmental friendliness, long-term solutions, anti-counterfeiting displays, and in food, cosmetics, polymers, bioactive molecules, and biochemistry ^{[6][7]}. Furthermore, in today's research, the design of novel luminous hybrid organic–nonorganic materials is critical ^{[8][9]}. The combination of phosphorescent dye as a sensitizer co-doped with a fluorescence emitter has made progress in developing luminescent materials for organic light-emitting diodes (OLEDs) in recent years ^{[10][11]}.

Fluorescence quenching or fluorescence enhancement is employed as an analytical technique ^{[12][13]}. The fluorescent labelling of the host molecule complex provides a useful tool for detecting the analyte's attachment to the host molecule ^{[14][15][16]}. For example, protein labelling using small molecule-based fluorescent probes is used in various biological experiments and is a valuable technique for determining the expression level and localization of a protein of interest in living cells ^[17].

Due to particular biological activity, crown ether's derivate and *N*-containing heterocyclic chemicals and their derivatives have been widely employed in agronomy and medicine ^[18]. Similar organic chemicals are of interest in pharmacology as effective tissue oxygenators and antidepressants, as well as in biotechnology; these compounds are employed for macromolecule binding ^{[19][20][21][22][23]}. *N*-containing heterocyclic compounds and macrocycle derivative crown ethers have remarkable photochemical, catalytic, and luminescent capabilities, indicating that they might be used to diagnose and cure various ailments. A few of their applications include photodynamic treatment and antimicrobial/antiparasitic activities against human pathogens and malarial parasites. Employment of fluorophores, including organic chromophores and crown ethers, with high selectivity, sensitivity, and stability constants while detecting tumor cells opens up new avenues for cancer research ^{[24][25]}.

Macrocyclic molecules, for example, crown ether, have been used in a wide range of chemical processes, including selective metal complexing agents and photo-induced electron transfer bio-mimetic research ^{[26][27][28][29]}. In contrast to the extensive coordination chemistry, little is known about crown ether coordination compounds' photoluminescence (PL). Crown ethers substituted with particular fluorescent dyes were the most commonly reported for PL. The use of such dye-substituted systems in sensing and analytical chemistry to detect the presence of particular metal cations was intensively investigated ^{[30][31][32]}.

A smart fluorescent probe with a crown ether moiety might be constructed as a sensor for metal anions, ions, and other biomolecules and then used to monitor biological processes in vivo [33]. The solvent effects of a crown ether complex containing a fluorescent anthracene unit are exceptional [34].

Supramolecular chemistry, inspired by nature's vast array of assemblies, has garnered significant attention in recent decades due to its diverse supra-structures, which consist of micelles, vesicles, and fibers, as well as its wide-ranging applications in sensors, drug delivery, luminescent materials, and bioimaging [35][36][37][38].

Fluorescence characteristics of *N*-containing heterocyclic compounds have recently received considerable interest. For example, fluorescent compounds known as quinolines have attracted the attention of scientists because of their use in high-tech applications [39]. Similarly, derivatives of the pyrazoloquinoline (PQ) family and quinoline are an example of fluorescent substances that may be of interest for several applications, including their use as oxidant scavengers and growth promoters [40][41]. These have also been found naturally in a wide range of foods and appear to be easily absorbed. More recently, heterocyclic azo compounds such as benzothiazole, pyrazole, and thiazole have been employed for electrochemical, biological, and nonlinear optics applications and structure–activity relationships for drug designing [SAR] [42][43][44]. Thiophene and thienopyrimidine derivatives have fluorescence features and are more efficient than other aromatic chemicals for anti-avian influenza virus (H5N1) action. Porphyrins are *N*-heterocyclic chemicals present in a wide variety of biological systems. Metalloporphyrins contain solely -pyrrolic substituents in biological systems and appear attached to proteins, creating supramolecular structures such as haemoglobin, myoglobin, cytochromes, catalases, and peroxidases, as well as chlorophylls and bacteriochlorophylls in reduced forms [45].

2. Applications of Heterocyclic Compounds

2.1. Anti-Mycobacterial Activity

Different symptoms such as respiratory issues, long-term coughs, and tuberculosis are treated by various plants in African and Asian countries. Many anti-tubercular drugs, with toxicity and side effects, are still used to treat tuberculosis. For treating *M. tuberculosis*, the synthesis of azo compounds was monitored and showed anti-tubercular activity. Maximum activity was shown by compounds 5-methyl-2-(5-methylbenzo[d]thiazol-2-yl)-4-(*p*-tolyl diazenyl)-1*H*-pyrazol-3(2*H*)-one (**1a**) and 5-methyl-2-(5-methylbenzo[d]thiazol-2-yl)-4-(*m*-tolyl diazenyl)-1*H*-pyrazol-3(2*H*)-one (**1b**) when compared to the copounds 4-((4-chlorophenyl) diazenyl)-5-methyl-2-(5-methylbenzo[d]thiazol-2-yl)-1*H*-pyrazol-3(2*H*)-one (**1c**) and 4-((4-bromophenyl) diazenyl)-5-methyl-2-(5-methylbenzo[d]thiazol-2-yl)-1*H*-pyrazol-3(2*H*)-one (**1d**) shown below in **Figure 1**, correspondingly. A previous study shows that the presence of a side chain to an azo dye along with a phenyl group substituent and a significantly enhanced electron-donating group ultimately decreased the growth of bacteria [5].




Figure 1. Structures of azo dye compounds (**1a–1d**) showing anti-mycobacterial activity.

2.2. Anticancer Activity

The photochemistry and the anti-tuberculosis activity of the in vitro azo compounds discussed above yielded good results, so their anticancer activity was also studied. An MTT test was performed for cell proliferation, and for this reason, different human cancer cell lines were used, such as chronic myeloid leukaemia (K562), lung carcinoma (A549), colon (HCT116), and T-lymphocyte (Jurkat) cell lines. **Table 1** shows their anticancer activity results. Data revealed that K562, Jurkat, and A549 cell lines containing various synthesized azo compounds displayed fair in vitro results ($IC_{50} > 50$). However, on the other hand, in comparison with other human cell lines, the HCT116 cell line showed relatively good activity in the presence of various compounds [46].

Table 1. Anticancer activities of azo compounds (**1a–d**).

Compounds	IC_{50} (μM)			
	HCT116	A549	Jurkat	K562
1a	34.65 ± 0.35	>50	>50	>50
1b	>50	>50	>50	>50
1c	43.33 ± 0.14	>50	>50	>50
1d	48.19 ± 0.31	>50	>50	>50

2.3. Therapeutic and Biological Applications

Various applications, such as anti-inflammatory, antibacterial, analgesic, antiviral, antipyretic, and anti-convulsant activities, belonged to 3-aminopyrroles derivatives, which are considered an essential family of compounds [47]. Thiophene compounds also play a significant role as agrochemicals [48][49], anti-avian influenza virus (H5N1), anti-tubercular, anti-breast cancer agents, AMPK activators, HIV, and multi-target kinase inhibitors [50].

The majority of roles, including serving as precursors for different biological molecules or connecting to various sulphur and nitrogen heterocycles, are imparted by some structural units combined to form a 2-aminothiophene product. Apart from this, UV-visible absorption and fluorescence of these compounds make them important for biological purposes. Thiophene derivatives can be used explicitly as valuable fluorescent dyes in confocal microscopy for bio-imaging [51].

2.4. Antiparasitic Activity of Metalloporphyrins and Their Role as Potentiometric Biosensors

Metalloporphyrins, known for their β -pyrrolic substitution, are important in forming useful supramolecules such as cytochromes, haemoglobin, peroxidases, myoglobin, and catalases [52][53]. The main reason porphyrins are gaining importance in the biological world day by day is their diverse functionality along with their remarkable structural features and positive properties in the field of photochemistry and spectroscopy. The use of metalloporphyrins as potentiometric sensors is common among all other functions—for example, Mn(III)-porphyrin derivatives are being used in the chloride ion measurement in samples of human serum [54].

The increase in antiparasitic activity of porphyrins is related to the presence of electrically charged substituents on these compounds. An ultimate decrease in the oxidative damage to the mosquitoes' larvae of genera *Culex*, *Aedes* [55][56], and *Anopheles* [57], while of adult flies of *Ceratitis capitates*, *Bactrocera oleae* species, and *Stomoxys calcitrans* [54][58] can be observed by porphyrin-based drugs. Photosensitization makes hematoporphyrin IX a powerful eco-friendly drug.

2.5. Antioxidant Activity

Disordered physiological processes such as neurodegenerative disorders are studied by reactive nitrogen and oxygen species or heterocyclic compounds [59]. Neuroprotection involves an option of antioxidant therapy, so antioxidants can be described as compounds capable of searching for free radicals. Discussing specific fluorescent heterocycles shown in **Figure 2** such as (3s,5s,7s)-*N*-(2,4-dinitrophenyl)adamantan-1-amine (**2a**), *N*-((3s,5s,7s)-adamantan-1-yl)-6-(dimethylamino)-naphthalene-2-sulfonamide (**2b**), 2-(adamantan-1-yl)-2*H*-isindole-1-carbonitrile (**2c**), provide a guide to the pharmacological industry as they are of great interest as antioxidant agents [60].




Figure 2. Structures of compounds (**2a–2c**) having antioxidant activities.

References

- Sanderson, M.J.; Smith, I.; Parker, I.; Bootman, M.D. Fluorescence microscopy. *Cold Spring Harb. Protoc.* 2014, 2014, 1042–1065.
- Schulman, S.; Sharma, A. Introduction to fluorescence spectroscopy. *Microchem. J.* 2000, 65, 353.
- McGown, L.B.; Nithipahkom, K. Molecular fluorescence and phosphorescence. *Appl. Spectrosc. Rev.* 2000, 35, 353–393.
- Derkowska-Zielinska, B.; Skowronski, L.; Biitseva, A.; Grabowski, A.; Naparty, M.K.; Smokal, V.; Kysil, A.; Krupka, O. Optical characterization of heterocyclic azo dyes containing polymers thin films. *Appl. Surf. Sci.* 2017, 421, 361–366.
- Maliyappa, M.R.; Keshavayya, J.; Mallikarjuna, N.M.; Krishna, P.M.; Shivakumara, N.; Sandeep, T.; Sailaja, K.; Nazrulla, M.A. Synthesis, characterization, pharmacological and computational studies of 4, 5, 6, 7-tetrahydro-1, 3-benzothiazole incorporated azo dyes. *J. Mol. Struct.* 2018, 1179, 630–641.
- Xu, S.; Chen, R.; Zheng, C.; Huang, W. Excited State Modulation for Organic Afterglow: Materials and Applications. *Adv. Mater.* 2016, 28, 9920–9940.
- Escribano, P.; Julián-López, B.; Planelles-Aragó, J.; Cordoncillo, E.; Viana, B.; Sanchez, C. Photonic and nanobiophotonic properties of luminescent lanthanide-doped hybrid organic-inorganic materials. *J. Mater. Chem.* 2008, 18, 23–40.
- Khattab, T.A.; Abd El-Aziz, M.; Abdelrahman, M.S.; El-Zawahry, M.; Kamel, S. Development of long-persistent photoluminescent epoxy resin immobilized with europium (II)-doped strontium aluminate. *Luminescence* 2020, 35, 478–485.
- Muhr, V.; Würth, C.; Kraft, M.; Buchner, M.; Baeumner, A.J.; Resch-Genger, U.; Hirsch, T. Particle-Size-Dependent Förster Resonance Energy Transfer from Upconversion Nanoparticles to Organic Dyes. *Anal. Chem.* 2017, 89, 4868–4874.
- Muhamad Sarih, N.; Myers, P.; Slater, A.; Slater, B.; Abdullah, Z.; Tajuddin, H.A.; Maher, S. White Light Emission from a Simple Mixture of Fluorescent Organic Compounds. *Sci. Rep.* 2019, 9, 11834.
- Girgis, A.S.; Basta, A.H.; El-Saied, H.; Mohamed, M.A.; Bedair, A.H.; Salim, A.S. Synthesis, quantitative structure–property relationship study of novel fluorescence active 2-pyrazolines and application. *R. Soc. Open Sci.* 2018, 5, 171964.
- Velic, D.; Knapp, M.; Köhler, G. Supramolecular inclusion complexes between a coumarin dye and β -cyclodextrin, and attachment kinetics of thiolated β -cyclodextrin to gold surface. *J. Mol. Struct.* 2001, 598, 49–56.
- Wagner, B.D.; Fitzpatrick, S.J. A comparison of the host-guest inclusion complexes of 1,8-ANS and 2,6-ANS in parent and modified cyclodextrins. *J. Incl. Phenom.* 2000, 38, 467–478.
- Kele, P.; Orbulescu, J.; Calhoun, T.L.; Gawley, R.E.; Leblanc, R.M. Coumaryl crown ether based chemosensors: Selective detection of saxitoxin in the presence of sodium and potassium ions. *Tetrahedron Lett.* 2002, 43, 4413–4416.
- Li, T.; Pang, H.; Wu, Q.; Huang, M.; Xu, J.; Zheng, L.; Wang, B.; Qiao, Y. Rigid Schiff Base Complex Supramolecular Aggregates as a High-Performance pH Probe: Study on the Enhancement of the Aggregation-Caused Quenching (ACQ) Effect via the Substitution of Halogen Atoms. *Int. J. Mol. Sci.* 2022, 23, 6259.
- Schönefeld, K.; Barann, A.; Vogel, K.; Feller, K.H.; Kunze, D.; Müller, P.; Weber, E. Fluorescence studies of crown ether complexes—Solvent effects regarding the inclusion properties of host-guest sensor complexes. *Int. J. Environ. Anal. Chem.* 2005, 85, 655–663.
- Prasad, P.K.; Motiei, L.; Margulies, D. Steps toward enhancing the fluorescence of small-molecule-based protein labels using supramolecular hosts. *Results Chem.* 2021, 3, 100134.
- Singh, S.K.; Singh, M.; Singh, S.K.; Gangwar, M.; Nath, G. Design, synthesis and mode of action of some benzothiazole derivatives bearing an amide moiety as antibacterial agents. *RSC Adv.* 2014, 4, 19013–19023.
- Ullah, F.; Khan, T.A.; Iltaf, J.; Anwar, S.; Khan, M.F.A.; Khan, M.R.; Ullah, S.; Rehman, M.F.U.; Mustaqeem, M.; Kotwica-Mojzych, K.; et al. Heterocyclic Crown Ethers with Potential Biological and Pharmacological Properties: From Synthesis

20. Cai, K.; Wang, F.; Lu, J.-Q.; Shen, A.-N.; Zhao, S.-M.; Zang, W.-D.; Gui, Y.-H.; Zhao, J.-Y. Nicotinamide Mononucleotide Alleviates Cardiomyopathy Phenotypes Caused by Short-Chain Enoyl-CoA Hydratase 1 Deficiency. *JACC Basic Transl. Sci.* 2022, 7, 348–362.
21. Wu, Z.; Li, C.; Zhang, F.; Huang, S.; Wang, F.; Wang, X.; Jiao, H. High-performance ultra-narrow-band green-emitting phosphor LaMgAl₁₁O₁₉:Mn²⁺ for wide color-gamut WLED backlight displays. *J. Mater. Chem. C* 2022, 10, 7443–7448.
22. Kityk, A.V. Absorption and fluorescence spectra of heterocyclic isomers from long-range-corrected density functional theory in polarizable continuum approach. *J. Phys. Chem. A* 2012, 116, 3048–3055.
23. Mao, G.; Orita, A.; Fenenko, L.; Yahiro, M.; Adachi, C.; Otera, J. Blue emitting fluorophores of phenyleneethynylenes substituted by diphenylethenyl terminal groups for organic light-emitting diodes. *Mater. Chem. Phys.* 2009, 115, 378–384.
24. Feng, Y.; Li, F.; Yan, J.; Guo, X.; Wang, F.; Shi, H.; Du, J.; Zhang, H.; Gao, Y.; Li, D.; et al. Pan-cancer analysis and experiments with cell lines reveal that the slightly elevated expression of DLGAP5 is involved in clear cell renal cell carcinoma progression. *Life Sci.* 2021, 287, 120056.
25. Qu, Y.-Y.; Zhao, R.; Zhang, H.-L.; Zhou, Q.; Xu, F.-J.; Zhang, X.; Xu, W.-H.; Shao, N.; Zhou, S.-X.; Dai, B.; et al. Inactivation of the AMPK-GATA3-ECHS1 Pathway Induces Fatty Acid Synthesis That Promotes Clear Cell Renal Cell Carcinoma Growth. *Cancer Res.* 2020, 80, 319–333.
26. Mohanty, J.; Pal, H.; Sapre, A.V. Excited singlet (S₁) state interactions of 2,2'- and 4,4'-biphenyldiols with chloroalkanes: Photoinduced dissociative electron transfer. *J. Chem. Phys.* 2002, 116, 8006–8014.
27. Yang, W.; Zhang, H.; Liu, Y.; Tang, C.; Xu, X.; Liu, J. Rh(III)-catalyzed synthesis of dibenzopyran-6-ones from aryl ketone O-acetyl oximes and quinones via C-H activation and C-C bond cleavage. *RSC Adv.* 2022, 12, 14435–14438.
28. Zhang, J.; Lv, J.; Wang, J. The crystal structure of (E)-1-(4-aminophenyl)-3-(p-tolyl)prop-2-en-1-one, C₁₆H₁₅NO. *Z. Krist.—New Cryst. Struct.* 2022, 237, 385–387.
29. Jarolímová, Z.; Vishe, M.; Lacour, J.; Bakker, E. Potassium ion-selective fluorescent and pH independent nanosensors based on functionalized polyether macrocycles. *Chem. Sci.* 2016, 7, 525–533.
30. Li, Y.P.; Yang, H.R.; Zhao, Q.; Song, W.C.; Han, J.; Bu, X.H. Ratiometric and selective fluorescent sensor for Zn²⁺ as an “off-on-off” switch and logic gate. *Inorg. Chem.* 2012, 51, 9642–9648.
31. Valeur, B.; Leray, I. Design principles of fluorescent molecular sensors for cation recognition. *Coord. Chem. Rev.* 2000, 205, 3–40.
32. Kapoor, S. Fluorescence properties of crown ethers with phenylbenzothiozole pendant group. *Chem. Phys. Lett.* 2005, 408, 290–294.
33. Li, J.; Yim, D.; Jang, W.; Yoon, J. Chem Soc Rev Recent progress in the design and applications of fluorescence probes containing crown ethers. *Chem. Soc. Rev.* 2016, 46, 2437–2458.
34. Erk, C. Cation Recognition with Fluorophore Crown Ethers 1. *Ind. Eng. Chem. Res.* 2000, 39, 3582–3588.
35. Lou, X.Y.; Yang, Y.W. Manipulating Aggregation-Induced Emission with Supramolecular Macrocycles. *Adv. Opt. Mater.* 2018, 6, 1800668.
36. Ji, X.; Dong, S.; Wei, P.; Xia, D.; Huang, F. A novel diblock copolymer with a supramolecular polymer block and a traditional polymer block: Preparation, controllable self-assembly in water, and application in controlled release. *Adv. Mater.* 2013, 25, 5725–5729.
37. Li, B.; He, T.; Fan, Y.; Yuan, X.; Qiu, H.; Yin, S. Recent developments in the construction of metallacycle/metallacage-cored supramolecular polymers: Via hierarchical self-assembly. *Chem. Commun.* 2019, 55, 8036–8059.
38. Całus, S.; Gondek, E.; Danel, A.; Jarosz, B.; Pokladko, M.; Kityk, A.V. Electroluminescence of 6-R-1,3-diphenyl-1H-pyrazoloquinoline-based organic light-emitting diodes (R = F, Br, Cl, CH₃, C₂H₃ and N(C₆H₅)₂). *Mater. Lett.* 2007, 61, 3292–3295.
39. Yan, H.; Cui, P.; Liu, C.; Yuan, S. Molecular Dynamics Simulation of Pyrene Solubilized in a Sodium Dodecyl Sulfate Micelle. *Langmuir* 2012, 28, 4931–4938.
40. Gąsiorowski, P.; Danel, K.S.; Matusiewicz, M.; Uchacz, T.; Kityk, A.V. From pirazoloquinolines to annulated azulene dyes: UVVIS spectroscopy and quantum chemical study. *J. Lumin.* 2010, 130, 2460–2468.
41. Stites, T.E.; Mitchell, A.E.; Rucker, R.B. Physiological importance of quinoenzymes and the O-quinone family of cofactors. *J. Nutr.* 2000, 130, 719–727.
42. Castro, M.C.R.; Schellenberg, P.; Belsley, M.; Fonseca, A.M.C.; Fernandes, S.S.M.; Raposo, M.M.M. Design, synthesis and evaluation of redox, second order nonlinear optical properties and theoretical DFT studies of novel bithiophene azo

dyes functionalized with thiadiazole acceptor groups. *Dye. Pigment.* 2012, 95, 392–399.

43. Beyzaei, H.; Aryan, R.; Moghaddam-Manesh, M.; Ghasemi, B.; Karimi, P.; Samareh Delarami, H.; Sanchooli, M. Evaluation and structure-activity relationship analysis of a new series of 4-imino-5H-pyrazolo pyrimidin-5-amines as potential antibacterial agents. *J. Mol. Struct.* 2017, 1144, 273–279.
44. Raposo, M.M.M.; Castro, M.C.R.; Fonseca, A.M.C.; Schellenberg, P.; Belsley, M. Design, synthesis, and characterization of the electrochemical, nonlinear optical properties, and theoretical studies of novel thienylpyrrole azo dyes bearing benzothiazole acceptor groups. *Tetrahedron* 2011, 67, 5189–5198.
45. Huang, X.; Groves, J.T. Oxygen Activation and Radical Transformations in Heme Proteins and Metalloporphyrins. *Chem. Rev.* 2018, 118, 2491–2553.
46. Maliyappa, M.R.; Keshavayya, J.; Mallikarjuna, N.M.; Pushpavathi, I. Novel substituted aniline based heterocyclic dispersed azo dyes coupling with 5-methyl-2-(6-methyl-1, 3-benzothiazol-2-yl)-2, 4-dihydro-3H-pyrazol-3-one: Synthesis, structural, computational and biological studies. *J. Mol. Struct.* 2019, 1205, 127576.
47. Khan, F.A.; Ali, G.; Rahman, K.; Khan, Y.; Ayaz, M.; Mosa, O.F.; Nawaz, A.; Hassan, S.S.U.; Bungau, S. Efficacy of 2-Hydroxyflavanone in Rodent Models of Pain and Inflammation: Involvement of Opioidergic and GABAergic Anti-Nociceptive Mechanisms. *Molecules* 2022, 27, 5431.
48. Mahmood, F.; Khan, J.A.; Mahnashi, M.H.; Jan, M.S.; Javed, M.A.; Rashid, U.; Sadiq, A.; Hassan, S.S.U.; Bungau, S. Anti-Inflammatory, Analgesic and Antioxidant Potential of New (2S,3S)-2-(4-isopropylbenzyl)-2-methyl-4-nitro-3-phenylbutanals and Their Corresponding Carboxylic Acids through In Vitro, In Silico and In Vivo Studies. *Molecules* 2022, 27, 4068.
49. Muhammad, I.; Luo, W.; Shoaib, R.M.; Li, G.L.; Hassan, S.S.U.; Yang, Z.H.; Xiao, X.; Tu, G.L.; Yan, S.K.; Ma, X.P.; et al. Guaiane-type sesquiterpenoids from *Cinnamomum migao* H. W. Li: And their anti-inflammatory activities. *Phytochemistry* 2021, 190, 112850.
50. Rashad, A.E.; Shamroukh, A.H.; Abdel-megeid, R.E.; Mostafa, A.; El-shesheny, R.; Kandeil, A.; Ali, M.A.; Banert, K. European Journal of Medicinal Chemistry Synthesis and screening of some novel fused thiophene and thienopyrimidine derivatives for anti-avian influenza virus (H5N1) activity. *Eur. J. Med. Chem.* 2010, 45, 5251–5257.
51. Baleiz, C. Synthesis and fluorescence properties of aminocyanopyrrole and aminocyanothiophene esters for biomedical and bioimaging applications. *J. Mol. Struct.* 2020, 1209, 127974.
52. Imran, M.; Ramzan, M.; Qureshi, A.K.; Khan, M.A.; Tariq, M. Emerging Applications of Porphyrins and Metalloporphyrins in Biomedicine and Diagnostic Magnetic Resonance Imaging. *Biosensors* 2018, 8, 95.
53. Zhang, X.; Qu, Y.; Liu, L.; Qiao, Y.; Geng, H.; Lin, Y.; Zhao, J. Homocysteine inhibits pro-insulin receptor cleavage and causes insulin resistance via protein cysteine-homocysteinylation. *Cell Rep.* 2021, 37, 109821.
54. Garcia, C.R.S.; Deda, D.K.; Iglesias, B.A.; Alves, E.; Araki, K. Porphyrin Derivative Nanoformulations for Therapy and Antiparasitic Agents. *Molecules* 2020, 25, 2080.
55. Awad, H.H.; El-tayeb, T.A.; Abd El-aziz, N.M.; Abdelkader, M.A. A Semi-field Study on the Effect of Novel Hematoporphyrin Formula on the Control of *Culex pipiens* Larvae. *J. Agric. Soc. Sci.* 2008, 85–88.
56. Lucantoni, L.; Magaraggia, M.; Lupidi, G.; Ouedraogo, R.K.; Esposito, F.; Fabris, C.; Jori, G.; Habluetzel, A. Novel, Mes-o-Substituted Cationic Porphyrin Molecule for Photo-Mediated Larval Control of the Dengue Vector *Aedes aegypti*. *J. Pntd.* 2011, 5, e1434.
57. Fabris, C.; Kossivi, R.; Coppellotti, O.; Dabiré, R.K.; Diabaté, A.; Di, P.; Guidolin, L.; Jori, G.; Lucantoni, L.; Lupidi, G.; et al. Acta Tropica Efficacy of sunlight-activatable porphyrin formulates on larvae of *Anopheles gambiae* M and S molecular forms and *An. arabiensis*: A potential novel biolarvicide for integrated malaria vector control. *Acta Trop.* 2012, 123, 239–243.
58. Ben Amor, T.; Jori, G. Sunlight-activated insecticides: Historical background and mechanisms of phototoxic activity. *Insect Biochem. Mol. Biol.* 2000, 30, 915–925.
59. Nishida, J.; Kawabata, J. DPPH Radical Scavenging Reaction of Hydroxy- and Methoxychalcones. *Biosci. Biotechnol. Biochem.* 2016, 8451, 193–202.
60. Joubert, J.; Van Dyk, S.; Green, I.R.; Malan, S.F. Bioorganic & Medicinal Chemistry Synthesis and evaluation of fluorescent heterocyclic aminoadamantanes as multifunctional neuroprotective agents. *Bioorg. Med. Chem.* 2011, 19, 3935–3944.

