

# Riboflavin Transporter Deficiency Neuronopathy

Subjects: Genetics & Heredity

Contributor: Nora Tang

Riboflavin transporter deficiency neuronopathy is a disorder that affects nerve cells (neurons). Affected individuals typically have hearing loss caused by nerve damage in the inner ear (sensorineural hearing loss) and signs of damage to other nerves.

Keywords: genetic conditions

---

## 1. Introduction

In addition to nerves in the inner ear, riboflavin transporter deficiency neuronopathy involves nerves found in the part of the brain that is connected to the spinal cord (the brainstem), specifically in a region of the brainstem known as the pontobulbar region. Damage to these nerves causes paralysis of the muscles controlled by them, a condition called pontobulbar palsy. Nerves in the pontobulbar region help control several voluntary muscle activities, including breathing, speaking, and moving the limbs. As a result of pontobulbar palsy, people with riboflavin transporter deficiency neuronopathy can have breathing problems; slurred speech; and muscle weakness in the face, neck, shoulders, and limbs. Affected individuals can also have muscle stiffness (spasticity) and exaggerated reflexes.

The age at which riboflavin transporter deficiency neuronopathy begins varies from infancy to young adulthood. When the condition begins in infancy, the first symptom is often breathing problems caused by nerve damage, which can be life-threatening. When the condition begins in children or young adults, sensorineural hearing loss usually occurs first, followed by signs of pontobulbar palsy.

If not treated, the signs and symptoms of riboflavin transporter deficiency neuronopathy worsen over time. Severe breathing problems and respiratory infections are the usual cause of death in people with this condition. Without treatment, affected infants typically survive less than one year. However, those who develop the condition after age 4 often survive more than 10 years.

Riboflavin transporter deficiency neuronopathy encompasses two conditions that were once considered distinct disorders: Brown-Vialetto-Van Laere syndrome (BVVLS) and Fazio-Londe disease. The two conditions have similar signs and symptoms, but Fazio-Londe disease does not include sensorineural hearing loss. Because these two conditions share a genetic cause and have overlapping features, researchers determined that they are forms of a single disorder.

## 2. Frequency

Riboflavin transporter deficiency neuronopathy is a rare condition. Approximately 100 cases have been reported in the scientific literature.

## 3. Causes

Riboflavin transporter deficiency neuronopathy is caused by mutations in the *SLC52A2* or *SLC52A3* gene. These genes provide instructions for making related proteins called riboflavin transporters: RFVT2 is produced from the *SLC52A2* gene, and RFVT3 is produced from the *SLC52A3* gene. Both proteins move (transport) a vitamin called riboflavin (also called vitamin B<sub>2</sub>) across the cell membrane. Riboflavin cannot be made by the body, so it must be obtained from the food a person eats. The RFVT3 protein is found at especially high levels in cells of the small intestine and is important for absorbing riboflavin during digestion so that the vitamin can be used in the body. The RFVT2 protein is found in cells of the brain and spinal cord and is important for ensuring that these tissues have enough riboflavin for proper functioning.

In the cells of the body, riboflavin is the core component of molecules called flavin adenine dinucleotide (FAD) and flavin mononucleotide (FMN). FAD and FMN are involved in many different chemical reactions and are required for a variety of cellular processes. One important role of these molecules is in the production of energy for cells. FAD and FMN are also

involved in the breakdown (metabolism) of carbohydrates, fats, and proteins.

Mutations in the *SLC52A2* or *SLC52A3* gene lead to an abnormal riboflavin transporter protein with impaired ability to transport riboflavin. Consequently, there is a reduction of riboflavin available in the body. However, it is unclear how these changes lead to the nerve problems characteristic of riboflavin transporter deficiency neuropathy.

### 3.3. The Genes Associated with Riboflavin Transporter Deficiency Neuropathy

- SLC52A2
- SLC52A3

## 4. Inheritance

Riboflavin transporter deficiency neuropathy usually follows an autosomal recessive pattern of inheritance, which means both copies of the gene in each cell have mutations. The parents of an individual with an autosomal recessive condition each carry one copy of the mutated gene, but they typically do not show signs and symptoms of the condition.

## 5. Other Names for This Condition

- Brown-Vialetto-Van Laere syndrome
- BVVLS
- Fazio-Londe disease
- Fazio-Londe syndrome
- pontobulbar palsy with deafness
- progressive bulbar palsy with sensorineural deafness
- riboflavin transporter deficiency

---

## References

1. Bosch AM, Abeling NG, Ijlst L, Knoester H, van der Pol WL, Stroomer AE, Wanders RJ, Visser G, Wijburg FA, Duran M, Waterham HR. Brown-Vialetto-Van Laere and Fazio Londe syndrome is associated with a riboflavin transporter defect mimicking mild MADD: a new inborn error of metabolism with potential treatment. *J Inher Metab Dis*. 2011 Feb;34(1):159-64. doi: 10.1007/s10545-010-9242-z.
2. Bosch AM, Stroek K, Abeling NG, Waterham HR, Ijlst L, Wanders RJ. The Brown-Vialetto-Van Laere and Fazio Londe syndrome revisited: natural history, genetics, treatment and future perspectives. *Orphanet J Rare Dis*. 2012 Oct 29;7:83. doi: 10.1186/1750-1172-7-83. Review.
3. Dipti S, Childs AM, Livingston JH, Aggarwal AK, Miller M, Williams C, Crow YJ. Brown-Vialetto-Van Laere syndrome; variability in age at onset and disease progression highlighting the phenotypic overlap with Fazio-Londe disease. *Brain Dev*. 2005 Sep;27(6):443-6.
4. Foley AR, Menezes MP, Pandraud A, Gonzalez MA, Al-Odaib A, Abrams AJ, Sugano K, Yonezawa A, Manzur AY, Burns J, Hughes I, McCullagh BG, Jungbluth H, Lim MJ, Lin JP, Megarbane A, Urtizberea JA, Shah AH, Antony J, Webster R, Broomfield A, Ng J, Mathew AA, O'Byrne JJ, Forman E, Scoto M, Prasad M, O'Brien K, Olpin S, Oppenheim M, Hargreaves I, Land JM, Wang MX, Carpenter K, Horvath R, Straub V, Lek M, Gold W, Farrell MO, Brandner S, Phadke R, Matsuura K, McGarvey ML, Scherer SS, Baxter PS, King MD, Clayton P, Rahman S, Reilly MM, Ouvrier RA, Christodoulou J, Züchner S, Muntoni F, Houlden H. Treatable childhood neuropathy caused by mutations in riboflavin transporter RFVT2. *Brain*. 2014 Jan;137(Pt 1):44-56. doi: 10.1093/brain/awt315.
5. Green P, Wiseman M, Crow YJ, Houlden H, Riphagen S, Lin JP, Raymond FL, Childs AM, Sheridan E, Edwards S, Josifova DJ. Brown-Vialetto-Van Laere syndrome, a pontobulbar palsy with deafness, is caused by mutations in c20orf54. *Am J Hum Genet*. 2010 Mar 12;86(3):485-9. doi: 10.1016/j.ajhg.2010.02.006.
6. Haack TB, Makowski C, Yao Y, Graf E, Hempel M, Wieland T, Tauer U, Ahting U, Mayr JA, Freisinger P, Yoshimatsu H, Imai K, Strom TM, Meitinger T, Yonezawa A, Prokisch H. Impaired riboflavin transport due to missense mutations in SLC5

2A2causes Brown-Vialetto-Van Laere syndrome. *J Inherit Metab Dis.* 2012Nov;35(6):943-8. doi: 10.1007/s10545-012-9513-y.

7. Johnson JO, Gibbs JR, Megarbane A, Urtizberea JA, Hernandez DG, Foley AR, Arepalli S, Pandraud A, Simón-Sánchez J, Clayton P, Reilly MM, Muntoni F, Abramzon Y, Houlden H, Singleton AB. Exome sequencing reveals riboflavin transporter mutations as a cause of motor neuron disease. *Brain.* 2012 Sep;135(Pt 9):2875-82. doi: 10.1093/brain/aws161.
8. Manole A, Houlden H. Riboflavin Transporter Deficiency Neuronopathy. 2015 Jun 11. In: Adam MP, Ardinger HH, Pagon RA, Wallace SE, Bean LJH, Stephens K, Amemiya A, editors. *GeneReviews®* [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2020. Available from <http://www.ncbi.nlm.nih.gov/books/NBK299312/>
9. Nabokina SM, Subramanian VS, Said HM. Effect of clinical mutations on functionality of the human riboflavin transporter-2 (hRFT-2). *Mol Genet Metab.* 2012 Apr;105(4):652-7. doi: 10.1016/j.ymgme.2011.12.021.
10. Oregon State University Linus Pauling Institute: Riboflavin

---

Retrieved from <https://encyclopedia.pub/entry/history/show/11951>