

Mixture Designs in Beverages/Foods/Pharmaceutical Health

Subjects: **Food Science & Technology**

Contributor: Diego Galvan

Design of Experiments (DoE) is a statistical tool used to plan and optimize experiments and is seen as a quality technology to achieve products excellence. Among the experimental designs (EDs), the mixture designs (MDs) stand out, being widely applied to improve conditions for processing, developing, or formulating novel products. Overall, the SOM indicated that Brazil presented the largest number of works using MDs. Among the continents, America and Asia showed a predominance in applications with the same amount of work. Comparing the MDs application areas, the analysis indicated that works are prevalent in food and beverage science in the American continent, while in Asia, health science prevails. MDs were more used to develop functional/nutraceutical products and the formulation of drugs for several diseases. However, we briefly describe some promising research fields in that MDs can still be employed.

design of experiments (DoE)

experimental designs (EDs)

mixture designs (MDs)

simplex lattice design (SLD)

simplex-centroid design (SCD)

1. Introduction

Currently, food and health science studies are becoming more complex with a considerable increase in the number of variables, requiring robust methods for simultaneous data analysis [1]. These analyzes can be facilitated using mathematical and statistical fundamentals, known as chemometric methods, which can be divided into the design of experiments, multivariate data analysis, and multivariate calibration [2][3][4].

Design of Experiments (DoE) is a statistical tool used to plan and optimize experiments and is seen as a quality technology to achieve products excellence [5][6][7][8][9]. Among the various experimental designs (EDs), the mixture designs (MDs) stand out. In MDs, two or more components are mixed in different proportions, and the characteristics of the resulting products are recorded. The responses are independent of physical states, depending only on the proportions of the ingredients present in the mixtures [7][8][10][11].

In MDs, the response depends only on the mixture and not on the mixture's total amount. The sum of a mixture's proportions in different components or ingredients is always 1% or 100% [8][12]. In an experiment with q components, the proportions of the ingredients may be denoted by x_1, x_2, \dots, x_q , where $x_i \geq 0$ for $i = 1, 2, \dots, q$ and $\sum_{i=1}^q x_i = 1$, where x_i represents the proportion of the i -th component. This equation removes a degree of freedom from the proportions and the factor space is, therefore, a $(q - 1)$ -dimensional regular simplex [2][8][9][13].

$$\text{Linear : } \hat{y} = \sum_{i=1}^q \beta_i x_i \quad (1)$$

$$\text{Quadratic : } \hat{y} = \sum_{i=1}^q \beta_i x_i + \sum_{i < j}^{q-1} \sum_j^q \beta_{ij} x_i x_j \quad (2)$$

$$\text{Special cubic : } \hat{y} = \sum_{i=1}^q \beta_i x_i + \sum_{i < j}^{q-1} \sum_j^q \beta_{ij} x_i x_j + \sum_{i < j}^{q-2} \sum_{j < k}^{q-1} \sum_k^q \beta_{ijk} x_i x_j x_k \quad (3)$$

$$\text{Full cubic : } \hat{y} = \sum_{i=1}^q \beta_i x_i + \sum_{i < j}^{q-1} \sum_j^q \beta_{ij} x_i x_j + \sum_{i < j}^{q-1} \sum_j^q \delta_{ij} x_i x_j (x_i - x_j) + \sum_{i < j}^{q-2} \sum_{j < k}^{q-1} \sum_k^q \beta_{ijk} x_i x_j x_k \quad (4)$$

The parameter β_i represents the expected response to the pure blend $x_i = 1$ and $x_j = 0$ when $j \neq i$. The term $\sum_{i=1}^q \beta_i x_i$ represents the linear blending portion. When curvature arises from nonlinear blending between component pairs, the parameters β_{ij} , which represent either synergistic or antagonistic blending, will be different from zero [2][8].

Developing a novel product requires mixing two or more ingredients, and specific characteristics or restrictions are often desired [14]. Therefore, product formulation is still a challenge that has dogged chemists, pharmacists, and food scientists. Conventionally, product development requires much work and experiments, which increase with the number of variables evaluated. Furthermore, when experiments are based on trial and error, without an experimental design, there is the possibility of presenting low reproducibility, robustness, versatility, as they are not statistically validated [15].

In this sense, the use of MDs makes it possible to find the best proportion of these components through predictive equations that allow the application of mathematical algorithms, making it possible to determine the optimal conditions for a formulation or an industrial process [2][5][6][7][8][13][16]. The application of these models aims at high quality, low cost, and optimizes single or multiple responses simultaneously (applying desirability functions), and solves challenges with restrictions [5][6][7][16].

Improvements in food, beverage, and health processing techniques have been increasingly recurrent due to the general expectation for higher quality/quantity products today. Jointly, factors that ensure food security to these demands. Thus, MDs can facilitate the development of novel products or techniques that seek to achieve these goals. Among some typical applications, we can list the development of foods and beverages with functional or nutraceutical properties [14][17][18][19], while in the pharmaceutical health area, they are used to formulate drugs for several diseases [15][20][21]. In addition, MDs can also be used for many other applications [5][22][23].

A general representation for the MDs is through the figure of an equilateral triangle for systems with three components. The vertices correspond to the pure components and the sides to the binary mixtures, while the points located inside the triangle represent the mixtures of three components, called the central or centroid point [2][8][9]. The variation of a given property with a mixture's composition is represented by a response surface drawn

above the triangle. Thus, the surface's representation can be done through a triangular diagram of contour lines [2] [12].

Over the years, many MDs types have been developed for a specific purpose with extensive application in science, engineering, and industry [2][12]. Among the most common/used are the simplex lattice design (SLD) [9][11][13], simplex-centroid design (SCD) [9][13][24], simplex axial design (SAD) [9][13][25] and extreme vertex design (EVD) [9][13][26][27][28]. For more details, please see the [Supplementary Material](#). Many works that employ designs for experiments with mixture followed the Scheffé regression models [5][6][7][13][16][29][30]. They can be extended to projects with more than three components [9][29]. The widespread models are represented in Equations (1)–(4).

2. Mixture Designs Applications

2.1. Application of Mixture Design around the World

Initially, all possible applications for MDs around the world were considered. According to the data set, **Figure 1** shows the countries and their continents that most used MDs in the beverage, food, pharmaceutical health, and other science in the last five years.




Figure 1. Application of mixture designs around the world between 2016 and 2020. Africa (a), America (b), Europe (c), and Asia (d).

2.2. Application of Mixture Design by Area

Overall, **Figure 2** shows that MDs have been increasingly applied in different areas of knowledge over the years, being more applied in the pharmaceutical health area, followed by the food and beverage areas. In this sense, researchers have applied MDs to improve processing conditions, new product development, or obtain unique sensory and physicochemical characteristics in products, among other numerous applications described briefly in the following sections.




Figure 2. Number of works by areas and year using MDs according to the search from January 2016 to February 2020.

2.3. Application of Mixture Design in Beverage

Table 1 shows some products that used MDs to improve the conditions for processing or formulating beverage products, classified into four subdivisions following careful previous analysis: alcoholic beverage, milk-based beverages, several beverages, and several juices.

Table 1. Application of mixture design in the beverage.

Papers	n _{ref}	References for Data Analysis
Application in Beverage	37	Refs.
Alcoholic beverage	1	[31]
Milk-based beverages	14	[32][33][34][35][36][37][38][39][40][41][42][43][44][45]
Several beverages	9	[18][46][47][48][49][50][51][52][53]
Several juices	13	[17][54][55][56][57][58][59][60][61][62][63][64][65]

In **alcoholic beverages**, simplex-lattice MDs was applied to craft beers to understand the sensory contribution of different hops (Cascade, Chinook, and Centennial) added to the American beer style. The study demonstrated that it is possible to achieve similar aroma profiles when dry-hopping beer with varying blends of Cascade, Chinook, and Centennial hops [31]. The use of MDs for these products has been scarce, demonstrating the possibility for future studies that seek to formulate alcoholic beverages with functional or antioxidant properties, as well as evaluate the effects of additions of spices in a beverage at different stages of fermentation, mixtures of raw materials, formulations of gluten-free or non-alcoholic beverages, among several other possible applications [14].

In **milk-based beverages**, the most frequent applications of MDs aim to improve the rheological, functional, sensory, and antioxidant properties of yogurts and fermented beverages based on milk. Probiotic products were most frequently evaluated, such as yogurt [34][36][37], fermented milk [39], and chocolate beverages [35]. Furthermore, other conventional products, such as yogurt [32][38][43] and dairy drink [45] formulation; yogurt [41] and milk chocolate [33] diet or light; fermented milk [40][44] and soy milk-based beverage [42] have been reported.

In this works, the authors use MDs for formulations of yogurt or fermented milk symbiotics, probiotics, or organic [34][37][39], with various applications. Such as investigate the influence of starter cultures mixtures for product development [36][43], evaluate the effects of different gums and their interactions on rheological, physicochemical, and sensory properties [32][35][40][44], formulate beverages with low/free sugar content [41][42], formulation of drinks with natural sweeteners [33], fortified with antioxidant [38], or for the formulation of low-cost milk drinks obtained from by-products [45].

In **several beverages**, the MDs were used to formulate some unconventional beverages, such as the development of a drink based on sweet potato peels, and a tea based on rooibos (*Aspalathus linearis*), white tea (*Camellia sinensis* var. *sinensis*), and roasted mate (*Ilex paraguariensis*) optimizing the antioxidant and antiproliferative activity of beverages [18][52]. As well as other formulated beverages, such as the development of banana, strawberry, and juçara smoothie [50], carbonated soft drinks formulated based on tagatose, sucrose, and stevia [47][49], coffee [48][51], peanut drink [46], and soursop [53] beverage formulation. Among the applications of MDs in several beverages, we highlight the authors' attempt to achieve improvements in sensory parameters, antioxidant, physical and chemical properties, and rheological of beverages.

In **several juices**, the applications of the MDs were in functional fruit juices [17][55][56][57][58][59][60][62][63][64], powders [54][61], and the quality of sugarcane [65] juice. In addition, the MDs have often been applied in formulations of probiotic and energetic functional juices to improve these products' sensory, antioxidant, and nutritional properties with potential benefits for consumers' health [54][56][60][61].

2.4. Application of Mixture Design in Food

In these subdivisions, several products are reported in which MDs were used for food formulations (**Table 2**). This group was classified as animal origin, bakery product, dairy frozen, fruit jelly, and vegetable origin according to careful analysis.

Table 2. Application of mixture design in food.

Papers	n _{ref}	References for Data Analysis
Application in Food	39	Refs.
Animal origin	10	[19][66][67][68][69][70][71][72][73][74]
Bakery product	16	[75][76][77][78][79][80][81][82][83][84][85][86][87][88][89][90]
Dairy frozen	3	[91][92][93]
Fruit jelly	5	[94][95][96][97][98]
Vegetable origin	5	[99][100][101][102][103]

In **animal origin** products, MDs have been used to formulate mortadella [68], poultry pâté [69], and beef patties [73] low-fat; sausage [67] and grilled beef [71] with aromatic herbal; sodium-reduced sausages [19][66]; prebiotic sausage formulation [72], gluten-free chicken nugget [70], and honeydew honey [74].

In general, part of the studies have used MDs for formulations of probiotic products of animal origin, such as the formulation of a prebiotic sausage with inulin, konjac, and starch [72] or formulation of a poultry pâté with inulin and lentil flour [69]. Other uses of the MDs were to try to reduce the fat or sodium content in these products. Such as formulating a low-fat mortadella using beef meat, pork meat, and pork back fat [68], formulation of sodium-reduced lean sausages with fish oil seeking partial replacement of NaCl by KCl and sodium tripolyphosphate [19][66], and the formulation of low-fat beef patties with the partial animal fat replacement by cold-pressed grape seed oil and pomegranate seed oil [73].

Other less conventional applications of MDs were for formulations of natural sausages with the addition of aromatic herbs to slow the oxidation of lipids and control the spoilage bacteria in the food [67], or to evaluate the inhibitory effect of spices/herbs and their mixtures on the formation of heterocyclic amines and mutagenic activity of grilled beef by MDs [71]. Moreover, there was only one report for the development of gluten-free industrialized animal products, in which the authors formulated a gluten-free pasta based on three Thai rice flour for use in a frozen battered chicken nugget [70].

In **bakery products**, MDs have been used in functional milk [85], chocolate [82], pecan nut [89][90], and sponge [86] cake formulations; gluten-free formulation of bread [75][81] and cakes [76]; noodle [88] and rice bread [87] formulations; cake [77] and staple bread [83] formulation; cassava crackers enriched [84]; flour [79] and whole-food [78] blend formulation and cereal recipe [80].

Among the most frequent applications of MDs in bakery products are formulations of gluten-free products, such as the formulation of bread using sorghum flour, rice flour, and millet flour [75]; a gluten-free bread based on chickpea flour [81], and a gluten-free cake enriched with protein [76]. Other applications of MDs extend the incorporation of products with different purposes, such as inserting jujube flour in the sponge cake [86], the addition of tikhur starch

as a substitute for semolina in the preparation of baked milk cake [85], formulation of chocolate cake with partial replacement of wheat flour with yacon and maca flour [82], formulation of cassava-fish crackers with a combination of ingredients such as cassava starch, high-quality cassava flour, and fish flour [84], formulation of rice noodles that include combinations of gelatinized corn starch, xanthan gum, and guar gum to improve the tensile strength of cooked noodles [88] and the formulation of a low-cost fiber-rich whole food from residues from the processing of orange juice [78].

In **dairy frozen**, the use of MDs was scarce. Among the applications, the use of MDs to optimize the combination of stabilizers (stone gum, carboxymethylcellulose, and guar gum) in the ice cream formulation [93], for the formulation of ice cream mix powder using milk protein, fat, sucrose, stabilizers, emulsifiers, and water [91], and formulation of antioxidants-rich ice cream containing aqueous extract rich in phenolic compounds and potential functional properties made of *Ilex paraguariensis*, *Melissa officinalis*, and *Cymbopogon citratus* [92].

In **fruit jelly**, the MDs were used for functional jelly formulations. Such as the formulation of a jelly enriched with apple puree, concentrated apple juice, and juice from sea-buckthorn berries added during the gelation of jelly [96] or the optimization of native Brazilian fruit jelly with jabuticaba, pitanga, and cambuci based on sensory and nutritional characteristics [94]. Other jelly formulations have been reported, such as jelly with Swazi indigenous fruits tincizi, tineyi, and umfomfo [95], jelly formulation of red fruit with blackberry, blueberry, and strawberry [97], and produce a pestil (fruit leather) from commercial pomegranate juice using xanthan gum, locust bean gum, and pregelatinized starch [98].

In **vegetable origin** products, the use of MDs went to tomato sauce formulated with tomato puree, onion puree, and extra virgin olive oil [103], elaboration of probiotic creamy sauce-based soymilk with okara flour fermented by *Lactobacillus acidophilus* [100]; oxidative stability of soybean oil with natural antioxidant from Tunisian aromatic plants [101][102] and replacement of salt NaCl by KCl and CaCl₂ in tomato soup using electronic tongue and mixture design [99].

2.5. Application of Mixture Design in Pharmaceutical Health

According to **Table 3**, the main applications of MDs in pharmaceutical health are for the development of novel products or the formulation of several drugs. This area was subdivided into five classes: drug formulation, encapsulation, extraction compounds, functional activity, and foams/or films-based.

Table 3. Application of mixture design in pharmaceutical health.

Papers	<i>n_{ref}</i>	References for Data Analysis
Application in Pharmaceutical Health	59	Refs.
Drug formulation	36	[15][20][21][104][105][106][107][108][109][110][111][112][113][114][115][116][117][118][119][120][121][122][123][124][125][126][127][128][129][130][131][132][133][134][135][136]

Papers	<i>n</i> _{ref}	References for Data Analysis
Encapsulation	6	[137] [138] [139] [140] [141] [142]
Extraction compounds	6	[143] [144] [145] [146] [147] [148]
Functional activity	9	[149] [150] [151] [152] [153] [154] [155] [156] [157]
Foams/films-based	2	[158] [159]

In **drug formulation**, the greatest number of studies using MDs stands out. The application of this method in developing pharmaceutical products has been more explored than in other areas. In this sense, the researchers have used MDs to optimize the composition of formulations and simultaneously estimate the main effects and the interaction of all variables in a drug formulation for several diseases, such as osteoporosis, cardiovascular, cancer, AIDS, dermatological, tuberculosis, gastrointestinal, etc.

The use of MDs has also been explored for the formulation of **encapsulants**. In this step, it is common to mix two or more components to improve specific properties of drugs, and a great way to evaluate the interaction between these encapsulating agents is through MDs. Among some applications, we can list the optimization of the ratio of materials to produce the cinnamon essential oil microcapsules by spray-drying with gum Arabic, maltodextrin, and inulin [\[137\]](#); or to evaluate the effect of encapsulating agents (gum Arabic, modified starch Capsul™ and maltodextrin DE 5) on anthocyanin retention in microcapsules produced by spray-drying of raw pomegranate juice [\[138\]](#). In addition, to formulate a bionanocomposites for entrapment of probiotic cells (*Bacillus coagulans*) using bacterial nanocellulose, pectin, and *Schizophyllum commune* [\[139\]](#); or for the formulation of Turkish oregano extract microcapsules prepared by spray-drying using different concentrations of maltodextrin and gum Arabic as encapsulating agent [\[140\]](#); or to check the performance of *Zingiber zerumbet* oil encapsulation [\[141\]](#); or encapsulation using carbohydrate, protein, coconut oil mixtures on the viability of probiotic cells (*Lactobacillus bulgaricus*) during spray-drying [\[142\]](#).

The **functional activity** can have several classes, where antioxidant and antimicrobial are the activities most combined with MDs recently. MDs have been used to evaluate the synergistic effects of antioxidant activity on mixtures of the essential oil from *Apium graveolens* L., *Thymus vulgaris* L., *Coriandrum sativum* L., and essential oils from *Ocimum basilicum* L., *Origanum majorana* L., and *Rosmarinus officinalis* L. [\[150\]](#)[\[151\]](#)[\[153\]](#). To maximize the antimicrobial activity of essential oils combined against *Escherichia coli* in milk [\[149\]](#) and *Salmonella typhimurium* [\[152\]](#), or to optimize strain mixture of *Lactobacillus* with the highest antimicrobial activity against common food-borne pathogenic bacteria (*Escherichia coli*, *Salmonella enteritidis*, *Listeria monocytogenes*, and *Bacillus cereus*) [\[154\]](#), the nematocidal activity of artemisia extract [\[155\]](#), maqui berry extract as an antioxidant/anti-inflammatory agent [\[156\]](#) and optimization of soy protein isolate, bovine whey protein and egg white protein hydrolysis with the protease Flavourzyme™ 500 L [\[157\]](#).

Certainly, the combination of essential oils generally expresses some interaction effect, and they can be synergistic, antagonistic, additive, or indifferent. In this sense, the MDs use has presented excellent perspectives

for future studies to evaluate the effectiveness of several types of functional activities of these products [14].

MDs have been applied to **extract compounds** from various medicinal plants, such as anthocyanins and phenolic compounds from jabuticaba skin and seed, by mixing solvents [143]; to optimize the activity of total phenolic and flavonoids compounds from mixture herbs *Cnestis palala*, *Urceola micrantha*, *Labisia pumila*, and *Microporus xanthopus* [145]; optimize the solvent mixture to extract total phenolic content and antioxidant capacity of camu-camu (*Myrciaria dubia*) seeds [147]; optimize the solvent proportions for extraction of aporphine alkaloids from the leaves of *Unonopsis duckei* [148]; optimize the extraction of curcuminoids from turmeric using ethyl lactate, ethanol, and water under mild conditions [146]; and optimization of essential oil extraction from Pitanga (*Eugenia uniflora* L.) leaves with petroleum ether, *n*-hexane, methanol, and ethanol [144].

In **foams/films-based**, the MDs were used in the formulation of edible films/coatings composite with different proportions of pectin, alginate, and whey protein concentrate for evaluated for physicochemical characteristics on the final product [159] and formulation of foam of cassava starch, peanut skin, and glycerol to improve mechanical flexural mechanical properties and water absorption capacity [158].

2.6. Application of Mixture Design in Other Areas

Table 4 presents a subdivision of the MDs' possible applications that can be used within the area, and these subdivisions were classified as applications related to the area: animal, bioenergy, biology, and materials.

Table4. Application of mixture design in other areas.

Papers	<i>n</i> _{ref}	References for Data Analysis
Application in Other Areas	23	Refs.
Animal	3	[160][161][162]
Bioenergy	9	[163][164][165][166][167][168][169][170][171]
Biology	4	[1][172][173][174]
Materials	7	[22][23][175][176][177][178][179]

According to **Table 4**, MD research in **bioenergy** aims to optimize biodiesel synthesis from a blend of five different oils [169]; investigate the effects of cellulose, xylan, and lignin constituents on biomass pyrolysis characteristics and bio-oil composition [165]; production of biomethane using municipal sludge wastes, grease trap waste, and meat processing waste [167]; investigate the ability of the oleaginous yeast *Debaryomyces etchellsii* strain for lipid production for biodiesel manufacture using different agro-industrial wastewaters (cheese whey, expired soft drinks and fresh olive mill) as substrates [163]; investigate the effect of light wavelength on *Dunaliella salina* to increase microalgae lipid productivity for biodiesel production [164]; optimization of viscosity and density of refined palm oil-Melaleuca Cajuputi oil binary blends for a novel biofuel [168]; formulation of a vermicomposting in organic matters with dairy manure vermicompost, straw, and peat using MDs in cucumber seedling experiment to evaluate the

compressed substrates [171]; optimization of concentrations of the solvents ethanol, acetone and water in the extraction of total phenolics present in cashew apple bagasse [166] and optimization of a method for the determination of metals from palm oil by flame atomic absorption spectrometry [170].

For the **materials** area, MDs have been used in glass formulation using iron phosphate base glass system that contained P_2O_5 , Fe_2O_3 , Al_2O_3 , Na_2O , and SO_3 [22]; hydraulic oil formulation with different materials based on specific restrictions applying multiresponse optimization [177]; analysis of the effects of nano-oil additives (using ZnO , Si_3N_4 and carbon nanotubes) on the wear properties of AISI 4140 steel material [178]; formulate and examine the mechanical properties of cotton shell particles integrated into glass-fiber-reinforced polymer composites [175]; optimization of bitumen formulations using asphaltic residue, vacuum residue, and three aromatic extracts (by-products) from the refining process of base oils [176]; formulation of alkali-activated cement mortars incorporating glass powder, slag, and calcium aluminate cement [23]; and investigate the arsenic (V) removal from waters using synthetic minerals using synthetic poorly crystallized aluminum hydroxide, calcined layered double hydroxide, and two-line ferrihydrite [179].

In **biology** applications, MDs propose substitutes of bacteriological agar manufacture in terms of texture by using some reported gelling agents used by food and agroindustry [172], metabolomic fingerprint investigation of reference and crossed coffees [1], secondary metabolites from *Mikania laevigata* leaves [173] and development of a chromatography method to determine drugs of abuse [174].

In the **animal** area, MDs were used for the development of essential oil-based phyto-formulations to control the cattle tick *Rhipicephalus microplus* using a mixture of the cinnamon (*Cinnamomum zeylanicum*), cumin (*Cuminum cyminum*), and allspice (*Pimenta dioica*) [160]; formulation of functional foods for animals mixing complex extracts of *Rhodiola crenulata*, *Astragalus membranaceus*, and *Panax quinquefolius* [161]; and development of an analytical chromatography method for the simultaneous investigation of veterinary drugs in poultry litter [162].

3. Behaviors, Trends, and Perspectives of Mixture Design Applications

Regarding the use of mixture design, the main perspective raised was the development of healthier products, such as the formulation of symbiotic products, which combine probiotics and prebiotics in the form of synergism, and products that aim to reduce the content of sodium, fat, sugar, or gluten from food. Given the greater concern with food quality and improved quality of life, consumers have increasingly chosen these products, which have leveraged and heated the market in this sector. Moreover, there is a growing interest in formulations of free gluten, sugar, or lactose products, especially for celiac, diabetic, and lactose intolerance prevalence.

Another emerging field of research has been studies related to allergenic substances in foods from animal or plant sources that can cause an overreaction of the immune system. In our search, there were no reports about works on this theme. In this sense, MDs' applications can be great tools for future formulations of these types of products.

We also observed promising prospects for studies to evaluate the substitution of chemical additives added to food and pharmaceutical products, such as antioxidants and synthetic stabilizers. Thus, the replacement by natural products such as herbs, fruits, and co-products from industrial processes are great paths to be followed, as they generate products with greater health benefits, add value to co-products, and have been the focus of intense research. Moreover, MDs are promising tools to assess the effects of synergistic interactions of functional activities, such as the antioxidant, antimicrobial and antiviral activity of essential oils and extracts of natural products.

Recently, the science and technology of the production of alcoholic beverages has been taken to another level. Consumers have shown a preference for functionalized drinks, such as the consumption of craft beers, one of the main reasons for this market's leverage [14]. This trend has been growing and opens the possibility of using MDs for formulations of craft beers and other alcoholic beverages.

Though drug formulation has been the main use of MDs, this sector still has an overall capacity for research in nanotechnology, such as the development/formulation of novel NLCs, SMEDDS, SNEOFs, S-SNEOFs, SNELS, microemulsions, nanoemulsions, microencapsulation, nanosuspensions, and nanocrystal-based products.

Trends in the use of MDs in other areas are growing, some promising fields for the application of these tools have been reported, such as the reuse or use of sources from renewable raw materials and industrial waste, formulations of growth media, development of novel analytical methods, nanotechnology, novel materials such as graphene, silicene, perovskite and metal-organic frameworks (MOFs) with specific characteristics for many applications.

Another general trend is the use of MDs to solve problems with multiple responses: restrictions of maximum, minimum, and target values for the responses that can be applied when seeking a formulation [5][6][14]. The strategy most used to optimize multiple responses makes use of the desirability function (D), proposed by Deringer and Suich [180], and other methods such as overlaid contour plots, constrained nonlinear optimization, and a Bayesian approach, which has also been employed [181]. From a practical point of view, multiresponse optimization makes the use of MDs more attractive. Among some promising examples of the application of the desirability function (D) present in the literature, we can mention the multiresponse optimization of the oxidative stability of biodiesel from natural [16] and synthetic [30] antioxidants, or a multiresponse optimization of the efficiency and cost of synthetic antioxidants added to biodiesel [6], or biodiesel formulation [5], or a multiresponse optimization of the effects of adding spices with antioxidant compounds in craft beer [14].

References

1. Guizellini, F.C.; Marcheafave, G.G.; Rakocevic, M.; Bruns, R.E.; Scarminio, I.S.; Soares, P.K. PARAFAC HPLC-DAD metabolomic fingerprint investigation of reference and crossed coffees. *Food Res. Int.* 2018, 113, 9–17.

2. Barros Neto, B.; Scarminio, I.S.; Bruns, R.E. *Como Fazer Experimentos: Pesquisa e Desenvolvimento na Ciência e na Indústria*, 4th ed.; Bookman: Porto Alegre, Brazil, 2010; ISBN 8577807134.
3. Ferreira, S.L.C. *Introdução às Técnicas de Planejamento de Experimentos*, 1st ed.; Vento Leste: Salvador, Brazil, 2015.
4. Galvan, D.; Aquino, A.; Effting, L.; Mantovani, A.C.G.; Bona, E.; Conte-Junior, C.A. E-sensing and nanoscale-sensing devices associated with data processing algorithms applied to food quality control: A systematic review. *Crit. Rev. Food Sci. Nutr.* 2021, 1–41.
5. Orives, J.R.; Galvan, D.; Coppo, R.L.; Rodrigues, C.H.F.; Angilelli, K.G.; Borsato, D. Multiresponse optimisation on biodiesel obtained through a ternary mixture of vegetable oil and animal fat: Simplex-centroid mixture design application. *Energy Convers. Manag.* 2014, 79, 398–404.
6. Orives, J.R.; Galvan, D.; Pereira, J.L.; Coppo, R.L.; Borsato, D. Experimental design applied for cost and efficiency of antioxidants in biodiesel. *J. Am. Oil Chem. Soc.* 2014, 91.
7. Jain, A.; Hurkat, P.; Jain, S.K. Development of liposomes using formulation by design: Basics to recent advances. *Chem. Phys. Lipids* 2019, 224, 1–16.
8. Calado, V.; Montgomery, D.C. *Planejamento de Experimentos Usando o Statistica*; Editora E-Papers: Rio de Janeiro, Brazil, 2003; ISBN 8587922831.
9. Montgomery, D.C. *Design and Analysis of Experiments*, 6th ed.; John Wiley and Sons: New York, NY, USA, 2005.
10. Galvan, D.; Orives, J.R.; Coppo, R.L.; Rodrigues, C.H.F.; Spacino, K.R.; Pinto, J.P.; Borsato, D.D. Study of oxidation kinetics of B100 biodiesel from soybean and pig fat: Activation energy determination. *Quim. Nova* 2014, 37, 244–248.
11. Scheffé, H. Experiments with Mixtures. *J. R. Stat. Soc. Ser. B* 1958, 20, 344–360.
12. Hill, T.; Lewicki, P. *Statistics: Methods and Applications: A Comprehensive Reference for Science, Industry, and Data Mining*, 1st ed.; StatSoft, Inc.: Tulsa, OK, USA, 2006; ISBN 1884233597.
13. Cornell, J.A. *Experiments with Mixtures: Designs, Models and the Analysis of Mixture Data*, 3rd ed.; Wiley: New York, NY, USA, 2002; ISBN 9780471393672.
14. Nunes Filho, R.C.; Galvan, D.; Effting, L.; Terhaag, M.M.; Yamashita, F.; de Toledo Benassi, M.; Spinosa, W.A. Effects of adding spices with antioxidants compounds in red ale style craft beer: A simplex-centroid mixture design approach. *Food Chem.* 2021, 365, 130478.
15. Das, S.K.; Khanam, J.; Nanda, A. Optimization of preparation method for ketoprofen-loaded microspheres consisting polymeric blends using simplex lattice mixture design. *Mater. Sci. Eng. C* 2016, 69, 598–608.

16. Coppo, R.L.; Pereira, J.L.; Silva, H.C.; Angilelli, K.G.; Rodrigues, P.R.P.; Galvan, D.; Borsato, D. Effect of natural antioxidants on oxidative stability of biodiesel from soybean oil. Applying simplex-centroid design. *J. Biobased Mater. Bioenergy* 2014, 8.

17. Ogundele, O.M.A.; Awolu, O.O.; Badejo, A.A.; Nwachukwu, I.D.; Fagbemi, T.N. Development of functional beverages from blends of Hibiscus sabdariffa extract and selected fruit juices for optimal antioxidant properties. *Food Sci. Nutr.* 2016, 4, 679–685.

18. Anastácio, A.; de Carvalho, I.S. Development of a beverage benchtop prototype based on sweet potato peels: Optimization of antioxidant activity by a mixture design. *Int. J. Food Sci. Nutr.* 2016, 67, 496–506.

19. Restrepo-Molina, D.A.; Arroyave-Maya, W.; González-Rodríguez, D.M.; Sepúlveda-Valencia, J.U.; Ciro-Velásquez, H.J. Design of a sodium-reduced preservative mixture for use in standard frankfurter sausages. *DYNA* 2019, 86, 17–24.

20. Bonda, A.F.; Rinaldi, M.; Segale, L.; Palugan, L.; Cerea, M.; Vecchio, C.; Pattarino, F. Nanonized itraconazole powders for extemporary oral suspensions: Role of formulation components studied by a mixture design. *Eur. J. Pharm. Sci.* 2016, 83, 175–183.

21. Ferreira-Nunes, R.; Gratieri, T.; Gelfuso, G.M.; Cunha-Filho, M. Mixture design applied in compatibility studies of catechin and lipid compounds. *J. Pharm. Biomed. Anal.* 2018, 149, 612–617.

22. Ray, C.S.; Samaranayake, V.A.; Mohammadkhah, A.; Day, T.E.; Day, D.E. Iron phosphate glass waste forms for vitrifying Hanford AZ102 low activity waste (LAW), part I: Glass formation model. *J. Non Cryst. Solids* 2017, 458, 149–156.

23. He, P.; Zhang, B.; Lu, J.X.; Poon, C.S. A ternary optimization of alkali-activated cement mortars incorporating glass powder, slag and calcium aluminate cement. *Constr. Build. Mater.* 2020, 240, 117983.

24. Scheffé, H. The Simplex-Centroid design for experiments with mixtures. *J. R. Stat. Soc. Ser. B Stat. Methodol.* 1963, 25, 235–251.

25. Cornell, J.A. Some comments on designs for cox's mixture polynomial. *Technometrics* 1975, 17, 25–35.

26. McLean, R.A.; Anderson, V.L. Extreme vertices design of mixture experiments. *Technometrics* 1966, 8, 447–454.

27. Piepel, G.F. Defining consistent constraint regions in mixture experiments. *Technometrics* 1983, 25, 97–101.

28. Myers, R.H.; Montgomery, D.C.; Anderson-Cook, C.M. *Response Surface Methodology: Process and Product Optimization Using Designed Experiments*, 2nd ed.; John Wiley & Sons, Inc.: New

York, NY, USA, 1995; ISBN 978-0-471-58100-0.

29. Piepel, G.F.; Szychowski, J.M.; Loeppky, J.L. Augmenting scheffé linear mixture models with squared and/or crossproduct terms. *J. Qual. Technol.* 2002, 34, 297–314.

30. Borsato, D.; Galvan, D.; Pereira, J.L.; Orives, J.R.; Angilelli, K.G.; Coppo, R.L. Kinetic and thermodynamic parameters of biodiesel oxidation with synthetic antioxidants: Simplex centroid mixture design. *J. Braz. Chem. Soc.* 2014, 25.

31. Lafontaine, S.R.; Shellhammer, T.H. Sensory Directed mixture study of beers dry-hopped with cascade, centennial, and chinook. *J. Am. Soc. Brew. Chem.* 2018, 76, 199–208.

32. Kashaninejad, M.; Razavi, S.M.A. The effects of different gums and their interactions on the rheological properties of instant camel yogurt: A mixture design approach. *J. Food Meas. Charact.* 2019, 13, 1299–1309.

33. Homayouni Rad, A.; Pirouzian, H.R.; Toker, O.S.; Konar, N. Application of simplex lattice mixture design for optimization of sucrose-free milk chocolate produced in a ball mill. *LWT Food Sci. Technol.* 2019, 115, 108435.

34. Karnopp, A.R.; Oliveira, K.G.; de Andrade, E.F.; Postingher, B.M.; Granato, D. Optimization of an organic yogurt based on sensorial, nutritional, and functional perspectives. *Food Chem.* 2017, 233, 401–411.

35. Oliveira, D.F.; Granato, D.; Barana, A.C. Development and optimization of a mixed beverage made of whey and water-soluble soybean extract flavored with chocolate using a simplex-centroid design. *Food Sci. Technol.* 2018, 38, 413–420.

36. Ghasempour, Z.; Moghaddas Kia, E.; Golbandi, S.; Ehsani, A. Effects of mixed starters on quality attributes of probiotic yogurt using statistical design approach. *Nutr. Food Sci.* 2019, 49, 158–169.

37. Bevilacqua, A.; Speranza, B.; Campaniello, D.; Sinigaglia, M.; Corbo, M.R.; Lamacchia, C. A preliminary report on the use of the design of experiments for the production of a synbiotic yogurt supplemented with gluten Friendly™ flour and *bifidobacterium infantis*. *Front. Microbiol.* 2019, 10, 1–11.

38. Hernández-Carranza, P.; Jattar-Santiago, K.Y.; Avila-Sosa, R.; Pérez-Xochipa, I.; Guerrero-Beltrán, J.A.; Ochoa-Velasco, C.E.; Ruiz-López, I.I. Antioxidant fortification of yogurt with red cactus pear peel and its mucilage. *CYTA J. Food* 2019, 17, 824–833.

39. Speranza, B.; Bevilacqua, A.; Campaniello, D.; Sinigaglia, M.; Musaico, D.; Corbo, M.R.; Lamacchia, C. The impact of Gluten Friendly flour on the functionality of an active drink: Viability of *Lactobacillus acidophilus* in a fermented milk. *Front. Microbiol.* 2018, 9, 2042.

40. Amaral, T.N.; Junqueira, L.A.; Prado, M.E.T.; Cirillo, M.A.; de Abreu, L.R.; Costa, F.F.; de Resende, J.V. Blends of *Pereskia aculeata* Miller mucilage, guar gum, and gum Arabic added to

fermented milk beverages. *Food Hydrocoll.* 2018, **79**, 331–342.

41. Carvalho, D.A.; Valente, G.D.F.S.; Assumpção, G.M.P. External preference map to evaluate the acceptance of light and diet yogurt prepared using natural sweeteners. *Cienc. Rural* 2018, **48**, 1–9.

42. Getu, R.; Tola, Y.B.; Neela, S. Optimization of soy milk, mango nectar and sucrose solution mixes for a better quality soy milk-based beverage. *Acta Sci. Pol. Technol. Aliment.* 2017, **16**, 379–391.

43. Dahlan, H.A.; Sani, N.A. The interaction effect of mixing starter cultures on homemade natural yogurt's pH and viscosity. *Int. J. Food Stud.* 2017, **6**, 239–243.

44. Sudha, A.; Priyanka Devi, K.S.; Sangeetha, V.; Sangeetha, A. Development of fermented millet sprout milk beverage based on physicochemical property studies and consumer acceptability data. *J. Sci. Ind. Res.* 2016, **75**, 239–243.

45. Quijano-Aviles, M.F.; Franco-Agurto, G.L.; Suárez-Quirumbay, K.B.; Barragán-Lucas, A.D.; Manzano-Santana, P.I. Linear programming formulation of a dairy drink made of cocoa, coffee and orange by-products. *Emir. J. Food Agric.* 2016, **28**, 554–559.

46. Gama, A.P.; Hung, Y.C.; Adhikari, K. Optimization of emulsifier and stabilizer concentrations in a model peanut-based beverage system: A mixture design approach. *Foods* 2019, **8**, 116.

47. Zendeboodi, F.; Yeganehzad, S.; Sadeghian, A. Optimizing the formulation of a natural soft drink based on biophysical properties using mixture design methodology. *J. Food Meas. Charact.* 2018, **12**, 763–769.

48. Marcheafave, G.G.; Pauli, E.D.; Tormena, C.D.; Mattos, L.E.; de Almeida, A.G.; Rakocevic, M.; Bruns, R.E.; Scarminio, I.S. Irrigated and CO₂ level effects on metabolism in *Coffea arabica* beans from mixture design—Near infrared fingerprints. *Microchem. J.* 2020, **152**, 104276.

49. Acevedo, W.; Capitaine, C.; Rodríguez, R.; Araya-Durán, I.; González-Nilo, F.; Pérez-Correa, J.R.; Agosin, E. Selecting optimal mixtures of natural sweeteners for carbonated soft drinks through multi-objective decision modeling and sensory validation. *J. Sens. Stud.* 2018, **33**, e12466.

50. Ribeiro, L.D.O.; Dos Santos, J.G.C.; Gomes, F.D.S.; Cabral, L.M.C.; Sá, D.D.G.C.F.; da Matta, V.M.; Freitas, S.P. Sensory evaluation and antioxidant capacity as quality parameters in the development of a banana, strawberry and juçara smoothie. *Food Sci. Technol.* 2018, **38**, 653–660.

51. Salamanca-Neto, C.A.R.; Marcheafave, G.G.; Scremen, J.; Barbosa, E.C.M.; Camargo, P.H.C.; Dekker, R.F.H.; Scarmínio, I.S.; Barbosa-Dekker, A.M.; Sartori, E.R. Chemometric-assisted construction of a biosensing device to measure chlorogenic acid content in brewed coffee beverages to discriminate quality. *Food Chem.* 2020, **315**, 126306.

52. Santos, J.S.; Deolindo, C.T.P.; Hoffmann, J.F.; Chaves, F.C.; do Prado-Silva, L.; Sant'Ana, A.S.; Azevedo, L.; do Carmo, M.A.V.; Granato, D. Optimized *Camellia sinensis* var. *sinensis*, *Ilex*

paraguaricensis, and Aspalathus linearis blend presents high antioxidant and antiproliferative activities in a beverage model. *Food Chem.* 2018, 254, 348–358.

53. Cheong, K.W.; Mirhosseini, H.; Tabatabaee Amid, B.; Sheikh Abdul Hamid, N.; Tan, C.P. The influence of main emulsion components on the physicochemical properties of soursop beverage emulsions: A mixture design approach. *J. Dispers. Sci. Technol.* 2018, 39, 934–942.

54. Kim, M.B.; Ko, J.Y.; Lim, S. Bin formulation optimization of antioxidant-rich juice powders based on experimental mixture design. *J. Food Process. Preserv.* 2017, 41, 1–10.

55. Curi, P.N.; de Almeida, A.B.; Tavares, B.D.S.; Nunes, C.A.; Pio, R.; Pasqual, M.; de Souza, V.R. Optimization of tropical fruit juice based on sensory and nutritional characteristics. *Food Sci. Technol.* 2017, 37, 308–314.

56. Siche, R.; Ávalos, C.; Arteaga, H.; Saldaña, E.; Vieira, T.M.F.S. Antioxidant capacity of binary and ternary mixtures of orange, grape, and starfruit juices. *Curr. Nutr. Food Sci.* 2015, 12, 65–71.

57. Ntsoane, M.L.; Sivakumar, D.; Mahajan, P.V. Optimisation of O₂ and CO₂ concentrations to retain quality and prolong shelf life of 'shelly' mango fruit using a simplex lattice mixture design. *Biosyst. Eng.* 2020, 192, 14–23.

58. Khalifa, I.; Li, M.; Mamet, T.; Li, C. Maltodextrin or gum Arabic with whey proteins as wall-material blends increased the stability and physicochemical characteristics of mulberry microparticles. *Food Biosci.* 2019, 31, 100445.

59. Sattar, S.; Ahmad, T.; Nisa, M.U.; Imran, M.; Holmes, M.; Maycock, J.; Nadeem, M.; Khan, M.K. Microwave processing impact on physicochemical and bioactive attributes of optimized peach functional beverage. *J. Food Process. Preserv.* 2019, 43, e13952.

60. Gao, Y.; Hamid, N.; Gutierrez-Maddox, N.; Kantono, K.; Kitundu, E. Development of a probiotic beverage using breadfruit flour as a substrate. *Foods* 2019, 8, 214.

61. De Lima, N.D.; Afonso, M.R.A.; da Costa, J.M.C.; Carvalho, J.D.G. Powdered beverage mix with acerola pulp, whey and maltodextrin. *Rev. Cienc. Agron.* 2019, 50, 578–583.

62. Khan, M.K.; Asif, M.N.; Ahmad, M.H.; Imran, M.; Arshad, M.S.; Hassan, S.; Khan, M.I.; Nisa, M.U.; Iqbal, M.M.; Muhammad, N. Ultrasound-assisted optimal development and characterization of stevia-sweetened functional beverage. *J. Food Qual.* 2019, 2019, 5916097.

63. Schiassi, M.C.E.V.; Lago, A.M.T.; de Souza, V.R.; dos Santos Meles, J.; de Resende, J.V.; Queiroz, F. Mixed fruit juices from Cerrado: Optimization based on sensory properties, bioactive compounds and antioxidant capacity. *Br. Food J.* 2018, 120, 2334–2348.

64. Yen, P.P.L.; Kitts, D.D.; Pratap Singh, A. Natural acidification with low-pH Fruits and incorporation of essential oil constituents for organic preservation of unpasteurized juices. *J. Food Sci.* 2018, 83, 2039–2046.

65. Andrade, D.F.; Guedes, W.N.; Pereira, F.M.V. Detection of chemical elements related to impurities leached from raw sugarcane: Use of laser-induced breakdown spectroscopy (LIBS) and chemometrics. *Microchem. J.* 2018, 137, 443–448.

66. Marchetti, L.; Argel, N.; Andrés, S.C.; Califano, A.N. Sodium-reduced lean sausages with fish oil optimized by a mixture design approach. *Meat Sci.* 2015, 104, 67–77.

67. Nurain, A.; Noriham, A.; Zainon, M.N. Optimisation of mechanically deboned chicken meat (MDCM) aromatic herbal sausage formulations using simplex-lattice mixture design. *Int. Food Res. J.* 2018, 25, 1799–1807.

68. Saldaña, E.; Siche, R.; da Silva Pinto, J.S.; de Almeida, M.A.; Selani, M.M.; Rios-Mera, J.; Contreras-Castillo, C.J. Optimization of lipid profile and hardness of low-fat mortadella following a sequential strategy of experimental design. *J. Food Sci. Technol.* 2018, 55, 811–820.

69. Momchilova, M.; Gradinarska, D.; Petrova, T.; Zsivanovits, G.; Bakalov, I.; Penov, N.; Yordanov, D. Inulin and lentil flour as fat replacers in meat-vegetable pâté—A mixture design approach. *Carpathian J. Food Sci. Technol.* 2019, 11, 5–14.

70. Pinkaew, P.; Naivikul, O. Development of gluten-free batter from three Thai rice cultivars and its utilization for frozen battered chicken nugget. *J. Food Sci. Technol.* 2019, 56, 3620–3626.

71. Sepahpour, S.; Selamat, J.; Khatib, A.; Manap, M.Y.A.; Abdull Razis, A.F.; Hajeb, P. Inhibitory effect of mixture herbs/spices on formation of heterocyclic amines and mutagenic activity of grilled beef. *Food Addit. Contam. Part A Chem. Anal. Control. Expo. Risk Assess.* 2018, 35, 1911–1927.

72. Safaei, F.; Abhari, K.; Khosroshahi, N.K.; Hosseini, H.; Jafari, M. Optimisation of functional sausage formulation with konjac and inulin: Using D-Optimal mixture design. *Foods Raw Mater.* 2019, 7, 177–184.

73. Velioglu, H.M. Low-fat beef patties with cold-pressed oils optimized by mixture design. *J. Food Nutr. Res.* 2016, 55, 89–100.

74. Da Silva, B.; Valdomiro Gonzaga, L.; Fett, R.; Oliveira Costa, A.C. Simplex-centroid design and Derringer's desirability function approach for simultaneous separation of phenolic compounds from Mimosa scabrella Bentham honeydew honeys by HPLC/DAD. *J. Chromatogr. A* 2019, 1585, 182–191.

75. Azarbad, H.R.; Mazaheri Tehrani, M.; Rashidi, H. Optimization of gluten-free bread formulation using sorghum, rice, and millet flour by D-optimal mixture design approach. *J. Agric. Sci. Technol.* 2019, 21, 101–115.

76. Bravo-Núñez, Á.; Sahagún, M.; Bravo-Núñez, A.; Gómez, M. Optimisation of protein-enriched gluten-free layer cakes using a mixture design. *Int. J. Food Sci. Technol.* 2019, 1–8.

77. Mohtarami, F. Effect of carrot pomace powder and dushab (traditional grape juice concentrate) on the physical and sensory properties of cakes: A combined mixtures design approach. *Curr. Nutr. Food Sci.* 2019, 15, 572–582.

78. Oduntan, A.O.; Arueya, G.L. Design, formulation, and characterization of a potential ‘whole food’ using fibre rich orange (*Citrus sinensis* Lin) pomace as base. *Bioact. Carbohydr. Diet. Fibre* 2019, 17, 100172.

79. Ohizua, E.R.; Adeola, A.A.; Idowu, M.A.; Sobukola, O.P.; Afolabi, T.A.; Ishola, R.O.; Ayansina, S.O.; Oyekale, T.O.; Falomo, A. Nutrient composition, functional, and pasting properties of unripe cooking banana, pigeon pea, and sweetpotato flour blends. *Food Sci. Nutr.* 2017, 5, 750–762.

80. Rytz, A.; Moser, M.; Lepage, M.; Mokdad, C.; Perrot, M.; Antille, N.; Pineau, N. Using fractional factorial designs with mixture constraints to improve nutritional value and sensory properties of processed food. *Food Qual. Prefer.* 2017, 58, 71–75.

81. Santos, F.G.; Fratelli, C.; Muniz, D.G.; Capriles, V.D. Mixture design applied to the development of chickpea-based gluten-free bread with attractive technological, sensory, and nutritional quality. *J. Food Sci.* 2018, 83, 188–197.

82. Tormena, M.M.L.; de Medeiros, L.T.; de Lima, P.C.; Possebon, G.; Fuchs, R.H.B.; Bona, E. Application of multi-block analysis and mixture design with process variable for development of chocolate cake containing yacon (*Smallanthus sonchifolius*) and maca (*Lepidium meyenii*). *J. Sci. Food Agric.* 2017, 97, 3559–3567.

83. Cherie, Z.; Ziegler, G.R.; Fekadu Gemedé, H.; Zewdu Woldegiorgis, A. Optimization and modeling of teff-maize-rice based formulation by simplex lattice mixture design for the preparation of brighter and acceptable injera. *Cogent Food Agric.* 2018, 4, 1443381.

84. Coker, O.J.; Sobukola, O.P.; Sanni, L.O.; Bakare, H.A.; Kajihausa, O.E.; Adebawale, A.R.A.; Tomlins, K. Quality attributes of cassava-fish crackers enriched with different flours: An optimization study by a simplex centroid mixture design. *J. Food Process Eng.* 2017, 40, 1–11.

85. Franklin, M.E.E.; Pushpadass, H.A.; Kamaraj, M.; Muthurayappa, M.; Battula, S.N. Application of D-optimal mixture design and fuzzy logic approach in the preparation of chhana podo (baked milk cake). *J. Food Process Eng.* 2019, 42, 1–10.

86. Hosseini, H.; Bolourian, S.; Shahidi, F. Extending the shelf-life of sponge cake by an optimized level of jujube fruit flour determined using custom mixture design. *Br. Food J.* 2019, 121, 3208–3232.

87. Locke, J.E.C.; González, L.C.; Loubes, M.A.; Tolaba, M.P. Optimization of rice bread formulation by mixture design and relationship of bread quality to flour and dough attributes. *LWT Food Sci. Technol.* 2019, 113, 108299.

88. Loubes, M.A.; Flores, S.K.; Tolaba, M.P. Effect of formulation on rice noodle quality: Selection of functional ingredients and optimization by mixture design. *LWT Food Sci. Technol.* 2016, 69, 280–286.

89. Maciel, L.G.; Ribeiro, F.L.; Teixeira, G.L.; Molognoni, L.; Nascimento dos Santos, J.; Larroza Nunes, I.; Mara Block, J. The potential of the pecan nut cake as an ingredient for the food industry. *Food Res. Int.* 2020, 127, 1–10.

90. Maciel, L.G.; Teixeira, G.L.; Block, J.M. Dataset on the phytochemicals, antioxidants, and minerals contents of pecan nut cake extracts obtained by ultrasound-assisted extraction coupled to a simplex-centroid design. *Data Br.* 2020, 28, 1–14.

91. Razak, A.; Rahman, A.; Amin, M.; Johari, M.; Parid, M. Optimisation of stabiliser combinations in instant ice cream mix powder formulation via mixture design methodology. *Int. Food Res. J.* 2019, 26, 1689–1698.

92. Gremski, L.A.; Coelho, A.L.K.; Santos, J.S.; Daguer, H.; Molognoni, L.; do Prado-Silva, L.; Sant'Ana, A.S.; da Silva Rocha, R.; da Silva, M.C.; Cruz, A.G.; et al. Antioxidants-rich ice cream containing herbal extracts and fructooligosaccharides: Manufacture, functional and sensory properties. *Food Chem.* 2019, 298, 1–10.

93. BahramParvar, M.; Tehrani, M.M.; Razavi, S.M.A.; Koocheki, A. Application of simplex-centroid mixture design to optimize stabilizer combinations for ice cream manufacture. *J. Food Sci. Technol.* 2015, 52, 1480–1488.

94. Curi, P.N.; de Almeida, A.B.; Pio, R.; Lima, L.C.D.O.; Nunes, C.A.; de Souza, V.R. Optimization of native Brazilian fruit jelly through desirability-based mixture design. *Food Sci. Technol.* 2019, 39, 388–395.

95. Dlamini, N.P.; Solomon, W.K. Optimization of blending ratios of jam from swazi indigenous fruits tincozi (*syzygium cordatum*), tineyi (*phyllodium zeyheri*) and umfomfo (*cephalanthus natalensis* oliv.) using mixture design. *Cogent Food Agric.* 2019, 5, 1–16.

96. Magomedov, G.O.; Zhuravlev, A.A.; Lobosova, L.A.; Zhurakhova, S.N. Optimization of prescription composition of jelly masses using the Scheffe's symplex plan. *Foods Raw Mater.* 2018, 6, 71–78.

97. Schiassi, M.C.E.V.; Salgado, D.L.; Meirelles, B.S.; Lago, A.M.T.; Queiroz, F.; Curi, P.N.; Pio, R.; de Souza, V.R. Berry Jelly: Optimization through desirability-based mixture design. *J. Food Sci.* 2019, 84, 1522–1528.

98. Tontul, I.; Topuz, A. Production of pomegranate fruit leather (pestil) using different hydrocolloid mixtures: An optimization study by mixture design. *J. Food Process Eng.* 2018, 41, 1–12.

99. Barnett, S.M.; Diako, C.; Ross, C.F. Identification of a Salt Blend: Application of the electronic tongue, consumer evaluation, and mixture design methodology. *J. Food Sci.* 2019, 84, 327–338.

100. Moraes Filho, M.L.; Busanello, M.; Prudencio, S.H.; Garcia, S. Soymilk with okara flour fermented by *Lactobacillus acidophilus*: Simplex-centroid mixture design applied in the elaboration of probiotic creamy sauce and storage stability. *LWT Food Sci. Technol.* 2018, 93, 339–345.

101. Meinhart, A.D.; Ferreira da Silveira, T.F.; Rosa de Moraes, M.; Petrarca, M.H.; Silva, L.H.; Oliveira, W.S.; Wagner, R.; André Bolini, H.M.; Bruns, R.E.; Filho, J.T.; et al. Optimization of frying oil composition rich in essential fatty acids by mixture design. *LWT Food Sci. Technol.* 2017, 84, 795–803.

102. Saoudi, S.; Chammem, N.; Sifaoui, I.; Jiménez, I.A.; Lorenzo-Morales, J.; Piñero, J.E.; Bouassida-Beji, M.; Hamdi, M.; LBazzocchi, I. Combined effect of carnosol, rosmarinic acid and thymol on the oxidative stability of soybean oil using a simplex centroid mixture design. *J. Sci. Food Agric.* 2017, 97, 3300–3311.

103. Yu, J.; Gleize, B.; Zhang, L.; Caris-Veyrat, C.; Renard, C.M.G.C. A D-optimal mixture design of tomato-based sauce formulations: Effects of onion and EVOO on lycopene isomerization and bioaccessibility. *Food Funct.* 2019, 10, 3589–3602.

104. Garg, B.; Katare, O.P.; Beg, S.; Lohan, S.; Singh, B. Systematic development of solid self-nanoemulsifying oily formulations (S-SNEOFs) for enhancing the oral bioavailability and intestinal lymphatic uptake of lopinavir. *Colloids Surf. B Biointerfaces* 2016, 141, 611–622.

105. Garg, B.; Beg, S.; Kaur, R.; Kumar, R.; Katare, O.P.; Singh, B. Long-chain triglycerides-based self-nanoemulsifying oily formulations (SNEOFs) of darunavir with improved lymphatic targeting potential. *J. Drug Target.* 2018, 26, 252–266.

106. De Godoi, K.R.R.; Basso, R.C.; Ming, C.C.; da Silva, V.M.; da Cunha, R.L.; Barrera-Arellano, D.; Ribeiro, A.P.B. Physicochemical and rheological properties of soybean organogels: Interactions between different structuring agents. *Food Res. Int.* 2019, 124, 239–251.

107. Granillo-Guerrero, V.G.; Villalobos-Espinosa, J.C.; Alamilla-Beltrán, L.; Téllez-Medina, D.I.; Hernández-Sánchez, H.; Dorantes-Álvarez, L.; Gutiérrez-López, G.F. Optimization of the formulation of emulsions prepared with a mixture of vitamins d and e by means of an experimental design simplex centroid and analysis of colocalization of its components. *Rev. Mex. Ing. Química* 2017, 16, 861–872.

108. Al Hagbani, T.; Altomare, C.; Salawi, A.; Nazzal, S. D-optimal mixture design: Formulation development, mechanical characterization, and optimization of curcumin chewing gums using oppanol® B 12 elastomer as a gum-base. *Int. J. Pharm.* 2018, 553, 210–219.

109. Hasa, D.; Perissutti, B.; Campisi, B.; Grassi, M.; Grabnar, I.; Golob, S.; Mian, M.; Voinovich, D. Quality improvement of melt extruded laminar systems using mixture design. *Eur. J. Pharm. Sci.* 2015, 75, 169–176.

110. Isailović, T.; Dordević, S.; Marković, B.; Randelovic, D.; Cekić, N.; Lukić, M.; Pantelić, I.; Daniels, R.; Savić, S. Biocompatible nanoemulsions for improved aceclofenac skin delivery: Formulation Approach using combined mixture-process experimental design. *J. Pharm. Sci.* 2016, 105, 308–323.

111. Kamboj, S.; Sharma, R.; Singh, K.; Rana, V. Aprepitant loaded solid preconcentrated microemulsion for enhanced bioavailability: A comparison with micronized Aprepitant. *Eur. J. Pharm. Sci.* 2015, 78, 90–102.

112. Kamboj, S.; Rana, V. Quality-by-design based development of a self-microemulsifying drug delivery system to reduce the effect of food on Nelfinavir mesylate. *Int. J. Pharm.* 2016, 501, 311–325.

113. Khurana, R.K.; Beg, S.; Burrow, A.J.; Vashishta, R.K.; Katare, O.P.; Kaur, S.; Kesharwani, P.; Singh, K.K.; Singh, B. Enhancing biopharmaceutical performance of an anticancer drug by long chain PUFA based self-nanoemulsifying lipidic nanomicellar systems. *Eur. J. Pharm. Biopharm.* 2017, 121, 42–60.

114. Lima, S.G.B.; Pinho, L.A.G.; Pereira, M.N.; Gratieri, T.; Sa-Barreto, L.L.; Gelfuso, G.M.; Cunha-Filho, M. Preformulation studies of finasteride to design matrix systems for topical delivery. *J. Pharm. Biomed. Anal.* 2018, 161, 273–279.

115. Malaquias, L.F.B.; Schulte, H.L.; Chaker, J.A.; Karan, K.; Durig, T.; Marreto, R.N.; Gratieri, T.; Gelfuso, G.M.; Cunha-Filho, M. Hot melt extrudates formulated using design space: One Simple process for both palatability and dissolution rate improvement. *J. Pharm. Sci.* 2018, 107, 286–296.

116. Mori, M.; Rossi, S.; Ferrari, F.; Bonferoni, M.C.; Sandri, G.; Chlapanidas, T.; Torre, M.L.; Caramella, C. Sponge-like dressings based on the association of chitosan and sericin for the treatment of chronic skin ulcers. I. design of experiments-assisted development. *J. Pharm. Sci.* 2016, 105, 1180–1187.

117. Mukherjee, R.; Sen, K.; Fontana, L.; Mao, C.; Chaudhuri, B. Quantification of moisture-induced cohesion in pharmaceutical mixtures. *J. Pharm. Sci.* 2019, 108, 223–233.

118. Patel, M.B.; Shaikh, F.; Patel, V.; Surti, N.I. Application of simplex centroid design in formulation and optimization of floating matrix tablets of metformin. *J. Appl. Pharm. Sci.* 2017, 7, 23–30.

119. Pires, F.Q.; Angelo, T.; Silva, J.K.R.; Sá-Barreto, L.C.L.; Lima, E.M.; Gelfuso, G.M.; Gratieri, T.; Cunha-Filho, M.S.S. Use of mixture design in drug-excipient compatibility determinations: Thymol nanoparticles case study. *J. Pharm. Biomed. Anal.* 2017, 137, 196–203.

120. Rao, S.; Barot, T.; Rajesh, K.S.; Jha, L.L. Formulation, optimization and evaluation of microemulsion based gel of Butenafine Hydrochloride for topical delivery by using simplex lattice mixture design. *J. Pharm. Investig.* 2016, 46, 1–12.

121. Riquelme, N.; Sepúlveda, C.; Arancibia, C. Influence of ternary emulsifier mixtures on oxidative stability of nanoemulsions based on avocado oil. *Foods* **2020**, *9*, 42.

122. Rossi, S.; Vigani, B.; Puccio, A.; Bonferoni, M.C.; Sandri, G.; Ferrari, F. Chitosan ascorbate nanoparticles for the vaginal delivery of antibiotic drugs in atrophic vaginitis. *Mar. Drugs* **2017**, *15*, 319.

123. Shah, N.; Seth, A.; Balaraman, R.; Sailor, G.; Java, A.; Gohil, D. Oral bioavailability enhancement of raloxifene by developing microemulsion using D-optimal mixture design: Optimization and in-vivo pharmacokinetic study. *Drug Dev. Ind. Pharm.* **2018**, *44*, 687–696.

124. Apeji, Y.E.; Oyi, A.R.; Isah, A.B.; Allagh, T.S.; Modi, S.R.; Bansal, A.K. Development and optimization of a starch-based co-processed excipient for direct compression using mixture design. *AAPS PharmSciTech* **2018**, *19*, 866–880.

125. Sisinthu, S.P.; Lynn Sarah, C.Y.; Rao, N.K. Optimization of coconut oil based self micro emulsifying drug delivery systems of olmesartan medoxomil by simplex centroid design. *Int. J. Appl. Pharm.* **2016**, *8*, 47–59.

126. Son, H.Y.; Chae, B.R.; Choi, J.Y.; Shin, D.J.; Goo, Y.T.; Lee, E.S.; Kang, T.H.; Kim, C.H.; Yoon, H.Y.; Choi, Y.W. Optimization of self-microemulsifying drug delivery system for phospholipid complex of telmisartan using D-optimal mixture design. *PLoS ONE* **2018**, *13*, 1–17.

127. Tibalinda, P.; Sempombe, J.; Shedafa, R.; Masota, N.; Pius, D.; Temu, M.; Kaale, E. Formulation development and optimization of Lamivudine 300 mg and Tenofovir Disoproxil Fumarate (TDF) 300 mg FDC tablets by D-optimal mixture design. *Helijon* **2016**, *2*, e00207.

128. Visetvichaporn, V.; Kim, K.H.; Jung, K.; Cho, Y.S.; Kim, D.D. Formulation of self-microemulsifying drug delivery system (SMEDDS) by D-optimal mixture design to enhance the oral bioavailability of a new cathepsin K inhibitor (HL235). *Int. J. Pharm.* **2020**, *573*, 1–9.

129. Winarti, L.; Martien, R.; Suwaldi; Hakim, L. An experimental design of SNEDDS template loaded with bovine serum albumin and optimization using D-optimal. *Int. J. Pharm. Clin. Res.* **2016**, *8*, 425–432.

130. Zhou, Z.; Dunn, C.; Khadra, I.; Wilson, C.G.; Halbert, G.W. Influence of physiological gastrointestinal surfactant ratio on the equilibrium solubility of BCS class II drugs investigated using a four component mixture design. *Mol. Pharm.* **2017**, *14*, 4132–4144.

131. Ashara, K.C.; Shah, K.V. Nuts and bolts of oil and temperature in topical preparations of salicylic acid. *Folia Med.* **2017**, *59*, 279–288.

132. Bennani, I.; Yachi, L.; Bennis, S.; Mefetah, H.; Cheikh, A.; Bouatia, M.; Rahali, Y. Oleogels based on vegetable oil and synthetic oil: Evaluation of the effect of the bentone on gelling using a mixture design. *Asian J. Pharm.* **2018**, *12*, 243–254.

133. Bennis, S.; Yachi, L.; Elalaoui, Y.; Bouatia, M.; Cherkaoui, N.; Laatiris, A.; Rahali, Y. Evaluation of the effect of the Organo-bentonite on gelification of Almond oil using a mixture design. *J. Mater. Environ. Sci.* 2017, 8, 2506–2512.

134. Chauhan, S.I.; Nathwani, S.V.; Soniwala, M.M.; Chavda, J.R. Development and characterization of multifunctional directly compressible Co-processed excipient by spray drying method. *AAPS PharmSciTech* 2017, 18, 1293–1301.

135. Cheong, A.M.; Tan, K.W.; Tan, C.P.; Nyam, K.L. Improvement of physical stability properties of kenaf (*Hibiscus cannabinus* L.) seed oil-in-water nanoemulsions. *Ind. Crop. Prod.* 2016, 80, 77–85.

136. Elkomy, M.H.; El Menshawe, S.F.; Abou-Taleb, H.A.; Elkarmalawy, M.H. Loratadine bioavailability via buccal transfersomal gel: Formulation, statistical optimization, in vitro/in vivo characterization, and pharmacokinetics in human volunteers. *Drug Deliv.* 2017, 24, 781–791.

137. Noghabi, M.S.; Molaveisi, M. Microencapsulation optimization of cinnamon essential oil in the matrices of gum Arabic, maltodextrin, and inulin by spray-drying using mixture design. *J. Food Process Eng.* 2019, 43, 1–13.

138. De Araujo Santiago, M.C.P.; Nogueira, R.I.; Paim, D.R.S.F.; Gouvêa, A.C.M.S.; Godoy, R.L.D.O.; Peixoto, F.M.; Pacheco, S.; Freitas, S.P. Effects of encapsulating agents on anthocyanin retention in pomegranate powder obtained by the spray drying process. *LWT Food Sci. Technol.* 2016, 73, 551–556.

139. Khorassani, A.C.; Shojaosadati, S.A. Bacterial nanocellulose-pectin bionanocomposites as prebiotics against drying and gastrointestinal condition. *Int. J. Biol. Macromol.* 2016, 83, 9–18.

140. Baranauskaite, J.; Ivanauskas, L.; Masteikova, R.; Kopustinskiene, D.; Baranauskas, A.; Bernatoniene, J. Formulation and characterization of Turkish oregano microcapsules prepared by spray-drying technology. *Pharm. Dev. Technol.* 2017, 22, 792–803.

141. Rosli, N.A.; Hasham, R.; Aziz, A. Design and physicochemical evaluation of nanostructured lipid carrier encapsulated zingiber zerumbet oil by d- optimal mixture design. *J. Teknol.* 2018, 3, 105–113.

142. Othman, N.Z.; Shadan, N.H.; Ramli, S.; Sarmidi, M.R. Study on the effects of carbohydrate-protein-coconut oil on the viability of *Lactobacillus bulgaricus* during spray drying, simulated gastrointestinal conditions and unrefrigerated storage by simplex-lattice mixture design. *Chem. Eng. Trans.* 2018, 63, 553–558.

143. Paludo, M.C.; Colombo, R.C.; Filho, J.T.; Hermosín-Gutiérrez, I.; Ballus, C.A.; Godoy, H.T. Optimizing the extraction of anthocyanins from the skin and phenolic compounds from the seed of jabuticaba fruits (*Myrciaria jabuticaba* o (Vell.) O. Berg) with ternary mixture experimental designs. *J. Braz. Chem. Soc.* 2019, 30, 1506–1514.

144. Araromi, D.O.; Alade, A.O.; Bello, M.O.; Bakare, T.; Akinwande, B.A.; Jameel, A.T.; Adegbola, S.A. Optimization of oil extraction from Pitanga (*Eugenia uniflora* L.) leaves using simplex centroid design. *Sep. Sci. Technol.* 2017, 52, 1341–1349.

145. Ismail, N.A.; Abu Bakar, M.F.; Kormin, F.; Linatoc, A.C.; Mohamed, M. Application of statistically simplex-centroid mixture design to optimize the TPC and TFC on the proportion of polyherbal formulation used by jakun women. *J. Eng. Appl. Sci.* 2018, 13, 6996–7002.

146. D'Archivio, A.A.; Maggi, M.A.; Ruggieri, F. Extraction of curcuminoids by using ethyl lactate and its optimisation by response surface methodology. *J. Pharm. Biomed. Anal.* 2018, 149, 89–95.

147. Fidelis, M.; do Carmo, M.A.V.; da Cruz, T.M.; Azevedo, L.; Myoda, T.; Miranda Furtado, M.; Boscacci Marques, M.; Sant'Ana, A.S.; Inês Genovese, M.; Young Oh, W.; et al. Camu-camu seed (*Myrciaria dubia*)—From side stream to an antioxidant, antihyperglycemic, antiproliferative, antimicrobial, antihemolytic, anti-inflammatory, and antihypertensive ingredient. *Food Chem.* 2020, 310, 1–14.

148. Rocha, P.R.N.; de Freitas, F.A.; Angolini, C.F.F.; Vasconcelos, L.S.F.; da Silva, A.L.B.; Costa, E.V.; da Silva, F.M.A.; Eberlin, M.N.; Bataglion, G.A.; Soares, P.K.; et al. Statistical mixture design investigation for extraction and quantitation of aporphine alkaloids from the leaves of *Unonopsis duckei* R.E. Fr. by HPLC–MS/MS. *Phytochem. Anal.* 2018, 29, 569–576.

149. Falleh, H.; Ben Jemaa, M.; Djebali, K.; Abid, S.; Saada, M.; Ksouri, R. Application of the mixture design for optimum antimicrobial activity: Combined treatment of *Syzygium aromaticum*, *Cinnamomum zeylanicum*, *Myrtus communis*, and *Lavandula stoechas* essential oils against *Escherichia coli*. *J. Food Process. Preserv.* 2019, 43, 1–11.

150. Crespo, Y.A.; Bravo Sánchez, L.R.; Quintana, Y.G.; Cabrera, A.S.T.; Bermúdez del Sol, A.; Mayancha, D.M.G. Evaluation of the synergistic effects of antioxidant activity on mixtures of the essential oil from *Apium graveolens* L., *Thymus vulgaris* L. and *Coriandrum sativum* L. using simplex-lattice design. *Helion* 2019, 5, 1–6.

151. Baj, T.; Baryluk, A.; Sieniawska, E. Application of mixture design for optimum antioxidant activity of mixtures of essential oils from *Ocimum basilicum* L., *Origanum majorana* L. and *Rosmarinus officinalis* L. *Ind. Crop. Prod.* 2018, 115, 52–61.

152. Fadil, M.; Fikri-Benbrahim, K.; Rachiq, S.; Ihssane, B.; Lebrazi, S.; Chraibi, M.; Haloui, T.; Farah, A. Combined treatment of *Thymus vulgaris* L., *Rosmarinus officinalis* L. and *Myrtus communis* L. essential oils against *Salmonella typhimurium*: Optimization of antibacterial activity by mixture design methodology. *Eur. J. Pharm. Biopharm.* 2018, 126, 211–220.

153. Ouedrhiri, W.; Balouiri, M.; Bouhdid, S.; Moja, S.; Chahdi, F.O.; Taleb, M.; Greche, H. Mixture design of *Origanum compactum*, *Origanum majorana* and *Thymus serpyllum* essential oils: Optimization of their antibacterial effect. *Ind. Crop. Prod.* 2016, 89, 1–9.

154. Yolmeh, M.; Khomeiri, M.; Ahmadi, Z. Application of mixture design to introduce an optimum cell-free supernatant of multiple-strain mixture (MSM) for *Lactobacillus* against food-borne pathogens. *LWT Food Sci. Technol.* 2017, 83, 298–304.

155. Perez-Roman, I.; Garcia-Rodriguez, J.J.; Kiekens, F.; Cordoba-Diaz, D.; Cordoba-Diaz, M. Enhanced nematocidal activity of a novel *Artemisia* extract formulated as a microemulsion. *Nat. Prod. Commun.* 2019, 14, 1–6.

156. Tenci, M.; Rossi, S.; Giannino, V.; Vigani, B.; Sandri, G.; Bonferoni, M.C.; Daglia, M.; Longo, L.M.; Macelloni, C.; Ferrari, F. An in situ gelling system for the local treatment of inflammatory bowel disease (IBD). the loading of maqui (*aristotelia chilensis*) berry extract as an antioxidant and anti-inflammatory agent. *Pharmaceutics* 2019, 11, 611.

157. Castro, R.J.S.; Inacio, R.F.; de Oliveira, A.L.R.; Sato, H.H. Statistical optimization of protein hydrolysis using mixture design: Development of efficient systems for suppression of lipid accumulation in 3T3-L1 adipocytes. *Biocatal. Agric. Biotechnol.* 2016, 5, 17–23.

158. Machado, C.M.; Benelli, P.; Tessaro, I.C. Constrained mixture design to optimize formulation and performance of foams based on cassava starch and peanut skin. *J. Polym. Environ.* 2019, 27, 2224–2238.

159. Chakravartula, S.S.N.; Soccio, M.; Lotti, N.; Balestra, F.; Dalla Rosa, M.; Siracusa, V. Characterization of composite edible films based on pectin/alginate/whey protein concentrate. *Materials* 2019, 12, 2454.

160. Díaz, E.L.; Padilla Camberos, E.; Castillo Herrera, G.A.; Estarrón Espinosa, M.; Espinosa Andrews, H.; Paniagua Buelnas, N.A.; Gutiérrez Ortega, A.; Martínez Velázquez, M. Development of essential oil-based phyto-formulations to control the cattle tick *Rhipicephalus microplus* using a mixture design approach. *Exp. Parasitol.* 2019, 201, 26–33.

161. Do Hong, H.; Kim, J.C.; Lim, T.G.; Song, Y.R.; Cho, C.W.; Jang, M. Mixing ratio optimization for functional complex extracts of *Rhodiola crenulata*, *Panax quinquefolius*, and *Astragalus membranaceus* using mixture design and verification of immune functional efficacy in animal models. *J. Funct. Foods* 2018, 40, 447–454.

162. Teglia, C.M.; Peltzer, P.M.; Seib, S.N.; Lajmanovich, R.C.; Culzoni, M.J.; Goicoechea, H.C. Simultaneous multi-residue determination of twenty one veterinary drugs in poultry litter by modeling three-way liquid chromatography with fluorescence and absorption detection data. *Talanta* 2017, 167, 442–452.

163. Arous, F.; Frikha, F.; Triantaphyllidou, I.E.; Aggelis, G.; Nasri, M.; Mechichi, T. Potential utilization of agro-industrial wastewaters for lipid production by the oleaginous yeast *Debaryomyces etchellsii*. *J. Clean. Prod.* 2016, 133, 899–909.

164. Bredda, E.H.; Da Silva, A.F.; Silva, M.B.; Da Rós, P.C.M. Mixture design as a potential tool in modeling the effect of light wavelength on *Dunaliella salina* cultivation: An alternative solution to increase microalgae lipid productivity for biodiesel production. *Prep. Biochem. Biotechnol.* 2019, 1, 1–11.

165. Fan, Y.; Cai, Y.; Li, X.; Jiao, L.; Xia, J.; Deng, X. Effects of the cellulose, xylan and lignin constituents on biomass pyrolysis characteristics and bio-oil composition using the Simplex Lattice Mixture Design method. *Energy Convers. Manag.* 2017, 138, 106–118.

166. Felix, A.C.S.; Alvarez, L.D.G.; Santana, R.A.; Valasques, G.L., Jr.; Bezerra, M.A.; de Oliveira Neto, N.M.; de Oliveira Lima, E.; de Oliveira Filho, A.A.; Franco, M.; do Nascimento, B.B., Jr. Application of experimental designs to evaluate the total phenolics content and antioxidant activity of cashew apple bagasse. *Rev. Mex. Ing. Quim.* 2018, 17, 165–175.

167. Kashi, S.; Satari, B.; Lundin, M.; Horváth, I.S.; Othman, M. Application of a mixture design to identify the effects of substrates ratios and interactions on anaerobic co-digestion of municipal sludge, grease trap waste, and meat processing waste. *J. Environ. Chem. Eng.* 2017, 5, 6156–6164.

168. Mat, S.C.; Idroas, M.Y.; Teoh, Y.H.; Hamid, M.F. Optimisation of viscosity and density of refined palm Oil-Melaleuca Cajuputi oil binary blends using mixture design method. *Renew. Energy* 2019, 133, 393–400.

169. Nouadjep, N.S.; Nso, E.; Gueguim Kana, E.B.; Kapseu, C. Simplex lattice mixture design application for biodiesel production: Formulation and characterization of hybrid oil as feedstock. *Fuel* 2019, 252, 135–142.

170. Valasques, G.S.; Dos Santos, A.M.P.; Da Silva, D.G.; Alves, J.P.S.; Bezerra, M.A. Use of constrained mixture design in the optimization of a method based on extraction induced by emulsion breaking for the determination of Ca, Mg, Mn, Fe and Zn from palm oil by flame atomic absorption spectrometry. *J. Braz. Chem. Soc.* 2018, 29, 2189–2196.

171. Yang, L.; Cao, H.; Yuan, Q.; Luoa, S.; Liu, Z. Component optimization of dairy manure vermicompost, straw, and peat in seedling compressed substrates using simplex-centroid design. *J. Air Waste Manag. Assoc.* 2018, 68, 215–226.

172. Sanchez-Cardozo, J.; Quintanilla-Carvajal, M.X.; Ruiz-Pardo, R.; Acosta-González, A. Evaluating gelling-agent mixtures as potential substitutes for bacteriological agar: An approach by mixture design. *DYNA* 2019, 86, 171–176.

173. Passari, L.M.Z.G.; Scarminio, I.S.; Marcheafave, G.G.; Bruns, R.E. Seasonal changes and solvent effects on fractionated functional food component yields from *Mikania laevigata* leaves. *Food Chem.* 2019, 273, 151–158.

174. Jacques, A.L.B.; dos Santos, M.K.; Limberger, R.P. Development and validation of a method using dried oral fluid spot to determine drugs of abuse. *J. Forensic Sci.* 2019, 64, 1906–1912.

175. Annamalai, M.; Ramasubbu, R. Optimizing the formulation of E-glass fiber and cotton shell particles hybrid composites for their mechanical behavior by mixture design analysis. *Mater. Tehnol.* 2018, 52, 207–214.

176. Varanda, C.; Portugal, I.; Ribeiro, J.; Silva, A.M.S.; Silva, C.M. Optimization of bitumen formulations using mixture design of experiments (MDOE). *Constr. Build. Mater.* 2017, 156, 611–620.

177. Khalilian, H.; Semnani, A.; Haddadi, H.; Nekoeinia, M. Multiresponse optimization of a hydraulic oil formulation by mixture design and response surface methods. *J. Tribol.* 2016, 138, 1–5.

178. Sahin, Y.B.; Celik, O.N.; Burnak, N.; Demirtas, E.A. Modeling and analysis of the effects of nano-oil additives on wear properties of AISI 4140 steel material using mixture design. *Proc. Inst. Mech. Eng. Part J J. Eng. Tribol.* 2016, 230, 442–451.

179. Dias, A.C.; Fontes, M.P.F.; Reis, C.; Bellato, C.R.; Fendorf, S. Simplex-Centroid mixture design applied to arsenic (V) removal from waters using synthetic minerals. *J. Environ. Manag.* 2019, 238, 92–101.

180. Derringer, G.; Suich, R. Simultaneous optimization of several response variables. *J. Qual. Technol.* 1980, 12, 214–219.

181. Piepel, G.F. 50 years of mixture experiment research: 1955–2004. In *Response Surface Methodology and Related Topics*; World Scientific: Singapore, 2006; pp. 283–327.

Retrieved from <https://encyclopedia.pub/entry/history/show/34444>