

White Sponge Nevus

Subjects: **Genetics & Heredity**

Contributor: Bruce Ren

White sponge nevus is a condition characterized by the formation of white patches of tissue called nevi (singular: nevus) that appear as thickened, velvety, sponge-like tissue.

genetic conditions

1. Introduction

White sponge nevus is a condition characterized by the formation of white patches of tissue called nevi (singular: nevus) that appear as thickened, velvety, sponge-like tissue. The nevi are most commonly found on the moist lining of the mouth (oral mucosa), especially on the inside of the cheeks (buccal mucosa). Affected individuals usually develop multiple nevi. Rarely, white sponge nevi also occur on the mucosae (singular: mucosa) of the nose, esophagus, genitals, or anus. The nevi are caused by a noncancerous (benign) overgrowth of cells.

White sponge nevus can be present from birth but usually first appears during early childhood. The size and location of the nevi can change over time. In the oral mucosa, both sides of the mouth are usually affected. The nevi are generally painless, but the folds of extra tissue can promote bacterial growth, which can lead to infection that may cause discomfort. The altered texture and appearance of the affected tissue, especially the oral mucosa, can be bothersome for some affected individuals.

2. Frequency

The exact prevalence of white sponge nevus is unknown, but it is estimated to affect less than 1 in 200,000 individuals worldwide.

3. Causes

Mutations in the *KRT4* or *KRT13* gene cause white sponge nevus. These genes provide instructions for making proteins called keratins. Keratins are a group of tough, fibrous proteins that form the structural framework of epithelial cells, which are cells that line the surfaces and cavities of the body and make up the different mucosae. The keratin 4 protein (produced from the *KRT4* gene) and the keratin 13 protein (produced from the *KRT13* gene) partner together to form molecules known as intermediate filaments. These filaments assemble into networks that provide strength and resilience to the different mucosae. Networks of intermediate filaments protect the mucosae from being damaged by friction or other everyday physical stresses.

Mutations in the *KRT4* or *KRT13* gene disrupt the structure of the keratin protein. As a result, keratin 4 and keratin 13 are mismatched and do not fit together properly, leading to the formation of irregular intermediate filaments that are easily damaged with little friction or trauma. Fragile intermediate filaments in the oral mucosa might be damaged when eating or brushing one's teeth. Damage to intermediate filaments leads to inflammation and promotes the abnormal growth and division (proliferation) of epithelial cells, causing the mucosae to thicken and resulting in white sponge nevus.

3.1 The genes associated with White sponge nevus

- *KRT13*
- *KRT4*

4. Inheritance

This condition is inherited in an autosomal dominant pattern, which means one copy of the altered gene in each cell can be sufficient to cause the disorder. However, some people who have a mutation that causes white sponge nevus do not develop these abnormal growths; this phenomenon is called reduced penetrance.

5. Other Names for This Condition

- Cannon's disease
- familial white folded mucosal dysplasia
- hereditary leukokeratosis
- hereditary mucosal leukokeratosis
- hereditary oral keratosis
- leukokeratosis of oral mucosa
- leukokeratosis, hereditary mucosal
- nevus of Cannon
- white folded gingivostomatosis
- white gingivostomatitis
- white sponge naevus
- white sponge nevus of Cannon
- white sponge nevus of mucosa
- WSN

References

1. Aghbali A, Pouralibaba F, Eslami H, Pakdel F, Jamali Z. White sponge nevus: a case report. *J Dent Res Dent Clin Dent Prospects*. 2009 Spring;3(2):70-2. doi:10.5681/joddd.2009.017.

2. Kimura M, Nagao T, Machida J, Warnakulasuriya S. Mutation of keratin 4 gene causing white sponge nevus in a Japanese family. *Int J Oral Maxillofac Surg.* 2013 May;42(5):615-8. doi: 10.1016/j.ijom.2012.10.030.
3. Marrelli M, Tatullo M, Dipalma G, Inchegolo F. Oral infection by *Staphylococcus aureus* in patients affected by White Sponge Nevus: a description of two cases occurred in the same family. *Int J Med Sci.* 2012;9(1):47-50.
4. Martelli H Jr, Pereira SM, Rocha TM, Nogueira dos Santos PL, Batista de Paula AM, Bonan PR. White sponge nevus: report of a three-generation family. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2007 Jan;103(1):43-7.
5. Nishizawa A, Nakajima R, Nakano H, Sawamura D, Takayama K, Satoh T, Yokozeki H. A de novo missense mutation in the keratin 13 gene in oral white sponge nevus. *Br J Dermatol.* 2008 Sep;159(4):974-5. doi:10.1111/j.1365-2133.2008.08716.x.
6. Rugg E, Magee G, Wilson N, Brandrup F, Hamburger J, Lane E. Identification of two novel mutations in keratin 13 as the cause of white sponge nevus. *Oral Dis.* 1999 Oct;5(4):321-4.
7. Shibuya Y, Zhang J, Yokoo S, Umeda M, Komori T. Constitutional mutation of keratin 13 gene in familial white sponge nevus. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2003 Nov;96(5):561-5.
8. Songu M, Adibelli H, Diniz G. White sponge nevus: clinical suspicion and diagnosis. *Pediatr Dermatol.* 2012 Jul-Aug;29(4):495-7. doi:10.1111/j.1525-1470.2011.01414.x.
9. Zhang JM, Yang ZW, Chen RY, Gao P, Zhang YR, Zhang LF. Two new mutations in the keratin 4 gene causing oral white sponge nevus in Chinese family. *Oral Dis.* 2009 Jan;15(1):100-5. doi: 10.1111/j.1601-0825.2008.01498.x.

Retrieved from <https://encyclopedia.pub/entry/history/show/12115>