

Protein One-Dimensional Sliding

Subjects: **Biophysics**

Contributor: Xiao-Wen Yang

One-dimensional (1D) sliding of DNA-binding proteins has been observed by numerous kinetic studies. It appears that many of these sliding events play important roles in a wide range of biological processes. However, one challenge is to determine the physiological relevance of these motions in the context of the protein's biological function. Here, we discuss methods of measuring protein 1D sliding by highlighting the single-molecule approaches that are capable of visualizing particle movement in real time. We also present recent findings that show how protein sliding contributes to function.

DNA-binding protein

one-dimensional sliding

single molecule

1. Introduction

The interactions between proteins and DNA are involved in most cellular functions of DNA, including DNA replication, DNA repair and recombination, and gene transcription and regulation. Many if not all protein–DNA interactions are essential for those processes. When the physical properties of genomic DNA have been well described by a worm-like chain model ^{[1][2]}, a number of biophysical studies have provided evidence of protein performing one-dimensional (1D) sliding along the DNA ^{[3][4][5][6]}. These highly dynamic protein–DNA interactions have served as models for understanding the macromolecular interactions in biology.

The concept of protein 1D sliding has been recognized for several decades. DNA/RNA polymerases and helicases that operate one nucleotide after another are early examples that obviously must move in 1D along a DNA strand ^{[7][8]}. Further investigations have found that many sequence-specific DNA-binding proteins, including a number of restriction endonucleases and transcription factors, also slide along the double-stranded DNA (dsDNA) to search for target sites ^{[9][10][11]}. While most studies have inspected protein 1D sliding using relatively simple systems, many complicated biological questions involve protein complex assembly and multiple reaction steps that could not be assessed with these systems. In this review, we have summarized recent studies of protein 1D sliding. Several modes of the sliding and the methods for measuring protein movement have been introduced and discussed here. Of particular interest, we highlight the observations that have ultimately revealed the biological functions of protein 1D sliding.

2. Modes of Protein 1D Sliding

The mechanisms of protein 1D sliding can be categorized into several modes. Probably the most familiar one for biologists is translocation. The majority of DNA/RNA polymerases and helicases hydrolyze nucleoside triphosphate

(NTP) and have been suggested to use the released energy for moving along DNA in a unidirectional manner [12][13][14][15]. This unidirectional movement is defined as translocation (**Figure 1 a**). Moreover, there are other proteins that also translocate along the DNA by NTP hydrolysis but do not couple the directional motions to the nucleotide incorporation or strand separation activity [16]. Those enzymes are referred as translocases. As it can be seen, the translocation is an active motion, and the proteins that perform translocation may be considered as molecular motors on DNA [17][18]. While the protein translocation activity certainly requires NTP hydrolysis, the reverse is not necessarily true—a protein that performs 1D sliding and contains NTPase activity may not translocate along the DNA (see Section 2.3).




Figure 1. Modes of protein 1D sliding. (a) Translocation: protein moves along DNA in a unidirectional way by NTP hydrolysis (NDP: nucleoside diphosphate, Pi: inorganic phosphate). (b) Facilitated diffusion: protein passively slides on DNA to search targets after nonspecific binding. (c) Sliding as a clamp: ring-like protein passively slides on DNA with intermittent DNA contacts. (d) Hopping or jumping: protein diffuses on DNA with a series of disassociation and rebinding events. (e) Intersegment transfer: protein moves from one binding site to another via the formation of a transient DNA loop. The double helix structures represent dsDNA and the arch or ring-like structures represent DNA-binding proteins. The black arrows indicate the directions of protein movements on DNA.

The continuous protein–DNA contacts without dissociation are usually necessary to support facilitated diffusion. However, there are proteins that obviously slide with intermittent DNA contacts. Members of this group include proteins that can form special conformations on DNA, such as a clamp encircling the DNA (**Figure 1 c**). A classic example is the DNA replication factor (β -clamp in prokaryotes, also known as proliferating cell nuclear antigen, or PCNA, in eukaryotes), which is a ring-shaped clamp and requires a clamp loader to load it onto the DNA [19][20]. Coincidentally, the DNA mismatch repair (MMR) proteins, MutS and MutL homologs, also form ATP-dependent ring-like clamps during MMR. These sliding clamps do not always interact with the DNA backbone but can maintain a relatively long lifetime (several minutes) during sliding [21][22][23][24]. It is now well established that the MutS and MutL sliding clamps are actually molecular switches that undergo Brownian motion along the DNA, while their ring-like structures prevent the proteins from disassociating from the DNA [21][24].

Another type of protein sliding with intermittent DNA contacts is hopping/jumping, where the protein diffuses with a series of disassociation and rebinding events ^[3] (**Figure 1 d**). Compared with the facilitated diffusion, one could imagine the protein may slide faster and is able to bypass obstacles during hopping/jumping. As a result, proteins combining facilitated diffusion and hopping/jumping seem to be more efficient when searching for target sites ^[25]. However, distinct from the sliding clamps that are topologically constrained to the DNA, it is now still unclear how the hopping/jumping occurs.

In some scenarios, large DNA molecules can form transient DNA loops. A few proteins are capable of moving between two sites on DNA via these intermediate “loops”, which is referred to as intersegmental transfer (**Figure 1 e**). Intersegmental transfer allows the protein to move from one site of DNA to another without dissociation, thus facilitating the target search ^[3]. However, intersegmental transfer is more likely to occur when the protein contains two or more DNA binding surfaces, since the two binding sites need to be accessed concurrently ^[3]. Therefore, this mode of sliding only relates to distant translocations on large DNA molecules by a limited subset of DNA-binding proteins ^{[26][27]}.

3. Approaches for Observing Protein 1D Sliding

In a typical single-molecule tracking experiment, the DNA is usually immobilized on the quartz surface via the biotin–neutravidin–biotin or digoxigenin– antibody interactions (**Figure 2 a**). DNA molecules can be stretched by laminar flow and linked to the surface at both ends, or they can be linked at a single end and stretched by a continuous laminar flow. A similar method such as “DNA curtains”, which uses a combination of nanofabricated surface patterns and fluid lipid bilayers to align hundreds of DNA molecules (**Figure 2 a**), has been developed recently ^[28]. However, some single-molecule experiments still suffer from the nonspecific protein binding on the surface, where the undesirable background signals significantly reduce the resolutions of the target proteins. Thus, a novel single-molecule tracking platform named “DNA skybridge” has been designed and applied to monitor proteins only on DNA (**Figure 2 a**) ^[29]. Although the quartz barrier used in the DNA skybridge platform has not been commercially available, it is expected that most single-molecule tracking systems can be replaced by this surface-condition-independent and high-throughput method in future. In some single-molecule tracking assays, C-trap optical tweezers combining confocal fluorescence microscope have been used ^[30]. Instead of DNA immobilization on a quartz surface, the C-trap captures a DNA molecule by optical traps (**Figure 2 a**), where the changes of DNA lengths and protein positions can be recorded simultaneously. While the C-trap instrument is now commercially available, one disadvantage is that only one molecule can be observed during a single experiment.




Figure 2. Single-molecule methods for observing protein 1D sliding. **(a)** An illustration of tracking a single protein molecule on DNA using TIRF or confocal microscopy. Four types of DNA immobilization methods are shown, including random immobilization, DNA curtains, DNA skybridges, and C-trap optical tweezers (from left to right). The gray rectangles represent the quartz surfaces and the yellow rectangle represents the lipid bilayers. The black spheres represent the dielectric beads. The fluorophore-labeled proteins (DNA-binding proteins with green dyes) on DNA are shown. **(b)** An illustration of monitoring the distance between DNA and protein using FRET (left) and PIFE (right). The fluorophore-labeled DNA substrates are immobilized on quartz slide surfaces. The DNA-binding proteins slide on DNA and cause FRET (left) or PIFE (right). **(c)** An illustration of detecting the forces associated with DNA–protein interactions using optical tweezers. Three types of experiment designs are shown, including surface-base assay, dumbbell-based assay using two optical traps to tether DNA double ends, and dumbbell-based assay using two optical traps to tether the DNA end and protein, respectively (from top to bottom). The black spheres represent the dielectric beads. **(d)** An illustration of detecting the forces associated with DNA–protein interactions using magnetic tweezers. A general set up (left) and a flow stretching set up (right) are shown. The black spheres represent the superparamagnetic beads. **(e)** An illustration of imaging the different states of DNA–protein interactions over time using AFM. The black box represents the detector which monitors the cantilever deflection.

To monitor the DNA–protein interactions in real time, the fluorophore-labeled proteins can be injected into the chamber after DNA immobilization. For single-molecule tracking, the Cyanine (Cy3 and Cy5) and Alexa Fluor dyes, which are small probes that have both high quantum yields and high photostabilities, are recommended. To conjugate the protein with a fluorophore, methods include Cys-maleimide chemistry [31], hydrazinyl-isopictetaspengler ligation [32], and sortase-mediated reaction [33], which only introduce small tags onto the proteins, and it has been shown that these can largely retain the biological activities of enzymes. In contrast, the commonly

used fluorescent proteins are not recommended due to their large sizes and short lifetimes under TIRF illumination [34].

Another type of force spectroscopy is referred to as magnetic tweezers, which utilize a similar concept as optical tweezers. Instead of using the optical traps, magnetic tweezers contain magnets that provide an external magnetic field to manipulate dozens of magnetic particles simultaneously. In a general magnetic tweezers set up, a superparamagnetic bead is tethered to one end of a DNA molecule, while the other end of the DNA is attached to the slide surface [35] (**Figure 2 d**, left). The external magnets are able to impart both twist and tension to the DNA, making magnetic tweezers an excellent method to study the motions of DNA translocases or topoisomerases [36][37][38]. Alternatively, a flow-stretching set-up has been designed, where the instrument combines the magnetic force and the drag force created by a laminar flow to hold the paramagnetic beads and stretch the DNA (**Figure 2 d**, right). Using the flow-stretching assay, a number of DNA molecules have been tracked in real time, and the DNA replication, unwinding, and excision events were studied [39][40][41]. Due to the large volume of the magnetic field, the magnetic force spectroscopy is able to monitor hundreds of DNA molecules at one time but with lower spatial and temporal resolutions.

Atomic force microscopy (AFM) generates images by scanning a small cantilever with a sharp tip over the surface of a sample. By placing the tip in contact with the molecule of interest and moving the surface with respect to the tip, force can be modulated precisely, thereby changing the cantilever deflection. Cantilever deflection is further monitored by a laser beam reflected from the sharp tip onto a position-sensing detector [42] (**Figure 2 e**). High-speed atomic force microscopy (HS-AFM) has been used to image the DNA–protein complex in real time, where the protein diffusion on DNA can be visualized [43]. Nevertheless, the main limitations of AFM come from the nonspecific interactions between proteins and the surface that may suppress the sliding [43].

4. Biological Significance of Protein 1D Sliding

Protein 1D sliding has been shown to be involved in many biological processes. Here, we summarize and discuss the significance of protein 1D sliding in biology.

One of the well-recognized functions of protein 1D sliding is to accelerate the target search process. For many DNA-binding proteins that specifically recognize particular sequences, such as transcription factors, endonucleases, and DNA methyltransferases, the target sites only constitute a minute fraction of the genome DNA (**Figure 3 a**). Thus, the target search should be highly efficient. Actually, it is the case. Some of the proteins locate their target sites much more rapidly than the theoretical calculated one that uses a general diffusion–collision mechanism [44]. The search process was later modeled as a two-step event, where nonspecific protein–DNA binding first occurs and is followed by a 1D sliding to the target site. For example, the *lacI* repressor [44][45], tumor suppressor p53 [11][46], and endonucleases EcoRV, BcnI, and Endonuclease V [9][47][48][49][50] have been reported to slide along the DNA to facilitate the target search. Interestingly, while the continuous protein–DNA contacts without dissociation were initially believed to be essential for the searching processes [11][47][48], further evidence have shown that the sliding was accompanied by occasional hopping/jumping events, which helped the enzymes to

locate their target sites [9][46][49][50][51][52][53][54]. The combination of sliding with continuous DNA contacts and hopping/jumping suggests a trade-off between speed and accuracy during the target search [52].




Figure 3. Biological functions of protein 1D sliding. (a) An illustration showing the process of an endonuclease targeting a sequence-specific site among the genome by facilitated diffusion. Arrows indicate the sliding of an endonuclease to a target site. (b) An illustration showing a helicase unwinding and rezipping on DNA. Black arrows indicate the directions of helicase translocation. (c) An illustration showing a DNA repair protein scanning along the DNA backbone and recognizing a damage site (red star). (d) An illustration showing the distant communications between a mismatch (red star) and an excision/incision site via MutS and MutL sliding clamps. Arrows indicate the sliding of MutS and MutL. (e) An illustration showing condensin binding to the DNA and extruding it as a loop to spatially organize the chromosomes.

As mentioned above (Section 2.3), β -clamp/PCNA forms a special ring-shaped conformation on DNA that provides a mobile platform to tether the DNA polymerase. While the replication sliding clamp itself usually adopts an intrinsically closed ring-like configuration, a sliding clamp loader belonging to the AAA+ ATPase family first binds ATP and recognizes the β -clamp/PCNA molecule to produce a cleft on the clamp [55]. Then, this complex binds a primer–template junction on DNA to stimulate the ATPase activity of the clamp loader [56][57]. ATP hydrolysis releases the loader from the β -clamp/PCNA and DNA, leading to the closure of the β -clamp/PCNA on substrate DNA [58][59]. Studies have demonstrated that the PCNA moves along DNA using two different sliding modes: it slides by tracking the DNA backbone in the rotational-coupled mode, while it moves at higher rates in the non-helical diffusion mode [60][61]. Most importantly, the replication sliding clamp exhibits a half-life of tens of minutes on DNA that enhances polymerase processivity by more than 1000-fold [62][63]. Other clamp-like proteins that have a long lifetime on DNA, such as the Mre11-Rad50-Nbs1 (MRN) complex, MutS, and MutL homologs, also have been shown to act as processivity factors during DNA repair [40][41][64][65].

DNA damage occurs at a rate of 10,000 to 1,000,000 lesions per cell per day [66]. To maintain the stability of the genome, repair proteins have to recognize DNA lesions efficiently. Many DNA repair proteins have been reported to

scan along the DNA for damage sites (**Figure 3 c**), including the base excision repair protein glycosylases [67][68][69][70], mismatch repair protein MutS homologs [71][72], and nucleotide excision repair protein Rad4, XPA, and UvrB [73][74]. Among these studies, it is generally believed that the 1D sliding is used to follow the DNA backbone for lesion recognition, where a damage site only slightly destabilizes the helical structure of a duplex DNA [71][70][72]. In other words, the scanning processes likely involve rotation-coupled diffusions along the DNA helix, while the proteins are in continuous contact with the duplex DNA [71][75].

References

1. Bustamante, C.; Marko, J.; Siggia, E.; Smith, S. Entropic elasticity of lambda-phage DNA. *Science* 1994, 265, 1599–1600.
2. Smith, S.B.; Cui, Y.; Bustamante, C. Overstretching B-DNA: The Elastic Response of Individual Double-Stranded and Single-Stranded DNA Molecules. *Science* 1996, 271, 795–799.
3. Berg, O.G.; Winter, R.B.; Von Hippel, P.H. Diffusion-driven mechanisms of protein translocation on nucleic acids. 1. Models and theory. *Biochemistry* 1981, 20, 6929–6948.
4. Khoury, A.M.; Lee, H.J.; Lillis, M.; Lu, P. Lac repressor-operator interaction: DNA length dependence. *Biochim. Biophys. Acta Gene Struct. Expr.* 1990, 1087, 55–60.
5. Park, C.S.; Wu, F.Y.; Wu, C.W. Molecular mechanism of promoter selection in gene transcription. II. Kinetic evidence for promoter search by a one-dimensional diffusion of RNA polymerase molecule along the DNA template. *J. Biol. Chem.* 1982, 257, 6950–6956.
6. Ricchetti, M.; Metzger, W.; Heumann, H. One-dimensional diffusion of Escherichia coli DNA-dependent RNA polymerase: A mechanism to facilitate promoter location. *Proc. Natl. Acad. Sci. USA* 1988, 85, 4610–4614.
7. Kabata, H.; Kurosawa, O.; Arai, I.; Washizu, M.; Margaron, S.A.; Glass, R.E.; Shimamoto, N. Visualization of Single Molecules of RNA Polymerase Sliding Along DNA. *Science* 1993, 262, 1561–1563.
8. Eggleston, A.; Rahim, N.A.; Kowalczykowski, S.C. A Helicase Assay Based on the Displacement of Fluorescent, Nucleic Acid-Binding Ligands. *Nucleic Acids Res.* 1996, 24, 1179–1186.
9. Bonnet, I.; Biebricher, A.; Porté, P.-L.; Loverdo, C.; Bénichou, O.; Voituriez, R.; Escudé, C.; Wende, W.; Pingoud, A.; Desbiolles, P. Sliding and jumping of single EcoRV restriction enzymes on non-cognate DNA. *Nucleic Acids Res.* 2008, 36, 4118–4127.
10. Rau, D.C.; Sidorova, N.Y. Diffusion of the Restriction Nuclease EcoRI along DNA. *J. Mol. Biol.* 2010, 395, 408–416.

11. Tafvizi, A.; Huang, F.; Leith, J.S.; Fersht, A.R.; Mirny, L.A.; van Oijen, A.M. Tumor Suppressor p53 Slides on DNA with Low Friction and High Stability. *Biophys. J.* 2008, 95, L01–L03.
12. Guajardo, R.; Sousa, R. A model for the mechanism of polymerase translocation. *J. Mol. Biol.* 1997, 265, 8–19.
13. Gelles, J.; Landick, R. RNA Polymerase as a Molecular Motor. *Cell* 1998, 93, 13–16.
14. Moore, K.J.; Lohman, T.M. Helicase-catalyzed DNA unwinding: Energy coupling by DNA motor proteins. *Biophys. J.* 1995, 68, 180S–185S.
15. Lohman, T.M.; Bjornson, K.P. Mechanisms of helicase-catalyzed DNA unwinding. *Annu. Rev. Biochem.* 1996, 65, 169–214.
16. Terakawa, T.; Bisht, S.; Eeftens, J.M.; Dekker, C.; Haering, C.H.; Greene, E.C. The condensin complex is a mechanochemical motor that translocates along DNA. *Science* 2017, 358, 672–676.
17. Bath, J.; Wu, L.J.; Errington, J.; Wang, J.C. Role of *Bacillus subtilis* SpoIIIE in DNA Transport Across the Mother Cell-Prespore Division Septum. *Science* 2000, 290, 995–997.
18. Aussel, L.; Barre, F.-X.; Aroyo, M.; Stasiak, A.; Stasiak, A.Z.; Sherratt, D. FtsK Is a DNA Motor Protein that Activates Chromosome Dimer Resolution by Switching the Catalytic State of the XerC and XerD Recombinases. *Cell* 2002, 108, 195–205.
19. Kong, X.-P.; Onrust, R.; O'Donnell, M.; Kuriyan, J. Three-dimensional structure of the β subunit of *E. coli* DNA polymerase III holoenzyme: A sliding DNA clamp. *Cell* 1992, 69, 425–437.
20. Hingorani, M.M.; O'Donnell, M. ATP Binding to the *Escherichia coli* Clamp Loader Powers Opening of the Ring-shaped Clamp of DNA Polymerase III Holoenzyme. *J. Biol. Chem.* 1998, 273, 24550–24563.
21. Liu, J.; Hanne, J.; Britton, B.M.; Bennett, J.; Kim, D.; Lee, J.-B.; Fishel, R. Cascading MutS and MutL sliding clamps control DNA diffusion to activate mismatch repair. *Nature*. 2016, 539, 583–587.
22. Gradia, S.; Acharya, S.; Fishel, R. The Human Mismatch Recognition Complex hMSH2-hMSH6 Functions as a Novel Molecular Switch. *Cell* 1997, 91, 995–1005.
23. Gradia, S.; Subramanian, D.; Wilson, T.; Acharya, S.; Makhov, A.; Griffith, J.; Fishel, R. hMSH2–hMSH6 Forms a Hydrolysis-Independent Sliding Clamp on Mismatched DNA. *Mol. Cell* 1999, 3, 255–261.
24. London, J.; Martín-López, J.; Yang, I.; Liu, J.; Lee, J.-B.; Fishel, R. Linker domain function predicts pathogenic MLH1 missense variants. *Proc. Natl. Acad. Sci. USA* 2021, 118, e2019215118.

25. Halford, S.E.; Marko, J.F. How do site-specific DNA-binding proteins find their targets? *Nucleic Acids Res.* 2004, 32, 3040–3052.
26. Doucleff, M.; Clore, G.M. Global jumping and domain-specific intersegment transfer between DNA cognate sites of the multidomain transcription factor Oct-1. *Proc. Natl. Acad. Sci. USA* 2008, 105, 13871–13876.
27. Suzuki, Y.; Gilmore, J.L.; Yoshimura, S.H.; Henderson, R.M.; Lyubchenko, Y.L.; Takeyasu, K. Visual Analysis of Concerted Cleavage by Type IIF Restriction Enzyme SfiI in Subsecond Time Region. *Biophys. J.* 2011, 101, 2992–2998.
28. Collins, B.E.; Ye, L.F.; Duzdevich, D.; Greene, E.C. DNA curtains: Novel tools for imaging protein-nucleic acid interactions at the single-molecule level. *Methods Cell Biol.* 2014, 123, 217–234.
29. Kim, D.; Rashid, F.; Cho, Y.; Zaher, M.S.; Cho, I.I.H.; Hamdan, S.M.; Jeong, C.; Lee, J.-B. DNA skybridge: 3D structure producing a light sheet for high-throughput single-molecule imaging. *Nucleic Acids Res.* 2019, 47, e107.
30. Qin, Z.; Bi, L.; Hou, X.-M.; Zhang, S.; Zhang, X.; Lu, Y.; Li, M.; Modesti, M.; Xi, X.-G.; Sun, B. Human RPA activates BLM's bidirectional DNA unwinding from a nick. *eLife* 2020, 9, e54098.
31. Roy, R.; Hohng, S.; Ha, T. A practical guide to single-molecule FRET. *Nat. Methods* 2008, 5, 507–516.
32. Liu, J.; Hanne, J.; Britton, B.M.; Shoffner, M.A.; Albers, A.E.; Bennett, J.; Zatezalo, R.; Barfield, R.M.; Rabuka, D.; Lee, J.-B.; et al. An Efficient Site-Specific Method for Irreversible Covalent Labeling of Proteins with a Fluorophore. *Sci. Rep.* 2015, 5, 16883.
33. Theile, C.S.; Witte, M.D.; Blom, A.E.M.; Kundrat, L.; Ploegh, H.L.; Guimaraes, C.P. Site-specific N-terminal labeling of proteins using sortase-mediated reactions. *Nat. Protoc.* 2013, 8, 1800–1807.
34. Seefeldt, B.; Kasper, R.; Seidel, T.; Tinnefeld, P.; Dietz, K.-J.; Heilemann, M.; Sauer, M. Fluorescent proteins for single-molecule fluorescence applications. *J. Biophotonics* 2008, 1, 74–82.
35. Saleh, O.A.; Péral, C.; Barre, F.-X.; Allemand, J.-F. Fast, DNA-sequence independent translocation by FtsK in a single-molecule experiment. *EMBO J.* 2004, 23, 2430–2439.
36. Strick, T.; Croquette, V.; Bensimon, D. Single-molecule analysis of DNA uncoiling by a type II topoisomerase. *Nature* 2000, 404, 901–904.
37. Harada, Y.; Ohara, O.; Takatsuki, A.; Itoh, H.; Shimamoto, N.; Kinosita, K. Direct observation of DNA rotation during transcription by Escherichia coli RNA polymerase. *Nature* 2001, 409, 113–115.

38. Dessinges, M.-N.; Lionnet, T.; Xi, X.G.; Bensimon, D.; Croquette, V. Single-molecule assay reveals strand switching and enhanced processivity of UvrD. *Proc. Natl. Acad. Sci. USA* 2004, 101, 6439–6444.
39. Lee, J.-B.; Hite, R.K.; Hamdan, S.; Xie, X.S.; Richardson, C.C.; van Oijen, A. DNA primase acts as a molecular brake in DNA replication. *Nature* 2006, 439, 621–624.
40. Jeon, Y.; Kim, D.; Martín-López, J.V.; Lee, R.; Oh, J.; Hanne, J.; Fishel, R.; Lee, J.-B. Dynamic control of strand excision during human DNA mismatch repair. *Proc. Natl. Acad. Sci. USA* 2016, 113, 3281–3286.
41. Liu, J.; Lee, R.; Britton, B.M.; London, J.A.; Yang, K.; Hanne, J.; Lee, J.-B.; Fishel, R. MutL sliding clamps coordinate exonuclease-independent *Escherichia coli* mismatch repair. *Nat. Commun.* 2019, 10, 5294.
42. Greenleaf, W.J.; Woodside, M.T.; Block, S.M. High-Resolution, Single-Molecule Measurements of Biomolecular Motion. *Annu. Rev. Biophys. Biomol. Struct.* 2007, 36, 171–190.
43. Shibata, M.; Nishimasu, H.; Kodera, N.; Hirano, S.; Ando, T.; Uchihashi, T.; Nureki, O. Real-space and real-time dynamics of CRISPR-Cas9 visualized by high-speed atomic force microscopy. *Nat. Commun.* 2017, 8, 1430.
44. Riggs, A.; Bourgeois, S.; Cohn, M. The lac repressor-operator interaction: III. Kinetic studies. *J. Mol. Biol.* 1970, 53, 401–417.
45. Wang, Y.M.; Austin, R.H.; Cox, E.C. Single Molecule Measurements of Repressor Protein 1D Diffusion on DNA. *Phys. Rev. Lett.* 2006, 97, 048302.
46. Tafvizi, A.; Huang, F.; Fersht, A.R.; Mirny, L.A.; van Oijen, A.M. A single-molecule characterization of p53 search on DNA. *Proc. Natl. Acad. Sci. USA* 2011, 108, 563–568.
47. Biebricher, A.; Wende, W.; Escudé, C.; Pingoud, A.; Desbiolles, P. Tracking of Single Quantum Dot Labeled EcoRV Sliding along DNA Manipulated by Double Optical Tweezers. *Biophys. J.* 2009, 96, L50–L52.
48. Klionsky, D.J.; Abdalla, F.C.; Abeliovich, H.; Abraham, R.T.; Acevedo-Arozena, A.; Adeli, K.; Agholme, L.; Agnello, M.; Agostinis, P.; Aguirre-Ghiso, J.A.; et al. Guidelines for the use and interpretation of assays for monitoring autophagy. *Autophagy* 2012, 8, 445–544.
49. Kostiuk, G.; Dikić, J.; Schwarz, F.W.; Sasnauskas, G.; Seidel, R.; Siksnys, V. The dynamics of the monomeric restriction endonuclease BcnI during its interaction with DNA. *Nucleic Acids Res.* 2017, 45, 5968–5979.
50. Ahmadi, A.; Rosnes, I.; Blicher, P.; Diekmann, R.; Schüttelpelz, M.; Glette, K.; Tørresen, J.; Bjørås, M.; Dalhus, B.; Rowe, A.D. Breaking the speed limit with multimode fast scanning of DNA by Endonuclease V. *Nat. Commun.* 2018, 9, 5381.

51. Terakawa, T.; Kenzaki, H.; Takada, S. p53 Searches on DNA by Rotation-Uncoupled Sliding at C-Terminal Tails and Restricted Hopping of Core Domains. *J. Am. Chem. Soc.* 2012, 134, 14555–14562.
52. Marklund, E.; Van Oosten, B.; Mao, G.; Amselem, E.; Kipper, K.; Sabantsev, A.; Emmerich, A.; Globisch, D.; Zheng, X.; Lehmann, L.C.; et al. DNA surface exploration and operator bypassing during target search. *Nature* 2020, 583, 858–861.
53. Subekti, D.R.G.; Murata, A.; Itoh, Y.; Takahashi, S.; Kamagata, K. Transient binding and jumping dynamics of p53 along DNA revealed by sub-millisecond resolved single-molecule fluorescence tracking. *Sci. Rep.* 2020, 10, 13697.
54. Cuculis, L.; Abil, Z.; Zhao, H.; Schroeder, C.M. TALE proteins search DNA using a rotationally decoupled mechanism. *Nat. Chem. Biol.* 2016, 12, 831–837.
55. Turner, J.; Hingorani, M.M.; Kelman, Z.; O'Donnell, M. The internal workings of a DNA polymerase clamp-loading machine. *EMBO J.* 1999, 18, 771–783.
56. Ellison, V.; Stillman, B. Opening of the Clamp: An intimate view of an ATP-driven biological machine. *Cell* 2001, 106, 655–660.
57. Hedglin, M.; Kumar, R.; Benkovic, S.J. Replication Clamps and Clamp Loaders. *Cold Spring Harb. Perspect. Biol.* 2013, 5, a010165.
58. Sakato, M.; O'Donnell, M.; Hingorani, M.M. A Central Swivel Point in the RFC Clamp Loader Controls PCNA Opening and Loading on DNA. *J. Mol. Biol.* 2012, 416, 163–175.
59. Kelch, B.A.; Makino, D.L.; O'Donnell, M.; Kuriyan, J. Clamp loader ATPases and the evolution of DNA replication machinery. *BMC Biol.* 2012, 10, 34.
60. Kochaniak, A.B.; Habuchi, S.; Loparo, J.J.; Chang, D.J.; Cimprich, K.A.; Walter, J.; van Oijen, A.M. Proliferating Cell Nuclear Antigen Uses Two Distinct Modes to Move along DNA. *J. Biol. Chem.* 2009, 284, 17700–17710.
61. De March, M.; Merino, N.; Barrera-Vilarmau, S.; Crehuet, R.; Onesti, S.; Blanco, F.J.; De Biasio, A. Structural basis of human PCNA sliding on DNA. *Nat. Commun.* 2017, 8, 13935.
62. Mizrahi, V.; Henrie, R.N.; Marlier, J.F.; Johnson, K.A.; Benkovic, S.J. Rate-limiting steps in the DNA polymerase I reaction pathway. *Biochemistry* 1985, 24, 4010–4018.
63. Leu, F.P.; O'Donnell, M. Interplay of Clamp Loader Subunits in Opening the β Sliding Clamp of *Escherichia coli* DNA Polymerase III Holoenzyme. *J. Biol. Chem.* 2001, 276, 47185–47194.
64. Lammens, K.; Bemeleit, D.J.; Möckel, C.; Clausing, E.; Schele, A.; Hartung, S.; Schiller, C.B.; Lucas, M.; Angermüller, C.; Söding, J.; et al. The Mre11:Rad50 Structure Shows an ATP-Dependent Molecular Clamp in DNA Double-Strand Break Repair. *Cell* 2011, 145, 54–66.

65. Myler, L.; Gallardo, I.F.; Soniat, M.; Deshpande, R.; Gonzalez, X.B.; Kim, Y.; Paull, T.T.; Finkelstein, I.J. Single-Molecule Imaging Reveals How Mre11-Rad50-Nbs1 Initiates DNA Break Repair. *Mol. Cell* 2017, 67, 891–898.
66. Alhmoud, J.F.; Woolley, J.F.; Al Moustafa, A.-E.; Malki, M.I. DNA Damage/Repair Management in Cancers. *Cancers* 2020, 12, 1050.
67. Nelson, S.R.; Dunn, A.R.; Kathe, S.D.; Warshaw, D.M.; Wallace, S.S. Two glycosylase families diffusively scan DNA using a wedge residue to probe for and identify oxidatively damaged bases. *Proc. Natl. Acad. Sci. USA* 2014, 111, E2091–E2099.
68. Rowland, M.M.; Schonhoft, J.D.; McKibbin, P.L.; David, S.S.; Stivers, J.T. Microscopic mechanism of DNA damage searching by hOGG1. *Nucleic Acids Res.* 2014, 42, 9295–9303.
69. Peng, S.; Wang, X.; Zhang, L.; He, S.; Zhao, X.S.; Huang, X.; Chen, C. Target search and recognition mechanisms of glycosylase AlkD revealed by scanning FRET-FCS and Markov state models. *Proc. Natl. Acad. Sci. USA* 2020, 117, 21889–21895.
70. Blainey, P.; van Oijen, A.; Banerjee, A.; Verdine, G.L.; Xie, X.S. A base-excision DNA-repair protein finds intrahelical lesion bases by fast sliding in contact with DNA. *Proc. Natl. Acad. Sci. USA* 2006, 103, 5752–5757.
71. Jeong, C.; Cho, W.-K.; Song, K.-M.; Cook, C.; Yoon, T.-Y.; Ban, C.; Fishel, R.; Lee, J.-B. MutS switches between two fundamentally distinct clamps during mismatch repair. *Nat. Struct. Mol. Biol.* 2011, 18, 379–385.
72. Gorman, J.; Chowdhury, A.; Surtees, J.A.; Shimada, J.; Reichman, D.R.; Alani, E.; Greene, E.C. Dynamic Basis for One-Dimensional DNA Scanning by the Mismatch Repair Complex Msh2-Msh6. *Mol. Cell* 2007, 28, 359–370.
73. Springall, L.; Hughes, C.D.; Simons, M.; Azinas, S.; Van Houten, B.; Kad, N.M. Recruitment of UvrBC complexes to UV-induced damage in the absence of UvrA increases cell survival. *Nucleic Acids Res.* 2018, 46, 1256–1265.
74. Beckwitt, E.C.; Jang, S.; Detweiler, I.C.; Kuper, J.; Sauer, F.; Simon, N.; Bretzler, J.; Watkins, S.C.; Carell, T.; Kisker, C.; et al. Single molecule analysis reveals monomeric XPA bends DNA and undergoes episodic linear diffusion during damage search. *Nat. Commun.* 2020, 11, 1356.
75. Blainey, P.; Luo, G.; Kou, S.C.; Mangel, W.F.; Verdine, G.L.; Bagchi, B.; Xie, X.S. Nonspecifically bound proteins spin while diffusing along DNA. *Nat. Struct. Mol. Biol.* 2009, 16, 1224–1229.

Retrieved from <https://encyclopedia.pub/entry/history/show/37876>