

# Enzyme Immobilization Techniques

Subjects: [Biochemical Research Methods](#)

Contributor: Mohammed K. Al-Sakkaf , Ibrahim Basfer , Mustapha Iddrisu , Salem A. Bahadi , Mustafa S. Nasser , Basim Abussaud , Qasem A. Drmash , Sagheer A. Onaizi

Researchers have explored the technique of enzyme immobilization as a means to overcome limitations that free enzymes encounter, including reduced performance, high costs, and impracticality for large-scale applications. Enzymatic treatment offers a sustainable and eco-friendly alternative to conventional physicochemical treatment methods, such as adsorption, coagulation, and advanced oxidation processes. Free enzymes are biodegradable, highly efficient, and selective biological catalysts that can operate under mild conditions, thus reducing energy consumption and minimizing the generation of harmful byproducts.

enzyme immobilization

wastewater treatment

## 1. Introduction

Environmental pollution has become a major global concern, with a wide range of contaminants, such as phenolic compounds and organic dyes, posing significant threats to water quality and ecosystem health [\[1\]\[2\]\[3\]\[4\]](#). The increasing release of these toxic and recalcitrant substances into the environment, originating from various industrial wastewater streams such as textile, pharmaceutical, and petrochemical industries, has caused widespread concern [\[5\]\[6\]](#). Contaminated water sources can lead to adverse effects on aquatic life and have potentially detrimental consequences for human health, as these pollutants may exhibit carcinogenic, mutagenic, or teratogenic effects [\[7\]](#). In particular, understanding the effect of organic dyes and phenolic compounds on human health is crucial for the development of effective policies and remediation strategies to minimize the risks associated with these contaminants [\[8\]](#). As the demand for clean water continues to increase due to population growth and industrialization, there is an urgent need to develop effective, efficient, and sustainable strategies for environmental remediation, focusing on the removal of these pollutants from wastewater [\[8\]\[9\]\[10\]](#).

Numerous conventional techniques have been employed to treat phenolic wastewater and organic dyes, each with its own advantages and disadvantages. For instance, adsorption is simple, flexible, and can be highly efficient with low capital cost, but the performance is adsorbent-dependent, and some adsorbents can be expensive [\[11\]\[12\]\[13\]\[14\]\[15\]\[16\]\[17\]](#). Distillation, while effective, can be energy-intensive and may not be suitable for all types of contaminants [\[18\]\[19\]\[20\]\[21\]](#). Chemical oxidation can be highly effective, but it often involves the use of harsh chemicals that can be harmful to the environment [\[22\]\[23\]\[24\]](#). Extraction is a fast and simple process, but it requires highly selective solvents, which can be expensive and harmful to the environment [\[25\]\[26\]\[27\]\[28\]](#). Membrane separation is easy to operate with high selectivity, but it can be expensive and suffers from issues such as membrane fouling [\[29\]\[30\]\[31\]](#).

[32]. Photocatalytic oxidation is a promising technique, but it often requires specific conditions to be effective [33][34][35][36].

Given these limitations of conventional techniques, there has been growing interest in environmental protection through the effective treatment of polluting streams [37][38][39][40], the utilization of sustainable and biodegradable materials [41][42][43][44][45][46][47][48], and the application of microorganisms for the biological treatment of wastewater contaminated with organic dyes, and phenolic compounds have also been explored [49][50]. Microorganisms utilize various enzymatic systems for the oxidative transformation of organic molecules, including laccases, ligninases, tyrosinases, monooxygenases, and dioxygenases [51]. Flavoenzymes, known as azoreductases, are present in both microorganisms and higher eukaryotes and are involved in the detoxification and biotransformation of azo dyes [52]. Furthermore, microorganisms utilize specific intracellular enzymes, namely oxidoreductases, to catalyze the metabolism of phenolic compounds [53]. These enzymatic systems offer several advantages over conventional techniques, including cost-effectiveness, sustainability, and the ability to operate under mild conditions [54][55][56]. This highlights the critical function of enzymes in the biodegradation processes of various organic pollutants, demonstrating the potential of harnessing these enzymatic systems for efficient and eco-friendly pollutant removal strategies [56]. As a result, a new approach has arisen in recent times where extracellular enzymes are utilized instead of whole microbial cells for the remediation of wastewater contaminated with organic substances.

Enzymes act as highly effective biological catalysts that enable specific reactions to occur. The lock and key model or the induced fit model can be used to explain their efficiency. By reducing the activation energy and stabilizing the transition state, enzymes enhance the reaction rate [57][58]. Enzymes possess desirable qualities such as high efficiency, high selectivity, and the ability to operate under milder conditions compared to other chemical catalysts. Enzymes offer a cost-effective advantage in that they operate under mild conditions, eliminating the need for expensive equipment that would otherwise be required for chemical catalysts to achieve extreme conditions such as high pressure or temperature. Their natural origin also makes them environmentally friendly due to their biodegradability and low environmental impact.

There has been growing interest in employing enzymes for the treatment of dye wastewater, and previous research has examined the use of various enzymes, including soybean peroxidase [59][60], horseradish peroxidase (HRP), lignin peroxidase (LiP) [61], and laccase [62][63], for their potential in treating dyes. Peroxidase, a member of the oxidoreductase enzyme family, can enable the oxidation of diverse substances in the presence of an oxidizing agent like chlorine, hydrogen peroxide, and potassium permanganate. A significant use of peroxidase is its ability to degrade aromatic compounds, especially synthetic dyes. This occurs when they are decomposed into individual components and the oxidative polymerization of phenolic compounds is triggered, leading to the creation of insoluble polymers [58][64]. Hydrogen peroxide plays a crucial role in the catalytic cycle of peroxidase enzymes. The reaction begins with the reaction between the Fe(III) state of peroxidase and hydrogen peroxide, leading to the formation of a high-oxidation-state intermediate consisting of a cation radical based on porphyrin and an Fe(IV) oxo ferryl center [65]. After the initial oxidation, the process consists of two reduction steps that bring the peroxidase back to its original state, compound II, with the production of free radicals. These free radicals then undergo

polymerization. However, a high concentration of hydrogen peroxide may inhibit the process, leading to a decrease in enzymatic activity [\[66\]](#).

The application of enzymes for dye wastewater treatment at an industrial scale is often hindered by various limitations such as elevated production expenses, reduced long-term operational stability, restricted reusability, and limited shelf life after the initial use [\[59\]\[67\]](#). In their crude form, enzymes may exhibit limited catalytic activity because of their vulnerability to inhibition, particularly in the case of complex dye wastewater [\[68\]](#). Heavy metals can have a detrimental effect on enzymatic activity, as different enzymes show varying degrees of sensitivity to these substances [\[58\]](#). In certain cases, heavy metals, such as mercury, can react with the reactive groups present in enzymes and render them incapable of catalyzing further reactions [\[69\]](#). As the complexity of wastewater effluent increases, it is expected that enzyme activity will decline, further underscoring the challenges associated with the use of enzymes in the treatment of wastewater [\[70\]\[71\]](#).

In recent years, the use of free enzymes has gained considerable attention among the various proposed methods for removing phenolic compounds [\[72\]\[73\]\[74\]\[75\]](#). Enzymatic treatment offers a sustainable and eco-friendly alternative to conventional physicochemical treatment methods, such as adsorption, coagulation, and advanced oxidation processes [\[76\]\[77\]](#). Free enzymes are biodegradable, highly efficient, and selective biological catalysts that can operate under mild conditions, thus reducing energy consumption and minimizing the generation of harmful byproducts [\[78\]\[79\]](#).

Researchers have explored the technique of enzyme immobilization as a means to overcome limitations that free enzymes encounter, including reduced performance, high costs, and impracticality for large-scale applications [\[56\]\[80\]\[81\]](#). Enzymes have limited operational stability, which can negatively impact their catalytic efficiency. Factors like temperature, pH, and exposure to harsh conditions or solvents can lead to enzyme denaturation, degradation, or aggregation, thereby limiting their effectiveness [\[82\]](#). Enzyme immobilization can improve the stability of enzymes as it offers a physical support system that safeguards them against destabilizing agents while maintaining their original structure [\[80\]\[83\]](#). The recovery and reuse of free enzymes after a reaction can be time-consuming and costly, but enzyme immobilization allows for easy separation from the reaction mixture and enables reuse for several reaction cycles, thereby decreasing the overall costs of enzyme usage [\[84\]\[85\]\[86\]](#). In some reactions, free enzymes can experience reduced catalytic activity due to mass transfer limitations, substrate and product inhibition, or poor substrate solubility in water. Immobilization can overcome these challenges by creating custom biocatalytic systems with improved mass transfer properties, enhanced enzyme-substrate interactions, and optimized reaction conditions [\[54\]](#). Moreover, free enzymes may exhibit low selectivity in certain reactions, especially when working with chiral compounds or complex substrate mixtures. Immobilization has the potential to enhance selectivity by enabling precise control over enzyme orientation and creating a microenvironment that promotes selective catalysis [\[87\]\[88\]](#).

According to Nguyen et al. [\[89\]](#), immobilized enzymes are more effective in eliminating phenolic compounds than free enzymes due to the synergistic effect of enzymatic reactions and pollutant adsorption on the solid support. While not examining the competition between the adsorption of pollutants and products or the influence of product

adsorption on enzyme activity, previous research has demonstrated that the adsorption capacity of the support medium deteriorates after multiple applications, even in the absence of enzymes. This suggests that irreversible adsorption of pollutants or products could be involved in the overall process, particularly when immobilized enzymes are recycled. Nguyen et al. [89] suggested that addressing the removal of irreversibly adsorbed pollutants or products could potentially enhance the effectiveness of the immobilized enzyme system.

There is a broad range of nanomaterials that are widely employed for the purpose of enzyme immobilization. This includes but is not limited to metal oxides [90], carbon dots [91], covalent organic frameworks (COFs) [92], graphene [93], CNTs [94], and MOFs [95]. Each of these nanomaterials possesses unique properties that make them suitable for enzyme immobilization. They often exhibit a high surface area to volume ratio, excellent conductivity, good chemical stability, and strong adsorption capabilities, which make them advantageous in improving the performance of immobilized enzymes [96].

Metal oxides like titanium dioxide and zinc oxide offer robustness, chemical stability, and biocompatibility, making them useful for enzyme immobilization [90][97]. Carbon dots, with their superior optical properties and biocompatibility, also have applications in this area [91]. COFs, due to their designable structures, large pore size, and high surface area, provide ideal platforms for enzyme immobilization [92]. However, this review specifically focuses on the unique advantages of graphene, CNTs, and MOFs for enzyme immobilization. Graphene and its derivatives, such as graphene oxide (GO), are widely used for enzyme immobilization due to their high surface area, excellent thermal and electrical conductivity, and strong  $\pi$ - $\pi$  stacking interactions, which allow effective enzyme adsorption and retention of their bioactivity [83][98]. CNTs offer similar benefits, with additional advantages coming from their tubular structure, which provides a protective environment for enzymes, enhancing their stability and reusability [99][100]. MOFs, with their highly ordered structures and large surface areas, offer unique possibilities for enzyme immobilization. Their pore size, shape, and functionality can be finely tuned, allowing the accommodation of a wide range of enzymes while preserving their activity and stability [101][102].

## 2. Enzyme Immobilization Techniques

Enzyme immobilization is a technique to improve enzyme stability and reusability while maintaining their activity. It refers to the physical or chemical confinement of enzymes in a distinct phase different from the substrate's phase [103]. These techniques can be classified into two broad categories: physical and chemical methods [103][104].

### 2.1. Physical Techniques

Physical immobilization, as the earliest form of immobilization, only involves physical interactions. In this method, neither the immobilizer nor the immobilization agent is changed, linked, or modified. This technique includes encapsulation, entrapment, and adsorption. These processes do not necessitate a covalent bond between the enzyme and the support, therefore maintaining the enzyme's native structure [105]. Adsorption involves enzymes interacting with a support material through forces such as hydrophobic interactions or salt bridges, while entrapment is a technique where enzymes are confined within gels or fibers using covalent or non-covalent bonds.

Similarly, encapsulation secures enzymes within semi-permeable capsules, allowing for the movement of small substrates or products while restricting the migration of larger enzymes [104][105][106].

### 2.1.1. Adsorption

Enzymes can be adsorbed onto support materials through interactions such as hydrophobic forces and salt bridges. Enzyme adsorption onto the support physically can be achieved by immersing the support material in the enzyme solution or by drying enzymes onto electrode surfaces. This immobilization method protects the adsorbed enzymes from factors such as proteolysis, aggregation, and interaction with hydrophobic surfaces [107]. Scientists have utilized eco-friendly materials as enzyme supports to promote sustainable practices. For instance, coconut fibers can retain high amounts of water and have strong cation exchange properties, microcrystalline cellulose has a strong binding capacity, and kaolin offers good enzyme retention through micro/mesoporous materials and chemical acetylation with thiol functionalization and large surface areas that are suitable for redox reactions [108][109][110][111][112][113]. Silanized molecular sieves have been found to be an effective support for enzyme adsorption, owing to the presence of silanols on the surface of the pores that allow for enzyme immobilization through the process of hydrogen bonding [114]. Modifications to the current support materials could potentially enhance enzyme immobilization. Prior investigations have delineated the water activity patterns of polypropylene hydrophobic granules-bound lipase, notably Accurel EP-100 [115]. It was observed that reducing the particle size of Accurel has a positive effect on reaction rates and enantiomeric ratios during biocatalysis [116].

To improve both process control and the cost-effectiveness of production, the immobilization of *Yarrowia lipolytica* lipase on supports like octadecyl-sepabeads and octyl-agarose through physical adsorption has been explored. As a result of this process, there were significant improvements in yields and a tenfold increase in stability when compared to free lipase. Octadecyl-sepabeads, which are hydrophobic in nature, enhance the affinity between the enzyme and support, explaining this observation [117]. After being adsorbed onto biodegradable poly (3-hydroxybutyrate-co-hydroxyvalerate), *Candida rugosa* lipase was able to retain 94% of its activity after four hours at 50 °C and could be reused for up to 12 cycles [118]. The supports were selected due to their flexible and less ordered nature when compared to polyhydroxybutyrate. Byssus threads activated with 1,4-butanediol diglycidyl ether provided a suitable matrix for immobilizing urease, leading to enhanced pH stability and maintaining 50% of the activity of the enzyme under dry conditions [119]. In recent years, biocompatible mesoporous silica nanoparticles (MSNs) have gained attention as an environmentally sustainable support for biocatalysis. The use of these supports not only reduces production costs but also avoids ethical concerns. Due to their durability and effectiveness, MSNs have been applied in energy-related biocatalytic processes [120]. **Table 1** presents the benefits, challenges, and solutions for overcoming the limitations of the adsorption technique.

**Table 1.** Common immobilization techniques, their advantages and disadvantages, and suggested approaches to overcome limitations.

Immobilization Technique	Advantages	Drawbacks	Approaches to Address the Limitations	Ref.
Adsorption	<ul style="list-style-type: none"> <li>Prevention of proteolysis</li> <li>Full activity retention</li> </ul>	<ul style="list-style-type: none"> <li>Non-targeted adsorption</li> <li>The expense of affinity binding</li> <li>The activity is affected by a slight shift in the reaction conditions</li> <li>The leaching of enzymes</li> </ul>	<ul style="list-style-type: none"> <li>Using a blocking agent to reduce interactions that aren't specific</li> <li>Specific pH for the charge difference between the silica support and the enzyme</li> <li>Pore size decrease following adsorption</li> </ul>	<a href="#">[121]</a> <a href="#">[122]</a> <a href="#">[123]</a>
Entrapment	<ul style="list-style-type: none"> <li>Moderate preparation circumstances</li> <li>Prevents direct contact with the environment outside</li> </ul>	<ul style="list-style-type: none"> <li>Limited mobility on mass transfer</li> <li>Leakage is the result of fewer physical restraints</li> </ul>	<ul style="list-style-type: none"> <li>Exact pore size selection based on enzyme size</li> <li>Further covalent fusion</li> </ul>	<a href="#">[124]</a> <a href="#">[125]</a> <a href="#">[126]</a>
Encapsulation	<ul style="list-style-type: none"> <li>Maintenance of enzymatic activity over prolonged periods</li> <li>Easy passage of small substrate molecules</li> <li>Large enzymes confined within the capsules</li> </ul>	<ul style="list-style-type: none"> <li>Difficulties in ensuring optimal diffusion of substrates and products</li> <li>Maintaining the structural integrity of the capsules under operational conditions</li> </ul>	<ul style="list-style-type: none"> <li>Development of capsules with improved stability, selectivity, and permeability</li> <li>Advances in materials science for better encapsulation materials and methods, such as 3D capsules</li> </ul>	<a href="#">[126]</a> <a href="#">[127]</a> <a href="#">[128]</a> <a href="#">[129]</a>
Covalent binding	<ul style="list-style-type: none"> <li>Reduced limitations of mass transfer</li> </ul>	<ul style="list-style-type: none"> <li>Specific binding site</li> </ul>	<ul style="list-style-type: none"> <li>Support and enzyme modification</li> </ul>	<a href="#">[130]</a> <a href="#">[131]</a>

Immobilization Technique	Advantages	Drawbacks	Approaches to Address the Limitations	Ref.
	<ul style="list-style-type: none"><li>Improved storage and stability of reaction</li><li>Stronger bonding</li></ul>	<ul style="list-style-type: none"><li>Denaturation of the enzyme's active site</li><li>Irreversible binding</li></ul>	<ul style="list-style-type: none"><li>Specific binding site</li></ul>	<a href="#">[132]</a>
Cross-linking	<ul style="list-style-type: none"><li>Aggregates may experience increased activity</li><li>Recyclability, higher loading capacity, and total activity retention</li></ul>	<ul style="list-style-type: none"><li>The cross-linking matrix's fragility</li><li>Agents that precipitate conflict</li><li>The pure enzyme is necessary for cross-linking enzyme crystals</li></ul>	<ul style="list-style-type: none"><li>The ideal aggregate size determined by the cross-linker-to-enzyme ratio</li><li>Stabilizing components for the structure</li><li>Using cross-linking enzyme crystals for enzymes that haven't been fully purified</li></ul>	<a href="#">[133]</a>

Effective entrapment has been realized with hybrid carriers made of alginate, gelatin, and calcium, which prevent the enzyme from leakage and offer increased mechanical stability [\[134\]](#). The implementation of nanostructured materials in enzyme immobilization, such as pristine materials and electrospun nanofibers, which are produced through a method known as electrospinning, has significantly impacted the field. Mesoporous silica entrapment has recently emerged as a highly promising technology in fields such as biomedicine, fine chemistry, biosensors, and biofuels. This is largely due to the material's unique properties, including a large surface area, uniform pore distribution, adjustable pore size, and high adsorption capacity. These features enable mesoporous silica to serve as an effective support material for various applications [\[135\]](#). Lipase and magnetite entrapment of nanoparticles simultaneously within biomimetic silica has been shown to increase activity with various silane additives [\[136\]](#). In the meantime, the selective binding and carrying properties of sol–gel matrices with supramolecular calixarene polymers have been used to entrap *C. rugosa* lipase [\[137\]](#)[\[138\]](#). In **Table 1**, entrapment's advantages, disadvantages, and strategies to tackle its limitations are summarized.

2.1.3. Encapsulation

The method of encapsulation immobilization entails the confinement of a variety of biomolecules within distinct polymeric structures [\[139\]](#). This process shares similarities with entrapment, as both techniques permit enzymes and cells to exist freely within a solution while remaining in a controlled environment. Encapsulation aims to secure delicate enzymes and cellular solutions within small vesicles with porous barriers, preventing larger enzymes from exiting or entering the capsules, while smaller substrates or products can traverse the semi-permeable barrier with ease [\[127\]](#). This method allows for the preservation of biological systems within a thin protective film, preventing



direct environmental exposure that could negatively affect the performance of the biocatalysts, hence, enabling the prolonged activity of these biocatalysts [140]. Various supportive materials, such as cellulose nitrate and nylon, are employed in the production of microcapsules that range in size from 10 to 100  $\mu\text{m}$  [141]. Furthermore, the process of ionotropic gelation of alginates and nanoporous silica-based sol–gel glasses has proven its efficacy in the field of enzyme encapsulation.

The simplicity of the encapsulation process distinguishes it, and advancements in material sciences have led to the improvement of this method, with benefits such as increased morphological stability, customizable physicochemical permeability, and reduced enzyme leakage [141]. The technique also offers the potential for co-immobilization, allowing for the possibility of immobilizing enzymes in any combination as required. Nevertheless, the method is not without its limitations. For example, issues related to diffusion can be significant, with the risk of membrane rupture if reaction products accumulate rapidly [128]. **Table 1** provides a summary of the benefits, limitations, and strategies to overcome the challenges associated with the encapsulation method.

## 2.2. Chemical Techniques

Chemical methods involve the formation of strong covalent bonds between the enzyme and the support, leading to higher stability and reusability. Chemical techniques include covalent binding, cross-linking, and affinity immobilization. Covalent binding attaches enzymes to supports through covalent bonds formed with specific amino acids in the enzyme's side chains [124]. Cross-linking forms covalent bonds between enzyme molecules using bifunctional or multifunctional agents. Affinity immobilization is a technique that utilizes the enzyme's specific binding properties to support materials under different physiological conditions [104][142].

### 2.2.1. Covalent Binding

Enzymes can be attached to supports through covalent binding, which relies on specific amino acids in the enzyme's side chains, such as arginine, aspartic acid, and histidine. The effectiveness of this process is largely determined by the reactivity and efficiency of the functional groups present in the support, such as imidazole, indolyl, and phenolic hydroxyl [133]. Utilizing surfaces modified with peptides for enzyme immobilization leads to enhanced specific activity and stability of the enzymes, as well as the regulated orientation of the proteins [143]. Using CNBr-activated agarose and CNBr-activated sepharose, which have carbohydrate moieties and glutaraldehyde as a spacer arm, is one method for covalently attaching enzymes to supports. According to studies, this immobilization strategy has proven to give the linked enzymes thermal stability [117][144]. Through covalent enzyme attachment, silica gel carriers modified with silanization and SBA-15 supports with Si-F-lined cage-like pores created highly stable and hyperactive biocatalysts [145]. The enhanced half-life and thermal stability of enzymes have been achieved via covalently attaching them to various supports such as mesoporous silica and chitosan [135][144]. Covalently linking enzymes to electrospun nanofiber leads to improved residual activity as a result of greater surface area and porosity. The implementation of nanodiametric supports has revolutionized biocatalyst immobilization [146][147][148][149][150]. Alcohol dehydrogenase was covalently bound to attapulgite nanofibers (hydrated magnesium silicate) due to their thermal endurance and varying nanosizes [151]. Cross-linked



enzyme aggregates have been developed by precipitating enzymes from aqueous solutions using organic solvents or ionic polymers [152]. The pharmaceutical industry has found covalent binding to magnetic nanoclusters to be useful in achieving varied orientations of immobilized enzymes. This approach has resulted in enhanced operational stability, durability, and reusability, making it a promising technique for enzyme immobilization [153]. One important function of cross-linking agents in enzyme immobilization is to maintain the enzymes' structural and functional integrity. Glutaraldehyde is a commonly used bifunctional cross-linker that can form stable covalent bonds both within and between enzyme subunits, thereby preserving the enzyme's activity and structure. It is also soluble in aqueous solvents, making it a convenient option for use in enzyme immobilization processes. **Table 1** outlines the advantages, drawbacks, and approaches to address the limitations associated with covalent binding.

### 2.2.2. Cross-Linking

Cross-linking is a method of immobilizing enzymes that do not require a support material and results in irreversible binding, preventing the enzyme from leaking into the substrate solution [142][154][155]. This immobilization technique, referred to as carrier-free immobilization, allows the enzymes to act as their carrier, thus resulting in a pure enzyme product and avoiding the drawbacks of using carriers [125][156]. The addition of carriers for enzyme immobilization may result in a decrease in activity, as the presence of non-catalytic components, referred to as ballast, can account for a significant proportion of the total mass, ranging from 90% to over 99%, ultimately leading to reduced space-time yields [152][156] and increased costs [152].

Cross-linking is a process of forming covalent bonds between enzyme molecules using bifunctional or multifunctional agents. One of the commonly used cross-linking agents is glutaraldehyde, owing to its affordability and large-scale availability [125][157]. For several decades, glutaraldehyde has been extensively utilized as a cross-linking agent to generate intermolecular cross-links between proteins, such as enzymes. The cross-linking of enzymes occurs through a reaction with free amino groups of lysine residues on neighboring enzyme molecules. This results in the formation of oligomers or polymers through both inter- and intramolecular aldol condensations, with the specific type of cross-linking dependent on the pH [152][158].

Cross-linked enzyme aggregates (CLEAs) are formed by precipitating enzymes with ammonium sulfate, acetone, or ethanol and then treating the aggregates with a cross-linking agent [125]. There are three methods for immobilizing enzymes, which are: (1) the blending of prepolymer and photosensitizer followed by gelling under near-UV radiation, (2) the freezing of enzyme-containing monomer solution into beads and subsequently polymerizing by gamma radiation, and (3) chemical polymerization via the combination of enzymes with acrylamide monomer and a cross-linking agent in a buffered aqueous solution. [159]. Lately, nanodiametric supports have induced significant advancements in biocatalyst immobilization [56][96][160][161]. The cross-linking immobilization of enzymes on electrospun nanofibers has been shown to improve residual activity, ascribed to the larger surface area and porosity of the substrate. CLEAs were employed to immobilize lysozyme on electrospun chitosan (CS) nanofibers, yielding a durable antibacterial material that can be used continuously [162]. **Table 1** provides an overview of the merits, drawbacks, and methods to address the challenges related to the cross-linking technique.

### 2.2.3. Affinity Immobilization

The affinity immobilization of enzymes involves utilizing their specific binding properties to support materials under different physiological conditions. There are two approaches to achieving this: first, by linking an affinity ligand specific to the target enzyme to the matrix, or second, by attaching the enzyme to a molecule that develops an affinity for the matrix <sup>[163]</sup>. The use of affinity adsorbents has not only been limited to the purification of enzymes but has also been extended to their simultaneous purification <sup>[164]</sup>. Sophisticated affinity matrices like chitosan-modified porous silica beads that are stable in alkali environments and multilayered concanavalin A attached to agarose are capable of immobilizing greater amounts of enzymes leading to better stability and efficiency <sup>[165][166]</sup>. The technique of bio affinity layering is an improvement over affinity immobilization, and it can significantly increase the capacity for enzyme binding and reuse. The non-covalent interactions, such as van der Waals forces, coulombic forces, and hydrogen bonding, among others, are utilized for this purpose <sup>[166][167]</sup>.

## References

1. Farhan Hanafi, M.; Sapawe, N. A Review on the Water Problem Associate with Organic Pollutants Derived from Phenol, Methyl Orange, and Remazol Brilliant Blue Dyes. *Mater. Today Proc.* 2020, 31, A141–A150.
2. Lellis, B.; Fávaro-Polonio, C.Z.; Pamphile, J.A.; Polonio, J.C. Effects of Textile Dyes on Health and the Environment and Bioremediation Potential of Living Organisms. *Biotechnol. Res. Innov.* 2019, 3, 275–290.
3. Asmaly, H.A.; Abussaud, B.; Ihsanullah; Saleh, T.A.; Bukhari, A.A.; Laoui, T.; Shemsi, A.M.; Gupta, V.K.; Atieh, M.A. Evaluation of Micro- and Nano-Carbon-Based Adsorbents for the Removal of Phenol from Aqueous Solutions. *Toxicol. Environ. Chem.* 2015, 97, 1164–1179.
4. Asmaly, H.A.; Ihsanullah; Abussaud, B.; Saleh, T.A.; Laoui, T.; Gupta, V.K.; Atieh, M.A. Adsorption of Phenol on Aluminum Oxide Impregnated Fly Ash. *Desalin. Water Treat.* 2016, 57, 6801–6808.
5. Ahmed, J.; Thakur, A.; Goyal, A. Industrial Wastewater and Its Toxic Effects. In *Biological Treatment of Industrial Wastewater*; The Royal Society of Chemistry: London, UK, 2021; pp. 1–14.
6. Muthukumaran, M. Advances in Bioremediation of Nonaqueous Phase Liquid Pollution in Soil and Water. In *Biological Approaches to Controlling Pollutants*; Elsevier: Amsterdam, The Netherlands, 2021; pp. 191–231.
7. Landrigan, P.J.; Stegeman, J.J.; Fleming, L.E.; Allemand, D.; Anderson, D.M.; Backer, L.C.; Brucker-Davis, F.; Chevalier, N.; Corra, L.; Czerucka, D.; et al. Human Health and Ocean Pollution. *Ann. Glob. Health* 2020, 86, 151.

8. Anku, W.W.; Mamo, M.A.; Govender, P.P. Phenolic Compounds in Water: Sources, Reactivity, Toxicity and Treatment Methods. In Phenolic Compounds—Natural Sources, Importance and Applications; InTech: Rijeka, Romania, 2017.
9. Manasa, R.L.; Mehta, A. Wastewater: Sources of Pollutants and Its Remediation. *Environ. Chem. A Sustain. World* 2020, 2, 197–219.
10. Akpor, O.B.; Ohiobor, G.O.; Olaolu, T.D.; Oghenerobor, B.; Akpor, G.O.; Ohiobor, T.; Debby, O. Heavy Metal Pollutants in Wastewater Effluents: Sources, Effects and Remediation. *Adv. Biosci. Bioeng.* 2014, 2, 37.
11. Ahmed, A.; Forster, M.; Jin, J.; Myers, P.; Zhang, H. Tuning Morphology of Nanostructured ZIF-8 on Silica Microspheres and Applications in Liquid Chromatography and Dye Degradation. *ACS Appl. Mater. Interfaces* 2015, 7, 18054–18063.
12. Gao, W.; Fatehi, P. Fly Ash Based Adsorbent for Treating Bleaching Effluent of Kraft Pulping Process. *Sep. Purif. Technol.* 2018, 195, 60–69.
13. Li, G.; Xu, Q.; Jin, X.; Li, R.; Dharmarajan, R.; Chen, Z. Enhanced Adsorption and Fenton Oxidation of 2,4-Dichlorophenol in Aqueous Solution Using Organobentonite Supported NZVI. *Sep. Purif. Technol.* 2018, 197, 401–406.
14. Abussaud, B.; Asmaly, H.A.; Ihsanullah; Saleh, T.A.; Gupta, V.K.; Laoui, T.; Atieh, M.A. Sorption of Phenol from Waters on Activated Carbon Impregnated with Iron Oxide, Aluminum Oxide and Titanium Oxide. *J. Mol. Liq.* 2016, 213, 351–359.
15. Bahadi, S.A.; Iddrisu, M.; Al-Sakkaf, M.K.; Elgzoly, M.A.A.; Drmosh, Q.A.; Al-Amrani, W.A.; Ahmed, U.; Zahid, U.; Onaizi, S.A. Optimization of Methyl Orange Adsorption on MgFeAl-LTH through the Manipulation of Solution Chemistry and Synthesis Conditions. *Emerg. Mater.* 2023.
16. Bahadi, S.A.; Iddrisu, M.; Al-Sakkaf, M.K.; Elgzoly, M.A.A.; Al-Amrani, W.A.; Ahmed, U.; Zahid, U.; Drmosh, Q.A.; Onaizi, S.A. Chemically versus Thermally Reduced Graphene Oxide: Effects of Reduction Methods and Reducing Agents on the Adsorption of Phenolic Compounds from Wastewater. *Emerg. Mater.* 2023.
17. Ismail, U.M.; Onaizi, S.A.; Vohra, M.S. Novel MgCuAl-Layered Triple Hydroxide for Aqueous Selenite and Selenate Treatment. *Emerg. Mater.* 2023.
18. Fortunato, L.; Elcik, H.; Blankert, B.; Ghaffour, N.; Vrouwenvelder, J. Textile Dye Wastewater Treatment by Direct Contact Membrane Distillation: Membrane Performance and Detailed Fouling Analysis. *J. Memb. Sci.* 2021, 636, 119552.
19. Criscuoli, A.; Zhong, J.; Figoli, A.; Carnevale, M.C.; Huang, R.; Drioli, E. Treatment of Dye Solutions by Vacuum Membrane Distillation. *Water Res.* 2008, 42, 5031–5037.

20. Jaradat, A.Q.; Gharaibeh, S.; Abu Irjei, M. The Application of Solar Distillation Technique as a Mean for Olive Mill Wastewater Management. *Water Environ. J.* 2018, 32, 134–140.
21. Crini, G.; Lichtfouse, E. Wastewater Treatment: An Overview. In *Green Adsorbents for Pollutant Removal*; Springer: Cham, Switzerland, 2018; pp. 1–21.
22. Shikuku, V.O.; Nyairo, W.N. Advanced Oxidation Processes for Dye Removal from Wastewater. In *Impact of Textile Dyes on Public Health and the Environment*; IGI Global: Hershey, PA, USA, 2022; pp. 205–238.
23. Loos, G.; Scheers, T.; Van Eyck, K.; Van Schepdael, A.; Adams, E.; Van der Bruggen, B.; Cabooter, D.; Dewil, R. Electrochemical Oxidation of Key Pharmaceuticals Using a Boron Doped Diamond Electrode. *Sep. Purif. Technol.* 2018, 195, 184–191.
24. Liu, Z.; Meng, H.; Zhang, H.; Cao, J.; Zhou, K.; Lian, J. Highly Efficient Degradation of Phenol Wastewater by Microwave Induced H<sub>2</sub>O<sub>2</sub>-CuOx/GAC Catalytic Oxidation Process. *Sep. Purif. Technol.* 2018, 193, 49–57.
25. Pandit, P.; Basu, S. Dye and Solvent Recovery in Solvent Extraction Using Reverse Micelles for the Removal of Ionic Dyes. *Ind. Eng. Chem. Res.* 2004, 43, 7861–7864.
26. Pandit, P.; Basu, S. Removal of Ionic Dyes from Water by Solvent Extraction Using Reverse Micelles. *Environ. Sci. Technol.* 2004, 38, 2435–2442.
27. Asrami, M.R.; Saien, J. Salting-out Effect on Extraction of Phenol from Aqueous Solutions by Ionic Liquid: Experimental Investigations and Modeling. *Sep. Purif. Technol.* 2018, 204, 175–184.
28. González, E.J.; Díaz, I.; Gonzalez-Miquel, M.; Rodríguez, M.; Sueiras, A. On the Behavior of Imidazolium versus Pyrrolidinium Ionic Liquids as Extractants of Phenolic Compounds from Water: Experimental and Computational Analysis. *Sep. Purif. Technol.* 2018, 201, 214–222.
29. Kadhim, R.J.; Al-Ani, F.H.; Al-Shaeli, M.; Alsalhy, Q.F.; Figoli, A. Removal of Dyes Using Graphene Oxide (GO) Mixed Matrix Membranes. *Membranes* 2020, 10, 366.
30. Ouyang, Z.; Huang, Z.; Tang, X.; Xiong, C.; Tang, M.; Lu, Y. A Dually Charged Nanofiltration Membrane by PH-Responsive Polydopamine for Pharmaceuticals and Personal Care Products Removal. *Sep. Purif. Technol.* 2019, 211, 90–97.
31. Zhang, Y.; Yu, W.; Li, R.; Xu, Y.; Shen, L.; Lin, H.; Liao, B.Q.; Wu, G. Novel Conductive Membranes Breaking through the Selectivity-Permeability Trade-off for Congo Red Removal. *Sep. Purif. Technol.* 2019, 211, 368–376.
32. Sunil, K.; Sherugar, P.; Rao, S.; Lavanya, C.; Balakrishna, G.R.; Arthanareeswaran, G.; Padaki, M. Prolific Approach for the Removal of Dyes by an Effective Interaction with Polymer Matrix Using Ultrafiltration Membrane. *J. Environ. Chem. Eng.* 2021, 9, 106328.

33. Badvi, K.; Javanbakht, V. Enhanced Photocatalytic Degradation of Dye Contaminants with TiO<sub>2</sub> Immobilized on ZSM-5 Zeolite Modified with Nickel Nanoparticles. *J. Clean. Prod.* 2021, 280, 124518.
34. Sirirerkratana, K.; Kemacheevakul, P.; Chuangchote, S. Color Removal from Wastewater by Photocatalytic Process Using Titanium Dioxide-Coated Glass, Ceramic Tile, and Stainless Steel Sheets. *J. Clean. Prod.* 2019, 215, 123–130.
35. Nguyen, D.C.T.; Cho, K.Y.; Oh, W.C. Mesoporous CuO-Graphene Coating of Mesoporous TiO<sub>2</sub> for Enhanced Visible-Light Photocatalytic Activity of Organic Dyes. *Sep. Purif. Technol.* 2019, 211, 646–657.
36. Lin, J.C.T.; Sopajaree, K.; Jitjanesuwan, T.; Lu, M.C. Application of Visible Light on Copper-Doped Titanium Dioxide Catalyzing Degradation of Chlorophenols. *Sep. Purif. Technol.* 2018, 191, 233–243.
37. Alkadhém, A.M.; Elgzoly, M.A.A.; Onaizi, S.A. Novel Amine-Functionalized Magnesium Oxide Adsorbents for CO<sub>2</sub> Capture at Ambient Conditions. *J. Environ. Chem. Eng.* 2020, 8, 103968.
38. Hezam, A.; Drmash, Q.A.; Ponnammá, D.; Bajiri, M.A.; Qamar, M.; Namratha, K.; Zare, M.; Nayan, M.B.; Onaizi, S.A.; Byrappa, K. Strategies to Enhance ZnO Photocatalyst's Performance for Water Treatment: A Comprehensive Review. *Chem. Rec.* 2022, 22, e202100299.
39. Al Lagtah, N.M.A.; Onaizi, S.A.; Albadarin, A.B.; Ghaith, F.A.; Nour, M.I. Techno-Economic Analysis of the Effects of Heat Integration and Different Carbon Capture Technologies on the Performance of Coal-Based IGCC Power Plants. *J. Environ. Chem. Eng.* 2019, 7, 103471.
40. Almarouf, H.S.; Nasser, M.S.; Al-Marri, M.J.; Khraisheh, M.; Onaizi, S.A. Demulsification of Stable Emulsions from Produced Water Using a Phase Separator with Inclined Parallel Arc Coalescing Plates. *J. Pet. Sci. Eng.* 2015, 135, 16–21.
41. Al-Sakkaf, M.K.; Onaizi, S.A. Crude Oil/Water Nanoemulsions Stabilized by Rhamnolipid Biosurfactant: Effects of Acidity/Basicity and Salinity on Emulsion Characteristics, Stability, and Demulsification. *Fuel* 2023, 344, 128052.
42. Lateef, S.A.; Ajumobi, O.O.; Onaizi, S.A. Enzymatic Desulfurization of Crude Oil and Its Fractions: A Mini Review on the Recent Progresses and Challenges. *Arab. J. Sci. Eng.* 2019, 44, 5181–5193.
43. Al-Sakkaf, M.K.; Onaizi, S.A. Rheology, Characteristics, Stability, and PH-Responsiveness of Biosurfactant-Stabilized Crude Oil/Water Nanoemulsions. *Fuel* 2022, 307, 121845.
44. Onaizi, S.A.; Alsulaimani, M.; Al-Sakkaf, M.K.; Bahadi, S.A.; Mahmoud, M.; Alshami, A. Crude Oil/Water Nanoemulsions Stabilized by Biosurfactant: Stability and PH-Switchability. *J. Pet. Sci. Eng.* 2021, 198, 108173.

45. Onaizi, S.A. Demulsification of Crude Oil/Water Nanoemulsions Stabilized by Rhamnolipid Biosurfactant Using Enzymes and PH-Swing. *Sep. Purif. Technol.* 2021, 259, 118060.
46. Onaizi, S.A.; He, L.; Middelberg, A.P.J. The Construction, Fouling and Enzymatic Cleaning of a Textile Dye Surface. *J. Colloid Interface Sci.* 2010, 351, 203–209.
47. Onaizi, S.A. Dynamic Surface Tension and Adsorption Mechanism of Surfactin Biosurfactant at the Air–Water Interface. *Eur. Biophys. J.* 2018, 47, 631–640.
48. Onaizi, S.A.; Malcolm, A.S.; He, L.; Middelberg, A.P.J. Directed Disassembly of an Interfacial Rubisco Protein Network. *Langmuir* 2007, 23, 6336–6341.
49. Singh, N.; Singh, J. An Enzymatic Method for Removal of Phenol from Industrial Effluent. *Prep. Biochem. Biotechnol.* 2002, 32, 127–133.
50. Ariaeenejad, S.; Motamedi, E.; Salekdeh, G.H. Highly Efficient Removal of Dyes from Wastewater Using Nanocellulose from Quinoa Husk as a Carrier for Immobilization of Laccase. *Bioresour. Technol.* 2022, 349, 126833.
51. Sariaslani, F.S.; Dalton, H. Microbial Enzymes for Oxidation of Organic Molecules. *Crit. Rev. Biotechnol.* 1989, 9, 171–257.
52. Misal, S.A.; Gawai, K.R. Azoreductase: A Key Player of Xenobiotic Metabolism. *Bioresour. Bioprocess.* 2018, 5, 17.
53. Gaur, G.; Gänzle, M.G. Conversion of (Poly)Phenolic Compounds in Food Fermentations by Lactic Acid Bacteria: Novel Insights into Metabolic Pathways and Functional Metabolites. *Curr. Res. Food Sci.* 2023, 6, 100448.
54. Basso, A.; Serban, S. Industrial Applications of Immobilized Enzymes—A Review. *Mol. Catal.* 2019, 479, 110607.
55. Jesionowski, T.; Zdarta, J.; Krajewska, B. Enzyme Immobilization by Adsorption: A Review. *Adsorption* 2014, 20, 801–821.
56. Alshabib, M.; Onaizi, S.A. A Review on Phenolic Wastewater Remediation Using Homogeneous and Heterogeneous Enzymatic Processes: Current Status and Potential Challenges. *Sep. Purif. Technol.* 2019, 219, 186–207.
57. Torres, J.A.; Nogueira, F.G.E.; Silva, M.C.; Lopes, J.H.; Tavares, T.S.; Ramalho, T.C.; Corrêa, A.D. Novel Eco-Friendly Biocatalyst: Soybean Peroxidase Immobilized onto Activated Carbon Obtained from Agricultural Waste. *RSC Adv.* 2017, 7, 16460–16466.
58. Gholami-Borujeni, F.; Mahvi, A.H.; Naseri, S.; Faramarzi, M.A.; Nabizadeh, R.; Alimohammadi, M. Application of Immobilized Horseradish Peroxidase for Removal and Detoxification of Azo Dye from Aqueous Solution. *Res. J. Chem. Environ.* 2011, 15, 217–222.

59. Silva, M.C.; Torres, J.A.; Vasconcelos De Sá, L.R.; Chagas, P.M.B.; Ferreira-Leitão, V.S.; Corrêa, A.D. The Use of Soybean Peroxidase in the Decolourization of Remazol Brilliant Blue R and Toxicological Evaluation of Its Degradation Products. *J. Mol. Catal. B Enzym.* 2013, 89, 122–129.
60. Chiong, T.; Lau, S.Y.; Lek, Z.H.; Koh, B.Y.; Danquah, M.K. Enzymatic Treatment of Methyl Orange Dye in Synthetic Wastewater by Plant-Based Peroxidase Enzymes. *J. Environ. Chem. Eng.* 2016, 4, 2500–2509.
61. Bhatia, D.; Sharma, N.R.; Singh, J.; Kanwar, R.S. Biological Methods for Textile Dye Removal from Wastewater: A Review. *Crit. Rev. Environ. Sci. Technol.* 2017, 47, 1836–1876.
62. Forootanfar, H.; Moezzi, A.; Aghaie-Khozani, M.; Mahmoudjanlou, Y.; Ameri, A.; Niknejad, F.; Ali Faramarzi, M. Synthetic Dye Decolorization by Three Sources of Fungal Laccase. *Iran. J. Environ. Health Sci. Eng.* 2012, 9, 27.
63. Onaizi, S.A.; Alshabib, M. The Degradation of Bisphenol A by Laccase: Effect of Biosurfactant Addition on the Reaction Kinetics under Various Conditions. *Sep. Purif. Technol.* 2021, 257, 117785.
64. Silva, M.C.; Torres, J.A.; Castro, A.A.; da Cunha, E.F.F.; Alves de Oliveira, L.C.; Corrêa, A.D.; Ramalho, T.C. Combined Experimental and Theoretical Study on the Removal of Pollutant Compounds by Peroxidases: Affinity and Reactivity toward a Bioremediation Catalyst. *J. Biomol. Struct. Dyn.* 2016, 34, 1839–1848.
65. Veitch, N.C. Horseradish Peroxidase: A Modern View of a Classic Enzyme. *Phytochemistry* 2004, 65, 249–259.
66. Jaiswal, N.; Pandey, V.P.; Dwivedi, U.N. Immobilization of Papaya Laccase in Chitosan Led to Improved Multipronged Stability and Dye Discoloration. *Int. J. Biol. Macromol.* 2016, 86, 288–295.
67. Effron, D.; De La Horra, A.M.; Defrieri, R.L.; Fontanive, V.; Palma, R.M. Effect of Cadmium, Copper, and Lead on Different Enzyme Activities in a Native Forest Soil. *Commun. Soil Sci. Plant Anal.* 2006, 35, 1309–1321.
68. Torres, J.A.; Silva, M.C.; Lopes, J.H.; Nogueira, A.E.; Nogueira, F.G.E.; Corrêa, A.D. Development of a Reusable and Sustainable Biocatalyst by Immobilization of Soybean Peroxidase onto Magnetic Adsorbent. *Int. J. Biol. Macromol.* 2018, 114, 1279–1287.
69. Wang, C.; Zhang, H.; Ren, D.; Li, Q.; Zhang, S.; Feng, T. Effect of Direct-Current Electric Field on Enzymatic Activity and the Concentration of Laccase. *Indian. J. Microbiol.* 2015, 55, 278–284.
70. Alshabib, M.; Onaizi, S.A. Enzymatic Remediation of Bisphenol A from Wastewaters: Effects of Biosurfactant, Anionic, Cationic, Nonionic, and Polymeric Additives. *Water Air Soil. Pollut.* 2020, 231, 428.



71. Alshabib, M.; Onaizi, S.A. Effects of Surface Active Additives on the Enzymatic Treatment of Phenol and Its Derivatives: A Mini Review. *Curr. Pollut. Rep.* 2019, 5, 52–65.
72. Sellami, K.; Couvert, A.; Nasrallah, N.; Maachi, R.; Tandjaoui, N.; Abouseoud, M.; Amrane, A. Bio-Based and Cost Effective Method for Phenolic Compounds Removal Using Cross-Linked Enzyme Aggregates. *J. Hazard. Mater.* 2021, 403, 124021.
73. Villegas, L.G.C.; Mashhadi, N.; Chen, M.; Mukherjee, D.; Taylor, K.E.; Biswas, N. A Short Review of Techniques for Phenol Removal from Wastewater. *Curr. Pollut. Rep.* 2016, 2, 157–167.
74. Onaizi, S.A. Statistical Analyses of the Effect of Rhamnolipid Biosurfactant Addition on the Enzymatic Removal of Bisphenol A from Wastewater. *Biocatal. Agric. Biotechnol.* 2021, 32, 101929.
75. Onaizi, S.A. Enzymatic Treatment of Phenolic Wastewater: Effects of Salinity and Biosurfactant Addition. In *Proceedings of the International Petroleum Technology Conference*, Virtual, 23 March–1 April 2021.
76. Chatterjee, S.; Kumari, S.; Rath, S.; Das, S. Prospects and Scope of Microbial Bioremediation for the Restoration of the Contaminated Sites. In *Microbial Biodegradation and Bioremediation*; Elsevier: Amsterdam, The Netherlands, 2022; pp. 3–31.
77. Kanaujiya, D.K.; Paul, T.; Sinharoy, A.; Pakshirajan, K. Biological Treatment Processes for the Removal of Organic Micropollutants from Wastewater: A Review. *Curr. Pollut. Rep.* 2019, 5, 112–128.
78. Ramirez, E.; de la Luz Asunción, M.; Rivalcoba, V.S.; Hernández, A.; Santos, C.V. Removal of Phenolic Compounds from Water by Adsorption and Photocatalysis. In *Phenolic Compounds—Natural Sources, Importance and Applications*; InTech: Rijeka, Romania, 2017.
79. Betancur-Ramírez, K.J.; Meneses-Jácome, A.; Ruiz-Colorado, A.A.; Gallego-Suárez, D. Life Cycle Assessment of an Alternative Enzymatic-Biological Treatment for Effluents from Industrial Processing of Potatoes. *J. Clean. Prod.* 2021, 324, 129151.
80. Sirisha, V.L.; Jain, A.; Jain, A. Enzyme Immobilization: An Overview on Methods, Support Material, and Applications of Immobilized Enzymes. *Adv. Food Nutr. Res.* 2016, 79, 179–211.
81. Lou, W.Y.; Fernández-Lucas, J.; Ge, J.; Wu, C. Enzyme or Whole Cell Immobilization for Efficient Biocatalysis: Focusing on Novel Supporting Platforms and Immobilization Techniques. *Front. Bioeng. Biotechnol.* 2021, 9, 59.
82. Saravanan, A.; Kumar, P.S.; Vo, D.V.N.; Jeevanantham, S.; Karishma, S.; Yaashikaa, P.R. A Review on Catalytic-Enzyme Degradation of Toxic Environmental Pollutants: Microbial Enzymes. *J. Hazard. Mater.* 2021, 419, 126451.

83. Xu, K.; Chen, X.; Zheng, R.; Zheng, Y. Immobilization of Multi-Enzymes on Support Materials for Efficient Biocatalysis. *Front. Bioeng. Biotechnol.* 2020, 8, 660.
84. Irfan, M.; Ghazanfar, M.; Ur Rehman, A.; Siddique, A. Strategies to Reuse Cellulase: Immobilization of Enzymes (Part II). In *Approaches to Enhance Industrial Production of Fungal Cellulases*; Springer: Berlin/Heidelberg, Germany, 2019; pp. 137–151.
85. Sepahvand, H.; Heravi, M.M.; Saber, M.; Hooshmand, S.E. Techniques and Support Materials for Enzyme Immobilization Using Ugi Multicomponent Reaction: An Overview. *J. Iran. Chem. Soc.* 2022, 19, 2115–2130.
86. Wang, Y.; Gu, Y.; Yang, S. Developing a Novel Strategy for Light-Triggered Reversible Enzyme Immobilization and Reuse of Support. *Alex. Eng. J.* 2022, 61, 6949–6957.
87. Rodrigues, R.C.; Ortiz, C.; Berenguer-Murcia, Á.; Torres, R.; Fernández-Lafuente, R. Modifying Enzyme Activity and Selectivity by Immobilization. *Chem. Soc. Rev.* 2013, 42, 6290–6307.
88. Lanie, M.; Ab, H. Enzymatic Strategies for Asymmetric Synthesis. *RSC Chem. Biol.* 2021, 2, 958–989.
89. Nguyen, L.N.; Hai, F.I.; Dosseto, A.; Richardson, C.; Price, W.E.; Nghiem, L.D. Continuous Adsorption and Biotransformation of Micropollutants by Granular Activated Carbon-Bound Laccase in a Packed-Bed Enzyme Reactor. *Bioresour. Technol.* 2016, 210, 108–116.
90. Isanapong, J.; Lohawet, K.; Kumnorkaew, P. Optimization and Characterization of Immobilized Laccase on Titanium Dioxide Nanostructure and Its Application in Removal of Remazol Brilliant Blue R. *Biocatal. Agric. Biotechnol.* 2021, 37, 102186.
91. Ponmudi, K.; Cherian, A.R.; Varghese, A. Carbon Dots as an Effective Material in Enzyme Immobilization for Sensing Applications. In *Carbon Dots in Analytical Chemistry Detection and Imaging*; Elsevier: Amsterdam, The Netherlands, 2023; pp. 241–253.
92. Oliveira, F.L.; França, A.d.S.; de Castro, A.M.; Alves de Souza, R.O.M.; Esteves, P.M.; Gonçalves, R.S.B. Enzyme Immobilization in Covalent Organic Frameworks: Strategies and Applications in Biocatalysis. *Chempluschem* 2020, 85, 2051–2066.
93. Zhang, J.; Zhang, J.; Zhang, F.; Yang, H.; Huang, X.; Liu, H.; Guo, S. Graphene Oxide as a Matrix for Enzyme Immobilization. *Langmuir* 2010, 26, 6083–6085.
94. Zhang, W.; Yang, Q.; Luo, Q.; Shi, L.; Meng, S. Laccase-Carbon Nanotube Nanocomposites for Enhancing Dyes Removal. *J. Clean. Prod.* 2020, 242, 118425.
95. Yuan, Y.; Cai, W.; Xu, J.; Cheng, J.; Du, K.S. Recyclable Laccase by Coprecipitation with Acidic Cu-Based MOFs for Bisphenol A Degradation in an Aqueous Environment. *Colloids Surf. B Biointerfaces* 2021, 204, 111792.

96. Nawaz, A.F.; Zafar, S.; Fatim, S.L.; Shahzadi, K.; Fatima, Z.; Siddique, I. Use of Nanomaterials for the Immobilization of Industrially Important Enzymes. *J. Nanotechnol. Res.* 2020, 3, 45–57.
97. Soares, A.M.B.F.; Gonçalves, L.M.O.; Ferreira, R.D.S.; de Souza, J.M.; Fangueiro, R.; Alves, M.M.M.; Carvalho, F.A.A.; Mendes, A.N.; Cantanhêde, W. Immobilization of Papain Enzyme on a Hybrid Support Containing Zinc Oxide Nanoparticles and Chitosan for Clinical Applications. *Carbohydr. Polym.* 2020, 243, 116498.
98. Ramakrishna, T.R.B.; Nalder, T.D.; Yang, W.; Marshall, S.N.; Barrow, C.J. Controlling Enzyme Function through Immobilisation on Graphene, Graphene Derivatives and Other Two Dimensional Nanomaterials. *J. Mater. Chem. B* 2018, 6, 3200–3218.
99. Wen, H.; Nallathambi, V.; Chakraborty, D.; Barton, S.C. Carbon Fiber Microelectrodes Modified with Carbon Nanotubes as a New Support for Immobilization of Glucose Oxidase. *Microchim. Acta* 2011, 175, 283–289.
100. Thakur, K.; Attri, C.; Seth, A. Nanocarriers-Based Immobilization of Enzymes for Industrial Application. *3 Biotech.* 2021, 11, 427.
101. Li, S.F.; Zhai, X.J.; Zhang, C.; Mo, H.L.; Zang, S.Q. Enzyme Immobilization in Highly Ordered Macro–Microporous Metal–Organic Frameworks for Rapid Biodegradation of Hazardous Dyes. *Inorg. Chem. Front.* 2020, 7, 3146–3153.
102. Feng, L.; Wang, K.Y.; Lv, X.L.; Yan, T.H.; Zhou, H.C. Hierarchically Porous Metal–Organic Frameworks: Synthetic Strategies and Applications. *Natl. Sci. Rev.* 2020, 7, 1743–1758.
103. Guisan, J.M.; López-Gallego, F.; Bolivar, J.M.; Rocha-Martín, J.; Fernandez-Lorente, G. The Science of Enzyme Immobilization. *Methods Mol. Biol.* 2020, 2100, 1–26.
104. Khan, M.R. Immobilized Enzymes: A Comprehensive Review. *Bull. Natl. Res. Cent.* 2021, 45, 207.
105. Imam, H.T.; Marr, P.C.; Marr, A.C. Enzyme Entrapment, Biocatalyst Immobilization without Covalent Attachment. *Green. Chem.* 2021, 23, 4980–5005.
106. Brena, B.; González-Pombo, P.; Batista-Viera, F. Immobilization of Enzymes and Cells, 3rd ed.; *Methods in Molecular Biology*; Humana Press: Totowa, NJ, USA, 2013; Volume 1051, pp. 15–31.
107. Spahn, C.; Minteer, S.D. Enzyme Immobilization in Biotechnology. *Recent. Pat. Eng.* 2008, 2, 195–200.
108. Dey, G.; Nagpal, V.; Banerjee, R. Immobilization of Alpha-Amylase from *Bacillus Circulans* GRS 313 on Coconut Fiber. *Appl. Biochem. Biotechnol.* 2002, 102–103, 303–313.
109. Rosales-Hernández, M.; Kispert, L.; Torres-Ramírez, E.; Ramírez-Rosales, D.; Zamorano-Ulloa, R.; Trujillo-Ferrara, J. Electron Paramagnetic Resonance Analyses of Biotransformation

- Reactions with Cytochrome P-450 Immobilized on Mesoporous Molecular Sieves. *Biotechnol. Lett.* 2007, 29, 919–924.
110. Karagulyan, H.K.; Gasparyan, V.K.; Decker, S.R. Immobilization of Fungal Beta-Glucosidase on Silica Gel and Kaolin Carriers. *Appl. Biochem. Biotechnol.* 2008, 146, 39–47.
111. Brígida, A.I.S.; Calado, V.M.A.; Gonçalves, L.R.B.; Coelho, M.A.Z. Effect of Chemical Treatments on Properties of Green Coconut Fiber. *Carbohydr. Polym.* 2010, 79, 832–838.
112. Huang, X.J.; Chen, P.C.; Huang, F.; Ou, Y.; Chen, M.R.; Xu, Z.K. Immobilization of *Candida Rugosa* Lipase on Electrospun Cellulose Nanofiber Membrane. *J. Mol. Catal. B Enzym.* 2011, 70, 95–100.
113. Mitchell, S.; Pérez-Ramírez, J. Mesoporous Zeolites as Enzyme Carriers: Synthesis, Characterization, and Application in Biocatalysis. *Catal. Today* 2011, 168, 28–37.
114. Díaz, J.F.; Balkus, K.J. Enzyme Immobilization in MCM-41 Molecular Sieve. *J. Mol. Catal. B Enzym.* 1996, 2, 115–126.
115. Persson, M.; Wehtje, E.; Adlercreutz, P. Immobilisation of Lipases by Adsorption and Deposition: High Protein Loading Gives Lower Water Activity Optimum. *Biotechnol. Lett.* 2000, 22, 1571–1575.
116. Sabbani, S.; Hedenström, E.; Nordin, O. The Enantioselectivity of *Candida Rugosa* Lipase Is Influenced by the Particle Size of the Immobilising Support Material Accurel. *J. Mol. Catal. B Enzym.* 2006, 42, 1–9.
117. Cunha, A.G.; Fernández-Lorente, G.; Bevilacqua, J.V.; Destain, J.; Paiva, L.M.C.; Freire, D.M.G.; Fernández-Lafuente, R.; Guisán, J.M. Immobilization of *Yarrowia Lipolytica* Lipase—A Comparison of Stability of Physical Adsorption and Covalent Attachment Techniques. *Appl. Biochem. Biotechnol.* 2008, 146, 49–56.
118. Cabrera-Padilla, R.Y.; Lisboa, M.C.; Fricks, A.T.; Franceschi, E.; Lima, A.S.; Silva, D.P.; Soares, C.M.F. Immobilization of *Candida Rugosa* Lipase on Poly(3-hydroxybutyrate-co-hydroxyvalerate): A New Eco-Friendly Support. *J. Ind. Microbiol. Biotechnol.* 2012, 39, 289–298.
119. Mishra, N.; Pithawala, K.; Bahadur, A. Byssus Thread: A Novel Support Material for Urease Immobilization. *Appl. Biochem. Biotechnol.* 2011, 165, 1568–1576.
120. Popat, A.; Hartono, S.B.; Stahr, F.; Liu, J.; Qiao, S.Z.; Lu, G.Q. Mesoporous Silica Nanoparticles for Bioadsorption, Enzyme Immobilisation, and Delivery Carriers. *Nanoscale* 2011, 3, 2801–2818.
121. Magner, E. Immobilisation of Enzymes on Mesoporous Silicate Materials. *Chem. Soc. Rev.* 2013, 42, 6213–6222.
122. Zucca, P.; Sanjust, E. Inorganic Materials as Supports for Covalent Enzyme Immobilization: Methods and Mechanisms. *Molecules* 2014, 19, 14139–14194.

123. Azodi, M.; Falamaki, C.; Mohsenifar, A. Sucrose Hydrolysis by Invertase Immobilized on Functionalized Porous Silicon. *J. Mol. Catal. B Enzym.* 2011, 69, 154–160.
124. Homaei, A.A.; Sariri, R.; Vianello, F.; Stevanato, R. Enzyme Immobilization: An Update. *J. Chem. Biol.* 2013, 6, 185–205.
125. Hanefeld, U.; Gardossi, L.; Magner, E. Understanding Enzyme Immobilisation. *Chem. Soc. Rev.* 2009, 38, 453–468.
126. Betancor, L.; Luckarift, H.R. Bioinspired Enzyme Encapsulation for Biocatalysis. *Trends Biotechnol.* 2008, 26, 566–572.
127. Kurzbaum, E.; Raizner, Y.; Kuc, M.E.; Kulikov, A.; Hakimi, B.; Kruh, L.I.; Menashe, O. Phenol Biodegradation by Bacterial Cultures Encapsulated in 3D Microfiltration-Membrane Capsules. *Environ. Technol.* 2019, 41, 2875–2883.
128. Maghraby, Y.R.; El-Shabasy, R.M.; Ibrahim, A.H.; Azzazy, H.M.E.S. Enzyme Immobilization Technologies and Industrial Applications. *ACS Omega* 2023, 8, 5184–5196.
129. Song, J.; He, W.; Shen, H.; Zhou, Z.; Li, M.; Su, P.; Yang, Y. Construction of Multiple Enzyme Metal–Organic Frameworks Biocatalyst via DNA Scaffold: A Promising Strategy for Enzyme Encapsulation. *Chem. Eng. J.* 2019, 363, 174–182.
130. Tran, D.N.; Balkus, K.J. Perspective of Recent Progress in Immobilization of Enzymes. *ACS Catal.* 2011, 1, 956–968.
131. Datta, S.; Christena, L.R.; Rajaram, Y.R.S. Enzyme Immobilization: An Overview on Techniques and Support Materials. *3 Biotech.* 2013, 3, 1–9.
132. Hartmann, M.; Jung, D. Biocatalysis with Enzymes Immobilized on Mesoporous Hosts: The Status Quo and Future Trends. *J. Mater. Chem.* 2010, 20, 844–857.
133. Singh, B.D. *Biotechnology Expanding Horizons*, 4th ed.; Kalyani Publishers: Delhi, India, 2012; ISBN 9789327222982.
134. Shen, Q.; Yang, R.; Hua, X.; Ye, F.; Zhang, W.; Zhao, W. Gelatin-Templated Biomimetic Calcification for  $\beta$ -Galactosidase Immobilization. *Process Biochem.* 2011, 46, 1565–1571.
135. Ispas, C.; Sokolov, I.; Andreescu, S. Enzyme-Functionalized Mesoporous Silica for Bioanalytical Applications. *Anal. Bioanal. Chem.* 2009, 393, 543–554.
136. Chen, G.C.; Kuan, I.C.; Hong, J.R.; Tsai, B.H.; Lee, S.L.; Yu, C.Y. Activity Enhancement and Stabilization of Lipase from *Pseudomonas Cepacia* in Polyallylamine-Mediated Biomimetic Silica. *Biotechnol. Lett.* 2011, 33, 525–529.
137. Tümtürk, H.; Karaca, N.; Demirel, G.; Şahin, F. Preparation and Application of Poly(N,N-Dimethylacrylamide-Co-Acrylamide) and Poly(N-Isopropylacrylamide-Co-Acrylamide)/Kappa-

- Carrageenan Hydrogels for Immobilization of Lipase. *Int. J. Biol. Macromol.* 2007, 40, 281–285.
138. Jegannathan, K.R.; Jun-Yee, L.; Chan, E.S.; Ravindra, P. Production of Biodiesel from Palm Oil Using Liquid Core Lipase Encapsulated in  $\kappa$ -Carrageenan. *Fuel* 2010, 89, 2272–2277.
139. Brena, B.; González-Pombo, P.; Batista-Viera, F. Immobilization of Enzymes: A Literature Survey. *Methods Mol. Biol.* 2013, 1051, 15–31.
140. Patil, J.S.; Kamalapur, M.V.; Marapur, S.C.; Kadam, D.V. Ionotropic gelation and polyelectrolyte complexation: The novel techniques to design hydrogel particulate sustained, modulated drug delivery system: A review. *Dig. J. Nanomater. Biostruct.* 2010, 5, 241–248.
141. Rother, C.; Nidetzky, B. Enzyme Immobilization by Microencapsulation: Methods, Materials, and Technological Applications. In *Encyclopedia of Industrial Biotechnology*; John Wiley & Sons, Inc.: Hoboken, NJ, USA, 2014; pp. 1–21.
142. Honda, T.; Miyazaki, M.; Nakamura, H.; Maeda, H. Immobilization of Enzymes on a Microchannel Surface through Cross-Linking Polymerization. *Chem. Commun.* 2005, 40, 5062–5064.
143. Fu, J.; Reinhold, J.; Woodbury, N.W. Peptide-Modified Surfaces for Enzyme Immobilization. *PLoS ONE* 2011, 6, e18692.
144. Hsieh, H.J.; Liu, P.C.; Liao, W.J. Immobilization of Invertase via Carbohydrate Moiety on Chitosan to Enhance Its Thermal Stability. *Biotechnol. Lett.* 2000, 22, 1459–1464.
145. Szymańska, K.; Bryjak, J.; Jarzębski, A.B. Immobilization of Invertase on Mesoporous Silicas to Obtain Hyper Active Biocatalysts. *Top. Catal.* 2009, 52, 1030–1036.
146. Kim, J.; Jia, H.; Wang, P. Challenges in Biocatalysis for Enzyme-Based Biofuel Cells. *Biotechnol. Adv.* 2006, 24, 296–308.
147. Huang, X.J.; Yu, A.G.; Xu, Z.K. Covalent Immobilization of Lipase from *Candida Rugosa* onto Poly(acrylonitrile-co-2-hydroxyethyl methacrylate) Electrospun Fibrous Membranes for Potential Bioreactor Application. *Bioresour. Technol.* 2008, 99, 5459–5465.
148. Sakai, S.; Liu, Y.; Yamaguchi, T.; Watanabe, R.; Kawabe, M.; Kawakami, K. Immobilization of *Pseudomonas Cepacia* Lipase onto Electrospun Polyacrylonitrile Fibers through Physical Adsorption and Application to Transesterification in Nonaqueous Solvent. *Biotechnol. Lett.* 2010, 32, 1059–1062.
149. Wu, L.; Yuan, X.; Sheng, J. Immobilization of Cellulase in Nanofibrous PVA Membranes by Electrospinning. *J. Memb. Sci.* 2005, 250, 167–173.
150. Ren, G.; Xu, X.; Liu, Q.; Cheng, J.; Yuan, X.; Wu, L.; Wan, Y. Electrospun Poly(vinyl Alcohol)/Glucose Oxidase Biocomposite Membranes for Biosensor Applications. *React. Funct. Polym.* 2006, 66, 1559–1564.

151. Hilal, N.; Kochkodan, V.; Nigmatullin, R.; Goncharuk, V.; Al-Khatib, L. Lipase-Immobilized Biocatalytic Membranes for Enzymatic Esterification: Comparison of Various Approaches to Membrane Preparation. *J. Memb. Sci.* 2006, 268, 198–207.
152. Sheldon, R.A. Characteristic Features and Biotechnological Applications of Cross-Linked Enzyme Aggregates (CLEAs). *Appl. Microbiol. Biotechnol.* 2011, 92, 467.
153. Yusdy; Patel, S.R.; Yap, M.G.S.; Wang, D.I.C. Immobilization of L-Lactate Dehydrogenase on Magnetic Nanoclusters for Chiral Synthesis of Pharmaceutical Compounds. *Biochem. Eng. J.* 2009, 48, 13–21.
154. Wang, A.; Wang, H.; Zhu, S.; Zhou, C.; Du, Z.; Shen, S. An Efficient Immobilizing Technique of Penicillin Acylase with Combining Mesocellular Silica Foams Support and P-Benzoquinone Cross Linker. *Bioprocess. Biosyst. Eng.* 2008, 31, 509–517.
155. Subramanian, A.; Kennel, S.J.; Oden, P.I.; Jacobson, K.B.; Woodward, J.; Doktycz, M.J. Comparison of Techniques for Enzyme Immobilization on Silicon Supports. *Enzym. Microb. Technol.* 1999, 24, 26–34.
156. Sheldon, R.A. Cross-Linked Enzyme Aggregates (CLEAs): Stable and Recyclable Biocatalysts. *Biochem. Soc. Trans.* 2007, 35, 1583–1587.
157. Górecka, E.; Jastrzębska, M. Immobilization Techniques and Biopolymer Carriers. *Biotechnol. Food Sci.* 2011, 75, 65–86.
158. Migneault, I.; Dartiguenave, C.; Bertrand, M.J.; Waldron, K.C. Glutaraldehyde: Behavior in Aqueous Solution, Reaction with Proteins, and Application to Enzyme Crosslinking. *Biotechniques* 2004, 37, 790–802.
159. Öztürk, B. Immobilization of Lipase from *Candida Rugosa* on Hydrophobic and Hydrophilic Supports; Izmir Institute of Technology: Urla, Türkiye, 2001.
160. Lee, C.K.; Au-Duong, A.N. Enzyme Immobilization on Nanoparticles: Recent Applications. *Emerg. Areas Bioeng.* 2017, 1, 67–80.
161. Sharifi, M.; Sohrabi, M.J.; Hosseinali, S.H.; Hasan, A.; Kani, P.H.; Talaei, A.J.; Karim, A.Y.; Nanakali, N.M.Q.; Salihi, A.; Aziz, F.M.; et al. Enzyme Immobilization onto the Nanomaterials: Application in Enzyme Stability and Prodrug-Activated Cancer Therapy. *Int. J. Biol. Macromol.* 2020, 143, 665–676.
162. Park, J.M.; Kim, M.; Park, H.S.; Jang, A.; Min, J.; Kim, Y.H. Immobilization of Lysozyme-CLEA onto Electrospun Chitosan Nanofiber for Effective Antibacterial Applications. *Int. J. Biol. Macromol.* 2013, 54, 37–43.
163. Sardar, M.; Roy, I.; Gupta, M.N. Simultaneous Purification and Immobilization of *Aspergillus Niger* Xylanase on the Reversibly Soluble Polymer Eudragit(TM) L-100. *Enzym. Microb. Technol.* 2000,



27, 672–679.

164. Ho, L.F.; Li, S.Y.; Lin, S.C.; Hsu, W.H. Integrated Enzyme Purification and Immobilization Processes with Immobilized Metal Affinity Adsorbents. *Process Biochem.* 2004, 39, 1573–1581.
165. Shi, Q.H.; Tian, Y.; Dong, X.Y.; Bai, S.; Sun, Y. Chitosan-Coated Silica Beads as Immobilized Metal Affinity Support for Protein Adsorption. *Biochem. Eng. J.* 2003, 16, 317–322.
166. Sardar, M.; Gupta, M.N. Immobilization of Tomato Pectinase on Con A–Seralose 4B by Bioaffinity Layering. *Enzym. Microb. Technol.* 2005, 37, 355–359.
167. Haider, T.; Husain, Q. Concanavalin A Layered Calcium Alginate-Starch Beads Immobilized Beta Galactosidase as a Therapeutic Agent for Lactose Intolerant Patients. *Int. J. Pharm.* 2008, 359, 1–6.

---

Retrieved from <https://encyclopedia.pub/entry/history/show/110601>