

# POIKTMP

Subjects: Genetics & Heredity

Contributor: Peter Tang

Hereditary fibrosing poikiloderma with tendon contractures, myopathy, and pulmonary fibrosis (abbreviated POIKTMP), is a disorder that affects many parts of the body, particularly the skin, muscles, lungs, and pancreas. Signs and symptoms vary among affected individuals.

Keywords: genetic conditions

---

## 1. Introduction

People with POIKTMP have patchy changes in skin coloring and small clusters of blood vessels just under the skin (telangiectases), a combination known as poikiloderma. These skin changes begin in infancy and occur primarily on the face. They can also have red, scaly skin patches and mild swelling (lymphedema) of the arms and legs; thickened skin on the palms of the hands and soles of the feet (palmoplantar keratoderma); and abnormal hardening (sclerosis) of tissues in the fingers and toes. People with this disorder usually have sparse scalp hair, and their eyelashes and eyebrows can also be sparse or absent. Affected individuals have a decreased ability to sweat (hypohidrosis), which impairs their ability to tolerate heat.

Reduced movement of joints (contractures) caused by shortening of the connective tissues that attach muscles to bone (tendons) usually develops during childhood in people with POIKTMP. These contractures often affect the calf, resulting in turning in (valgus deformity) of the feet. Contractures can also affect the elbows and wrists. In addition, people with POIKTMP usually develop muscle weakness (myopathy) in the arms and legs, and medical imaging shows abnormal fatty tissue in the muscles.

Adults with POIKTMP can develop a condition called pulmonary fibrosis, in which scar tissue forms in the lungs. Pulmonary fibrosis eventually causes difficulty breathing and can be life-threatening within a few years after symptoms begin.

In addition to the skin, muscle, and lung problems that give this condition its name, people with POIKTMP can also have a shortage (deficiency) of enzymes produced by the pancreas to aid in the digestion of fats. This deficiency can lead to diarrhea and poor absorption of fats and fat-soluble vitamins. Liver problems, short stature, and delayed puberty can also occur in affected individuals. Intellectual development is not affected by this disorder.

## 2. Frequency

The prevalence of POIKTMP is unknown. At least 25 affected individuals have been described in the medical literature. POIKTMP is thought to be under-diagnosed because affected individuals may have only one or a few features of the disorder, and health care providers may not recognize these features as part of POIKTMP.

## 3. Causes

POIKTMP is caused by mutations in the *FAM111B* gene. This gene provides instructions for making a protein whose function is not well understood. The FAM111B protein, which is found in many parts of the body, contains a functional region called a peptidase domain. Similar proteins containing such a domain are able to break down other proteins. However, the types of proteins the FAM111B protein interacts with and the roles it plays in the body are unknown.

The *FAM111B* gene mutations that cause POIKTMP result in production of an abnormal FAM111B protein from one copy of the gene in each cell. Because most of the *FAM111B* mutations identified in people with POIKTMP result in changes in the peptidase domain, researchers think that the mutations alter the protein's function, and that these changes in FAM111B function underlie the varied signs and symptoms of POIKTMP.

### 3.1. The gene associated with Hereditary fibrosing poikiloderma with tendon contractures, myopathy, and pulmonary fibrosis

- FAM111B

## 4. Inheritance

This condition is inherited in an autosomal dominant pattern, which means one copy of the altered gene in each cell is sufficient to cause the disorder.

In about half of cases, an affected person inherits the mutation from one affected parent. Other cases result from new mutations in the gene and occur in people with no history of the disorder in their family.

## 5. Other Names for This Condition

- hereditary sclerosing poikiloderma with tendon and pulmonary involvement
- HFP
- POIKTMP

---

## References

1. Khumalo NP, Pillay K, Beighton P, Wainwright H, Walker B, Saxe N, Mayosi BM, Bateman ED. Poikiloderma, tendon contracture and pulmonary fibrosis: a new autosomal dominant syndrome? *Br J Dermatol*. 2006 Nov;155(5):1057-61.
2. Küry S, Mercier S, Shaboodien G, Besnard T, Barbarot S, Khumalo NP, Mayosi BM, Bézieau S. CUGC for hereditary fibrosing poikiloderma with tendon contractures, myopathy, and pulmonary fibrosis (POIKTMP). *Eur J Hum Genet*. 2016 May;24(5). doi:10.1038/ejhg.2015.205.
3. Mercier S, Küry S, Salort-Campana E, Magot A, Agbim U, Besnard T, Bodak N, Bou-Hanna C, Bréhéret F, Brunelle P, Caillon F, Chabrol B, Cormier-Daire V, David A, Eymard B, Faivre L, Figarella-Branger D, Fleurence E, Ganapathi M, Gherardi R, Goldenberg A, Hamel A, Igual J, Irvine AD, Israël-Biet D, Kannengiesser C, Labois C, Le Caignec C, Mahé JY, Mallet S, MacGowan S, McAleer MA, McLean I, Méni C, Munnich A, Mussini JM, Nagy PL, Odel J, O'Regan GM, Péréon Y, Perrier J, Piard J, Puzenat E, Sampson JB, Smith F, Soufir N, Tanji K, Thauvin C, Ulane C, Watson RM, Khumalo NP, Mayosi BM, Barbarot S, Bézieau S. Expanding the clinical spectrum of hereditary fibrosing poikiloderma with tendon contractures, myopathy and pulmonary fibrosis due to FAM111B mutations. *Orphanet J Rare Dis*. 2015 Oct 15;10:135. doi: 10.1186/s13023-015-0352-4.
4. Mercier S, Küry S, Shaboodien G, Houniet DT, Khumalo NP, Bou-Hanna C, Bodak N, Cormier-Daire V, David A, Faivre L, Figarella-Branger D, Gherardi RK, Glen E, Hamel A, Labois C, Le Caignec C, Lindenbaum P, Magot A, Munnich A, Mussini JM, Pillay K, Rahman T, Redon R, Salort-Campana E, Santibanez-Koref M, Thauvin C, Barbarot S, Keavney B, Bézieau S, Mayosi BM. Mutations in FAM111B cause hereditary fibrosing poikiloderma with tendon contracture, myopathy, and pulmonary fibrosis. *Am J Hum Genet*. 2013 Dec 5;93(6):1100-7. doi:10.1016/j.ajhg.2013.10.013.
5. Otsu U, Moriwaki S, Iki M, Nozaki K, Horiguchi Y, Kiyokane K. Early blistering, poikiloderma, hypohidrosis, alopecia and exocrine pancreatic hypofunction: a peculiar variant of Rothmund-Thomson syndrome? *Eur J Dermatol*. 2008 Nov-Dec;18(6):632-4. doi: 10.1684/ejd.2008.0509.
6. Seo A, Walsh T, Lee MK, Ho PA, Hsu EK, Sidbury R, King MC, Shimamura A. FAM111B Mutation Is Associated With Inherited Exocrine Pancreatic Dysfunction. *Pancreas*. 2016 Jul;45(6):858-62. doi: 10.1097/MPA.0000000000000529.
7. Takeichi T, Nanda A, Yang HS, Hsu CK, Lee JY, Al-Ajmi H, Akiyama M, Simpson MA, McGrath JA. Syndromic inherited poikiloderma due to a de novo mutation in FAM111B. *Br J Dermatol*. 2017 Feb;176(2):534-536. doi: 10.1111/bjd.14845.