

Arabidopsis LSH8 ABA Signaling

Subjects: Cell Biology

Contributor: Jinpeng Zou

LSH is a family of transcription factors with diversified functions, the members of which, in turn, are named *LSH1-LSH10*. *LIGHT-DEPENDENT SHORT HYPOCOTYLS 1* (*LSH1*) was first found in Arabidopsis in 2004, the overexpression of which can enhance the light response of Arabidopsis thaliana seedlings and show an obvious short hypocotyl phenotype. All members of the LSH family have a highly conserved Domain of Unknown Function 640 (DUF640) domain, which is also called the Arabidopsis *LSH1* and Oryza G1 (ALOG) domain in the Pfam protein database.

Keywords: Arabidopsis ; LSH8 ; ABA ; seed germination ; proteomics ; ARPs

1. Introduction

As a sessile organism, plants need to undergo a complex internal regulation mechanism and environmental signal regulation to survive in adverse and changeable environments ^[1]. Phytohormone ABA is an important signaling regulator that plays a crucial role in mediating seed germination and maturation, seedling growth, stomatal movement, flowering and stress responses ^{[1][2]}. For example, ABA can regulate seed dormancy to prevent premature germination of seeds under stress conditions so that the seeds are able to germinate under suitable conditions, improving the germination rate ^[3]. These important functions of ABA are derived from the sophisticated regulatory network of ABA ^[4].

Current research demonstrates that the ABA signaling network in Arabidopsis includes five important components: ABA receptors with PYR1-like (PYL) components, negative regulator type 2C protein phosphatases (PP2C), positive regulator SNF1-related protein kinase 2 (SnRK2), transcription factors of basic leucine zippers (bZIP) and ABA-responsive genes ^[5]. The signal transduction of ABA in plants occurs in the following pathways. When ABA is deficient, PP2C with phosphatase activity dephosphorylates SnRK2 to inhibit the expression of downstream ABA-responsive genes activated by SnRK2, while in the presence of ABA, the complex of ABA binding to PYR/PYL/RCAR receptors inhibits the phosphatase activity of PP2C, from which SnRK2 is released. The released SnRK2 phosphorylates the downstream transcription factors ABI3/ABI4/ABI5 and ABA-response element binding factors (ABFs), thereby activating the expression of ABA-responsive genes ^{[4][6][7][8][9][10][11]}. Numerous previous studies have shown that a large number of transcription factors in the ABA signaling pathway play an indispensable role. For example, ABI3 is a B3-type transcription factor, and ABI5 is a bZIP transcription factor, both of which mediate ABA-induced inhibition of seed germination and initial seedling growth to participate in ABA signal transduction at the seedling stage ^{[12][13][14]}. Their loss-of-function mutation leads to the weakening of the inhibitory effect of ABA on seed germination ^{[13][15]}. Exogenous ABA significantly induces the expression of ABI5, the overexpression lines of which show a hypersensitivity phenotype to ABA during seed germination and early seedling development ^[16]. Furthermore, ABI5, as a bZIP transcription factor, can bind to the ABA binding response element (ABRE) in the promoter region of the target genes to activate the target genes' expression ^[17]. Additionally, ABF1, ABF2/AREB1, ABF3, ABF4/AREB2 and other transcription factors have been reported to play important roles in the ABA signaling network ^{[18][19]}, and after the phosphorylation caused by activated SnRK2, ABF/AREB directly binds to the promoter of stress response genes (such as *RD29A* and *RD29B*) to stimulate its transcriptional activity under stress conditions ^{[19][20][21]}. These reports suggest that a large number of transcription factors play an important role in the highly complex signaling network of ABA.

LSH is a family of transcription factors with diversified functions, the members of which, in turn, are named *LSH1-LSH10*. *LIGHT-DEPENDENT SHORT HYPOCOTYLS 1* (*LSH1*) was first found in Arabidopsis in 2004 ^[22], the overexpression of which can enhance the light response of Arabidopsis thaliana seedlings and show an obvious short hypocotyl phenotype. All members of the LSH family have a highly conserved Domain of Unknown Function 640 (DUF640) domain, which is also called the Arabidopsis *LSH1* and Oryza G1 (ALOG) domain in the Pfam protein database ^{[22][23]}. The DUF640/ALOG domain contains four all- α helices, the additional insertion of a zinc ribbon and a nuclear location signal (NLS) ^[24]. Proteins with the DUF640/ALOG domain comprise a class of specific transcription factors in plants, with characteristics of binding DNA sequence specificity, transcriptional regulation activity, nuclear localization and homodimer formation, and

control plant growth and development in many aspects. Therefore, transcription factor proteins with such a domain often have specific functions [23][25][26][27]. Studies have found that *LSH1* inhibits hypocotyl length in *Arabidopsis thaliana* in a light-dependent manner. The expression of *LSH3* and *LSH4* in the cells of various lateral organs, such as the cotyledon, the leaf and the flower organ, inhibits the differentiation of the boundary organ [25][28][29]. *LSH9* interacts with the temperature sensor ELF3 to regulate hypocotyl elongation [30][31]. In addition, proteins of the LSH family can regulate inflorescence structure and flower organ development in other plant species [32]. *LSH* family genes extensively participate in different biological processes in plants, but whether its family genes participate in plant stress response remains unknown.

2. *LSH8* Regulates Seed Germination and the Elongation of Primary and Lateral Root

LSH family genes are reported to be expressed in hypocotyl and flower organs. They are important for the growth and development of plants, but their function in the hormone signaling network remains unknown. To understand the function of *LSH8* in the ABA signaling pathway, we obtained 35S::LUC-*LSH8* overexpression lines *LSH8*-#2 and *LSH8*-#5, *lsh8* mutant lines *lsh8-1* (SALK_024841) and *lsh8-2* (CS845710). However, ABA inhibition on the seed germination of *lsh8* mutant lines *lsh8-1* and *lsh8-2* were obviously attenuated. Thus, *LSH8* overexpression lines were recognized as ABA sensitive and *lsh8* mutant lines insensitive (**Figure 1A**). As the seed germination is under the joint regulation of the hormones ABA and GA, during which GA promotes seed germination, presenting the opposite effect of ABA, we used GA biosynthesis inhibitor paclobutrazol (PAC) to verify the *LSH8* phenotype and found that under the PAC treatment condition, the seed germination of different genotypes was identical to that under ABA treatment. At the same time, we applied an appropriate amount of GA on ABA treatment conditions, the result of which showed that GA could weaken the inhibitory effect of ABA on seed germination (**Figure 1A**). To summarize, the above results indicate that ABA and GA simultaneously participate in the process of seed germination. More importantly, we identified a new positive regulator *LSH8* in the ABA signaling pathway. Further statistical results of germination rate showed that under either ABA or PAC treatment, the germination rate of *lsh8* mutant lines was significantly higher than that of Col4, while the germination rate of *LSH8* overexpression lines was significantly lower. Additionally, under ABA treatment with a moderate amount of GA, the seed germination rate of *LSH8* overexpression lines and *lsh8* mutant lines was improved (**Figure 1B**).




Figure 1. ABA phenotype of *LSH8* overexpression and *lsh8* mutant lines. **(A)** Germination phenotypes of *LSH8* overexpression lines (*LSH8*-#2 and *LSH8*-#5), *lsh8* mutant lines (*lsh8*-1 and *lsh8*-2) and wild type (Col4). Seeds were germinated and grown on 1/2 MS (mock) and 1/2 MS containing 0.5 μ M ABA, 30 μ M PAC, 0.5 μ M ABA + 50 μ M GA for 5 d, respectively. Scale bar: 1 cm. **(B)** Statistical analysis of germination rate described in **(A)**. Data represent mean \pm SD of at least 64 seeds. **(C)** Comparison of root length among genotypes on 1/2 MS with or without 30 μ M ABA, respectively. Scale bar: 1 cm. **(D)** Statistical analysis of the differences in root length among the genotypes shown in **(C)**. Data are shown as mean \pm SD ($n > 10$). Asterisks in **(B,D)** indicate statistically significant differences compared with wild-type Col4: *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$ (Student's *t*-test).

Since the *LSH8*-mediated ABA signaling pathway participates in the regulation of seed germination, whether *LSH8* is involved in ABA-mediated seedling root elongation is another issue of concern. After growing on 1/2 MS medium without ABA for 4 days, the seeds of different genotypes were transferred to 1/2 MS medium with ABA for another 4 days. The root length phenotype and root length measurements showed that when compared with wild-type Col4, the primary roots of the overexpression lines *LSH8*-#2 and *LSH8*-#5 were significantly shorter, but the primary roots of mutant lines *lsh8*-1

and *Lsh8-2* were significantly longer, and their lateral roots were obviously increasing (**Figure 1C,D**), showing that *LSH8* promotes ABA's effect on inhibiting the primary root elongation and lateral root development.

3. The Prediction of Upstream Element of *LSH8* and the Expression of *LSH8* Regulated by ABA

In the ABA signaling pathway, ABREs can be recognized by specific transcription factors to activate the expression of ABA downstream related response genes. Most of the promoter region of ABA response genes contains conserved G-box-like *cis*-elements, ABREs (PyACGTGG/TC). Previous studies have shown that genes successfully activated and expressed by ABA require multiple ABREs or one ABRE bound to several coupling elements (CEs). Analyzing the data of PlantCare, we found that the promoter region of *LSH8* contained two ABREs and one G-box element.

With ePlant (<http://bar.utoronto.ca/eplant> (accessed on 25 August 2020)), we predicted that *LSH8* was localized in the nucleus. Further protoplast subcellular localization experiment showed that *LSH8* was localized in the nucleus (**Figure 2A**). Additionally, to analyze the *LSH8* expression in different tissues, we detected the root, the stem, the leaf, the flower and the silique of Arabidopsis by qPCR, the result of which showed that the highest expression of *LSH8* happened in the flower and the silique and the lowest in the stem (**Figure 2B**). Previous studies have found that *PhLSH7b* of petunia, a homologous gene of *LSH8* in Arabidopsis, regulates plant flowering, but the function of *LSH8* in Arabidopsis is still unknown. To study the function of *LSH8*, we checked the public microarray data (<http://bar.utoronto.ca/efp/cgi-bin/efpWeb.cgi> (accessed on 13 September 2020)) and found that the expression of *LSH8* may be affected by ABA during seed germination. To confirm this expression pattern, we detected the changes in the transcriptional level and protein level of *LSH8* in wild-type Col4 under ABA treatment. The results showed that the transcriptional level of *LSH8* started to have a gradual decrease after 1.5 h of ABA treatment (**Figure 2C**). With additional Luciferase assay and Western blot detection, we found that the protein level of *LSH8* from *LSH8* overexpression lines also showed a downtrend under ABA treatment (**Figure 2D,E**). The above results indicated that ABA inhibited the expression of *LSH8* at both transcriptional and protein levels.




Figure 2. Expression pattern analysis of *LSH8*. **(A)** Subcellular localization analysis of GFP and GFP-*LSH8* in Arabidopsis wild-type Col4 protoplasts. The channels from left to right are GFP, Chloroplast, Bright and Merged channels, respectively. Scale bar: 10 μm. **(B)** Tissue-specific expression of *LSH8*. Various tissues of wild-type Col4 were grown under normal conditions, and the *LSH8* expression was determined by qPCR. Data are shown as mean ± SD (*n* = 3). **(C)** qPCR analysis of *LSH8* transcriptional level under ABA treatment. Five-day-old Col4 seedlings were treated with 100 μM ABA for 0–5 h. Data are shown as mean ± SD (*n* = 3). **(D)** LUC signals in 5-d-old LUC-*LSH8* overexpressing seedlings treated with 100 μM ABA. Signals were detected every 10 min, and the detecting period is 5 h. Data are shown as mean ± SD (*n* = 3). **(E)** Immunoblot analyzing the ABA-induced decline of *LSH8* protein in the LUC-*LSH8* overexpressing lines. Whole

seedlings of 5-day-old LUC-*LSH8* overexpression lines were treated with 100 μ M ABA for 5 h. The expression of LUC-*LSH8* fusion protein was detected by immunoblotting with an anti-LUC antibody. HSP90 was used as loading control. Asterisks in (C) indicate statistically significant differences compared with normal conditions (0 h): ns, $p > 0.05$; *, $p < 0.05$; **, $p < 0.01$ (Student's *t*-test).

References

- Hauser, F.; Waadt, R.; Schroeder, J.I. Evolution of abscisic acid synthesis and signaling mechanisms. *Curr. Biol.* 2011, 21, R346–R355. [Google Scholar] [CrossRef]
- Finkelstein, R.R.; Gampala, S.S.; Rock, C.D. Absciscic acid signaling in seeds and seedlings. *Plant Cell* 2002, 14, S15–S45. [Google Scholar] [CrossRef]
- Verslues, P.E.; Zhu, J.K. New developments in abscisic acid perception and metabolism. *Curr. Opin. Plant Biol.* 2007, 10, 447–452. [Google Scholar] [CrossRef]
- Finkelstein, R. Absciscic Acid synthesis and response. *Arab. Book* 2013, 11, e0166. [Google Scholar] [CrossRef]
- Yang, W.; Zhang, W.; Wang, X. Post-translational control of ABA signalling: The roles of protein phosphorylation and ubiquitination. *Plant Biotechnol. J.* 2017, 15, 4–14. [Google Scholar] [CrossRef]
- Umezawa, T.; Sugiyama, N.; Mizoguchi, M.; Hayashi, S.; Myouga, F.; Yamaguchi-Shinozaki, K.; Ishihama, Y.; Hirayama, T.; Shinozaki, K. Type 2C protein phosphatases directly regulate abscisic acid-activated protein kinases in Arabidopsis. *Proc. Natl. Acad. Sci. USA* 2009, 106, 17588–17593. [Google Scholar] [CrossRef]
- Vlad, F.; Rubio, S.; Rodrigues, A.; Sirichandra, C.; Belin, C.; Robert, N.; Leung, J.; Rodriguez, P.L.; Laurière, C.; Merlot, S. Protein phosphatases 2C regulate the activation of the Snf1-related kinase OST1 by abscisic acid in Arabidopsis. *Plant Cell* 2009, 21, 3170–3184. [Google Scholar] [CrossRef] [PubMed]
- Bhaskara, G.B.; Nguyen, T.T.; Verslues, P.E. Unique drought resistance functions of the highly ABA-induced clade A protein phosphatase 2Cs. *Plant Physiol* 2012, 160, 379–395. [Google Scholar] [CrossRef] [PubMed]
- Fujita, Y.; Yoshida, T.; Yamaguchi-Shinozaki, K. Pivotal role of the AREB/ABF-SnRK2 pathway in ABRE-mediated transcription in response to osmotic stress in plants. *Physiol. Plant* 2013, 147, 15–27. [Google Scholar] [CrossRef] [PubMed]
- Furihata, T.; Maruyama, K.; Fujita, Y.; Umezawa, T.; Yoshida, R.; Shinozaki, K.; Yamaguchi-Shinozaki, K. Absciscic acid-dependent multisite phosphorylation regulates the activity of a transcription activator AREB1. *Proc. Natl. Acad. Sci. USA* 2006, 103, 1988–1993. [Google Scholar] [CrossRef] [PubMed]
- Zhu, J.K. Abiotic Stress Signaling and Responses in Plants. *Cell* 2016, 167, 313–324. [Google Scholar] [CrossRef]
- Giraudat, J.; Hauge, B.M.; Valon, C.; Smalle, J.; Parcy, F.; Goodman, H.M. Isolation of the Arabidopsis ABI3 gene by positional cloning. *Plant Cell* 1992, 4, 1251–1261. [Google Scholar]
- Finkelstein, R.R.; Lynch, T.J. The Arabidopsis abscisic acid response gene ABI5 encodes a basic leucine zipper transcription factor. *Plant Cell* 2000, 12, 599–609. [Google Scholar] [CrossRef]
- Söderman, E.M.; Brocard, I.M.; Lynch, T.J.; Finkelstein, R.R. Regulation and function of the Arabidopsis ABA-insensitive 4 gene in seed and abscisic acid response signaling networks. *Plant Physiol.* 2000, 124, 1752–1765. [Google Scholar] [CrossRef] [PubMed]
- Dai, M.; Xue, Q.; McCray, T.; Margavage, K.; Chen, F.; Lee, J.H.; Nezames, C.D.; Guo, L.; Terzaghi, W.; Wan, J.; et al. The PP6 phosphatase regulates ABI5 phosphorylation and abscisic acid signaling in Arabidopsis. *Plant Cell* 2013, 25, 517–534. [Google Scholar] [CrossRef]
- Lopez-Molina, L.; Mongrand, S.; Chua, N.H. A postgermination developmental arrest checkpoint is mediated by abscisic acid and requires the ABI5 transcription factor in Arabidopsis. *Proc. Natl. Acad. Sci. USA* 2001, 98, 4782–4787. [Google Scholar] [CrossRef] [PubMed]
- Shen, Q.; Zhang, P.; Ho, T.H. Modular nature of abscisic acid (ABA) response complexes: Composite promoter units that are necessary and sufficient for ABA induction of gene expression in barley. *Plant Cell* 1996, 8, 1107–1119. [Google Scholar] [PubMed]
- Joshi, R.; Wani, S.H.; Singh, B.; Bohra, A.; Dar, Z.A.; Lone, A.A.; Pareek, A.; Singla-Pareek, S.L. Transcription Factors and Plants Response to Drought Stress: Current Understanding and Future Directions. *Front. Plant Sci.* 2016, 7, 1029. [Google Scholar] [CrossRef] [PubMed]
- Yoshida, T.; Fujita, Y.; Sayama, H.; Kidokoro, S.; Maruyama, K.; Mizoi, J.; Shinozaki, K.; Yamaguchi-Shinozaki, K. AREB1, AREB2, and ABF3 are master transcription factors that cooperatively regulate ABRE-dependent ABA signaling in

- lved in drought stress tolerance and require ABA for full activation. *Plant J. Cell Mol. Biol.* 2010, 61, 672–685. [Google Scholar] [CrossRef] [PubMed]
20. Uno, Y.; Furihata, T.; Abe, H.; Yoshida, R.; Shinozaki, K.; Yamaguchi-Shinozaki, K. Arabidopsis basic leucine zipper transcription factors involved in an abscisic acid-dependent signal transduction pathway under drought and high-salinity conditions. *Proc. Natl. Acad. Sci. USA* 2000, 97, 11632–11637. [Google Scholar] [CrossRef] [PubMed]
 21. Narusaka, Y.; Nakashima, K.; Shinwari, Z.K.; Sakuma, Y.; Furihata, T.; Abe, H.; Narusaka, M.; Shinozaki, K.; Yamaguchi-Shinozaki, K. Interaction between two cis-acting elements, ABRE and DRE, in ABA-dependent expression of Arabidopsis rd29A gene in response to dehydration and high-salinity stresses. *Plant J. Cell Mol. Biol.* 2003, 34, 137–148. [Google Scholar] [CrossRef]
 22. Zhao, L.; Nakazawa, M.; Takase, T.; Manabe, K.; Kobayashi, M.; Seki, M.; Shinozaki, K.; Matsui, M. Overexpression of LSH1, a member of an uncharacterised gene family, causes enhanced light regulation of seedling development. *Plant J. Cell Mol. Biol.* 2004, 37, 694–706. [Google Scholar] [CrossRef]
 23. Yoshida, A.; Suzaki, T.; Tanaka, W.; Hirano, H.Y. The homeotic gene long sterile lemma (G1) specifies sterile lemma identity in the rice spikelet. *Proc. Natl. Acad. Sci. USA* 2009, 106, 20103–20108. [Google Scholar] [CrossRef]
 24. Chen, F.; Zhou, Q.; Wu, L.; Li, F.; Liu, B.; Zhang, S.; Zhang, J.; Bao, M.; Liu, G. Genome-wide identification and characterization of the ALOG gene family in Petunia. *BMC Plant Biol.* 2019, 19, 600. [Google Scholar] [CrossRef] [PubMed]
 25. Takeda, S.; Hanano, K.; Kariya, A.; Shimizu, S.; Zhao, L.; Matsui, M.; Tasaka, M.; Aida, M. CUP-SHAPED COTYLEDON1 transcription factor activates the expression of LSH4 and LSH3, two members of the ALOG gene family, in shoot organ boundary cells. *Plant J. Cell Mol. Biol.* 2011, 66, 1066–1077. [Google Scholar] [CrossRef] [PubMed]
 26. Iyer, L.M.; Aravind, L. ALOG domains: Provenance of plant homeotic and developmental regulators from the DNA-binding domain of a novel class of DIRS1-type retrotransposons. *Biol. Direct* 2012, 7, 39. [Google Scholar] [CrossRef] [PubMed]
 27. Peng, P.; Liu, L.; Fang, J.; Zhao, J.; Yuan, S.; Li, X. The rice TRIANGULAR HULL1 protein acts as a transcriptional repressor in regulating lateral development of spikelet. *Sci. Rep.* 2017, 7, 13712. [Google Scholar] [CrossRef]
 28. Hibara, K.; Takada, S.; Tasaka, M. CUC1 gene activates the expression of SAM-related genes to induce adventitious shoot formation. *Plant J. Cell Mol. Biol.* 2003, 36, 687–696. [Google Scholar] [CrossRef]
 29. Kamiuchi, Y.; Yamamoto, K.; Furutani, M.; Tasaka, M.; Aida, M. The CUC1 and CUC2 genes promote carpel margin meristem formation during Arabidopsis gynoecium development. *Front. Plant Sci.* 2014, 5, 165. [Google Scholar] [CrossRef]
 30. Press, M.O.; Queitsch, C. Variability in a Short Tandem Repeat Mediates Complex Epistatic Interactions in Arabidopsis thaliana. *Genetics* 2017, 205, 455–464. [Google Scholar] [CrossRef] [PubMed]
 31. Hayes, S.; Schachtschabel, J.; Mishkind, M.; Munnik, T.; Arisz, S.A. Hot topic: Thermosensing in plants. *Plant Cell Environ.* 2021, 44, 2018–2033. [Google Scholar] [CrossRef]
 32. Yoshida, A.; Sasao, M.; Yasuno, N.; Takagi, K.; Daimon, Y.; Chen, R.; Yamazaki, R.; Tokunaga, H.; Kitaguchi, Y.; Sato, Y.; et al. TAWAWA1, a regulator of rice inflorescence architecture, functions through the suppression of meristem phase transition. *Proc. Natl. Acad. Sci. USA* 2013, 110, 767–772. [Google Scholar] [CrossRef] [PubMed]