# **Microbe-Associated Molecular Patterns**

#### Subjects: Plant Sciences

Contributor: Benedict C. Offor , Ian A. Dubery , Lizelle A. Piater

Plants depend on both preformed and inducible defence responses to defend themselves against biotic stresses stemming from pathogen attacks. In this regard, plants perceive pathogenic threats from the environment through pattern recognition receptors (PRRs) that recognise microbe-associated molecular patterns (MAMPs), and so induce plant defence responses against invading pathogens. Close to thirty PRR proteins have been identified in plants, however, the molecular mechanisms underlying MAMP perception by these receptors/receptor complexes are not fully understood. As such, knockout (KO) of genes that code for PRRs and co-receptors / defence-associated proteins is a valuable tool to study plant immunity. The loss of gene activity often causes changes in the phenotype of the model plant, allowing in vivo studies of gene function and associated biological mechanisms. Here, we review the functions of selected PRRs, brassinosteroid insensitive 1 (BRI1) associated receptor kinase 1 (BAK1) and other associated defence proteins that have been identified in plants, and also outline KO lines generated by T-DNA insertional mutagenesis as well as the effect on MAMP perception - and triggered immunity (MTI). In addition, we further review the role of membrane raft domains in flg22-induced MTI in Arabidopsis, due to the vital role in the activation of several proteins that are part of the membrane raft domain theory in this regard.

BAK1 innate immunity KO MAMPs membrane raft MTI PRRs T-DNA

# 1. Well-Studied MAMP Perception System(s)

## **1.1. Bacterial MAMPs**

### 1.1.1. Flagellin Perception

Flagellin is the main protein component of bacterial flagella, that acts as a MAMP in both plants and animals. Plants have a notably sensitive perception system for the highly conserved domain in the N-terminus of eubacterial flagellin <sup>[1]</sup>. The perception thereof, and particularly the N-terminus flagellin peptides, flg22 and flg15 from *Pseudomonas syringae* pv. *tabaci*, lead to the alkalinisation of the culture medium in suspension-cultured cells of *Lycopersicon peruvianum* (a wild relative of tomato), *A. thaliana*, potato and tobacco respectively <sup>[1]</sup>. Flagellin from the rice-incompatible strain, N1141 of *P. avenae*, was also shown to induce the hypersensitive cell death and accumulation of EL2 mRNA (elicitor-responsive gene) in cultured rice cells <sup>[2]</sup>. As mentioned, FLS2 is a LRR-RLK involved in the perception of the bacterial flagellin and immune responses in *A. thaliana* <sup>[3][4]</sup>. Furthermore, the recognition of bacterial flg22 by Arabidopsis FLS2 induces a FLS2-BAK1 complex formation and triggers defence signalling <sup>[4][5]</sup>. By using chemical cross-linking and immunoprecipitation techniques, Chinchilla et al. <sup>[6]</sup> showed that the specificity of flagellin perception and immune responses is mediated by the binding of FLS2 to flg22 in *A.* 

*thaliana*. The *fls2* mutants carrying T-DNA insertion in the flagellin receptor gene *FLS2* were more susceptible to *Pst* DC3000 compared to the wildtype (WT) <sup>[Z]</sup>. Upon flg22 treatment, *FLS2* mutants (*fls2-101*), generated by the insertion of T-DNA at the third *FLS2* exon, showed impaired binding of the flg22 and reduced seedling growth inhibition <sup>[8]</sup>.

As stated, flg22 induces a FLS2-BAK1 complex in a ligand-dependent manner, and leads to defence responses in *A. thaliana* <sup>[4]</sup>. This study showed that although *BAK1* mutants (*bak1-3* and *bak1-4* with T-DNA inserted in 5<sup>th</sup> intron and 9<sup>th</sup> exon, respectively) were defective in an oxidative burst generation, they did not affect the binding of flg22 to FLS2. Furthermore, flg22 induces recruitment of the U-box E3 ubiquitin ligases (PUB12 and PUB13), previously mentioned, to the FLS2-BAK1 complex and this causes FLS2 degradation and the concomitant attenuation of immune signalling in Arabidopsis <sup>[9]</sup>. Here, there was a decrease in FLS2 ubiquitination and *Pst* DC3000 infection in mutant plants (*pub12-2, pub13* and *pub12/13* double knockout), compared to WT controls. A recent study has furthermore shown that flagellin-sensing 3 (FLS3) directly and specifically binds flgII-28 (a second flagellin epitope distinct from flg22), which enhances immune responses in tomato <sup>[10]</sup>. Finally, it is also important to note that BAK1 is involved in the defence signalling activities of both FLS2 and FLS3 <sup>[4][10]</sup>.

#### 1.1.2. EF-Tu Perception

Elongation factor-thermo unstable (EF-Tu), a prokaryotic elongation factor involved in the synthesis of proteins, is a bacterial MAMP that is recognised by the LRR-RLK, EFR, in Arabidopsis [11]. N. benthamiana normally lacks EF-Tu responsiveness, but achieved the ability to recognise the MAMP when EFR was transiently expressed in the plant [11]. Kunze et al. [12] showed that EFR binds directly to the acetylated N-terminus epitope of elf18 (the first amino acids of EF-Tu) and elicits innate immunity in Arabidopsis and other Brassicaceae species. The first acetylated 12 N-terminal amino acids (elf12) were, however, not able to elicit an immune response, but rather acted as a specific antagonist of elf18 [12]. Interestingly, even though EF-Tu induces MTI in rice, the elf18 peptide failed to trigger an immune response. However, another epitope, EFa50 (50-amino acid from central region of EF-Tu comprising Lys176 to Gly222) induced MTI responses, although the related PRR is still unknown <sup>[13]</sup>. Rice leaves treated with EFa50 induced early defence responses such as increased  $H_2O_2$  and callose deposition, and triggered resistance to coinfection with pathogenic bacteria. In this regard, Arabidopsis EFR T-DNA insertion mutants (efr-1 and efr-2), with both T-DNAs inserted in 1<sup>st</sup> exon, were insensitive to EF-Tu and susceptible to the bacterium Agrobacterium tumefaciens [11] (Table 1). In fact, efr mutants did not show typical growth inhibition as for the WT, increased oxidative burst and ethylene biosynthesis or induced resistance to Pst DC3000 upon EF-Tu treatment. Jeworutzki et al. [14] reported involvement of EFR and FLS2 in the Ca<sup>2+</sup>-associated opening of plasma membrane anion channels during early bacterial flagellin and EF-Tu defence signalling in Arabidopsis mesophyll cells. Using electrophysiological approaches, they showed that the efr-1 was defective in the membrane potential depolarisation, which is indispensable for cytosolic calcium influx in response to elf18 treatment.

#### 1.1.3. LPS Perception

LPS is an amphipathic molecule found on the outer membrane of Gram-negative bacteria and protects the organism against antimicrobial compounds found in the environment <sup>[15][16]</sup>. This MAMP is involved in the bacterial

adhesion and induction of defence-related responses in both mammals and plants <sup>[17]</sup>. In the former, LPS recognition is orchestrated by lipopolysaccharide binding protein (LBP), before recruitment into a complex comprising soluble myeloid differentiation protein 2 (MD-2), membrane attached cluster of differentiation 14 (CD14), and the transmembrane Toll-like receptor 4 (TLR4), thereby leading to mammalian defence activation against LPS <sup>[18][19]</sup>. LBP and bactericidal/permeability-increasing protein (BPI) thus play a vital role in regulation of immune responses against LPS in mammals <sup>[19][20]</sup>. As such, the mechanism by which LPS is recognised has been well-studied in animals, however, in plants, the mechanism of perception and recognised moiety/epitope(s) of LPS is still debateable.

In this regard, LPS, as well as its lipid A moiety from *Burkholderia cepacia, Xanthomonas campestris and P. syringae*, trigger the upregulation of genes involved in immunity and defence <sup>[21][22][23]</sup>, with lipid A speculated to be the major elicitor in *A. thaliana* <sup>[24]</sup>. LPS has furthermore been shown to trigger defence responses, such as the generation of nitrogen oxide (NO), ROS, elevation of cytoplasmic Ca<sup>2+</sup> concentration, stomatal closure and the expression of pathogenesis-related (PR) genes in Arabidopsis, tobacco and rice suspension cells <sup>[25][26][27][28][29]</sup>. Ranf et al. <sup>[25]</sup> earlier reported that LPS from *Xanthomonas* and *Pseudomonas* are sensed by the bulb-type (B-type) lectin S-domain (SD)-1 RLK lipooligosaccharide-specific reduced elicitation (LORE) and that *LORE* mutants showed defects in the LPS-induced elevation of cytosolic calcium, ROS and defence gene (*AtFRK1*) expression. Additionally, Shang-Guan et al. <sup>[24]</sup> reported the existence of two ROS production phases, characterised by a weak initial and second stronger ROS generation in *A. thaliana*. T-DNA insertion *LORE* mutants, however, showed little or no difference in second phase ROS production compared with WT, suggesting another LPS receptor(s). In contradiction to the original LORE study, a recent report implicated bacterial 3-hydroxy fatty acids in LORE-dependent induction of immune response in Arabidopsis <sup>[30]</sup>. Here, LORE could not sense LPS that was repurified to remove free 3-hydroxy fatty acids, indicating that LORE is not the receptor for LPS.

Using *B. cepacia* LPS-affinity capture strategies, Vilakazi et al. <sup>[31]</sup> and Baloyi et al. <sup>[32]</sup> identified BAK1 and other defence response proteins associated with the plasma membrane fraction in LPS-treated *A. thaliana*. Additionally, Arabidopsis LBP/BPI related-1 (AtLBR-1) and LBP/BPI related-2 (AtLBR-2) were shown to bind to both rough and smooth LPS, and regulate the expression of the *pathogenesis-related* 1 (*PR1*) gene <sup>[23]</sup>. *AtLBR* T-DNA single mutants (*Atlbr-1*, *Atlbr-2-1*, *Atlbr2-2*) and (*Atlbr*-DKO) double-knockouts, generated by crossing *Atlbr-1* and *Atlbr-2-1* plants, were defective in ROS generation and in upregulation of LPS-induced *PR1* expression. A transcriptome analysis revealed that AtLBR-2 plays an indispensable role in the upregulation of 65 genes associated with defence responses upon *Pseudomonas* LPS treatment <sup>[33]</sup>. Furthermore, there was a defect in the upregulation of defence-related genes and salicylic acid (SA)-mediated signalling in the *Atlbr-2-1* mutants, compared to the WT after *Pseudomonas* LPS treatment.

Lastly, a recent study has reported OsCERK1, the chitin co-receptor, as an LPS receptor/co-receptor in rice, but not in *A. thaliana* <sup>[34]</sup>. This indicates a significant difference between LPS perception in rice and Arabidopsis.

#### **1.1.4. Peptidoglycan Perception**

Rice chitin elicitor-binding protein (CEBiP) homologs in Arabidopsis, LYM1 and LYM3 RLPs, bind in a ligandspecific manner to PGNs, heteropolymers that are part of the building blocks of the cell walls of Gram-negative and Gram-positive bacteria [35]. There was a significant expression of the immune marker gene, flagellin-induced receptor kinase FRK1, upon treatment of Arabidopsis with PGNs from Gram-negative Pst DC3000 [35]. Homozygous T-DNA insertional mutants of LYM1 and LYM3 showed a strongly reduced PGN-inducible marker gene expression and were more susceptible to Pst DC3000 bacterial infection than the WT (Table 1). The same authors further reported the involvement of AtCERK1 in the LYM1/LYM3 perception of PGN and subsequent immune response in A. thaliana. Thus, AtCERK1 is the additional protein that provides the cytoplasmic kinase domain that is lacking in the LYM1/LYM3 RLPs needed for the downstream transphosphorylation of PGN signalling. The study suggests that Arabidopsis senses PGNs in a LYM1/LYM3 and AtCERK1-dependent manner, similar to the chitin perception system in rice that uses OsCEBIP and OsCERK1 [36]. Furthermore, the direct interaction of PGN with AtLYM1 and AtLYM3 has been reported, but not with AtCERK1 [35]. In rice, OsCERK1 associates with LysM motif-containing proteins (LYP4 and LYP6) in PGN-induced defence responses [37][38][39]. Importantly, the putative ability of CERK1 to participate in the recognition and signalling of more than one MAMP supports the hypothesis of the capability of one PRR/co-receptor to recognise more than one MAMP, which favours the generation of transgenic crop plants with enhanced/altered recognition capabilities [40].

# 1.2. Fungal MAMPs

### 1.2.1. Chitin Perception

Chitin is a  $\beta(1 \rightarrow 4)$ -linked polymer of N-acetylglucosamine, a major structural component in the exoskeleton of arthropods and cell walls of fungi [41]. Chitin and its fragments (chitooligosaccharides) are MAMPs recognised by plants PRRs, which elicit defence responses such as the oxidative burst, protein phosphorylation, transcriptional activation of defence-related genes and phytoalexin biosynthesis [41][42][43]. Kaku et al. [42] isolated and characterised a chitin elicitor binding protein (OsCEBiP), an RLP involved in the perception of chitin oligosaccharides in cultured rice cells. It was observed that OsCEBiP has two LysM motifs and a C-terminal transmembrane domain, however no intracellular kinase domain, suggesting that it requires additional proteinpartner(s) to perform a signal transduction role. In rice, the LysM receptor, OsCEBiP, binds to the chitin oligosaccharide and forms a complex with OsCERK1 to trigger defence response [36]. Conversely, in *A. thaliana*. the LysM receptor, AtCERK1, directly binds to chitin, dimerises and triggers immune responses [44]. In the extracellular domain, OsCERK1 has one LysM motif, whereas AtCERK1 has three LysM motifs [36][45] that mediate the binding of chitin [46][47]. There are five genes encoding LysM receptor-like kinases (LYKs), which are comprised of LYK1 (CERK1) and LYK2 - LYK5, and three genes encoding LysM receptor-like proteins (LYPs) in the A. thaliana genome [48]. In this regard, LysM receptor-like kinase 1 (LysM RLK1) is involved in chitin signalling and fungal resistance in Arabidopsis <sup>[49]</sup>. A T-DNA insertion knockout mutant of LysM RLK1 blocked the induction of chitooligosaccharide-responsive genes (CRGs) by chitooligosaccharides and increased susceptibility to fungal pathogens, but not bacterial pathogens. This was reversed when the mutant was complemented with the WT LysM *RLK1* gene using the cauliflower mosaic virus (CaMV) 35S promoter [44][49]. CERK1 KO mutants were unable to

respond to chitin oligosaccharide elicitors when compared to the WT that exhibited rapid generation of ROS in Arabidopsis <sup>[44]</sup>.

LYK4 is another LysM-RLK involved in chitin signalling and plant immunity in *A. thaliana*, and possibly in the chitin recognition receptor complex <sup>[50]</sup>. *LYK4* mutants were defective in the activation of chitin-responsive genes and were also susceptible to bacterial and fungal pathogen infection. Cao et al. <sup>[51]</sup> showed that AtLYK5 binds chitin with higher affinity and forms a chitin-induced complex with AtCERK1, thereby triggering immunity in *A. thaliana*. The higher affinity of AtLYK5 for chitin, compared to AtCERK1, also suggests the former to be the major chitin binding protein in Arabidopsis. These authors <sup>[51]</sup> further reported involvement of AtLYK5 in chitin-induced AtCERK1 phosphorylation and homodimerisation <sup>[51]</sup>. While *Atlyk5-2* was significantly impaired in chitin signal responses, *Atlyk4*/*Atlyk5-2* double mutant resulted in a complete loss of chitin response, indicating an overlap of signal function between AtLYK5 and AtLYK4.

The LysM domain-containing glycosylphosphatidylinositol-anchored protein 2 (LYM2), one of the three CEBiP homologs in Arabidopsis, binds chitin and mediates a reduction in molecular flux via the plasmodesmata <sup>[52]</sup>. AtCERK1 is not involved in the chitin-mediated regulation of plasmodesmata flux, thereby suggesting the presence of an alternative novel disease resistance mechanism in Arabidopsis [52][53]. Shinya et al. [54] showed that the Arabidopsis LYM2 can recognise chitin oligosaccharides in a similar way as the rice OsCEBiP, but does not participate in chitin signalling. The KO mutant of LYM2 (lym2-1) was shown to be incapable of chitin-induced plasmodesmata flux, and susceptible to fungal pathogens (Botrytis cenerea and Alternaria brassicicola), but exhibited chitin-induced MAPK activation and an oxidative burst when compared to the WT [52][54]. LvsM RLK1interacting kinase 1 (LIK1) interacts with CERK1 and regulates chitin-induced MTI in Arabidopsis [55]. LIK1 mutants (lik1-1, lik1-2, lik1-3 and lik1-4), with T-DNA insertions located in intron 2, 13, and exon 18, respectively, showed an enhanced response to both chitin and flagellin elicitors. Furthermore, the mutants were defective in the expression of genes involved in jasmonic acid (JA) and ethylene (ET) signalling pathways, that have been shown to mediate resistance to necrotrophic pathogens <sup>[56]</sup>. In *A. thaliana* a powdery mildew-resistant kinase 1 (PMRK1), a RLK that is localised in the plasma membrane, is responsible for early chitin-induced defence signals against fungal pathogens [57]. PMRK1 KO mutant (pmrk1) was more susceptible to both Golovinomyces cichoracearum and Plectosphaerella cucumerina (Table 1). In fact, these numerous identified PPRs linked to chitin perception suggest the complexity of plant chitin defence signalling.

# 2. BAK1 and Other Associated Proteins in MAMP Signalling

Following MAMP perception, PRRs trigger downstream events involving protein association/dissociation. BAK1 was initially identified as a co-receptor in BRI1-mediated brassinosteroid (BR) signalling, which modulates plant growth and development <sup>[58][59]</sup>. Studies have shown that BAK1 and related somatic embryogenesis receptor kinases (SERK) proteins associate with other LRR-RLKs or LRR-RLPs, and regulate plant growth and immunity <sup>[4]</sup> [60][61].

Table 2 outlines different BAK1 and other associated proteins, and the implication of their KOs in plant MTI. In *A. thaliana*, both FLS2 and EFR form a complex with the co-receptor BAK1 to elicit immune responses immediately upon flg22 or elf18 perception, respectively <sup>[4][5][62][63]</sup>. Plants carrying *BAK1* mutants (*bak1-3* and *bak1-4*), generated by T-DNA insertion, displayed abnormal early and late flagellin-triggered responses <sup>[4][5]</sup>. In this regard, there was a significant reduction in the oxidative burst triggered by elf26 in *BAK1* mutants, indicating that EF-Tu is also affected by the mutation in BAK1 <sup>[4]</sup>. Interestingly, *BAK1* mutants were not completely impaired to flg22 or elf18 perception, indicating that BAK1 was not the only rate-limiting component and therefore suggests additional regulatory protein(s), such as BKK1, that are part of the FLS2 and EFR receptor complexes <sup>[4][5][60]</sup>. BAK1-disrupted *N. benthamiana* plants displayed decreased induction of MTI responses by the csp22 peptide (part of bacterial cold-shock protein) and INF1 (an oomycete elicitor) <sup>[5]</sup>. Furthermore, Arabidopsis *BAK1* KO mutants *brassicicola* <sup>[64]</sup>. These results suggest a central role for BAK1 in modulating other PRRs besides FLS2 and EFR in plant defence signalling. The exact mechanism by which BAK1 mediates defence signalling is, however, not resolved. A recent study also showed BAK1 involvement in the tomato FLS3 recognition of flgII-28 (another flagellin epitope) and resulting immune response signalling <sup>[10]</sup>.

BAK1 and BAK1-LIKE1 (BKK1) have dual physiological roles by positively regulating a BR-dependent plant growth pathway, and negatively regulating a BR-independent cell-death <sup>[65]</sup>. Here, cell death-control mediated by BAK1 and BKK1 is SA-dependent <sup>[66]</sup>. Upon flagellin perception, BIK1 as a RLCK, associates with the FLS2-BAK1 receptor complex to initiate plant innate immunity and cell death <sup>[67]</sup>. There was a significant loss of flg22-induced resistance to *Pst* DC3000 infection in *BIK1* mutant seedlings, however, the mutation did not affect flg22-induced FLS2 and BAK1 association. On the other hand, *BIK1* mutants were susceptible to necrotrophic pathogens but were resistant to a virulent bacterial pathogen *Pst* DC3000 <sup>[68]</sup>. Chen et al. <sup>[69]</sup> demonstrated that the *bik1* mutant displayed a strong SA-dependent resistance to *Plasmodiophora brassicae*, an obligate biotroph protist that induces gall formation in cruciferous plants. *Bak1-4 bik1* double mutants exhibited increased expression of plant defence genes and cell death phenotypes compared to *BIK1* single mutant <sup>[70]</sup>, highlighting the cooperativity of BIK1 and BAK1 influence in plant immunity.

BIR2, a novel LRR-RLK, interacts with BAK1 in a kinase-dependent manner, and negatively regulates BAK1dependent MAMP-triggered immune signalling <sup>[71]</sup>. Upon ligand binding to FLS2, BAK1 is released from BIR2 and recruited to the FLS2 complex. Therefore, BIR2 inhibits autoimmune cell-death responses by keeping BAK1 under control. Gao et al. <sup>[72]</sup> showed that BIR1, a BAK1-interacting RLK, negatively regulates multiple plant resistance signalling responses, and suppresses cell death in Arabidopsis. *BIR1* KO mutants (*bir1-1*) showed activation of constitutive defence responses and extensive cell death. However, the LRR-RLK SUPPRESSOR OF BIR1-1 (SOBIR1) and BAK1 function as co-receptors for LRR-RLPs, BAK1, and not SOBIR1, acts a co-receptor for LRR-RLKs <sup>[63]</sup>. SOBIR1, a co-receptor/adaptor for LRR-RLPs recruits BAK1 to SOBIR1-RLP23 and SOBIR1-RLP30 complex upon nlp20 and Sclerotinia culture filtrate elicitor1 (SCFE1) perception, respectively, in Arabidopsis <sup>[73][74]</sup>. Here, *SOBIR1* mutant (*sobir1-12*) was more susceptible to fungal *Sclerotinia sclerotiorum* and *B. cineria* <sup>[75]</sup>. The dissociation of BIR1 upon MAMP recognition by PRRs allows BAK1 to form an active complex with SOBIR1, which triggers downstream cell death and defence signalling <sup>[76]</sup>. The Arabidopsis malectin-like LRR-RLK, IMPAIRED OOMYCETE SUSCEPTIBILITY1 (IOS1) associated with PRRs FLS2, EFR and CERK1 in BAK1-dependent and -independent MTI responses <sup>[72]</sup>. Arabidopsis *IOS1* mutant (*ios1-2*) showed perturbations in the latter, including defective chitin responses and delayed upregulation of the PTI marker gene *FLG22-INDUCED RECEPTOR-LIKE KINASE1* (*FRK1*), as well as reduced downy mildew infection <sup>[74]</sup>. The malectin-like RLK FERONIA (FER), facilitates the ligand-induced complex formation of PRRs in Arabidopsis <sup>[78][79]</sup>. As such, the EFR/FLS2-BAK1 complex formation has been shown to be promoted by FER and inhibited by Rapid Alkalinization Factor 23 (RALF23) <sup>[77]</sup>. Furthermore, a *FER* mutant (*fer-4*) showed diminished ligand-induced EFR/FLS2 complex formation, with the co-receptor BAK1. In addition, AtFER is involved in the negative regulation of jasmonic acid (JA) and coronatine (COR) signalling <sup>[79]</sup>. In support, BAK1 and other defence-responsive proteins were identified in *A. thaliana* plasma membranes after *B. cepacia* and *E. coli* LPS treatments <sup>[31][32]</sup>. Here, proteins identified were similar to some previously implicated proteins upon flg22 elicitation, suggesting that LPS perception and signalling could likely resemble that of flg22.

# References

- 1. Georg Felix; Juliana D. Duran; Sigrid Volko; Thomas Boller; Plants have a sensitive perception system for the most conserved domain of bacterial flagellin. *The Plant Journal* **1999**, *18*, 265-276, 10.1046/j.1365-313x.1999.00265.x.
- Fang-Sik Che; Yoshihiro Nakajima; Noriko Tanaka; Megumi Iwano; Tomomi Yoshida; Seiji Takayama; Ikuo Kadota; Akira Isogai; Flagellin from an Incompatible Strain ofPseudomonas avenaeInduces a Resistance Response in Cultured Rice Cells. *Journal of Biological Chemistry* 2000, 275, 32347-32356, 10.1074/jbc.m004796200.
- 3. L Gómez-Gómez; T Boller; FLS2: an LRR receptor-like kinase involved in the perception of the bacterial elicitor flagellin in Arabidopsis.. *Molecular Cell* **2000**, *5*, 1003–1011.
- Delphine Chinchilla; Cyril Zipfel; Silke Robatzek; Birgit Kemmerling; Thorsten Nürnberger; Alexandra M.E. Jones; Georg Felix; Thomas Boller; A flagellin-induced complex of the receptor FLS2 and BAK1 initiates plant defence. *Nature* 2007, 448, 497-500, 10.1038/nature05999.
- Antje Heese; Dagmar R. Hann; Selena Gimenez-Ibanez; Alexandra M.E. Jones; Kai He; Jia Li; Julian I. Schroeder; Scott C. Peck; John P. Rathjen; The receptor-like kinase SERK3/BAK1 is a central regulator of innate immunity in plants. *Proceedings of the National Academy of Sciences* 2007, 104, 12217-12222, 10.1073/pnas.0705306104.
- Delphine Chinchilla; Zsuzsa Bauer; Martin Regenass; Thomas Boller; Georg Felix; The Arabidopsis receptor kinase FLS2 binds flg22 and determines the specificity of flagellin perception. *Plant Cell* 2006, *18*, 465-476, 10.1105/tpc.105.036574.
- 7. Cyril Zipfel; Silke Robatzek; Lionel Navarro; Edward J. Oakeley; Alexandra M.E. Jones; Georg Felix; Thomas Boller; Bacterial disease resistance in Arabidopsis through flagellin perception.

Nature 2004, 428, 764-767, 10.1038/nature02485.

- F. Mark Dunning; Wenxian Sun; Kristin L. Jansen; Laura Helft; Andrew Bent; Identification and Mutational Analysis of Arabidopsis FLS2 Leucine-Rich Repeat Domain Residues That Contribute to Flagellin Perception. *The Plant Cell* 2007, 19, 3297-3313, 10.1105/tpc.106.048801.
- Lu, D.; Lin, W.; Gao, X.; Wu, S.; Cheng, C.; Avila, J.; Heese, A.; Devarenne, T.P.; He, P.; Shan, L. Direct ubiquitination of pattern recongnition receptor FLS2 attenuates plant innate immunity. Science 2011, 332, 1439–1442.
- Sarah R. Hind; Susan Strickler; Patrick C. Boyle; Diane M. Dunham; Zhilong Bao; Inish M. O'doherty; Joshua A. Baccile; Jason S. Hoki; Elise G. Viox; Christopher Clarke; et al.Boris A VinatzerFrank C. SchroederGregory B. Martin Tomato receptor FLAGELLIN-SENSING 3 binds flgII-28 and activates the plant immune system. *Nature Plants* **2016**, *2*, 1-8, 10.1038/nplants.2016. 128.
- 11. Cyril Zipfel; Gernot Kunze; Delphine Chinchilla; Anne Caniard; Alexandra M.E. Jones; Thomas Boller; Georg Felix; Perception of the Bacterial PAMP EF-Tu by the Receptor EFR Restricts Agrobacterium-Mediated Transformation. *Cell* **2006**, *125*, 749-760, 10.1016/j.cell.2006.03.037.
- 12. Kunze, G; The N terminus of bacterial Elongation Factor Tu elicits innate immunity in Arabidopsis plants. *Plant Cell* **2004**, *16*, 3496–3507.
- 13. Takehito Furukawa; Hiroaki Inagaki; Ryota Takai; Hiroyuki Hirai; Fang-Sik Che; Two Distinct EF-Tu Epitopes Induce Immune Responses in Rice and Arabidopsis. *Molecular Plant-Microbe Interactions*® **2014**, *27*, 113-124, 10.1094/mpmi-10-13-0304-r.
- Elena Jeworutzki; M. Rob G. Roelfsema; Uta Anschütz; Elżbieta Król; J. Theo M. Elzenga; Georg Felix; Thomas Boller; Rainer Hedrich; D. Becker; Early signaling through the Arabidopsis pattern recognition receptors FLS2 and EFR involves Ca2+-associated opening of plasma membrane anion channels. *The Plant Journal* 2010, *62*, 367-378, 10.1111/j.1365-313x.2010.04155.x.
- 15. Alba Silipo; Gitte Erbs; Tomonori Shinya; J. Maxwell Dow; Michelangelo Parrilli; Rosa Lanzetta; Naoto Shibuya; Mari-Anne Newman; Antonio Molinaro; Glyco-conjugates as elicitors or suppressors of plant innate immunity. *Glycobiology* **2009**, *20*, 406-419, 10.1093/glycob/cwp201.
- 16. Chaeyeong Lee; Mohamed Mannaa; Namgyu Kim; Juyun Kim; Yeounju Choi; Soo Hyun Kim; Boknam Jung; Hyun-Hee Lee; Jungkwan Lee; Young-Su Seo; et al. Stress Tolerance and Virulence-Related Roles of Lipopolysaccharide in Burkholderia glumae.. *Plant Pathol. J* 2019, 35, 445-458.
- 17. Stefanie Ranf; Immune Sensing of Lipopolysaccharide in Plants and Animals: Same but Different. *PLOS Pathogens* **2016**, *12*, e1005596, 10.1371/journal.ppat.1005596.
- 18. Thorsten Nurnberger; Frédéric Brunner; Birgit Kemmerling; Lizelle Piater; Innate immunity in plants and animals: striking similarities and obvious differences.. *Immunological Reviews* **2004**,

198, 249-266, 10.1111/j.0105-2896.2004.0119.x.

- Weiss, J; Bactericidal/permeability-increasing protein (BPI) and lipopolysaccharide-binding protein (LBP): Structure, function and regulation in host defence against Gram-negative bacteria. *Biochem. Soc. Trans.* 2003, 31, 785–790.
- 20. Peter Elsbach; Jerrold Weiss; Role of the bactericidal/permeability-increasing protein in host defence. *Current Opinion in Immunology* **1998**, *10*, 45-49, 10.1016/s0952-7915(98)80030-7.
- Ntakadzeni E Madala; Maria R Leone; Antonio Molinaro; Ian A. Dubery; Deciphering the structural and biological properties of the lipid A moiety of lipopolysaccharides from Burkholderia cepacia strain ASP B 2D, in Arabidopsis thaliana. *Glycobiology* **2010**, *21*, 184-194, 10.1093/glycob/cwq14 6.
- Ntakadzeni E Madala; Antonio Molinaro; Ian A. Dubery; Distinct carbohydrate and lipid-based molecular patterns within lipopolysaccharides from Burkholderia cepacia contribute to defenseassociated differential gene expression in Arabidopsis thaliana. *Innate Immunity* 2011, 18, 140-154, 10.1177/1753425910392609.
- Lizasa, S.; Lizasa, E.; Matsuzaki, S.; Tanaka, H.; Kodama, Y.; Watanabe, K.; Nagano, Y. Arabidopsis LBP/BPI related-1 and -2 bind to LPS directly and regulate PR1 expression. Sci. Rep. 2016, 6, 1–10.
- Keke Shang-Guan; Min Wang; Nang Myint Phyu Sin Htwe; Ping Li; Yaoshen Li; Fan Qi; Dawei Zhang; Min Cao; Chanhong Kim; Haiyong Weng; Haiyan Cen; Ian M. Black; Parastoo Azadi; Russell W. Carlson; Gary Stacey; Yan Liang; Lipopolysaccharides Trigger Two Successive Bursts of Reactive Oxygen Species at Distinct Cellular Locations. *Plant Physiology* **2018**, *176*, 2543-2556, 10.1104/pp.17.01637.
- 25. Stefanie Ranf; Nicolas Gisch; Milena Schäffer; Tina Illig; Lore Westphal; Yuriy A Knirel; Patricia M. Sanchez Carballo; Ulrich Zähringer; Ralph Hückelhoven; Justin Lee; Dierk Scheel; A lectin S-domain receptor kinase mediates lipopolysaccharide sensing in Arabidopsis thaliana. *Nature Immunology* **2015**, *16*, 426-433, 10.1038/ni.3124.
- 26. Dana Zeidler; Ulrich Zähringer; Isak Gerber; Ian A. Dubery; Thomas Hartung; Wolf Bors; Peter Hutzler; Jörg Durner; From The Cover: Innate immunity in Arabidopsis thaliana: Lipopolysaccharides activate nitric oxide synthase (NOS) and induce defense genes. *Proceedings of the National Academy of Sciences* **2004**, *101*, 15811-15816, 10.1073/pnas.04045 36101.
- Isak B. Gerber; Dana Zeidler; Ian A. Dubery; Early perception responses of Nicotiana tabacum cells in response to lipopolysaccharides from Burkholderia cepacia. *Planta* 2004, *218*, 647-657, 1 0.1007/s00425-003-1142-0.

- Yoshitake Desaki; Ayako Miya; Balakrishnan Venkatesh; Shinji Tsuyumu; Hisakazu Yamane; Hanae Kaku; Eiichi Minami; Naoto Shibuya; Bacterial Lipopolysaccharides Induce Defense Responses Associated with Programmed Cell Death in Rice Cells. *Plant And Cell Physiology* 2006, 47, 1530-1540, 10.1093/pcp/pcl019.
- 29. Maeli Melotto; William Underwood; Jessica Koczan; Kinya Nomura; Sheng Yang He; Plant Stomata Function in Innate Immunity against Bacterial Invasion. *Cell* **2006**, *126*, 969-980, 10.101 6/j.cell.2006.06.054.
- 30. Bart Thomma; Kutschera A; Dawid C; Gisch N; Schmid C; Raasch L; Gerster T; Schäffer M; Smakowska-Luzan E; Belkhadir Y; Vlot Ac; Chandler Ce; Schellenberger R; Schwudke D; Ernst Rk; Dorey S; Hückelhoven R; Hofmann T; Ranf S; F1000Prime recommendation of Bacterial medium-chain 3-hydroxy fatty acid metabolites trigger immunity in Arabidopsis plants.. *F1000 -Post-publication peer review of the biomedical literature* **2019**, *364*, , 10.3410/f.735538698.79355 9797.
- 31. Cornelius S. Vilakazi; Ian A. Dubery; Lizelle Piater; Identification of lipopolysaccharide-interacting plasma membrane-type proteins in Arabidopsis thaliana. *Plant Physiology and Biochemistry* **2017**, *111*, 155-165, 10.1016/j.plaphy.2016.11.025.
- Nwaxigombe M. Baloyi; Ian A. Dubery; Lizelle Piater; Proteomic analysis of Arabidopsis plasma membranes reveals lipopolysaccharide-responsive changes. *Biochemical and Biophysical Research Communications* 2017, 486, 1137-1142, 10.1016/j.bbrc.2017.04.016.
- Sayaka Iizasa; Ei'Ichi Iizasa; Keiichi Watanabe; Yukio Nagano; Transcriptome analysis reveals key roles of AtLBR-2 in LPS-induced defense responses in plants.. *BMC Genomics* 2017, 18, 995, 10.1186/s12864-017-4372-4.
- 34. Yoshitake Desaki; Yusuke Kouzai; Yusuke Ninomiya; Ryosuke Iwase; Yumi Shimizu; Keito Seko; Antonio Molinaro; Eiichi Minami; Naoto Shibuya; Hanae Kaku; Yoko Nishizawa; OsCERK1 plays a crucial role in the lipopolysaccharide-induced immune response of rice. *New Phytologist* 2017, 217, 1042-1049, 10.1111/nph.14941.
- 35. Roland Willmann; Heini M. Lajunen; Gitte Erbs; Mari-Anne Newman; Dagmar Kolb; Kenichi Tsuda; Fumiaki Katagiri; Judith Fliegmann; Jean-Jacques Bono; Julie V. Cullimore; Anna K. Jehle; Friedrich Götz; Andreas Kulik; Antonio Molinaro; Volker Lipka; Andrea A Gust; Thorsten Nurnberger; Arabidopsis lysin-motif proteins LYM1 LYM3 CERK1 mediate bacterial peptidoglycan sensing and immunity to bacterial infection.. *Proceedings of the National Academy of Sciences* 2011, 108, 19824-9, 10.1073/pnas.1112862108.
- 36. Takeo Shimizu; Takuto Nakano; Daisuke Takamizawa; Yoshitake Desaki; Naoko Ishii-Minami; Yoko Nishizawa; Eiichi Minami; Kazunori Okada; Hisakazu Yamane; Hanae Kaku; Naoto Shibuya; Two LysM receptor molecules, CEBiP and OsCERK1, cooperatively regulate chitin elicitor signaling in rice. *The Plant Journal* **2010**, *64*, 204-214, 10.1111/j.1365-313X.2010.04324.x.

- 37. Yoshitake Desaki; Kana Miyata; Maruya Suzuki; Naoto Shibuya; Hanae Kaku; Plant immunity and symbiosis signaling mediated by LysM receptors. *Innate Immunity* **2017**, *24*, 92-100, 10.1177/175 3425917738885.
- 38. Bing Liu; Jian-Feng Li; Ying Ao; Jinwang Qu; Zhangqun Li; Jianbin Su; Yang Zhang; Jun Liu; Dongru Feng; Kangbiao Qi; Yanming He; Jinfa Wang; Hong-Bin Wang; Lysin motif-containing proteins LYP4 and LYP6 play dual roles in peptidoglycan and chitin perception in rice innate immunity.. *The Plant Cell* **2012**, *24*, 3406-19, 10.1105/tpc.112.102475.
- 39. Ying Ao; Zhangqun Li; Dongru Feng; Feng Xiong; Jun Liu; Jian-Feng Li; Menglong Wang; Jinfa Wang; Bing Liu; Hong-Bin Wang; OsCERK1 and OsRLCK176 play important roles in peptidoglycan and chitin signaling in rice innate immunity. *The Plant Journal* **2014**, *80*, 1072-1084, 10.1111/tpj.12710.
- 40. Xi Jiang; Han Bao; Hans Merzendorfer; Qing Yang; Immune Responses of Mammals and Plants to Chitin-Containing Pathogens.. *Single Molecule and Single Cell Sequencing* **2019**, *1142*, 61-81, 10.1007/978-981-13-7318-3\_4.
- 41. Wolfgang Knogge; Dierk Scheel; LysM receptors recognize friend and foe.. *Proceedings of the National Academy of Sciences* **2006**, *103*, 10829-30, 10.1073/pnas.0604601103.
- Kaku, H.; Nishizawa, Y.; Ishii-Minami, N.; Akimoto-Tomiyama, A.; Dohmae, N.; Takio, K.; Minami, E.; Shibuya, N. Plant cells recognize chitin fragments for defence signaling through a plasma membrane receptor. PNAS 2006, 103, 11086–11091.
- 43. Akira Yamada; N Shibuya; Osamu Kodama; Tadami Akatsuka; Induction of Phytoalexin Formation in Suspension-cultured Rice Cells byN-Acetyl-chitooligosaccharides. *Bioscience, Biotechnology, and Biochemistry* **1993**, *57*, 405-409, 10.1271/bbb.57.405.
- Miya, A.; Albert, P.; Shinya, T.; Desaki, Y.; Ichimura, K.; Shirasu, K.; Narusaka, Y.; Kawakami, N.; Kaku, H.; Shibuya, N; et al. CERK1, a LysM receptor kinase, is essential for chitin elicitor signaling in Arabidopsis. *PNAS* 2007, *104*, 1–6.
- 45. Tingting Liu; Zixu Liu; Chuanjun Song; Yunfei Hu; Zhifu Han; J. She; Fangfang Fan; Jiawei Wang; Changwen Jin; Junbiao Chang; Jian-Min Zhou; Jijie Chai; Chitin-Induced Dimerization Activates a Plant Immune Receptor. *Science* **2012**, *336*, 1160-1164, 10.1126/science.1218867.
- 46. Elena K. Petutschnig; Alexandra M.E. Jones; Liliya Serazetdinova; Ulrike Lipka; Volker Lipka; The Lysin Motif Receptor-like Kinase (LysM-RLK) CERK1 Is a Major Chitin-binding Protein in Arabidopsis thaliana and Subject to Chitin-induced Phosphorylation\*\*. *Journal of Biological Chemistry* 2010, 285, 28902-28911, 10.1074/jbc.M110.116657.
- Meritxell Antolín-Llovera; Martina Ried; Andreas Binder; Martin Parniske; Receptor Kinase Signaling Pathways in Plant-Microbe Interactions. *Annual Review of Phytopathology* 2012, 50, 451-473, 10.1146/annurev-phyto-081211-173002.

- 48. Tanaka, K.; Nguyen, C.T.; Liang, Y.; Cao, Y.; Stacey, G; Role of LysM receptors in chitin-triggered plant innate immunity. *Plant Signal. Behav.* **2013**, *8*, e22598.
- 49. Jinrong Wan; Xue-Cheng Zhang; David Neece; Katrina M. Ramonell; Steve Clough; Sung Yong Kim; Minviluz G. Stacey; Gary Stacey; A LysM Receptor-Like Kinase Plays a Critical Role in Chitin Signaling and Fungal Resistance in Arabidopsis[W][OA]. *The Plant Cell* **2008**, *20*, 471-481, 10.11 05/tpc.107.056754.
- 50. Jinrong Wan; Kiwamu Tanaka; Xue-Cheng Zhang; Geon Hui Son; Laurent Brechenmacher; Tran Hong Nha Nguyen; Gary Stacey; LYK4, a lysin motif receptor-like kinase, is important for chitin signaling and plant innate immunity in Arabidopsis.. *Plant Physiology* **2012**, *160*, 396-406, 10.110 4/pp.112.201699.
- 51. Yangrong Cao; Yan Liang; Kiwamu Tanaka; Cuong Nguyen; Robert P Jedrzejczak; Andrzej Joachimiak; Gary Stacey; The kinase LYK5 is a major chitin receptor in Arabidopsis and forms a chitin-induced complex with related kinase CERK1. *eLife* **2014**, *3*, 10.7554/eLife.03766.
- 52. Christine Faulkner; Elena Petutschnig; Yoselin Benitez Alfonso; Martina Beck; Silke Robatzek; Volker Lipka; Andrew J. Maule; LYM2-dependent chitin perception limits molecular flux via plasmodesmata. *Proceedings of the National Academy of Sciences* **2013**, *110*, 9166-9170, 10.10 73/pnas.1203458110.
- 53. Narusaka, Y.; Shinya, T.; Narusaka, M.; Motoyama, N.; Shimada, H.; Murakami, K; Presence of LYM2 dependent but CERK1 independent disease resistance in Arabidopsis. *Plant Signal. Behav.* 2013, 8, e25345.
- 54. Tomonori Shinya; Noriko Motoyama; Asahi Ikeda; Miyuki Wada; Kota Kamiya; Masahiro Hayafune; Hanae Kaku; Naoto Shibuya; Functional Characterization of CEBiP and CERK1 Homologs in Arabidopsis and Rice Reveals the Presence of Different Chitin Receptor Systems in Plants. *Plant And Cell Physiology* **2012**, 53, 1696-1706, 10.1093/pcp/pcs113.
- 55. Le, M.H.; Cao, Y.; Zhang, X.; Stacey, G; LIK1, a CERK1-interacting kinase, regulates plant immune responses in Arabidopsis. *PLoS ONE* **2019**, *9*, e102245.
- Jane Glazebrook; Contrasting Mechanisms of Defense Against Biotrophic and Necrotrophic Pathogens. *Annual Review of Phytopathology* 2005, *43*, 205-227, 10.1146/annurev.phyto.43.0402 04.135923.
- 57. Justin Ray; Xin Yang; Feng Kong; Tingwei Guo; Fengyan Deng; Steven Clough; Katrina Ramonell; A novel receptor-like kinase involved in fungal pathogen defence in Arabidopsis thaliana. *Journal of Phytopathology* **2018**, *166*, 506–515, 10.1111/jph.12711.
- 58. Jia Li; Jiangqi Wen; Kevin A Lease; Jason T Doke; Frans E. Tax; John Walker; BAK1, an Arabidopsis LRR Receptor-like Protein Kinase, Interacts with BRI1 and Modulates Brassinosteroid Signaling. *Cell* **2002**, *110*, 213-222, 10.1016/s0092-8674(02)00812-7.

- 59. Kyoung Hee Nam; Jianming Li; BRI1/BAK1, a Receptor Kinase Pair Mediating Brassinosteroid Signaling. *Cell* **2002**, *110*, 203-212, 10.1016/s0092-8674(02)00814-0.
- Milena Roux; Benjamin Schwessinger; Catherine Albrecht; Delphine Chinchilla; Alexandra M.E. Jones; Nick Holton; Frederikke Gro Mailnovsky; Mahmut Tor; Sacco De Vries; Cyril Zipfel; The Arabidopsis Leucine-Rich Repeat Receptor–Like Kinases BAK1/SERK3 and BKK1/SERK4 Are Required for Innate Immunity to Hemibiotrophic and Biotrophic Pathogens[W]. *The Plant Cell* 2011, 23, 2440-2455, 10.1105/tpc.111.084301.
- Dressano, K.; Ceciliato, P.H.O.; Silva, A.L.; Guerrero-Abad, J.C.; Bergonci, T.; Ortiz-Morea, F.A.; Bürger, M.; Silva-Filho, M.C.; Moura, D.S; BAK1 is involved in AtRALF1-induced inhibition of root cell. *PLoS Genet* **2017**, *13*, e1007053.
- 62. Yadong Sun; Lei Li; Alberto P. Macho; Zhifu Han; Zehan Hu; Cyril Zipfel; Jian-Min Zhou; Jijie Chai; Structural Basis for flg22-Induced Activation of the Arabidopsis FLS2-BAK1 Immune Complex. *Science* **2013**, *342*, 624-628, 10.1126/science.1243825.
- 63. Thomas W.H. Liebrand; Harrold A. Van Den Burg; Matthieu H. A. J. Joosten; Two for all: receptorassociated kinases SOBIR1 and BAK1. *Trends in Plant Science* **2014**, *19*, 123-132, 10.1016/j.tpla nts.2013.10.003.
- 64. Birgit Kemmerling; Anne Schwedt; Patricia Rodriguez; Sara Mazzotta; Markus Frank; Synan F. AbuQamar; Tesfaye Mengiste; Shigeyuki Betsuyaku; Jane E. Parker; Carsten Müssig; Bart P. H. J. Thomma; Catherine Albrecht; Sacco C. De Vries; Heribert Hirt; Thorsten Nürnberger; The BRI1-Associated Kinase 1, BAK1, Has a Brassinolide-Independent Role in Plant Cell-Death Control. *Current Biology* **2007**, *17*, 1116-1122, 10.1016/j.cub.2007.05.046.
- Kai He; Xiaoping Gou; Tong Yuan; Honghui Lin; Tadao Asami; Shigeo Yoshida; Scott D. Russell; Jia Li; BAK1 and BKK1 Regulate Brassinosteroid-Dependent Growth and Brassinosteroid-Independent Cell-Death Pathways. *Current Biology* 2007, *17*, 1109-1115, 10.1016/j.cub.2007.05. 036.
- 66. Junbo Du; Yang Gao; Yanyan Zhan; Shasha Zhang; Yujun Wu; Yao Xiao; Bo Zou; Kai He; Xiaoping Gou; Guojing Li; Honghui Lin; Jia Li; Nucleocytoplasmic trafficking is essential for BAK1and BKK1-mediated cell-death control. *The Plant Journal* **2016**, *85*, 520-531, 10.1111/tpj.13125.
- Dongping Lu; Shujing Wu; Xiquan Gao; Yulan Zhang; Libo Shan; Ping He; A receptor-like cytoplasmic kinase, BIK1, associates with a flagellin receptor complex to initiate plant innate immunity. *Proceedings of the National Academy of Sciences* 2009, 107, 496-501, 10.1073/pnas.0 909705107.
- 68. Jun Liu; Sufen Chen; Lijuan Chen; Qi Zhou; Menglong Wang; Dongru Feng; Jian-Feng Li; Jinfa Wang; Hong-Bin Wang; Bing Liu; BIK1 cooperates with BAK1 to regulate constitutive immunity and cell death in Arabidopsis. *Journal of Integrative Plant Biology* **2017**, *59*, 234-239, 10.1111/jip b.12529.

- 69. Thierry Halter; Julia Imkampe; Sara Mazzotta; Michael Wierzba; Sandra Postel; Christoph Bücherl; Christian Kiefer; Mark Stahl; Delphine Chinchilla; Xiaofeng Wang; Thorsten Nurnberger; Cyril Zipfel; Steven Clouse; Jan Willem Borst; Sjef Boeren; Sacco C. De Vries; Frans E. Tax; Birgit Kemmerling; The Leucine-Rich Repeat Receptor Kinase BIR2 Is a Negative Regulator of BAK1 in Plant Immunity. *Current Biology* **2014**, *24*, 134-143, 10.1016/j.cub.2013.11.047.
- 70. Paola Veronese; Hirofumi Nakagami; Burton Bluhm; Synan F. AbuQamar; Xi Chen; John Salmeron; Robert A. Dietrich; Heribert Hirt; Tesfaye Mengiste; The Membrane-Anchored BOTRYTIS-INDUCED KINASE1 Plays Distinct Roles in Arabidopsis Resistance to Necrotrophic and Biotrophic PathogensW<sup>II</sup>. *The Plant Cell* **2005**, *18*, 257-273, 10.1105/tpc.105.035576.
- 71. Tao Chen; Kai Bi; Zhangchao He; Zhixiao Gao; Ying Zhao; Yanping Fu; Jiasen Cheng; Jiatao Xie; Daohong Jiang; Arabidopsis Mutant bik1 Exhibits Strong Resistance to Plasmodiophora brassicae. *Frontiers in Physiology* **2016**, *7*, 1517, 10.3389/fphys.2016.00402.
- 72. Minghui Gao; Xia Wang; Ngmei Wang; Fang Xu; Xiaojun Ding; Zu-Wang Wang; Dongling Bi; Yu Ti Cheng; She Chen; Xin Li; Yuelin Zhang; Regulation of Cell Death and Innate Immunity by Two Receptor-like Kinases in Arabidopsis. *Cell Host & Microbe* **2009**, *6*, 34-44, 10.1016/j.chom.2009.0 5.019.
- 73. Isabell Albert; Hannah Böhm; Markus Albert; Christina E. Feiler; Julia Imkampe; Niklas Wallmeroth; Caterina Brancato; Tom Raaymakers; Stan Oome; Heqiao Zhang; Elżbieta Król; Christopher Grefen; Andrea A Gust; Jijie Chai; Rainer Hedrich; Guido Van Den Ackerveken; Thorsten Nurnberger; An RLP23–SOBIR1–BAK1 complex mediates NLP-triggered immunity. *Nature Plants* **2015**, *1*, 15140, 10.1038/nplants.2015.140.
- 74. Yanan Liu; Xingchuan Huang; Meng Li; Ping He; Yuelin Zhang; Loss-of-function of Arabidopsis receptor-like kinase BIR1 activates cell death and defense responses mediated by BAK1 and SOBIR1. *New Phytologist* **2016**, *212*, 637-645, 10.1111/nph.14072.
- 75. Zhang, W.; Fraiture, M.; Kolb, D.; Löffelhardt, B.; Desaki, Y.; Boutrot, F.F.G.; Tör, M.; Zipfel, C.; Gust, A.A.; Brunner, F; et al. Arabidopsis receptor-like protein 30 and receptor-like kinase suppresor of bir1-1/evershed mediate innate immunity to necrotrophic fungi. *Plant Cell* **2013**, *25*, 4227–4241.
- 76. Yu-Hung Yeh; Dario Panzeri; Yasuhiro Kadota; Yi-Chun Huang; Pin-Yao Huang; Chia-Nan Tao; Milena Roux; Hsiao-Chiao Chien; Tzu-Chuan Chin; Po-Wei Chu; Cyril Zipfel; Laurent Zimmerli; The Arabidopsis Malectin-Like/LRR-RLK IOS1 Is Critical for BAK1-Dependent and BAK1-Independent Pattern-Triggered Immunity. *The Plant Cell* **2016**, *28*, 1701-1721, 10.1105/tpc.16.00 313.
- 77. Martin Stegmann; Jacqueline Monaghan; Elwira Smakowska-Luzan; Hanna Rovenich; Anita Lehner; Nicholas Holton; Youssef Belkhadir; Cyril Zipfel; The receptor kinase FER is a RALF-

regulated scaffold controlling plant immune signaling. *Science* **2017**, *355*, 287-289, 10.1126/scien ce.aal2541.

- 78. Yu Xiao; Martin Stegmann; Zhifu Han; Thomas DeFalco; Katarzyna Parys; Li Xu; Youssef Belkhadir; Cyril Zipfel; Jijie Chai; Mechanisms of RALF peptide perception by a heterotypic receptor complex.. *Nature* **2019**, *572*, 270-274, 10.1038/s41586-019-1409-7.
- 79. Hongqing Guo; Trevor M. Nolan; Gaoyuan Song; Sanzhen Liu; Zhouli Xie; Jiani Chen; Patrick S Schnable; Justin W. Walley; Yanhai Yin; FERONIA Receptor Kinase Contributes to Plant Immunity by Suppressing Jasmonic Acid Signaling in Arabidopsis thaliana. *Current Biology* **2018**, *28*, 3316-3324.e6, 10.1016/j.cub.2018.07.078.

Retrieved from https://encyclopedia.pub/entry/history/show/8082