

Nematode Communities and Soil Health under Climate Change

Subjects: **Ecology**

Contributor: David Pires , Valeria Orlando , Raymond L. Collett , David Moreira , Sofia R. Costa , Maria L. Inácio

Soil health is intimately intertwined with ecosystem services. Climate change negatively impacts ecosystem functioning, by altering carbon and nitrogen biogeochemical cycles and shifting nutrient bioavailability, thus hampering food production and exacerbating biodiversity loss. Soil ecosystem services are provided by belowground biota, and as the most abundant metazoans on Earth, nematodes are key elements of soil food webs and reliable bioindicators of soil health.

abiotic stress

beneficial nematodes

ecosystem services

food webs

functional ecology

soil health

soil microfauna

1. Introduction

Soil is a complex system and a crucial component of sustainability ^[1]. Soil health is broadly defined as “the capacity of soil to function as a vital living system, within ecosystem and land-use boundaries, to sustain plant and animal productivity, maintain or enhance water and air quality, and promote plant and animal health” ^[2]. In other words, the concept highlights the ability of soil to perform important agricultural and ecological functions, including productivity, adaptability to management and inputs, and resilience against biotic and abiotic stressors. It must also exhibit robust resistance to degradation processes and the ability to rebound from disturbances due to its inherent resilience, defined by Holling as “a measure of the persistence of systems and of their ability to absorb change and disturbance and still maintain the same relationships between populations or state variables” ^{[3][4]}. Logically, soil health and ecosystem services are intimately intertwined. Ecosystem services encompass a wide range of benefits obtained from ecosystems, including (i) provisioning services (e.g., food and water), (ii) regulating services (e.g., natural disaster regulation, pest and pathogen control, and soil conservation), (iii) supporting services (e.g., nutrient cycling and pedogenesis), and (iv) cultural services (e.g., spiritual and recreational benefits) ^[5]. Soil is responsible for net primary production, it sustains plant and animal life, promotes water quality regulation, remediates pollution, intervenes in nutrient cycling, while providing physical stability and support ^{[6][7]}. Furthermore, it enhances the environment overall, by moderating climate at local, regional, and global scales ^[8]. In order to support sustainability, managing soil health must take into account that (i) improving multiple soil ecosystem services requires a multifunctional approach; (ii) enhancing one soil service can have favorable effects on some services but unfavorable outcomes on others; (iii) soil health management must ensure the long-term sustainability of soil services ^[9].

Climate change has a direct impact on the biological, chemical, and physical properties of soil, as it leads to shifts in temperature regimes and precipitation patterns [10][11]. Consequently, carbon (C), nitrogen (N), and hydrology cycles are likely to suffer the backlash [10], and the presence of multiple environmental stressors caused by human footprint were found to hamper soil ecosystem services across biomes [12]. Semiarid tropical regions of the world are particularly vulnerable, and with soils acting as important C reservoirs, a severe depletion of organic C will inevitably affect soil health [10]. Moreover, the decomposition of soil organic matter is thermosensitive [13][14], and climate change could lead to organic C-exhausted soils in response to altered C and N biogeochemical cycles and shifts in nutrient bioavailability, further exacerbating biodiversity loss [10].

Nematodes are part of the soil microfauna and represent approximately 80% of all multicellular animals of the terrestrial biosphere [6][15]. They are highly adaptable and successful animals, having colonized nearly all ecosystems on the planet [16][17][18]. According to their feeding habits, soil-dwelling nematodes can be assigned to one of five trophic groups: bacterivores, fungivores, herbivores, omnivores, and predators [19]. Nematode families and genera have been classified into a colonizer–persister scale (*c-p*), and given a rating from 1 to 5, indicative of their life strategy [20]. The *c-p* 1 group is made up of colonizers (*r*-strategists) like opportunistic bacterial feeders that rapidly increase in numbers under favorable conditions, exhibiting a short life cycle, high colonization ability, and tolerance to disturbance [20]. On the other hand, the *c-p* 5 group consists of persisters (*K*-strategists), such as some herbivores, omnivores, and predators, with a low reproduction rate, long life cycle, low colonization ability, and high sensitivity to disturbance [20]. Due to their rapid and taxon-specific response to environmental changes, nematodes are valuable bioindicators [21][22]. Nematofauna diversity is largely influenced by factors such as soil texture, soil moisture, and food availability [21]. However, the response of nematodes to environmental stress varies among trophic groups, with those having shorter generation times and/or high fecundity showing a positive response, while those with longer generation times and/or lower fecundity being more sensitive [23][24]. Free-living nematodes (bacterivores, fungivores, omnivores, and predators), widely referred to as beneficial nematodes, outnumber herbivores in terms of abundance and diversity, and they play critical roles in ecosystem functioning, occupying key ecological niches in belowground food webs, and are involved in C sequestration, energy transfer, and nutrient mineralization, increasing their availability to plants and, thus, improving soil fertility [25][26][27][28]. Soil nematodes directly or indirectly contribute to (i) human well-being, by driving key processes to food production; (ii) climate regulation, by intervening in the short and long-term fluxes and flows of C in and out of soils; and (iii) support terrestrial life and diversity, through processes like decomposition, nutrient cycling, and regulation of pests and pathogens (**Figure 1**) [29]. Healthy soils typically have a high abundance and diversity of free-living nematodes in complex food webs with long chains and feedback loops, and a low proportion of herbivores [23][30][31]. Indeed, a nematode community analysis can provide invaluable information on the status of soils: (i) a high ratio of bacterial- to fungal-feeding nematodes indicates that organic matter is predominantly decomposed by bacteria and that rapid nutrient cycling is occurring; (ii) a predominance of fungivores indicates that nutrient cycling is relatively slow, as the decomposition channel is dominated by fungi; (iii) low population densities of omnivores and predators suggest disturbance, such as excessive fertilizer inputs, tillage, or the presence of pollutants; (iv) high numbers of omnivores and predators indicate that the system is biologically complex and resilient, and has some natural ability to suppress plant-parasitic nematodes and other soil-borne pathogens [32]. In agricultural systems, plant-parasitic

nematodes are problematic, but in a broader ecological context, they are fundamental in shaping aboveground vegetation communities and contributors to plant performance in natural ecosystems by plant–soil feedbacks [33][34]. Upon feeding on their hosts, herbivorous nematodes alter root exudation patterns, and indirectly modify the rhizobiome (the microbial diversity attached to and influenced by roots), thus curbing nutrient availability to plants, especially of N and phosphorous (P), and contributing to plant community dynamics [30][35][36]. On the other hand, while the contributions of beneficial nematodes to plant performance remain largely unknown, they can have positive effects on plants by stimulating microbe-induced C sequestration, and keeping pests and pathogens at bay [37]. Nematodes are aquatic animals that require water to move, feed, and reproduce, and climate extremes are anticipated to shift the structure of nematode communities and their roles in ecosystems [38][39][40]. However, the impacts of climate change on nematode abundance and functional groups have not been consensual, displaying significant variation across different studies [41][42][43][44][45].




Figure 1. Illustrative representation of the main ecosystem services provided by soil-dwelling nematodes.

2. Nematode Community Dynamics under Climate Change

A demographic explosion is driving unprecedented food demand and pushing natural ecosystems to fragmentation. The effects of anthropogenic activities in natural systems are known to reduce aboveground biodiversity, hindering ecosystem services and, consequently, their contributions to human well-being [46], but data of such impacts on belowground taxa are scarce, perhaps because they are difficult to assess. Nevertheless, nematode community

structure is a fast and reliable predictor of ecological disturbance, giving us insights into the status of the whole system.

2.1. Temperature

The soil nematode community structure seems to be particularly susceptible to warming, favoring some groups in detriment of others, but these negative effects can be limited by aboveground vegetation that counters them to some extent. Both parameters also affect vegetation, with an anticipated indirect effect on nematode communities [47], although these confounding effects are not assessed here. The main effects of increasing temperature on nematode communities are summarized in **Table 1**.

Table 1. Effect of increasing temperature on soil nematode communities.

Factor	Effects on Soil Nematodes	Reference
High temperature	Reduction in nematode diversity and genera richness	[48]
	Reduction in herbivores, with a short-term decrease in bacterivores and fungivores who recover over time, and relative tolerance of omnivores–predators	[49]

2.2. Water Stress

Drought is a key limitation of soil nematode abundance, even in the short term, and has persistent effects. Nevertheless, the soil nematode community structure recovers gradually as precipitation increases, even though these positive effects are not straightforward, and water uptake by plants can further stress an already fragile system. The main effects of water stress on nematode communities are summarized in **Table 2**.

1. Brevik, E.C.; Cerda, A.; Mataix-Solera, J.; Pereg, L.; Quinlan, J.N.; Six, J.; Van Oost, K. The Interdisciplinary Nature of SOIL. SOIL 2015, 1, 117–129.

2. Doran, J.W.; Zeiss, M.R. Soil Health and Sustainability: Managing the Biotic Component of Soil Quality. Appl. Soil. Ecol. 2000, 15, 3–11.

Factor(s)	Effects on Soil Nematodes	Reference
Long-term increase in mean annual precipitation	Decline in total nematode abundance due to predation pressure, and increase in nematode abundance	[50]
	High proportion of plant-parasitic nematodes	[51]
	Decrease in free-living nematode diversity and evenness	[52]

Singh, 24. 4, 1–23. g.

Factor(s)	Effects on Soil Nematodes	Reference
Water scarcity	Reduction in total nematode abundance, and decrease in relative abundance of bacterivores, fungivores, and predators; relative abundance of herbivores unaffected	[53]
	Lower bacterivore and fungivore abundance, marginal increase in omnivores and decrease in fungivores, higher maturity and structure index, and lower prey:predator ratio	[54]

Change. In *Climate Change and Soil Interactions*; Prasad, M.N.V., Pietrzykowski, M., Eds.; Elsevier: Amsterdam, The Netherlands, 2020; pp. 751–767. ISBN 978-0-12-818032-7.

2.3. Land Use

11. Tripathi, A.; Pandey, V.; Ranjan, M.R. Climate Change and Its Impact on Soil Properties. In *Climate Change and the Microbiome*; Choudhary, D.K., Mishra, A., Varma, A., Eds.; Springer: Cham, Switzerland, 2021; Volume 63, pp. 139–153. ISBN 978-3-030-76863-8.

12. Rilling, M.C.; van der Heijden, M.G.A.; Berghuis, M.; Jous, Y.B.; Riedo, J.; Sanz-lazaro, A.; Moreno Jimenez, E.; Romero, E.; Tedersoo, L.; Delgado-Baquerizo, M. Increasing the Number of Stressors Reduces Soil Ecosystem Services Worldwide. *Nat. Clim. Change* 2023, 13, 478–483.

13. Conant, R.T.; Ryan, M.G.; Agren, G.I.; Birge, H.E.; Davidson, E.A.; Eliasson, P.E.; Evans, S.E.; Frey, S.D.; Giardina, C.P.; Hopkins, F.M.; et al. Temperature and Soil Organic Matter Decomposition Rates - Synthesis of Current Knowledge and a Way Forward. *Glob. Chang. Biol.* 2011, 17, 3392–3404.

Table 3. Effects of different land uses on soil nematode communities.

Factor(s)	Effects on Soil Nematodes	Reference
Intensive land use	Increase in bacterivores, fungivores, and herbivores	[55]
Soil type	Lower biodiversity and C/N ratios in cultivated soils, resulting in a reduction in nematode abundance and diversity; increase in abundance of trophic groups, total abundance, and diversity of nematodes in soils with higher pH and C and N contents	[56]
Land use conversion	Negative impacts on soil taxa across a large spatial scale, but agronomic practices limit the climatic constraints on belowground biodiversity	[57]

19. Yeates, G.W.; Bongers, T.; De Goede, R.G.; Freckman, D.W.; Georgieva, S.S. Feeding Habits in Soil Nematode Families and Genera-an Outline for Soil Ecologists. *J. Nematol.* 1993, 25, 315–

Factor(s)	Effects on Soil Nematodes	Reference
Monoculture	Increase in fungivores, and decrease in bacterivores, herbivores, and omnivores–predators	[58]
Monoculture, tillage, and manure	Increase in bacterivores and fungivores, and decrease in omnivores–predators	[59]
Vegetation succession	Positive effect on nematode abundance, diversity, complexity of community structure, and diversity-weighted abundance	[60]
Livestock grazing	Decline in bacterivores, herbivores, and omnivores–predators abundance, lower total nematode abundance, and no detrimental effect on fungivore abundance	[61]
Land degradation	Lower nematode trophic diversity	[62]
Intensive land use	Generalists equally abundant in all ecosystems, with specialists dominating agricultural landscapes and less abundant in low disturbed soils; highest richness and diversity in grasslands and dairy farms, with low abundances in shrubland–woodland habitats	[63]
Plant resource-use strategies	Plants with acquisitive strategies promoted nematode abundance, but fewer opportunistic nematodes in the rhizosphere of acquisitive plants compared to conservative plants	[64]
Agricultural practices	(i) Conventional practices decrease abundance, trophic structure, and taxonomic richness of nematode communities; (ii) agroecological practices enhance the functional and taxonomic diversity of nematodes: total nematode abundance and absolute abundance of fungivores, herbivores, and omnivores–predators; reduction in herbivore abundance in crop rotation; increase in omnivores–predators in cover crops; increase in bacterivores and fungivores in organic fertilization; reduction in nematode abundance and food web structure in monoculture and pesticide application, while copiotrophic nematodes are favored	[65]

30. Pineda, F., Sánchez, C., Morán, C., Gómez, A., Pacheco, C., Barón, C., & Robard, A.; Oktaba, L.; Filser, J.; Kindlmann, P.; et al. Global Distribution of Soil Fauna Functional Groups and Their Estimated Litter Consumption across Biomes. Sci. Rep. 2022, 12, 17362.

Factor(s)	Effects on Soil Nematodes	References
Spatial distance and environmental filtering	Plant type altered β -diversity of nematodes; spatial turnover was the primary process driving total β -diversity of the nematode community	[66]
Vegetation restoration with varying degrees of degradation	Strong effects on soil nematodes observed under low degradation	[67]
Crop-tree thinning	Increase in abundance of soil nematodes, along with the relative abundance of herbivores in all systems; increase in proportion of stress-tolerant enrichment and general opportunists	[68]
Harvest frequency and legume density	Legume addition and density were drivers of total nematode abundance, especially bacterivores, while improving metabolic activities of total nematodes, bacterivores, and omnivores–predators; positive effects of legume addition subsided after increased harvesting frequency	[69]
Agricultural practices	Plant diversity enhanced the activity of beneficial nematodes; positive effects of nematodes on plant growth and function associated with higher values in soil pH and cation contents	[70]

44. Blankinship, J.C.; Nieklaus, P.A.; Hungate, B.A. A Meta-Analysis of Responses of Soil Biota to Global Change. *Oecologia* 2011, 165, 553–565.

45. Cesarz, S.; Reich, P.B.; Scheu, S.; Ruess, L.; Schaefer, M.; Eisenhauer, N. Nematode Functional Guilds, Not Trophic Groups, Reflect Shifts in Soil Food Webs and Processes in Response to Interacting Global Change Factors. *Pedobiologia* 2015, 58, 23–32.

2.4. Nutrient Enrichment

The structure of nematode communities can be adversely affected by mineral fertilization, resulting in simplified soil food webs with fewer omnivores–predators and more herbivores and bacterivores, impairing ecosystem functions.

46. Wall, D.H.; Nielsen, U.N.; Six, J. Soil Biodiversity and Human Health. *Nature* 2015, 528, 69–76.

47. De Deyn, G.B.; Raaijmakers, C.E.; van Ruijven, J.; Berendse, F.; van der Putten, W.H. Plant Species Identity and Diversity Effects on Different Trophic Levels of Nematodes in the Soil Food Web. *Oikos* 2004, 106, 576–586.

On the other hand, organic amendments can increase nematode diversity and promote a higher energy transfer in the system. Although the above studies mainly report on crop nutrition management, they are informative of the nematode community response to nutrient quantities and balance in soils. Global change has been promoting the deposition of N and other nutrients to both agricultural and natural ecosystems, mostly accompanying urbanization.

48. Wang, J.; Li, W.; Zhang, X.; Li, X.; Li, P.; Shi, X.; Wu, H.; Pan, C. Change in Soil Nematode Abundance and Composition under Elevated and Canopy Warming in a Paddy Field (Plant) due to climate effects. *2019*, *445*, 425–437. effects of nutrient enrichment on nematode communities are summarized in **Table 4**.

49. Liu, Y.; Wang, W.; Liu, P.; Zhou, H.; Chen, Z.; Suonan, J. Plant-Soil Mediated Effects of Long-Term Warming on Soil Nematodes of Alpine Meadows on the Qinghai–Tibetan Plateau. *Biology* **2022**, *11*, 1596. **Table 4.** Effects of nutrient enrichment on soil nematode communities.

Factor(s)	Effects on Soil Nematodes	Reference
Long-term N enrichment	Plant removal dwindled nematode taxon richness and abundance of bacterivores and herbivores; the abundance of fungivores and omnivores–predators increased under the same conditions	[71]
Long-term N fertilization	Greater nematode abundance in fertilized plots, while richness, diversity, and ecological maturity were lower; enriched food web mostly driven by bacterivores and herbivores, with persisting effects overtime	[72]
High N deposition	Decrease in most nematode trophic groups and community diversity under understory addition of N compared to canopy addition of N	[73]
Short-term N and P enrichment under soil acidification	Nematode variables, including community structure, were largely unaffected by short-term nutrient enrichment under soil acidification	[74]
Long-term organic amendments and mineral fertilization	Positive effect on the abundance of most functional guilds by organic amendments, which enhanced the energy transfer among nematode communities, while increasing the relative allocation of energy flux to bacterivores and fungivores and decreasing the relative allocation to herbivores	[75]
Liming, P, and zinc inputs	P input significantly increased nematode diversity and genera; bacterivores and herbivores were the most abundant trophic groups, and predators the least	[76]

59. Yogaswara, D.A.; Kasmara, H.; Hermawan, W. Using Nematode Community to Evaluate Banana Soil Food Web in Mekargalih, Cianjur, West Java. *Pertanika J. Trop. Agric. Sci.* **2021**, *44*, 465–

Factor(s)	Effects on Soil Nematodes	Reference
	common; nematode biodiversity was unaffected by liming, and nematode diversity and maturity were reduced in the absence of liming	

and Decouples Their Relationships with Ecosystem Functions in Grasslands. Glob. Chang. Biol. 2020, 26, 960–970.

2.5. Combined Stressors

62. Han, X.; Li, Y.; Du, X.; Li, Y.; Wang, Z.; Jiang, S.; Li, O. Effect of Grassland Degradation on Soil In general, the greater the number of disturbances, whether directly or indirectly caused by human activities, the Quality and Soil Biotic Community in a Semi-Arid Temperate Steppe. Ecol. Process. 2020, 9, 63. longer the recovery time for the system and the greater the impact on soil nematode communities, which compared to their benefits to humankind, although manipulative studies and experimental research has been elucidating the impacts of each independent community, observational studies and complex systems suggest strong interactive effects among them. It is therefore anticipated that a large sampling effort at appropriate scales, combined with an in-depth local and remote-sensing characterization of sampled ecosystems can provide key knowledge that can be translated into management and policies for climate change adaptation and mitigation. The Root Traits Mediate Functional Guilds of Soil Nematodes in an Ex-Arable Field. Soil. Biol. Biochem. 2020, 151, 108038.

63. Vazquez, C.; Goede, R.G.M.; Korthals, G.W.; Rutgers, M.; Schouten, A.J.; Creamer, R. The combined effect of multiple stressors on nematode communities is summarized in **Table 5**. Biochem. 2020, 151, 108038.

64. Zhang, C.; Wang, J.; Ren, Z.; Hu, Z.; Tian, S.; Fan, W.; Chen, X.; Griffiths, B.S.; Hu, F.; Liu, M. Root Traits Mediate Functional Guilds of Soil Nematodes in an Ex-Arable Field. Soil. Biol. Biochem. 2020, 151, 108038.

65. Puissant, J.; Villenave, C.; Chauvin, C.; Plassard, C.; Blanchard, E. Trap 1 Quantification of the Global Impact of Agricultural Practices on Soil Nematodes: A Meta-Analysis. Soil. Biol. Biochem. 2020, 151, 108038.

Factors	Effects on Soil Nematodes	Reference
Soil moisture, P addition, and aboveground vegetation	Plant type and water availability had a greater impact on nematode abundance and community composition; drought was detrimental to the total density of nematodes and functional guilds; bacterivores, herbivores, and omnivores were significantly more abundant in soils with legumes	[77]
Biotic (microbial biomass and competition) and abiotic variables (moisture, salinity, and elevation)	Spatial segregation between two competing bacterivore species, with contrasting responses to abiotic factors: one best adapted to high salinity, lower temperatures, and low moisture environments, while the other thrives at higher temperatures, higher soil moisture, and lower salinity	[78]

Technique. Sustainability 2021, 13, 7181.

7	Factors	Effects on Soil Nematodes	Reference	Of
7	Climatic, soil, and historical factors	Current factors, particularly climate, are more influential than historical factors in shaping nematode diversity patterns on a broader scale	[79]	Shi-arid
7	Liming treatments	Interacting nematode and microbial communities minimally impacted by liming, with an increase in omnivores and predators, who keep bacterivores under control; stronger interaction in the presence of an abundant microbial community	[80]	Shi-arid
7	Increasing aridity across a large spatial scale	Decline in total and relative nematode abundance of each functional guild under increasing aridity; taxonomic richness of total nematode community and functional guilds decreased under moisture scarcity; at the dry end of the aridity gradient, richness of bacterivores was higher, while herbivores declined steadily; richness of fungivores and omnivores–predators remained relatively stable up to a certain point, before dropping steeply	[81]	Shi-arid
7	Drought and fertilization	Drought favored bacterivores and fungivores, and likely had detrimental effects on higher trophic levels; fertilization caused a prominent increase in bacterivores and an equally significant drop in fungivores	[82]	Shi-arid
7	Elevated CO ₂ and N, warming, and drought	Increase in nematode density at elevated N and ambient CO ₂ , and ambient N and elevated CO ₂	[83]	Shi-arid
8	Warming and precipitation	Decrease in nematode abundance, especially of bacterivores and herbivores (with minor effects on fungivores), under artificial warming, but the nematode community diversity and functions remained stable; decrease in nematode abundance, especially of bacterivores and omnivores–predators, under reduced precipitation, with fungivores and herbivores relatively insensitive to water stress; increase in nematode abundance and community diversity with water availability	[84]	Shi-arid

82. Siebert, J.; Sünneemann, M.; Auge, H.; Berger, S.; Cesarz, S.; Ciobanu, M.; Guerrero-Ramírez, N.R.; Eisenhauer, N. The Effects of Drought and Nutrient Addition on Soil Organisms Vary across Taxonomic Groups, but Are Constant across Seasons. *Sci. Rep.* 2019, 9, 639.

Factors	Effects on Soil Nematodes	Reference
Ecological and edaphic factors	Reduced nematode abundance and diversity with increasing altitude, with bacterivores consistently the dominant group; nematode diversity was mostly influenced by temperature and moisture; decrease in nematode abundance with increasing soil acidity; nematode diversity and richness were directly proportional to nutrient (N and P) levels	[85]
N deposition under reduced water availability	Reduced nematode abundance and diversity under N addition and reduced water input; synergistic effects of N addition and reduced water input on soil nematode communities at higher trophic levels; sole addition of N was more detrimental to the nematode community	[86]
Returning agricultural residues	Nematode diversity was lower in treatments with conventional chemical NPK fertilizers; positive correlation between omnivore–predator abundance and ecosystem multifunctionality and soil fertility	[87]

1.; Rich, 184–
atode
, China.
Wijaya,
des.
nd
rest.
nd

88. Ferris, H. Form and Function: Metabolic Footprints of Nematodes in the Soil Food Web. Eur. J. Soil. Biol. 2010, 46, 97–104.

89. Zhang, X.; Ferris, H.; Mitchell, J.; Liang, W. Ecosystem Services of the Soil Food Web after Long-Term Application of Agricultural Management Practices. Soil. Biol. Biochem. 2017, 111, 36–43.

3. Nematode Contributions to Soil Health

The adaptability of nematodes, as evidenced by frequent shifts in habitat transition, along with their ability to withstand environmental changes, has significant implications for soil functions on a global scale. Indeed, the ability of nematode functional guilds to perform ecosystem services depends on the soil condition [25] that can be measured according to their metabolic footprints [88]. However, land use (including soil properties), agricultural management practices, and climate all contribute to shaping nematode communities and, thus, affect nematode contributions to soil health through modification of their abundance, functional groups, and metabolic footprints, leading to shifts in ecosystem processes and services [89][90][91].

90. Wood, J.R.; Holdaway, R.J.; Orwin, K.H.; Morse, C.; Bonner, K.I.; Davis, C.; Boisthede, N.; Dickie, I.A. No Single Driver of Biodiversity. Divergent Responses of Multiple Taxa across Land Use Types. Ecosphere 2017, 8, e01997.

91. Bender, S.F.; Wagg, C.; van der Heijden, M.G.A. An Underground Revolution: Biodiversity and Soil Ecological Engineering for Agricultural Sustainability. Trends Ecol. Evol. 2016, 31, 440–452.

92. Adams, B.J.; Arif, W.; Farris, P.; Barden, R.D.; et al. A Global Database of Soil Nematode Abundance and Functional Group Composition. Sci. Data 2020, 7, 108.

93. Karuri, H. Nematode Community Structure and Functional Guilds Differ in Tea Fields and Tropical Forest. Geoderma 2021, 392, 115006.

94. Jiang, Y.; Wang, Z.; Liu, H.; Han, Y.; Wang, Y.; Wang, Q.; Liu, T. Nematodes and Their Bacterial Prey Improve Phosphorus Acquisition by Wheat. New Phytol. 2023, 237, 974–986.

95. Zheng, J.; Dini-Andreote, F.; Luan, L.; Gelsen, S.; Xue, J.; Li, H.; Sun, B.; Jiang, Y. Nematode Predation and Competitive Interactions Affect Litter Decomposition, nutrient cycling, and plant nutrient uptake. Indeed, understanding the susceptibilities and ecological preferences of soil nematodes can help predict changes

- in ecosystems services. To determine which factors best explain a soil health trait, Martin et al. [31] explored the incorporation of nematode communities in the assessment of belowground biota by using an exploratory factor analysis of permanganate oxidizable C, soil protein, and mineralizable C based on data from two longstanding Resistance and Resilience under Copper or Heat Stress. *Soil. Ecol. Lett.* 2020, 2, 220–229.
96. Chen, X.; Xue, W.; Xue, J.; Griffiths, B.S.; Liu, M. Contribution of Bacterivorous Nematodes to Soil analysis of permanganate oxidizable C, soil protein, and mineralizable C based on data from two longstanding experiments across many different fields. Fungivores were found to be highly integrated with soil biological health indicators of permanganate oxidizable C and acid phosphatase, suggesting that this functional guild may be essential for supplying essential ecosystem services, especially C cycling and P mineralization. To determine the Climate Variables Influence the Nematode Communities in the Gaotinga Dry Forest. *Appl. Soil. Ecol.* 2020, 150, 103474.
97. da Silva, J.V.C.d.; Hirschfeld, M.N.C.; Cares, J.F.; Esteves, A.M. Land Use, Soil Properties and Climate Variables Influence the Nematode Communities in the Gaotinga Dry Forest. *Appl. Soil. Ecol.* 2020, 150, 103474.
98. Bastida, F.; Eldridge, D.J.; Abades, S.; Alfaro, P.D.; Gallardo, A.; García-Velázquez, L.; García, C.; Han, S.C.; Pérez, C.A.; Santos, F.; et al. Climatic Vulnerabilities and Ecological Preferences of Soil Invertebrates across Biomes. *Mol. Ecol.* 2020, 29, 752–761.
99. Majdi, N.; Trautspurger, W.; Fueser, H.; Gansfort, B.; Laffaille, P.; Maire, A. Effects of a Broad Range of Experimental Temperatures on the Population Growth and Body-Size of Five Species of Free-Living Nematodes. *J. Biol.* 2019, 80, 21–36.
100. Andreazzi, W.S.; Pisco, A.F.; C. W. Ankomplex, K.E.; Chua, S.; de Tommaso, C.; M. Guan, P.; C. S. Chai, P.; A. S. Salas, C. E. M. Wad, D. H. Light, B. Size, Structure of Soil Fauna along Geographic and Temporal Gradients of Precipitation in Grasslands. *Soil Biol. Biochem.* 2020, 140, 107638.
101. Wilschut, R.A.; Geisen, S.; Martens, H.; Kostenko, O.; Hollander, M.; Hooven, F.C.; Weser, C.; Snoek, L.B.; Bloem, J.; Cakovic, D.; et al. Latitudinal Variation in Soil Nematode Communities microbiome structure, but this enhancement varied according to nematode feeding behavior: nematodes that had under Climate Warming-related Range-expanding and Native Plants. *Glob. Chang. Biol.* 2019, 25, 2714–2726.
102. Wang, X.; Xiao, S.; Yang, X.; Liu, Z.; Zhou, X.; Du, G.; Zhang, L.; Guo, A.; Chen, S.; Nielsen, N.P. Dominant Plant Species Influence Nematode Richness by Moderating Understory Diversity and bioavailability, by interacting with phosphate-solubilizing bacteria, thus enhancing plant performance and providing key ecosystem services. *Soil Biol. Biochem.* 2019, 137, 107566.
103. Bennett, J.A.; Koch, A.M.; Forsythe, J.; Johnson, N.C.; Tilman, D.; Klironomos, J. Resistance of Soil Biota and Plant Growth to Disturbance Increases with Plant Diversity. *Ecol. Lett.* 2020, 23, 119–128.
104. Wu, L.; Chen, H.; Chen, D.; Wang, S.; Wu, Y.; Wang, B.; Liu, S.; Yue, L.; Yu, J.; Bai, Y. Soil Biota Diversity and Plant Diversity Both Contributed to Ecosystem Stability in Grasslands. *Ecol. Lett.* 2023, 26, 858–868.

Resistance and diversity in soil biota. <https://encyclopedia.pub/entry/show/47443>

Resilient and diverse soil biota contribute to efficient decomposition, leading to more nutrient mineralization that readily become available to plants, thus reducing the need for fertilizer application. Moreover, a natural regulation of pests and diseases performed by soil nematodes will reduce our dependence on pesticides. Ultimately, with reduced farmer intervention and corresponding anthropogenic impact, systems will be less disturbed and soils healthier.

4. How Nematodes Promote Soil Resilience

As highly adaptable animals, with diverse roles in ecosystem functioning, the physiological and life history traits of nematodes make them less susceptible to environmental changes compared to larger fauna higher-up in the food web. Indeed, these characteristics could prove useful for the resistance and resilience of soils to natural and anthropogenic changes.

To better comprehend the role of bacterivores in maintaining functional stability of ecosystems under disturbance, Chen et al. [96] studied their contributions to promoting soil resistance and resilience under copper and heat stress. The relative shifts in two dominant bacterivore genera, *Acrobeloides* and *Protorhabditis*, responded differently to disturbance. *Protorhabditis* exhibited greater resistance and resilience to copper stress compared to *Acrobeloides*, while both genera displayed higher resilience only by the end of the experiment under heat stress. Indeed, bacterivores showed a positive effect on soil resilience under thermal stress starting at 28 days. The increase in relative abundance of bacterivores did not significantly affect soil resistance in terms of microbiota but it improved soil resilience to copper stress. The differences in responses of soil function to disturbance highlight the role of bacteria-feeding nematodes in promoting ecosystem stability under stress. To determine the effects of soil properties, rainfall, and temperature on soil nematodes, da Silva et al. [97] analyzed the changes in nematode community structure under contrasting types of land use in a seasonally dry tropical forest in Brazil. Nematofauna composition in the secondary forest differed in abundance and richness compared to agricultural systems, being expectedly higher in the former and lower in the latter, with bacterivores and omnivore–predators more susceptible to the type of land use. The variation in taxonomic composition among the studied sites was strongly related to soil properties, monthly mean rainfall, and temperature, which accounted for 65.42% of the total variation. These results further indicate that anthropogenic activities, expressed by the conversion of native vegetation to cropping systems, which modify soil characteristics, as well as climate variables, negatively affect the structure and composition of nematode communities. Nevertheless, changes in nematode community composition and structure can be reversed by allowing the fields to undergo secondary forest regeneration after abandonment. Seeking to identify the major ecological predictors of soil invertebrate diversity, Bastida et al. [98] surveyed 83 locations in six continents, from polar to arid climates, to study three soil invertebrates: nematodes, arachnids, and rotifers. Different ecosystem types such as forest, grasslands, and shrublands were included in the survey and nematodes were the most abundant, accounting for 43% of all taxa surveyed. Aridity was detrimental to the diversity of nematodes, whereas forest, plant richness, and annual net primary productivity were positively correlated. These findings exposed potential vulnerabilities of soil invertebrates to climate change in locations where hotter temperatures may occur in the future. Moreover, deforestation processes and increase in aridity may reduce nematode diversity, providing evidence of the importance of vegetation and climate for the diversity of soil invertebrates. Considering an increasing likelihood of extreme climatic events, Majdi et al. [99] exposed five species of free-living bacterivorous nematodes to a wide range of temperatures under controlled conditions, and their population growth rates and body-size distributions were measured. Body size at maturity was inversely proportional to temperature, mature females were laying a smaller number of eggs at higher temperatures, and a prevalence of early juvenile stages resulted in reduced body-mass structure with increasing temperature. Additionally, closely related species like *Plectus acuminatus* and *P. cf. velox* had very different thermal tolerance ranges, with the population growth of most tested species declining between 25 and 30 °C, and *A. nanus* exhibiting

the broadest thermal tolerance range. This study demonstrated how thermic stress can induce changes in the growth and size–structure of bacterivores. To investigate the community-weighted mean body mass of soil nematodes, Andriuzzi et al. [100] studied the role of water availability in the body size of these invertebrates across a gradient of precipitation in North American grasslands, ranging from arid to semiarid and mesic conditions. An increase in nematode community-weighted mean mass from arid to mesic conditions was observed, but no effects were reported at the arid site. When grouping community-weighted mean mass by feeding habits, only plant-parasitic nematodes showed a positive response to water input in semiarid and mesic conditions. This suggests that aridity acts as a buffer against large-bodied nematodes, limiting community body size shifts in response to extreme events, either drought or rainfall. Aiming to study the latitudinal variation in soil nematode communities under climate warming-related range-expanding and native plants, Wilschut et al. [101] showed that the composition of soil nematode communities changes across a latitudinal gradient, but not their richness or abundance, with plant species identity (both range-expanding and native plant species) being the strongest predictor of this shift. These findings further indicate that this variation is less dependent on soil characteristics, such as pH and soil moisture. In addition, plant species that expand their range due to climate warming may experience advantages by being free from nematode herbivory in their new habitat. A plant removal experiment was set up by Wang et al. [102] to better understand how dominant vegetation changes impact nematode assemblages. Edaphic properties, especially soil C and N content, were the primary drivers of nematode community structure and community-weighted mean biomass, with no observable short-term effects resulting from vegetation removal. However, long-term effects on nematode assemblages are expectable due to nutrient flow mediated by shifts in vegetation composition. To characterize and explore the relationship between soil biota and plant diversity and productivity, Bennett et al. [103] carried out a long-term experiment. Plant species richness had a positive effect on fungi, including increased arbuscular mycorrhizal fungi, while reducing plant-parasitic nematodes. Overall, soil biota resistance to disturbance increased with plant diversity, highlighting the importance of plant species richness for belowground communities. To evaluate the impacts of various measures of trophic diversity, climate, and soil environmental factors across three spatial scales, Wu et al. [104] conducted a field survey on the stability of ecosystems on the Mongolian Plateau. Soil biota diversity, including α - and β -diversity, positively contributed to ecosystem stability, with soil nematode diversity and trophic groups associated with higher ecosystem stability. The relatively low abundance of herbivores may have contributed to enhanced plant performance by increasing root exudation, which stimulated microbial activity and nutrient availability. The positive association of soil biota diversity with ecosystem stability was similar to that of plant diversity in some cases. Similarly, an increase in the abundance of higher trophic levels such as omnivores–predators and microbial-feeding nematodes may have resulted in improved nutrient transfer to plants, leading to enhanced plant productivity and maintaining ecosystem stability through top–down effects.

Severe anthropogenic impacts often lead to simplified soil food webs, with limited top–down control by omnivores–predators, ultimately compromising ecosystem functioning and impairing their natural ability to mitigate the effects of climate change. It is therefore crucial to restore the complexity of soil food webs to enhance soil resilience to climate change.