Uranyl Carbonate Minerals

Subjects: Materials Science, Characterization & Testing Contributor: Vladislav Gurzhiy

Uranyl carbonates are one of the largest groups of secondary uranium(VI)-bearing natural phases being represented by 40 minerals approved by the International Mineralogical Association, overtaken only by uranyl phosphates and uranyl sulfates. Uranyl carbonate phases form during the direct alteration of primary U ores on contact with groundwaters enriched by CO2, thus playing an important role in the release of U to the environment. The presence of uranyl carbonate phases has also been detected on the surface of "lavas" that were formed during the Chernobyl accident.

uranyl carbonate mineral crystal structure topology structural complexity

1. Introduction of Uranyl Carbonate Minerals

Uranyl carbonate phases play a very important role in all processes related to the nuclear fuel cycle. under the influence of groundwaters enriched with CO2, which can be derived from the dissolution of host carbonate rocks or from the atmosphere ^{[1][2][3][4][5]}. In dissolved form, uranyl carbonates can play an important role in U release to the environment. And of course, it should not be forgotten that uranyl-carbonate mineralization has been described among the alteration products of the "lavas" that were formed during the accident at the fourth reactor of the Chernobyl nuclear power plant in 1986 ^{[6][7]}.

There are 40 uranyl carbonate mineral species approved by the International Mineralogical Association as of 1 November 2020, thus making this group one of the most representative among secondary uranium minerals, coming third only after phosphates and sulfates [8][9]. Despite a fairly large number of known compounds, the structural diversity is not as great as one might expect. It is also of interest that about a quarter of approved minerals still have their crystal structures undetermined. The amount of synthetic structurally characterized uranyl carbonates is inferior to natural phases but can give an idea of the crystallization conditions present in the environment.

Current work is devoted to reviewing the topological diversity and growth conditions of natural uranyl carbonates and their synthetic analogs. Information-based complexity measures have been performed to determine contributions of various substructural building blocks and particular topological types into the complexity of the whole structure, which is related to the stability of a crystalline compound.

2. Synthetic Uranyl Carbonates

First of all, it should be noted that the number of synthetic phases for which the structures were determined is significantly inferior to the structurally characterized natural uranyl carbonates in the ratio of 19:32. Whereas for other groups of U(VI)-bearing compounds this proportion is usually opposite ^{[8][10][11]}. The first structurally characterized synthetic uranyl carbonate, to our knowledge, was one of the simplest phases **4a**, sodium-bearing Na₄(UO₂)(CO₃)₃ ^[12]. It is of interest that the first crystal structure of the natural uranyl carbonate, rutherfordine (**41**), was reported the year before ^[13]. The papers of K. Mereiter from the TU Wien (Austria) should be certainly noted in the first row among the works devoted to the synthesis and structural studies of synthetic uranyl carbonates. Then, the studies carried out by the A.M. Fedoseev and I.A. Charushnikova from the Frumkin Institute of Physical chemistry and Electrochemistry RAS (Russian Federation) and by V.N. Serezhkin from the Samara State University (Russian Federation) should be mentioned. A substantial portion of the synthetic uranyl carbonate compounds was synthesized and studied by P. C. Burns and his colleagues from the University of Notre Dame (USA), who significantly contribute to the studies of uranyl carbonate minerals as well.

All synthetic experiments can be roughly divided into two groups. Moreover, the majority of inorganic uranyl carbonates were synthesized by evaporation at room temperature and only a few of them were obtained from hydrothermal conditions. Uranyl nitrate hexahydrate was usually used as the source of U. But in some experiments, more specific reagents were used: UO₂ powder (for **1** and **2**), UO₂(CO₃) (for **7a**), Ag₄[UO₂(CO₃)₃] (for 8a), α -UO₂MoO₄(H₂O)₂ (for 10), and Na₄UO₂(CO₃)₃ (for 21). Potassium, sodium, cesium, or thallium carbonates were used as the source of CO₃ ions within the synthetic experiments. Compounds **36** and **10** can be considered as exceptions due to the usage of guanidine carbonate and carbamide in the respective syntheses. Several protocols of synthetic experiments deserve special attention. Thus, compound 1 [14] was formed as the result of the dissolution of UO₂ powder in the solution of K_2CO_3 and H_2O_2 at room temperature. Later it was filtered through a 0.45 µm polyamide syringe filter, and an additional 1.5 mL of 35 wt% H₂O₂ was added. Afterward it was transferred to a borosilicate scintillation vial and layered with methanol. The compound **2**^[15] was obtained similarly, except for the scintillation vial step. The crystals of 4a [16] were obtained by hydrothermal synthesis at 135 °C in a sealed silica glass tube at about 20 MPa. The compound **7a** ^[17] was synthesized by evaporation at room temperature, but before being left to evaporate, the dissolution of the precipitate was achieved via heating the solution over a steam bath. The crystals of **7b** ^[18] were obtained from the solution that was stirred for five days. The compound **11** ^[19] was synthesized by slow addition of uranyl nitrate solution to the solution of $Tl_2(CO_3)_2$ at the temperature of c.a. 57 °C. The solvent was removed in a vacuum desiccator for two months from the resulting yellow-green solution, and crystals were then taken from the precipitate. To obtain the crystals of **36** ^[20], the initial solution was stirred vigorously for several days, afterwards, it was centrifuged and the supernatant was removed via pipet. The crystals were obtained from the precipitate, which was slowly cooled to 5 °C under a CO₂ atmosphere.

Nine of uranyl carbonate minerals have synthetic analog, which was also obtained mostly by evaporation at room temperature, except for the crystals of **7** ^[21], which were formed during the two months of evaporation at vacuum-desiccator. The compound **4a** ^[22] was synthesized by hydrothermal reaction at 220 °C. The crystals of **41** ^[23] were obtained by purging the solution of UO₃ with 70 kPa CO₂ at the glove box for 24 h.

Very special attention can be paid to compound **4a** ^[22], which may be the same sodium uranyl carbonate that was found among alteration products of the "lavas" resulting from the nuclear accident of the Chernobyl nuclear power plant ^{[6][7]}.

3. Topological Analysis

The majority of uranyl carbonate crystal structures are based on finite clusters, which are represented by only two topological types (<u>Table 1</u>). The topological variety of layered uranyl carbonate complexes is significantly larger; however, the amount of compounds, which structures are based on the 2D units, is much lower. There are only ten uranyl carbonate compounds known with a layered structure, and all of them are natural phases.

The crystal structures of two synthetic K-bearing uranyl carbonates **1** ^[14] and **2** ^[15] differing only in the hydration state and one uranyl carbonate, templated by guanidinium molecules **3** ^[24], are based on the finite clusters of the *c*c0-1:2-9 topological type (Figure 1a,b). In terms of polyhedral representation, these clusters can be described as a hexagonal bipyramid that shares two of its equatorial edges with CO₃ groups spaced by one non-shared equatorial edge. There is a peroxide molecule arranged in the equatorial plane spaced from a carbonate group by another non-shared edge. Such topological type was also described in the structures of two uranyl nitrate compounds, pure inorganic ^[25] and organically templated ^[26], in which the peroxide group was replaced by two H₂O molecules. It should be noted that this topological type has an isomer, if two equatorial CO₃ groups are *trans*-arranged, being spaced by two non-shared edges, in contrast to the *cis*-arrangement in the structures of **1–3**. The *trans*-isomer is one of the most common types of uranyl nitrate finite clusters ^[27], while it has not been observed in the structures of uranyl carbonates at all.



Figure 1. Finite clusters in the crystal structures of natural and synthetic uranyl carbonates and their graphical representations (see Table 1 and text for details). Legend: see Figure 1; peroxide molecule is indicated by red bond (a); see Section 3.3 for details.

Uranyl tricarbonate cluster (UTC), which is shown in Figure 1c, is the most common structural unit among the natural and synthetic uranyl carbonate phases. There are 39 compounds known (Table 1), whose structures are based on these finite clusters, and is in sum 2.5 times more than the amount of all other structurally characterized uranyl carbonates (**15**). The topology of UTC belongs to the *cc*0-1:3-2 type (Figure 1d). This topology can be obtained from the previous *cis-cc*0-1:2-9 type by the replacement of the peroxide molecule by the third CO_3 group, resulting in the formation of a triangular cluster with the uranyl hexagonal bipyramid arranged in its core and ideal -6*m*2 point group symmetry.

No.	Chemical Formula	emical Formula Mineral Name		a, Å/ α, °	b, Ål β, °	c, Å/ y, °	Ref.				
Finite Clusters											
cc0-1:2-9											
1	K ₄ [(UO ₂ (CO ₃) ₂ (O ₂)](H ₂ O)		P2 ₁ /n	6.9670(14)/90	9.2158(18)/91.511(3)	18.052(4)/90	[<u>14]</u>				
2	K ₄ [(UO ₂ (CO ₃) ₂ (O ₂)](H ₂ O) _{2.5}		P2 ₁ /n	6.9077(14)/90	9.2332(18)/91.310(4)	21.809(4)/90	[<u>15</u>]				
3	$(CN_{3}H_{6})_{4}[UO_{2}(CO_{3})_{2}(O_{2})]\cdot H_{2}O$		Pca2 ₁	15.883(1)/90	8.788(2)/90	16.155(1)/90	[<u>24</u>]				
	cc0-1:3-2										
4	Na ₄ (UO ₂)(CO ₃) ₃	Čejkaite	Сс	9.2919(8)/90	16.0991(11)/91.404(5)	6.4436(3)/90	[<u>28</u>]				
4a	Na ₄ (UO ₂)(CO ₃) ₃		P-3c	9.3417/90	9.3417/90	12.824/120	[<u>22]</u> [<u>16]</u> [<u>12]</u>				
4b	Na ₄ (UO ₂)(CO ₃) ₃	Cejkaite old model	P-1	9.291(2)/90.73(2)	9.292(2)/90.82(2)	12.895(2)/120.00(1)	[<u>29</u>]				
5	K ₃ Na(UO ₂)(CO ₃) ₃		P-62c	9.29(2)/90	9.29(2)/90	8.26(2)/120	[<u>30</u>]				
6	K ₃ Na(UO ₂)(CO ₃) ₃ (H ₂ O)	Grimselite	P-62c	9.2507(1)/90	9.2507(1)/90	8.1788(1)/120	[<u>23]</u> [<u>31]</u> [<u>32]</u> [<u>33]</u>				
6a	Rb ₆ Na ₂ ((UO ₂)(CO ₃) ₃) ₂ (H ₂ O)	Rb analogue of Grimselite	P-62c	9.4316(7)/90	9.4316(7)/90	8.3595(8)/120	[<u>34</u>]				

Table 1. Crystallographic characteristics of natural and synthetic uranyl carbonates.

No.	Chemical Formula	Mineral Name	Sp.Gr.	a, Å/ α, °	b, Ål β, °	с, Å/ у, °	Ref.
7	K ₄ (UO ₂)(CO ₃) ₃	Agricolaite	C2/c	10.2380(2)/90	9.1930(2)/95.108(2)	12.2110(3)/90	[<u>35</u>]
7a	K ₄ UO ₂ (CO ₃) ₃		C2/c	10.247(3)/90	9.202(2)/95.11(2)	12.226(3)/90	[<u>21</u>]
7b	K ₄ (UO ₂)(CO ₃) ₃		C2/c	10.240(7)/90	9.198(4)/95.12(4)	12.222(12)/90	[<u>17</u>]
58	Rb ₄ (UO ₂)(CO ₃) ₃		C2/c	10.778(5)/90	9.381(2)/94.42(3)	12.509(3)/90	[<u>18</u>]
8	Cs ₄ UO ₂ (CO ₃) ₃ (H ₂ O) ₆		P2 <u>1</u> /n	11.1764(4)/90	9.5703(4)/ 96.451(2)	18.5756(7)/90	[<u>36</u>]
8a	Cs ₄ (UO ₂ (CO ₃) ₃)(H ₂ O) ₆		P2 ₁ /n	18.723(3)/90	9.647(2)/96.84(1)	11.297(2)/90	[<u>37</u>]
9	Cs ₄ (UO ₂ (CO ₃) ₃)		C2/c	11.5131(9)/90	9.6037(8)/93.767(2)	12.9177(10)/90	[<u>38</u>]
10	(NH ₄) ₄ (UO ₂ (CO ₃) ₃)		C2/c	10.679(4)/90	9.373(2)/96.43(2)	12.850(3)/90	[<u>39</u>]
11	Tl ₄ ((UO ₂)(CO ₃) ₃)		C2/c	10.684(2)/90	9.309(2)/94.95(2)	12.726(3)/90	[<u>19</u>]
12	Mg ₂ (UO ₂)(CO ₃) ₃ (H ₂ O) ₁₈	Bayleyite	P2 ₁ /a	26.560(3)/90	15.256(2)/92.90(1)	6.505(1)/90	[<u>40]</u> [<u>41</u>]
13	CaMg(UO ₂)(CO ₃) ₃ (H ₂ O) ₁₂	Swartzite	P2 ₁ /m	11.080(2)/90	14.634(2)/99.43(1)	6.439(1)/90	[<u>40]</u> [<u>42]</u> [<u>43</u>]
14	Ca ₂ (UO ₂)(CO ₃) ₃ (H ₂ O) ₁₁	Liebigite	Bba2	16.699(3)/90	17.557(3)/90	13.697(3)/90	[<u>40]</u> [<u>44]</u> [<u>45]</u> [<u>46]</u>
15	Ca ₉ (UO ₂) ₄ (CO ₃) ₁₃ ·28H ₂ O	Markeyite	Pmmn	17.9688(13)	18.4705(6)	10.1136(4)	[<u>47</u>]
16	Ca ₈ (UO ₂) ₄ (CO ₃) ₁₂ ·21H ₂ O	Pseudomarkeyite	P2 ₁ /m	17.531(3)	18.555(3)	9.130(3)/103.95(3)	[<u>48</u>]
17	Sr ₂ UO ₂ (CO ₃) ₃)(H ₂ O) ₈		P2 ₁ /c	11.379(2)/90	11.446(2)/93.40 (1)	25.653(4)/90	[<u>49</u>]
18	Na ₆ Mg(UO ₂) ₂ (CO ₃) ₆ ·6H ₂ O	Leószilárdite	C2/m	11.6093(21)/90	6.7843(13)/91.378(3)	15.1058(28)/90	[<u>50</u>]
19	Na ₂ Ca(UO ₂)(CO ₃) ₃ (H ₂ O) _{5.3}	Andersonite	R-3m	17.8448(4)/90	17.8448(4)/90	23.6688(6)/120	[23] [40] [51] [52] [53] [54] [55] [56] [57] [58]

No.	Chemical Formula	Mineral Name	Sp.Gr.	a, Ål α, °	b, Ål β, °	с, Å/ у, °	Ref.
							[<u>59</u>] [<u>60</u>] [<u>61</u>]
20	Na ₂ Ca ₈ (UO ₂) ₄ (CO ₃) ₁₃ ·27H ₂ O	Natromarkeyite	Pmmn	17.8820(13)	18.3030(4)	10.2249(3)	[<u>48</u>]
21	Ca ₃ Na _{1.5} (H ₃ O) _{0.5} (UO ₂ (C O ₃) ₃) ₂ (H ₂ O) ₈		Pnnm	18.150(3)/90	16.866(6)/90	18.436(3)/90	[<u>62</u>]
22	K ₂ Ca(UO ₂)(CO ₃) ₃ .6H ₂ O	Braunerite	P2 ₁ /c	17.6725(12)/90	11.6065(5)/101.780(8)	29.673(3)/90	[<u>63</u>]
23	K ₂ Ca ₃ [(UO ₂)(CO ₃) ₃] ₂ ·8(H ₂ O)	Linekite	Pnnm	17.0069(5)/90	18.0273(5)/90	18.3374(5)/90	[<u>64</u>]
23a	K ₂ Ca ₃ ((UO ₂ (CO ₃) ₃) ₂ (H ₂ O) ₆		Pnnm	17.015(2)/90	18.048(2)/90	18.394(2)/90	[<u>65</u>]
24	SrMg(UO ₂)(CO ₃) ₃ (H ₂ O) ₁₂	Swartzite-(Sr)	P2 ₁ /m	11.216(2)/90	14.739(2)/99.48(1)	6.484(1)/90	[<u>40]</u> [<u>42</u>]
25	Na _{0.79} Sr _{1.40} Mg _{0.17} (UO ₂ (C O ₃) ₃)(H ₂ O) _{4.66}		Pa-3	20.290(3)/90	20.290(3)/90	20.290(3)/90	[<u>66</u>]
26	MgCa ₅ Cu ₂ (UO ₂) ₄ (CO ₃) ₁₂ (H ₂ O) ₃₃	Paddlewheelite	Pc	22.052(4)/90	17.118(3)/90.474 (2)	19.354(3)/90	[<u>67</u>]
27	Na ₈ [(UO ₂)(CO ₃) ₃](SO ₄) ₂ ·3H ₂ O	Ježekite	P-62m	9.0664(11)/90	9.0664(11)/90	6.9110(6)/120	[<u>68</u>]
28	NaCa ₃ (UO ₂)(CO ₃) ₃ (SO ₄)F(H ₂ O) ₁₀	Schröckingerite	P-1	9.634(1)/91.41(1)	9.635(1)/92.33(1)	14.391(2)/120.26(1)	(40) (69) (70) (71) (72)
29	MgCa ₄ F ₂ [UO ₂ (CO ₃) ₃] ₂ (H ₂ O) _{17.29}	Albrechtschraufite	P-1	13.569(2)/115.82(1)	13.419(2)/107.61(1)	11.622(2)/92.84(1)	[<u>73]</u> [<u>74</u>]
30	Ca ₅ (UO ₂ (CO ₃) ₃) ₂ (NO ₃) ₂ (H ₂ O) ₁₀		P2 ₁ /n	6.5729(9)/90	16.517(2)/90.494(3)	15.195(2)/90	[<u>75</u>]
31	Ca ₆ (UO ₂ (CO ₃) ₃) ₂ C _{I4} (H ₂ O) ₁₉		P4/mbm	16.744(2)/90	16.744(2)/90	8.136(1)/90	[<u>75</u>]
32	Ca ₁₂ (UO ₂ (CO ₃) ₃) ₄ C _{I8} (H ₂ O) ₄₇		Fd-3	27.489(3)/90	27.489(3)/90	27.489(3)/90	[<u>75</u>]
33	Nd ₂ Ca[(UO ₂)(CO ₃) ₃](CO ₃) ₂ (H ₂ O) _{10.5}	Shabaite-(Nd)	P-1	8.3835(5)/90.058(3)	9.2766(12)/89.945(4)	31.7519(3)/90.331(4)	[<u>76</u>] [<u>77</u>]
34	[C(NH ₂) ₃] ₄ [UO ₂ (CO ₃) ₃]		R3	12.3278(1)/90	12.3278(1)/90	11.4457(2)/120	[<u>78</u>]
35	(C ₄ H ₁₂ N) ₄ [UO ₂ (CO ₃) ₃]·8H ₂ O		P2 ₁ /n	10.5377(18)/90	12.358(2)/99.343(4)	28.533(5)/90	[<u>79</u>]

No.	Chemical Formula	Mineral Name	Sp.Gr.	a, Å/ α, °	b, Ål β, °	c, Å/ y, °	Ref.				
			CC	:0-1:2-10							
36	[C(NH ₂) ₃] ₆ [(UO ₂) ₃ (CO ₃) ₆] (H ₂ O) _{6.5}		P-1	6.941(2)/95.63(2)	14.488(2)/98.47(2)	22.374 (2)/101.88(2)	[<u>20</u>]				
	Nanoclusters										
37	Mg ₈ Ca ₈ (UO ₂) ₂₄ (CO ₃) ₃₀ O ₄ (OH) ₁₂ (H ₂ O) ₁₃₈	Ewingite	I4 ₁ /acd	35.142(2)/90	35.142(2)/90	47.974(3)/90	[<u>80</u>]	-			
				Layers							
			5 ⁴ 4 ²	3 ⁴ (β-U ₃ O ₈)							
38	CaU(UO ₂) ₂ (CO ₃)O ₄ (OH) (H ₂ O) ₇	Wyartite	P212121	11.2706(8)/90	7.1055(5)/90	20.807(1)/90	[<u>81]</u> [<u>82]</u> [<u>83</u>]	-			
38a	Ca(U(UO ₂) ₂ (CO ₃) _{0.7} O ₄ (OH) _{1.6}) (H ₂ O) _{1.63}	Wyartite dehydrated	Pmcn	11.2610(6)/90	7.0870(4)/90	16.8359(10)/90	[<u>84</u>]				
			6 ¹ 5 ² 4 ² 3 ² ((phosphuranylite)							
39	Ca(UO ₂) ₃ (CO	Fontanite	P2 ₁ /n	6.968(3)/90	17.276(7)/90.064(6)	15.377(6)/90	[<u>85</u>] [<u>86</u>]	triurany			
			6 ¹ 5 ² 4 ² 3	3 ⁶ (roubaultite)				er urany			
40	Cu ₂ (UO ₂) ₃ (CO ₃) ₂ O ₂ (OH) ₂ (H ₂ O) ₄	Roubaultite	P-1	7.767(3)/92.16(4)	6.924(3)/90.89(4)	7.850(3)/93.48(4)	[<u>87]</u> [<u>88</u>]	ו a large ע of the			
			6 ¹ 3 ² -I ((rutherfordine)				e of this			
41	(UO ₂)(CO ₃)	Rutherfordine	Imm2	4.840(1)/90	9.273(2)/90	4.298(1)/90	[<u>23]</u> [<u>89]</u> [<u>13]</u> [<u>90]</u> [<u>91</u>]	n2 point			
42	Ca(H ₂ O) ₃ [(UO ₂) ₃ (CO ₃) _{3.6} O _{0.2}]	Sharpite	Cmcm	4.9032(4)	15.6489(11)	22.0414(18)	[<u>92</u>]	ll known			
42a	Ca(UO ₂) ₆ (CO ₃) ₅ (OH) ₄ .6H ₂ O	Sharpite	[<mark>80</mark>] Orth	21.99(2)	15.63(2)	4.487(4)	[<u>93</u>] [<u>94</u>]	e urany			
			6 ¹ 3 ² -II (widenmannite)				-teriinked			
43	3 Pb ₂ [(UO ₂)(CO ₃) ₂]	Widenmannite	Pmmn	4.9350(7)/90	9.550(4)/90	8.871(1)/90	[<u>31]</u> [<u>95</u>]	- building			

pentagonal pipyramids, 6 *cis*-isomer of the uranyl picarbonate unit (*cc*0-1:2-9), and 6 OTC complex (Figure 1h–j). Moreover, linkage of all building units occurs only through the carbonate groups. Mg and Ca ions as well as H_2O molecules are arranged both inside and in between the U-bearing nanoclusters.

The crystal structures of wyartite (**38**) ^{[81][82][83][84]} and its dehydrated form are based on the similar layered complexes that belong to the so-called β -U₃O₈-sheet anion topology (Figure 3a,b). Topology has the 5⁴4²3⁴ ring symbol and consists of infinite chains of edge-sharing pentagons that are linked with the neighbor chains through the common vertices and separated by the chains of squares and triangles. The crystal structures of **38** and **38a** are remarkable for being the only U(V)-bearing natural phases. Pentagons are occupied by the U⁶⁺ ions, squares correspond to the irregular U⁵⁺O₇ polyhedra, whereas triangles are vacant. Carbonate groups share edges with the U⁵⁺-centered polyhedra and are arranged towards the interlayer space.

No.	Chemical Formula	Mineral Name	Sp.Gr.	a, Å/ α, °	b, Ål β, °	с, Å/ у, °	Ref.			
							[<u>96</u>] [<u>97</u>]			
	Layers of Miscellaneous Topology									
44	Y ₂ (UO ₂) ₄ (CO ₃) ₃ O ₄ ·14H ₂ O	Kamotoite-(Y)	P21/n	12.3525(5)	12.9432 (5)/99.857(3)	19.4409(7)	[<u>98]</u> [<u>99</u>]			
45	$\begin{array}{c} [(Y_{4,22}Nd_{3.78}) \\ (H_2O)_{25}(UO_2)_{16}O_8 \\ (OH)_8(CO_3)_{16}](H_2O)_{14} \end{array}$	Bijvoetite-(Y)	B2 ₁	21.234(3)/90	12.958(2)/90.00(2)	44.911(7)/90	[<u>100</u>] [<u>101</u>]	-		
46	Ca(UO ₂)(CO ₃) ₂ ·5H ₂ O	Meyrowitzite	P2 ₁ /n	12.376(3)	16.0867(14)/107.679(13)	20.1340(17)	[<u>102</u>]	-		
		Mine	erals with	Undefined Structures						
47	Cu ₂ (Ce,Nd,La) ₂ (UO ₂)(CO ₃) ₅ (OH) ₂ ·1.5H2O	Astrocyanite-(Ce)	Hex	14.96(2)/90	14.96(2)/90	26.86(4)/120	[<u>103</u>]	-		
48	(UO ₂)(CO ₃)·H ₂ O	Blatonite	Hex or Trig	15.79(1)/90	15.79(1)/90	23.93(3)/120	[<u>104</u>]	-		
49	(UO ₂)(CO ₃)·nH ₂ O	Joliotite	Orth	8.16	10.35	6.32	[<u>31</u>]	-		
50	CaGd ₂ (UO ₂) ₂₄ (CO ₃) ₈ Si ₄ O ₂₈ . 60H ₂ O	Lepersonnite-(Gd)	Pnnm or Pnn2	16.23(3)/90	38.74(9)/90	11.73(3)/90	[<u>100</u>]	ir anior		
51	Ca(UO ₂)(CO ₃) ₂ ·3H ₂ O	Metazellerite	Pbn2 ₁ or Pbnm	9.718(5)	18.226(9)	4.965(4)	[<u>105</u>]	_		
52	(UO ₂) ₂ (CO ₃)(OH) ₂ ·4H ₂ O	Oswaldpeet	P21/c	4.1425(6)	14.098(3)/103.62(1)	18.374(5)	[<u>106</u>]	respond		
53	Ca ₃ Mg ₃ (UO ₂) ₂ (CO ₃) ₆ (OH) ₄ . 18H ₂ O	Rabbitite	Mon	[<u>114]</u> 32.6(1)	1 2 2 23.8(1)/~90	2 9.45(5)	[<u>107</u>]	topology		
54	Ca(UO ₂) ₃ (CO ₃)(OH) ₆ ·3H ₂ O	Urancalcarite	Pbnm or Pbn2 ₁ ,	15.42(3)	16.08(4)	6.970(6)	[<u>108</u>]	ng edge		
55	Ca ₂ Cu(UO ₂)(CO ₃) ₄ ·6H ₂ O	Voglite	P2 ₁ or P2 ₁ /m	25.97	24.50/104.0	10.70	[<u>109]</u> [<u>110]</u>	topology		
56	Ca(UO ₂)(CO ₃) ₂ ·5H ₂ O	Zellerite	Pmn2 ₁ or Pmnm	11.220(15)	19.252(16)	4.933(16)	[<u>105</u>] [<u>111</u>]	ariety o		
57	CaZn ₁₁ (UO ₂)(CO ₃) ₃ (OH) ₂₀ . 4H ₂ O	Znucalite	Orth	10.72(1)	[<u>8][10]</u> 25.16(1)	6.325(4)	[<u>112</u>]			

Another anion topology that consists of hexagons, pentagons, squares and triangles with the 6¹5²4²3⁶ ring symbol (Figure 2e,f) was described in the structure of roubaultite (**40**) ^[87][88]. Roubaultite anion topology contains the same infinite chains of edge-sharing pentagon dimers linked by edge-sharing hexagons that were observed in the phosphuranylite topology. But in the structure of **40**, these chains are separated by infinite chains of edge-sharing squares decorated with trimers of edge-sharing triangles on the sides. All hexagons and pentagons are also occupied by the uranyl ions, as I was realized in the phosphuranylite topology. Squares are occupied by Cucentered slightly distorted octahedra. The middle triangle from each trimer is occupied by the carbonate group, leaving the other two triangles vacant.

The simplest uranyl carbonate, at least according to the chemical composition, rutherfordine $(UO_2)(CO_3)$ ^{[23][89][13]} ^{[90][91]}, has a layered structure. The anion topology is also rather simple; it consists of parallel chains of edge-sharing hexagons separated by hourglass dimers of edge-sharing triangles (Figure 3g,h). All hexagons are occupied by the uranyl ions, and one triangle from each dimer is occupied by the CO₃ group, keeping the second

No.	Chemical Formula	Mineral Name	Sp.Gr.	a, Å/ α, °	b, Å/ β, °	с, Å/ у, °	Ref.	ig to the
57a	CaZn ₁₂ (UO ₂)(CO ₃) ₃ (OH) ₂₂ ·	Znucalite	Tricl	12.692(4)/89.08(2)	25.096(6)/91.79(2)	11.685(3)/90.37(3)	[<u>113</u>]	complex.
		Jonbeu us bein	9 1011110		ioudies of rutheric	nume topology.	Luon	module

represents a triple band of edge-sharing uranyl hexagonal bipyramids, a part of the triangular spaces between which are occupied by carbonate groups. These modules are arranged in the structure at approximate right angles to each other and are linked by the Ca-centered polyhedra, which are arranged on the crests of sawtooth waves (Figure 3a,b). Despite the curvature of such zigzag layers (in contrast to flat layered structure in rutherfordine) and the arrangement of Ca ions in the centers of square antiprisms, projection of such corrugated layers onto the (010) plane corresponds to the rutherfordine topology, with the equatorial arrangement of Ca polyhedra ligands having hexagon shape.



Figure 3. The crystal structures of sharpite (**a**,**b**) and meyrowitzite (**c**) with simplified topological representation (**d**). Legend: see <u>Figure 1</u>; Ca polyhedra = light blue.

The crystal structure of widenmannite (43) $^{[31][95][96][97]}$ is based on layered complexes, the topology of which consists of hexagons and triangles with the same $6^{1}3^{2}$ ring symbol as was found in the structure of rutherfordine. However, the arrangement of polygons in both structures is different. Widenmannite anion topology is built by the hexagons linked by vertex-sharing to other six hexagons, while all of its six edges are shared with triangles, thus forming trihexagonal tiling, which was used by Johannes Kepler in his book $^{[115]}$ and is also known under the kagome pattern name. In the ideal structure of the widenmannite (Figure 2i,j) each second row of hexagonal bipyramids should be vacant, but in the real structure the disorder with partial occupancy of the U sites takes place, which results in the occupation of all hexagons by the uranyl ions and half of the triangles oriented in the same direction are occupied by carbonate groups, keeping another half vacant.

The crystal structures of two *REE*-bearing minerals kamotoite-(Y) (**44**) ^{[98][99]} and bijvoetite-(Y) (**45**) ^{[100][101]} are based on highly remarkable and very rare layered complexes that have not been observed in any other natural or synthetic compound. The topology of the 2D complex is based on infinite chains of alternating dimers of edge-sharing uranyl hexagonal and pentagonal bipyramids (Figure 2k,I). Dimers of pentagonal bipyramids are arranged along the chain's extension, while dimers of hexagonal bipyramids are arranged perpendicularly. Each hexagonal bipyramid shares two of its oblique equatorial edges, not taking part in the linkage between U polyhedra, with CO_3 groups. These chains are linked into the 2D structure via irregular Y³⁺- or Nd³⁺-centered coordination polyhedra through the 6th non-shared equatorial edge from one chain and two O atoms of two carbonate groups from the neighbor chain. It should be noted that the resulting U- and *REE*-bearing layers are electroneutral, so the 3D structure formation is provided by the H-bonding system, which involves H₂O molecules from the coordination sphere of *REE* atoms and from the interlayer space.

The final to date topological type, which has been described in the structures of natural and synthetic uranyl carbonates, is observed in the structure of calcium uranyl carbonate mineral meyrowitzite (**46**) ^[102]. The structure of the layered complex is composed of UTC clusters sharing apical vertices of CO₃ groups with uranyl pentagonal bipyramids (Figure 3c). This structural type, unlike the rest of the topologies described herein, is the least dense in terms of the interconnection of U coordination polyhedra, and its topology will become clearer if graphical representation is used for illustration (Figure 3d). Thus, if uranyl pentagonal bipyramid is represented by grey nods, U hexagonal bipyramids by white nods, and the line between nods appearing if pentagonal bipyramid shares an equatorial O atom with the CO₃ group from the UTC cluster, the resulting graph of the complex will correspond to one of the most common *cc*1-1:2-4 topological type among the U(VI) bearing structures in general (e.g., ^{[8][10]}). Since the concept of graphical representation is violated, this description is more appropriate to use not as a direct interpretation of the topology, but as an approximate model.

4. Structural and Topological Complexity

Structural complexity measures have been implemented in several stages and the results of calculations are listed in Figure 4 and Figure 5 and Table 2. At the first stage, the topological complexity (**TI**), according to the maximal point (for finite clusters) or layer symmetry group has been determined, as these are the basic structure building units. At the second stage, the structural complexity (**SI**) of the U-bearing complexes has been calculated taking into account its real symmetry. The next informational contribution comes from the stacking (**LS**) of finite clusters and layered complexes (if more than one complex is within the unit cell). The fourth contribution to the total structural complexity is derived by the interstitial structure (**IS**). The last portion of information comes from the interstitial H bonding system (**H**). It should be noted that the H atoms related to the U-bearing clusters and layers were considered as a part of those complexes, but not within the contribution of the H-bonding system. Complexity parameters for the whole structure have been determined using *ToposPro* software [116].



Figure 4. Ladder diagrams showing contributions to structural complexity (bits per unit cell) (**a**) and normalized contributions (in %) (**b**) for the structures based on uranyl carbonate finite clusters. Legend: TI = topological information; SI = structural information; LS = layer stacking; IS = interstitial structure; HB = hydrogen bonding.



Figure 5. Ladder diagrams showing contributions to structural complexity (bits per unit cell) (**a**) and normalized contributions (in %) (**b**) for the structures based on uranyl carbonate layers. Legend: see Figure 4.

References

- 1. Alwan, A.K.; Williams, P.A. The aqueous chemistry of uranium minerals. Part 2. Minerals of the liebigite group. Mineral. Mag. 1980, 43, 665–667.
- 2. Clark, D.L.; Hobart, D.E.; Neu, M.P. Actinide carbonate complexes and their importance in actinide environmental chemistry. Chem. Rev. 1995, 95, 25–48.
- Plášil, J. Oxidation–hydration weathering of uraninite: The current state-of-knowledge. J. Geosci. 2014, 59, 99–114.
- Stefaniak, E.A.; Alsecz, A.; Frost, R.; Mathe, Z.; Sajo, I.E.; Torok, S.; Worobiec, A.; Van Grieken, R. Combined SEM/EDX and micro-Raman spectroscopy analysis of uranium minerals from a former uranium mine. J. Hazard Mater. 2009, 168, 416–423.
- 5. Driscoll, R.J.P.; Wolverson, D.; Mitchels, J.M.; Skelton, J.M.; Parker, S.C. A Raman spectroscopic study of uranyl minerals from Cornwall, UK. RSC Adv. 2014, 4, 59137–59149.
- Teterin, Y.A.; Baev, A.S.; Bogatov, S.A. X-ray photoelectron study of samples containing reactor fuel from "lava" and products growing on it which formed at Chernobyl NPP due to the accident. J. Electron Spectrosc. Relat. Phenom. 1994, 68, 685–694.
- 7. Burakov, B.E.; Strykanova, E.E.; Anderson, E. Secondary uranium minerals on the surface of Chernobyl "Lava". Mat. Res. Soc. Symp. Proc. 1996, 465, 1309–1311.
- 8. Lussier, A.J.; Lopez, R.A.K.; Burns, P.C. A revised and expanded structure hierarchy of natural and synthetic hexavalent uranium compounds. Can. Mineral. 2016, 54, 177–283.
- 9. Gurzhiy, V.V.; Plášil, J. Structural complexity of natural uranyl sulfates. Acta Crystallogr. 2019, B75, 39–48.
- Gurzhiy, V.V.; Kuporev, I.V.; Kovrugin, V.M.; Murashko, M.N.; Kasatkin, A.V.; Plášil, J. Crystal chemistry and structural complexity of natural and synthetic uranyl selenites. Crystals 2019, 9, 639.
- Krivovichev, S.V.; Burns, P.C. Actinide compounds containing hexavalent cations of the VI group elements (S, Se, Mo, Cr, W). In Structural Chemistry of Inorganic Actinide Compounds; Krivovichev, S.V., Burns, P.C., Tananaev, I.G., Eds.; Elsevier: Amsterdam, The Netherland, 2007; pp. 95–182.
- 12. Douglass, M. Tetrasodium uranyl tricarbonate, Na4UO2(CO3)3. Anal. Chem. 1956, 28, 1635.

- 13. Christ, C.L.; Clark, J.R.; Evans, H.T., Jr. Crystal structure of rutherfordine, UO2CO3. Science 1955, 121, 472–473.
- Goff, G.S.; Brodnax, L.F.; Cisneros, M.R.; Peper, S.M.; Field, S.E.; Scott, B.L.; Runde, W.H. First identification and thermodynamic characterization of the ternary U(VI) species, UO2(O2) (CO3)2(4-), in UO2-H2O2-K2CO3 solutions. Inorg. Chem. 2008, 47, 1984–1990.
- Zehnder, R.; Peper, S.; Brian, L.; Runde, S.; Runde, W. Tetrapotassium dicarbonatodioxoperoxouranium(VI) 2.5-hydrate, K4[U(CO3)2O2(O2)]·2.5H2O. Acta Crystallogr. 2005, C61, 3–5.
- 16. Císařová, I.; Skála, R.; Ondruš, P.; Drábek, M. Trigonal Na4[UO2(CO3)3]. Acta Crystallogr. 2001, E37, 32–34.
- Han, J.-C.; Rong, S.-B.; Chen, Q.-M.; Wu, X.-R. The determination of the crystal structure of tetrapotassium uranyl tricarbonate by powder X-ray diffraction method. Chin. J. Chem. 1990, 4, 313–318.
- 18. Chernorukov, N.G.; Mikhailov, Y.N.; Knyazev, A.V.; Kanishcheva, A.S.; Zamkovaya, E.V. Synthesis and crystal structure of rubidium uranyltricarbonate. Russ. J. Coord. Chem. 2005, 31, 364–367.
- 19. Mereiter, K. Structure of Thallium Tricarbonatodioxouranat (VI). Acta Crystallogr. Cryst. Struct. Commun. 1986, C42, 1682–1684.
- 20. Allen, P.G.; Bucher, J.J.; Clark, D.L.; Edelstein, N.M.; Ekberg, S.A.; Gohdes, J.W.; Hudson, E.A.; Kaltsoyannis, N.; Lukens, W.W.; Neu, M.P.; et al. Multinuclear NMR, Raman, EXAFS, and X ray diffraction studies of uranyl carbonate complexes in near-neutral aqueous solution. X-ray structure of (C(NH2)3)6((UO2)3(CO3)6).6.5(H2O). Inorg. Chem. 1995, 34, 4797–4807.
- 21. Anderson, A.; Chieh, C.; Irish, D.E.; Tong, J.P.K. An X-Ray crystallographic, Raman, and infrared spectral study of crystalline potassium uranyl carbonate, K4UO2(CO3)3. Can. J. Chem. 1980, 58, 1651–1658.
- 22. Li, Y.; Krivovichev, S.V.; Burns, P.C. The crystal structure of Na4(UO2)(CO3)3 and its relationship to schröckingerite. Mineral. Mag. 2001, 65, 297–304.
- 23. Kubatko, K.-A.; Helean, K.B.; Navrotsky, A.; Burns, P.C. Thermodynamics of uranyl minerals: Enthalpies of formation of rutherfordine, UO2CO3, andersonite, Na2CaUO2(CO3)3(H2O)5, and grimselite, K3NaUO2(CO3)3H2O. Amer. Mineral. 2005, 90, 1284–1290.
- 24. Mikhailov, Y.N.; Lobanova, G.M.; Shchelokov, R.N. X-ray structural study of the guanidinium uranyl-peroxo-dicarbonate hydrate (CN3H6)4UO2O2(CO3)2·H2O. Zh. Neorg. Khim. 1981, 26, 718–722.
- 25. Shuvalov, R.R.; Burns, P.C. A monoclinic polymorph of uranyl dinitrate trihydrate, [UO2(NO3)2(H2O)2]·H2O. Acta Crystallogr. 2003, C59, 71–73.

- 26. Thuéry, P. Uranyl Ion Complexes with Cucurbit[n]urils (n = 6, 7, and 8): A new family of uranylorganic frameworks. Cryst. Growth Des. 2008, 8, 4132–4143.
- 27. Gurzhiy, V.V.; Kornyakov, I.V.; Tyumentseva, O.S. Uranyl nitrates: Byproducts of the synthetic experiments or key indicators of the reaction progress? Crystals 2020, 10, 1122.
- Plášil, J.; Fejfarová, K.; Dušek, M.; Škoda, R.; Rohlíček, J. Actinides in Geology, Energy, and the Environment. Revision of the symmetry and the crystal structure of čejkaite, Na4(UO2)(CO3)3. Am. Mineral. 2013, 98, 549–553.
- 29. Ondruš, P.; Skála, R.; Veselovský, F.; Sejkora, J.; Vitti, C. Čejkaite, the triclinic polymorph of Na4(UO2)(CO3)3—A new mineral from Jachymov, Czech Republic. Am. Min. 2003, 88, 686–693.
- 30. Mazzi, F.; Rinaldi, F. La struttura cristallina del K3Na(UO2)(CO3)3. Period. Mineral. 1961, 30, 1– 21.
- 31. Walenta, K. Widenmannit und Joliotit, zwei neue Uranylkarbonatmineralien aus dem Schwarzwald. Schweiz. Mineral. Petrogr. Mitt. 1976, 56, 167–185.
- 32. Li, Y.; Burns, P. The crystal structure of synthetic grimselite, K3Na[(UO2)(CO3)3](H2O). Can. Mineral. 2001, 39, 1147–1151.
- Plášil, J.; Fejfarová, K.; Skála, R.; Škoda, R.; Meisser, N.; Hloušek, J.; Císařová, I.; Dušek, M.; Veselovský, F.; Čejka, J.; et al. The crystal chemistry of the uranyl carbonate mineral grimselite, (K,Na)3Na[(UO2)(CO3)3](H2O), from Jáchymov, Czech Republic. Mineral. Mag. 2012, 76, 443– 453.
- 34. Kubatko, K.A.; Burns, P.C. The Rb analogue of grimselite, Rb6Na2((UO2)(CO3)3)2 (H2O). Acta Crystallogr. Cryst. Struct. Commun. 2004, C60, 25–26.
- 35. Skála, R.; Ondruš, P.; Veselovský, F.; Císařová, I.; Hloušek, J. Agricolaite, a new mineral of uranium from Jáchymov, Czech Republic. Mineral. Petrol. 2011, 103, 169–175.
- 36. Charushnikova, I.A.; Fedoseev, A.M.; Perminov, V.P. Synthesis and Crystal Structure of Cesium Actinide(VI) Tricarbonate Complexes Cs4AnO2(CO3)3·6H2O, An(VI) = U., Np, Pu. Radiochemistry 2016, 58, 578–585.
- 37. Mereiter, K. Structure of cesium tricarbonatodioxouranate(VI) hexahydrate. Acta Crystallogr. Cryst. Struct. Commun. 1988, C44, 1175–1178.
- 38. Krivovichev, S.V.; Burns, P.C. Synthesis and crystal structure of Cs4(UO2(CO3)3). Radiochemistry 2004, 46, 12–15.
- Serezhkin, V.N.; Soldatkina, M.A.; Boiko, N.V. Refinement of the crystal-structure of (NH4)4UO2(CO3)3. J. Struct. Chem. 1983, 24, 770–774.

- 40. Axelrod, J.M.; Grimaldi, F.S.; Milton, C.; Murata, K.J. The uranium minerals from the Hillside mine, Yavapai County, Arizona. Am. Mineral. 1951, 36, 1–22.
- 41. Mayer, H.; Mereiter, K. Synthetic bayleyite, Mg2[UO2(CO3)3]·18H2O: Thermochemistry, crystallography and crystal structure. Tschermaks Mineral. Petrogr. Mitt. 1986, 35, 133–146.
- 42. Mereiter, K. Synthetic swartzite, CaMg[UO2(CO3)3]·12H2O, and its strontium analogue, SrMg[UO2(CO3)3]·12H2O: Crystallography and crystal structures. Neues Jahrb. Mineral. Mon. 1986, 1986, 481–492.
- 43. Amayri, S.; Arnold, T.; Foerstendorf, H.; Geipel, G.; Bernhard, G. Spectroscopic characterization of synthetic becquerelite, Ca[UO2)6O4(OH)6]·8H2O, and swartzite, CaMg[UO2(CO3)3]·12H2O. Can. Mineral. 2004, 42, 953–962.
- 44. Vochten, R.; Van Haverbeke, L.; Van Springel, K. Synthesis of liebigite and andersonite, and study of their thermal behavior and luminescence. Can. Mineral. 1993, 31, 167–171.
- 45. Smith, J.L. Two new minerals—medjidite (sulphate of uranium and lime)—Liebigite (carbonate of uranium and lime). Am. J. Sci. Arts 1848, 5, 336–338.
- 46. Mereiter, K. The crystal structure of liebigite, Ca2UO2(CO3)3·~11H2O. Tschermaks Mineral. Petrogr. Mitt. 1982, 30, 277–288.
- 47. Kampf, A.R.; Plášil, J.; Kasatkin, A.V.; Marty, J.; Čejka, J. Markeyite, a new calcium uranyl carbonate mineral from the Markey mine, San Juan County, Utah, USA. Mineral. Mag. 2018, 82, 1089–1100.
- Kampf, A.R.; Olds, T.A.; Plášil, J.; Burns, P.C.; Marty, J. Natromarkeyite and pseudomarkeyite, two new calcium uranyl carbonate minerals from the Markey mine, San Juan County, Utah, USA. Mineral. Mag. 2020, 84, 753–765.
- 49. Mereiter, K. Structure of strontium tricarbonatodioxouranate(VI) octahydrate. Acta Crystallogr. Cryst. Struct. Commun. 1986, C42, 1678–1681.
- 50. Olds, T.; Sadergaski, L.; Plášil, J.; Kampf, A.; Burns, P.; Steele, I.; Marty, J.; Carlson, S.; Mills, O. Leószilárdite, the first Na,Mg-containing uranyl carbonate from the Markey Mine, San Juan County, Utah, USA. Mineral. Mag. 2017, 81, 1039–1050.
- Gurzhiy, V.V.; Krzhizhanovskaya, M.G.; Izatulina, A.R.; Sigmon, G.E.; Krivovichev, S.V.; Burns, P.C. Structure refinement and thermal stability studies of the uranyl carbonate mineral Andersonite, Na2Ca[(UO2)(CO3)3]·(5+x)H2O. Minerals 2018, 8, 586.
- 52. Coda, A.; Della Giusta, A.; Tazzoli, V. The structure of synthetic andersonite, Na2Ca[UO2(CO3)3].xH2O (x = 5.6). Acta Cryst. 1981, B37, 1496–1500.
- 53. Mereiter, K. Neue kristallographische Daten ueber das Uranmineral Andersonit. Anz. Österr. Akad. Wiss. Mathemat. Naturwiss. Kl. 1986, 123, 39–41.

- 54. Plášil, J.; Čejka, J. A note on the molecular water content in uranyl carbonate mineral andersonite. J. Geosci. 2015, 60, 181–187.
- 55. Coda, A. Ricerche sulla struttura cristallina dell'Andersonite. Atti Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Nat. Ser. 1963, 34, 299–304.
- 56. Čejka, J.; Urbanec, Z.; Čejka, J., Jr. To the crystal chemistry of andersonite. Neu. Jb. Mineral. Mh. 1987, 11, 488–501.
- 57. De Neufville, J.P.; Kasdan, A.; Chimenti, R.J.L. Selective detection of uranium by laser-induced fluorescence: A potential remote-sensing technique. 1: Optical characteristics of uranyl geologic targets. Appl. Opt. 1981, 20, 1279–1296.
- Amayri, S.; Arnold, T.; Reich, T.; Foerstendorf, H.; Geipel, G.; Bernhard, G.; Massanek, A. Spectroscopic characterization of the uranium carbonate andersonite Na2Ca[UO2(CO3)3]·6H2O. Environ. Sci. Technol. 2004, 38, 6032–6036.
- 59. Frost, R.L.; Carmody, O.; Ertickson, K.L.; Weier, M.L.; Čejka, J. Molecular structure of the uranyl mineral andersonite—A Raman spectroscopic study. J. Molec. Struct. 2004, 703, 47–54.
- 60. Čejka, J. To the chemistry of andersonite and thermal composition of dioxo-tricarbonatouranates. Coll. Czech. Chem. Commun. 1969, 34, 1635–1656.
- 61. Čejka, J.; Urbanec, Z. Thermal and infrared spectrum analyses of natural and synthetic andersonites. J. Therm. Anal. 1988, 33, 389–394.
- 62. Vochten, R.; van Haverbeke, L.; van Springel, K.; Blaton, N.; Peeters, M. The structure and physicochemical characteristics of a synthetic phase compositionally intermediate between liebigite and andersonite. Can. Mineral. 1994, 32, 553–561.
- 63. Plášil, J.; Mereiter, K.; Kampf, A.R.; Hloušek, J.; Škoda, R.; Čejka, J.; Němec, I.; Ederová, J. Braunerite IMA 2015-123. CNMNC Newsletter No. 31. Mineral. Mag. 2016, 80, 692.
- 64. Plášil, J.; Čejka, J.; Sejkora, J.; Hloušek, J.; Škoda, R.; Novák, M.; Dušek, M.; Císařová, I.; Němec, I.; Ederová, J. Línekite, K2Ca3[(UO2)(CO3)3]2.8H2O, a new uranyl carbonate mineral from Jáchymov, Czech Republic. J. Geosci. 2017, 62, 201–213.
- 65. Kubatko, K.-A.; Burns, P. The crystal structure of a novel uranyl tricarbonate, K2Ca3[(UO2) (CO3)3)]2(H2O)6. Can. Mineral. 2004, 42, 997–1003.
- Effenberger, H.; Mereiter, K. Structure of a cubic sodium strontium magnesium tricarbonatodioxouranate(VI) hydrate. Acta Crystallogr. Cryst. Struct. Commun. 1988, C44, 1172– 1175.
- 67. Olds, T.; Plášil, J.; Kampf, A.; Dal Bo, F.; Burns, P. Paddlewheelite, a New Uranyl Carbonate from the Jáchymov District, Bohemia, Czech Republic. Minerals 2018, 8, 511.

- Plášil, J.; Hloušek, J.; Kasatkin, A.V.; Belakovskiy, D.I.; Čejka, J.; Chernyshov, D. Ježekite, Na8[(UO2)(CO3)3](SO4)2·3H2O, a new uranyl mineral from Jáchymov, Czech Republic. J. Geosci. 2015, 60, 259–267.
- 69. Schrauf, A. Schröckingerit, ein neues Mineral von Joachimsthal. Tschermaks Mineral. Petrogr. Mitt. 1873, 1, 137–138.
- 70. Jaffe, H.W.; Sherwood, A.M.; Peterson, M.J. New data on schroeckingerite. Am. Mineral. 1948, 33, 152–157.
- 71. Smith, D.K. An X-ray crystallographic study of schroeckingerite and its dehydration product. Am. Mineral. 1959, 44, 1020–1025.
- 72. Mereiter, K. Crystal structure and crystallographic properties of a schröckingerite from Joachimsthal. Tschermaks Mineral. Petrogr. Mitt. 1986, 35, 1–18.
- 73. Mereiter, K. The crystal structure of albrechtschraufite, MgCa4F2[(UO2)(CO3)3]2·17H2O. Acta Crystallogr. 1984, A40, 247.
- 74. Mereiter, K. Description and crystal structure of albrechtschaufite, MgCa4F2[UO2(CO3)3]2·17-18H2O. Mineral. Petrol. 2013, 107, 179–188.
- 75. Li, Y.; Burns, P.C. New structural arrangements in three ca uranyl carbonate compounds with multiple anionic species. J. Solid State Chem. 2002, 166, 219–228.
- Plášil, J.; Škoda, R. Crystal structure of the (REE)–uranyl carbonate mineral shabaite-(Nd). J. Geosci. 2017, 62, 97–105.
- 77. Deliens, M.; Piret, P. La shabaïte-(Nd), Ca(TR)2(U02)(C03)4(OH)2.6H20, nouvelle espéce minérale de Kamoto, Shaba, Zaïre. Eur. J. Mineral. 1989, 1, 85–88.
- Fedosseev, A.M.; Gogolev, A.V.; Charushnikova, I.A.; Shilov, V.P. Tricarbonate complex of hexavalent Am with guanidinium: Synthesis and structural characterization of [C(NH2)3]4[AmO2(CO3)3]·2H2O, comparison with [C(NH2)3]4[AnO2(CO3)3] (An=U, Np, Pu). Radiochim. Acta 2011, 99, 679–686.
- 79. Reed, W.A.; Oliver, A.G.; Rao, L. Tetrakis(tetramethylammonium) tricarbonatodioxidouranate octahydrate. Acta Crystallogr. 2011, 67, 301–303.
- 80. Olds, T.; Plášil, J.; Kampf, A.; Simonetti, A.; Sadergaski, L.; Chen, Y.-S.; Burns, P. Ewingite: Earth's most complex mineral. Geology 2017, 45, 1007–1010.
- Bullemin, C.; Protas, J. Ianthinite et wyartite. Bull. Société Française Minéralogie Cristallogr. 1959, 82, 80–86.
- 82. Burns, P.C.; Finch, R.J. Wyartite: Crystallographic evidence for the first pentavalent-uranium mineral. Am. Mineral. 1999, 84, 1456–1460.

- 83. Frost, R.L.; Henry, D.A.; Erickson, K. Raman spectroscopic detection of wyartite in the presence of rabejacite. J. Raman Spectrosc. 2004, 35, 255–260.
- 84. Hawthorne, F.C.; Finch, R.J.; Ewing, R.C. The crystal structure of dehydrated wyartite, Ca(CO3) [U5+(U6+O2)2O4(OH)](H2O)3. Can. Mineral. 2006, 44, 1379–1385.
- 85. Deliens, M.; Piret, P. La fontanite, carbonate hydraté d'uranyle et de calcium, nouvelle espèce minérale de Rabejac, Hérault, France. Eur. J. Mineral. 1992, 4, 1271–1274.
- 86. Hughes, K.A.; Burns, P.C. A new uranyl carbonate sheet in the crystal structure of fontanite, Ca[(UO2)3(CO3)2O2](H2O)6. Am. Mineral. 2003, 88, 962–966.
- 87. Cesbron, F.; Pierrot, R.; Verbeek, T. La roubaultite Cu2(UO2)3(OH)10.5H2O, une nouvelle espèce minérale. Bull. Société Française Minéralogie Cristallogr. 1970, 93, 550–554.
- 88. Ginderow, D.; Cesbron, F. Structure de la roubaultite Cu2(UO2)3(CO3)2O2(OH)2·4H2O. Acta Crystallogr. 1985, 41, 654–657.
- 89. Marckwald, W. Ueber Uranerze aus Deutsch-Ostafrika. Zent. Mineral. Geol. Paläontologie 1906, 1906, 761–763.
- 90. Frondel, C.; Meyrowitz, R. Studies of uranium minerals (XIX): Rutherfordine, diderichite, and clarkeite. Am. Mineral. 1956, 41, 127–133.
- 91. Finch, R.J.; Cooper, M.A.; Hawthorne, F.C.; Ewing, R.C. Refinement of the crystal structure of rutherfordine. Can. Mineral. 1999, 37, 929–938.
- 92. Plášil, J. A unique structure of uranyl-carbonate mineral sharpite: A derivative of the rutherfordine topology. Z. Krist. Cryst. Mater. 2018, 233, 579–586.
- 93. Melón, M.J. La sharpite, nouveau carbonate d'uranyle du Congo belge. Bull. Séances l'Institut R. Colonial Belg. 1938, 9, 333–336.
- 94. Cejka, J.; Mrazek, Z.; Urbanec, Z. New data on sharpite, a calcium uranyl carbonate. Neues Jahrb. Mineral. Monatsh. 1984, 1984, 109–117.
- 95. Elton, N.J.; Hooper, J.J. Widenmannite from Cornwall, England: The second world occurrence. Mineral. Mag. 1995, 59, 745–749.
- 96. Plášil, J.; Čejka, J.; Sejkora, J.; Škácha, P.; Goliáš, V.; Jarka, P.; Laufek, F.; Jehlička, J.; Němec,
 I.; Strnad, L. Widenmannite, a rare uranyl lead carbonate: Occurrence, formation and
 characterization. Miner. Mag. 2010, 74, 97–110.
- 97. Plášil, J.; Palatinus, L.; Rohlíček, J.; Houdková, L.; Klementová, M.; Goliáš, V.; Škácha, P. Crystal structure of lead uranyl carbonate mineral widenmannite: Precession electron-diffraction and synchrotron powder-diffraction study. Am. Mineral. 2014, 99, 276–282.

- 98. Deliens, M.; Piret, P. La kamototïte-(Y), un nouveau carbonate d'uranyle et de terres rares de Kamoto, Shaba, Zaïre. Bull. Minéralogie 1986, 109, 643–647.
- 99. Plášil, J.; Petříček, V. Crystal structure of the (REE)-uranyl carbonate mineral kamotoite-(Y). Mineral. Mag. 2017, 81, 653–660.
- 100. Deliens, M.; Piret, P. Bijvoetite et lepersonnite, carbonates hydrates d'uranyle et de terres rares de Shinkolobwe, Zaire. Can. Mineral. 1982, 22, 231–238.
- 101. Li, Y.; Burns, P.C.; Gault, R.A. A new rare-earth element uranyl carbonate sheet in the structure of bijvoetite-(Y). Can. Mineral. 2000, 38, 153–162.
- 102. Kampf, A.R.; Plášil, J.; Olds, T.A.; Nash, B.P.; Marty, J.; Belkin, H.E. Meyrowitzite, Ca(UO2) (CO3)2·5H2O, a new mineral with a novel uranyl-carbonate sheet. Am. Mineral. 2019, 104, 603–610.
- 103. Deliens, M.; Piret, P. L'astrocyanite-(Ce), Cu2(TR)2(UO2)(CO3)5(OH)2·1,5 H2O, nouvelle espèce minérale de Kamoto, Shaba, Zaïre. Eur. J. Mineral. 1990, 2, 407–411.
- 104. Vochten, R.; Deliens, M. Blatonite, UO2CO3·H2O, A new uranyl carbonate monohydrate from San Juan County, Utah. Can. Mineral. 1998, 36, 1077–1081.
- 105. Coleman, R.G.; Ross, D.R.; Meyrowitz, R. Zellerite and metazellerite, new uranyl carbonates. Am. Mineral. 1966, 51, 1567–1578.
- 106. Vochten, R.; Deliens, M.; Medenbach, O. Oswaldpeetersite, (UO2)2CO3(OH)2·4H2O, a new basic uranyl carbonate mineral from the Jomac uranium mine, San Juan County, Utah, U.S.A. Can. Mineral. 2001, 39, 1685–1689.
- 107. Thompson, M.E.; Weeks, A.D.; Sherwood, A.M. Rabbittite, a new uranyl carbonate from Utah. Am. Mineral. 1955, 40, 201–206.
- 108. Deliens, M.; Piret, P. L'urancalcarite, Ca(UO2)3CO3(OH)6.3H2O, nouveau minéral de Shinkolobwe, Shaba, Zaïre. Bull. Minéralogie 1984, 107, 21–24.
- 109. Vogl, J.F. Drei neue Mineral-Vorkommen von Joachimsthal. Jahrb. Kais. Königlichen Geol. Reichsanst. 1853, 4, 220–223.
- 110. Piret, P.; Deliens, M. New crystal data for Ca, Cu, UO2 hydrated carbonate: Voglite. J. Appl. Crystallogr. 1979, 12, 616.
- 111. Frost, R.L.; Dickfos, M.J.; Cejka, J. Raman spectroscopic study of the uranyl carbonate mineral zellerite. J. Raman Spectrosc. 2008, 39, 582–586.
- 112. Chiappero, P.J.; Sarp, H. Nouvelles données sur la znucalite et seconde occurrence: Le Mas d'Alary, Lodève (Hérault, France). Arch. Sci. 1993, 46, 291–301.

- 113. Ondruš, P.; Veselovský, F.; Rybka, R. Znucalite, Zn12(UO2)Ca(CO3)3(OH)22·4H2O, a new mineral from Príbram, Czechoslovakia. Neues Jahrb. Mineral. Mon. 1990, 1990, 393–400.
- 114. Demartin, F.; Diella, V.; Donzelli, S.; Gramaccioli, C.M.; Pilati, T. The importance of accurate crystal structure determination of uranium minerals. I. Phosphuranylite KCa(H3O)3(UO2)7(PO4)4O4·8H2O. Acta Crystallogr. 1991, B47, 439–446.
- 115. Kepler, J. Harmonices Mundi Libri V; Forni: Bologna, Italy, 1619.
- 116. Blatov, V.A.; Shevchenko, A.P.; Proserpio, D.M. Applied topological analysis of crystal structures with the program package ToposPro. Cryst. Growth. Des. 2014, 14, 3576–3586.

Retrieved from https://encyclopedia.pub/entry/history/show/26547