

Sirtuins in Kidney Diseases

Subjects: **Biochemistry & Molecular Biology**

Contributor: Yu Ah Hong , Ji Eun Kim , Minjee Jo , Gang-Jee Ko

Sirtuins (SIRT) are class III histone deacetylases (HDACs) that play important roles in aging and a wide range of cellular functions. Sirtuins are crucial to numerous biological processes, including proliferation, DNA repair, mitochondrial energy homeostasis, and antioxidant activity. Mammals have seven different sirtuins, SIRT1–7, and the diverse biological functions of each sirtuin are due to differences in subcellular localization, expression profiles, and cellular substrates. In this review, we summarize research advances into the role of sirtuins in the pathogenesis of various kidney diseases including acute kidney injury, diabetic kidney disease, renal fibrosis, and kidney aging along with the possible underlying molecular mechanisms. The available evidence indicates that sirtuins have great potential as novel therapeutic targets for the prevention and treatment of kidney diseases.

sirtuins

kidney

acute kidney injury

diabetic nephropathy

chronic kidney disease

aging kidney

1. Introduction

Sirtuins (SIRT) are a family of nicotinamide adenine dinucleotide (NAD)⁺-dependent class III histone deacetylases (HDACs) that use the coenzyme NAD⁺ to deacetylate lysine residues in histone and non-histone proteins. Due to their ability to target post-translational acyl modifications of various cellular substrates, sirtuins are crucial to numerous biological processes including proliferation, DNA repair, mitochondrial energy homeostasis, and antioxidant activity [1]. Silent information regulator 2 (SIR2) was the first sirtuin discovered and was originally found in *Saccharomyces cerevisiae* [2]. SIR2 is critical for transcriptional silencing in budding *S. cerevisiae* and extension of the lifespan of mother cells through suppression of toxic recombinant DNA circle formation, as well as other processes including the management of molecules damaged by oxidative stress [3]. Seven different sirtuin proteins localized to different subcellular compartments have been identified in mammals [4]. SIRT1 and SIRT2 are distributed in both the nucleus and cytoplasm. SIRT3, SIRT4, and SIRT5 are mainly present in the mitochondria, while SIRT6 and SIRT7 are primarily found in the nucleus [5]. The diverse biological functions of the various sirtuins are attributed to differences in subcellular localization, expression profiles, and cellular substrates.

2. Molecular Targets of Sirtuins

SIRT1, as the first sirtuin identified in mammals, has been the most widely investigated. Although SIRT1 is mainly present in the nucleus, it can translocate into the cytoplasm under specific conditions, such as ischemic stress or embryonic development [6][7]. In the kidney, SIRT1 is widely expressed in tubular cells and podocytes [8]. At the

cellular level, SIRT1 may play important roles in several biological processes, including energetic homeostasis [9], apoptosis [10], mitochondrial biogenesis [11], and autophagy [12]. SIRT1 mediates the longevity effect of caloric restriction through regulation of glucose and lipid metabolism [13]. Caloric restriction increases the NAD⁺/NADH ratio and subsequently SIRT1 activity [14]. Using the coenzyme NAD⁺, SIRT1 enhances chromatin silencing and transcriptional repression through deacetylation of histones [15]. SIRT1 can directly deacetylate histone lysines H4K16 (H4 Lys16), H3K9 (H3 Lys9), H3K14 (H3 Lys14), and H1K26 (H1 Lys26) upon recruitment to chromatin [16]. SIRT1 acts as the key transcriptional modulator of cell survival via regulation of p53 [17], nuclear factor- κ B (NF- κ B) p65 [18], signal transducer and activator of transcription 3 (STAT3) [19], and the Forkhead box class O (FoxO) family [20]. SIRT1 regulates the cellular response to hypoxic stress through deacetylation of hypoxia-induced factor 1 α (HIF-1 α) [21][22]. SIRT1 also activates peroxisome proliferator-activated receptor gamma (PPAR γ) through deacetylation of PPAR γ coactivator-1 α (PGC-1 α) [23][24]. In addition, SIRT1 regulates cellular homeostasis through its activation of adenosine monophosphate-activated protein kinase (AMPK) via liver kinase B1 (LKB1) and inhibition of the mammalian target of rapamycin (mTOR) [25][26]. SIRT1 also accelerates deacetylation and nuclear translocation of β -catenin and regulates the Wnt/ β -catenin transcriptional pathway [27].

SIRT2 is a cytoplasmic sirtuin that is co-localized with microtubules and deacetylates the major component of microtubules [28]. However, SIRT2 has also been observed in the nucleus and mitochondria, and its translocation to the nucleus may indicate an important epigenetic role [29]. SIRT2 has been shown to deacetylate several substrates including histone lysines H4K16 [30], H3K56 (H3 Lys56) [31], α -tubulin [28], PR-Set7 [32], phosphoenolpyruvate carboxykinase 1 (PEPCK1) [33], NF- κ B p65 [34], and FoxO family proteins [35][36], and is also involved in the regulation of the cell cycle, DNA repair, and stress responses [30]. SIRT2 regulates binding of p65 to the promoters of anti-inflammatory chemokines such as C-X-C motif chemokine ligand 2 (CXCL2) and CC motif chemokine ligand 2 (CCL2) [34]. SIRT2 also regulates insulin-induced protein kinase B (PKB/AKT) activation via constitutive phosphoinositide 3-kinase (PI3K) activity, and this regulation is involved in AMPK-dependent SIRT2 phosphorylation [37]. In addition, SIRT2 is a central regulator of the defense mechanism against reactive oxidative species (ROS) through FoxO3a deacetylation [36]. In the kidney, SIRT2 is mainly expressed in proximal epithelial tubular cells [34].

SIRT3 is the best characterized among mitochondrial sirtuins. SIRT3 is normally localized inside the mitochondria, but can translocate into the nucleus under stressful conditions, along with overexpression of SIRT5 [38]. SIRT3 is a NAD⁺-dependent deacetylase in mitochondria that regulates energy metabolism; the anti-oxidative defense system; lipid metabolism; and longevity via deacetylation of various substrates including acetyl-coenzyme A synthetase 2 (ACSS2) [39], glutamate dehydrogenase (GDH) [40], and superoxide dismutase 2 (SOD2) [41]. SIRT3 also reduces levels of ROS through regulation of NF- κ B [42], AMPK [43], PGC-1 α [44], and FoxO3a [45]. SIRT3 regulates the acetylation status of multiple proteins to improve function, including those associated with fatty acid oxidation, ketogenesis, amino acid catabolism, and urea and tricarboxylic acid cycles, in addition to mitochondrial regulation [46]. In the kidney, SIRT3 has been described as a crucial regulator of mitochondrial dynamics in proximal epithelial tubular cells [47].

SIRT4, the second mitochondrial sirtuin, has been observed in the mitochondrial matrix where it binds to proteins in similar pathways to those regulated by SIRT3. SIRT4 exhibits NAD⁺-dependent adenosine diphosphate (ADP)-ribosyltransferase activity and inhibits GDH activity through ADP-ribosylation [48]. SIRT4 is also a lysine deacetylase and can remove multiple derivatives of methylglutaryl from lysine residues to regulate leucine metabolism [49]. However, reduction of SIRT4 under conditions of nutrient deprivation coordinates the use of fatty acids as an energy source, suggesting that SIRT4 might antagonize the dietary restriction-mediated effects of SIRT1 and SIRT3. SIRT4 also suppresses fatty acid oxidation through inhibition of malonyl CoA decarboxylase (MCD) [50], peroxisome proliferator-activated receptor α (PPAR α) [51], and AMPK signaling [52]. SIRT4 also acts as a tumor suppressor, regulating the cellular metabolic response to DNA damage via repression of glutamine metabolism [53]. However, little is known about the role of SIRT4 in the kidney [54].

SIRT5, the third mitochondrial sirtuin, was thought to localize primarily to mitochondria [55]. However, several studies have demonstrated that SIRT5 is also present in the cytosol, peroxisomes, and nucleus [56][57][58]. SIRT5 was initially described as a mitochondrial deacetylase, but it has minimal deacetylase activity [59]. Recent studies have reported that SIRT5 shows distinct affinity for negatively charged acyl lysine modifications, and acts as a prominent cellular desuccinylase, demalonylase, and deglutarylase, thereby regulating various metabolic pathways [60]. Pathway analysis identified multiple target pathways of SIRT5 including fatty acid oxidation, branched-chain amino acid catabolism, the citric acid cycle, adenosine triphosphate (ATP) synthesis, oxidative phosphorylation (OXPHOS), stress responses, ketogenesis, and pyruvate metabolism [61]. The absence of SIRT5 led to hypersuccinylation of mitochondrial proteins in the kidney [62][63] and increased blood ammonia levels [64], suggesting that SIRT5 may play roles in renal and liver metabolism. A recent study revealed that SIRT5 is highly expressed in proximal epithelial tubular cells [65]. However, the function of SIRT5 in the kidney remains poorly understood.

SIRT6 is predominantly a nuclear protein and deacetylates diverse proteins, including H3K9 and H3K56 [66]. SIRT6 exhibits both deacetylase and mono-ADP-ribosyltransferase activities [67]. Like other sirtuins, SIRT6 regulates longevity via deacetylation of several transcription factors associated with DNA repair, glucose and lipid metabolism, cellular senescence, and inflammation [68][69][70][71]. SIRT6 also functions as a corepressor of HIF-1 α , suppressing glucose uptake and glycolysis [70]. In the kidney, SIRT6 plays an important role in podocyte injury and renal fibrosis in podocytes and proximal epithelial tubular cells [72][73][74].

SIRT7 is localized to the nucleus and has NAD⁺-dependent HDAC activity [75]. SIRT7 appears to exhibit relatively weak and substrate-specific deacetylase activity. The known deacetylation substrates of SIRT7 are only H3K18 (H3 Lys18), p53, polymerase-associated factor 53 (PAF53), the nucleolar protein nucleophosmin (NPM1), and GA-binding protein- β -1 (GABP- β 1) [76]. SIRT7 deficiency is associated with a high mutation rate, increased sensitivity to DNA damage, and apoptosis. These findings suggest that SIRT7 regulates genome stability through its effects on transcriptional regulation, DNA replication, and DNA repair [77][78]. In the kidney, SIRT7 is expressed in proximal tubules and collecting ducts [79]. However, the role of SIRT7 in kidney injury remains unclear.

3. The Role of Sirtuins in Renal Disease

3.1. Sirtuins in Acute Kidney Injury

Interest in the role of sirtuins in the pathophysiology of various renal diseases has grown recently. Given the tubular expression of sirtuins and their modulating effects on oxidative stress, inflammation, and mitochondrial dysfunction, they are likely to be involved in the pathogenesis of acute kidney injury (AKI) [80].

3.2. Sirtuins in the Aging Kidney

Aging is a multifactorial process characterized by progressive decline in physiological function. The kidney is a typical target organ of age-associated tissue damage, and the increased incidence of chronic kidney disease (CKD) in elderly is an emerging health problem worldwide [81]. Various sirtuins have been demonstrated to mitigate kidney aging. SIRT1 expression was found to be reduced in aging kidneys, and this change was associated with changes in the expression of other target molecules such as PGC-1 α /estrogen-related receptor-1 α (ERR-1 α), PPAR α , Klotho, and HIF-1 α [82][83][84]. Recently, podocyte-specific reduction of SIRT1 was found to accelerate kidney injury in aging mice [85]. Therefore, SIRT1 is believed as a potential target for treatment of kidney aging.

Caloric restriction has been shown to extend lifespan and may affect numerous cellular aspects of kidney aging. Kume et al. demonstrated that long-term caloric restriction from one to two years in mice promoted SIRT1 expression in aging kidneys, which resulted in attenuation of hypoxia-induced kidney injury via SIRT1-mediated deacetylation of FoxO3a and activation of autophagy. Even short-term activation of SIRT1 through caloric restriction promoted autophagy and reduced mitochondrial oxidative damage in 25-month-old rats [86]. SIRT1 and its target proteins may play an important role in renoprotection of aging kidneys, which is accomplished through stimulation of autophagy. Pharmacologically induced SIRT1 activation significantly reduced tubulointerstitial fibrosis and improved renal function through enhancement of Nrf2/HO-1 signaling and AMPK/PGC-1 α signaling [87].

Along with other sirtuins, SIRT3 has been reported to act as an essential regulator of cell senescence. SIRT3 is associated with renin–angiotensin–aldosterone system (RAAS) activation, which is known to play a role in kidney aging. In kidneys of aged mice, angiotensin II (Ang II) type 1 receptor (AT₁R) deletion upregulated nicotinamide phosphoribosyltransferase (Nampt) and SIRT3 and resulted in markedly prolonged lifespan. Ang II treatment downregulated SIRT3 expression in tubular epithelial cells, and this effect was inhibited by AT₁ antagonist administration. These findings suggest a biochemical link between Ang II and SIRT3 through AT₁R in aging kidneys [88]. However, a recent study reported conflicting results: this study found that expression of SIRT1 and Nampt expression, but not SIRT3, was significantly reduced in the kidneys of aged mice with AT₁R-associated protein (ATRAP) deletion [89]. Although the role of SIRT3 in kidney aging via the Ang II-AT₁R signaling pathway remains unclear, SIRT3 deficiency is known to cause severe renal fibrosis in aging kidneys associated with increased transforming growth factor- β 1 (TGF- β 1) expression and hyperacetylation of glycogen synthase kinase-3 β (GSK-3 β), resulting in phosphorylation of Smad3, c-Jun, and β -catenin [90]. SIRT6 activation due to caloric restriction also attenuated age-associated kidney injury through inhibition of the proinflammatory NF- κ B signaling pathway [91]. Together, these findings indicate that sirtuins play a role in attenuating tissue injury in aging kidneys, likely via

attenuation of oxidative stress and inflammation, and it supports that the investigation of sirtuins should be done as therapeutic targets for kidney aging.

3.3. Sirtuins in Diabetic Kidney Disease

Caloric restriction not only slows aging and increases lifespan, but also increases insulin sensitivity [92][93]. In a clinical study, fasting glucose levels and insulin resistance improved after a 12-week intensive weight reduction program based on caloric restriction among obese individuals with advanced diabetic nephropathy (DN), and also led to improvement in kidney function [94]. Dietary restriction in diabetic rat models increased SIRT1 expression in the kidneys and improved renal function including albuminuria, creatinine clearance, and renal histology [95][96]. Thus, caloric restriction activates sirtuins that may be beneficial in preventing the progression of DN.

Several studies have suggested that SIRT1 decreases mitochondrial oxidative stress and apoptosis through modulation of p53 [97], the AMPK/PGC-1 α pathway [98][99][100], the Nrf2 pathway [101], and the FoxO family [102], thereby providing protection against DN. SIRT1 also restrained renal inflammation and fibrosis under hyperglycemic conditions through HIF-1 α signaling in mesangial cells [103]. Podocyte-specific SIRT1 deletion in diabetic mice led to proteinuria and podocyte injury, and these changes were associated with renal inflammation due to hyperacetylation of STAT3/NF- κ B [104]. In addition, SIRT1 modulated angiogenesis through downregulation of vascular endothelial growth factor (VEGF) and Flk-1 (VEGFR-2) expression in high glucose (HG)-treated podocytes and endothelial cells, but these effects were attenuated by the genetic elimination of SIRT1 [105].

SIRT1 may participate in the crosstalk between podocytes and tubular cells in DN. Podocyte-specific SIRT1 deletion caused severe mesangial expansion and podocyte loss [104], while SIRT1 overexpression in podocytes attenuated renal damage in diabetic mice [106]. SIRT1 deletion in proximal tubules also increased albuminuria, which upregulated the tight junction protein claudin-1, in streptozotocin (STZ)-induced diabetic mice [107]. Exposure of podocytes to medium obtained from proximal tubular cells cultured with HG downregulated SIRT1 and upregulated claudin-1 expression. These effects were abolished in podocytes exposed to medium from proximal tubular cells overexpressing SIRT1, even under HG conditions [107]. Based on these findings, Hasegawa et al. proposed a functional relationship between proximal tubules and podocytes, referred to as 'proximal tubule–podocyte communication' [107].

Recent research has demonstrated a role for SIRT1 in proximal tubule–podocyte communication in association with sodium–glucose cotransporter 2 (SGLT2). In diabetic kidneys, high glucose levels around the proximal tubules may trigger glucose transporter 2 (GLUT2)-mediated intracellular glucose uptake via SGLT2 upregulation, causing a concomitant decrease in SIRT1. An SGLT2 inhibitor recovered SIRT1 expression in diabetic mice and HG-treated proximal tubular cells [108], as well as AMPK phosphorylation [109]. Interplay between AMPK/SIRT1 signaling and sodium transport mechanisms in the kidney may partially explain the role of the SGLT2 inhibitor in ameliorating the development of DN [110].

The associations of sirtuins other than SIRT1 with DN have also been explored. SIRT3 overexpression suppressed HG-induced apoptosis by reducing ROS accumulation through modulation of Akt/FoxO signaling in proximal tubular cells [111]. On the other hand, SIRT3 suppression was associated with activation of TGF- β /Smad3 signaling and increased HIF-1 α accumulation, which subsequently caused abnormal glycolysis and kidney fibrosis in diabetic mice and proximal tubular cells [112]. SIRT4 overexpression led to downregulated expression of apoptosis-related proteins such as NADPH oxidase 1 (NOX1), Bcl-2-associated X protein (Bax), and phosphorylated p38, along with upregulated expression of Bcl-2, which was associated with attenuation of the inflammatory response in HG-simulated podocytes [113].

SIRT6 deletion exacerbated podocyte injury in diabetic mice, and SIRT6 overexpression with HG treatment protected against podocyte injury through epigenetic regulation of Notch1 and Notch4 transcription due to deacetylation of H3K9. SIRT6 was also found to regulate the immune response by activating M2 macrophages, which are protective against podocyte injury, in STZ-induced diabetic mice [114]. In a recent study, selective deletion of Nampt in proximal tubule cells of STZ-induced diabetic mice led to downregulation of SIRT6, which was accompanied by thickening of the tubular basement membrane, type IV collagen deposition, enhanced renal fibrosis, and albuminuria. Selective deletion of SIRT6 in the proximal tubules of diabetic mice caused a phenotype similar to that of Nampt knockout mice. Therefore, the Nampt–Sirt6 axis in proximal tubules was suggested to be a key player in the fibrogenic extracellular matrix remodeling associated with DN [115].

3.4. Sirtuins in Chronic Kidney Disease

Renal tubular fibrosis is a major pathognomonic phenomenon in CKD [116], and sirtuins have been demonstrated to play an important role in tubular fibrosis. SIRT1 knockout mice were found to have prominent tubular fibrosis in a model of unilateral ureter obstruction (UUO) [117]. Suppression of SIRT1 expression in mouse renal medullary interstitial cells resulted in substantial reduction of cellular resistance to oxidative stress [117]. In addition, SIRT1 activation inhibited tubular fibrosis in a 5/6 nephrectomy model and a UUO model [118][119]. SIRT1 expression was also found to be involved in the pathogenesis of chronic renal allograft dysfunction and chronic cyclosporine A (CsA) nephropathy. In rat kidneys with chronic allograft dysfunction, decreased SIRT1 was associated with mononuclear cell infiltration and interstitial fibrosis due to upregulation of inflammatory cytokines [120]. In a mouse model of chronic CsA nephropathy, SIRT1 expression was reduced according to the degree of tubulointerstitial fibrosis through the Nrf2 and PI3K/Akt/FoxO1 pathways [121].

Various mechanisms have been suggested to underlie the pathogenetic link between SIRT1 and the development of renal fibrosis. Endothelial SIRT1 expression may play an important role, as SIRT1 deletion in the endothelium of mice caused spontaneous interstitial fibrosis without glomerular involvement, even at a young age. Moreover, tubulointerstitial fibrosis after long-term folic acid treatment was aggravated in mice with endothelium-specific SIRT1 deletion [122]. Endothelial SIRT1 depletion also enhanced the senescence of pericapillary tubular endothelial cells, which manifested as impaired endothelial proliferation and increased expression of molecules in the Notch1 signaling pathway [123]. A role for sirtuins in the epithelial-to-mesenchymal transition (EMT) during the development of renal fibrosis has also been suggested [124]. SIRT1 up-regulation by resveratrol treatment ameliorated renal

fibrosis in proximal tubular cells treated with TGF- β in addition to a mouse model of UUO; this was found to be due to inhibition of the EMT through deacetylation of Smad4 and inhibition of matrix metalloproteinase-7 (MMP-7) [125].

Consistent with observations in AKI, pharmacological inhibition of SIRT2 resulted in reduction of renal interstitial fibrosis in UUO models [126][127], which was accompanied by decreases in expression of epidermal growth factor receptor (EGFR), platelet-derived growth factor receptor- β (PDGFR- β), STAT3 [126], and E3-ubiquitin ligase murine double-minute 2 (MDM2) [127]. In contrast, SIRT3 plays an important role in the endothelial-to-mesenchymal transition (EndoMT) associated with the vascular pathology of renal fibrosis [128]. EndoMT is a novel mechanism of renal fibrosis and is characterized by a phenotypic transition from vascular endothelial cells to myofibroblasts [129]. Honokiol, a pharmaceutical SIRT3 activator, decreased renal inflammation and fibrosis through regulation of mitochondrial dynamics via the NF- κ B/TGF- β 1/Smad signaling pathway [130]. SIRT6 knockout aggravated TGF- β -induced fibrosis in mouse tubular epithelial cells, while pharmacological inhibition of SIRT6 deacetylase activity by OSS_128167 induced kidney fibrosis in a mouse model of UUO through modulation of the Wnt/ β -catenin signaling pathway. Therefore, various sirtuins appear to be involved in kidney fibrosis and related processes.

References

1. Finkel, T.; Deng, C.X.; Mostoslavsky, R. Recent progress in the biology and physiology of sirtuins. *Nature* 2009, 460, 587–591.
2. Kim, S.; Benguria, A.; Lai, C.Y.; Jazwinski, S.M. Modulation of life-span by histone deacetylase genes in *Saccharomyces cerevisiae*. *Mol. Biol. Cell* 1999, 10, 3125–3136.
3. Kaeberlein, M.; McVey, M.; Guarente, L. The SIR2/3/4 complex and SIR2 alone promote longevity in *Saccharomyces cerevisiae* by two different mechanisms. *Genes Dev.* 1999, 13, 2570–2580.
4. Smith, B.C.; Hallows, W.C.; Denu, J.M. Mechanisms and molecular probes of sirtuins. *Chem. Biol.* 2008, 15, 1002–1013.
5. Haigis, M.C.; Sinclair, D.A. Mammalian sirtuins: Biological insights and disease relevance. *Annu. Rev. Pathol.* 2010, 5, 253–295.
6. Tong, C.; Morrison, A.; Mattison, S.; Qian, S.; Bryniarski, M.; Rankin, B.; Wang, J.; Thomas, D.P.; Li, J. Impaired SIRT1 nucleocytoplasmic shuttling in the senescent heart during ischemic stress. *FASEB J.* 2013, 27, 4332–4342.
7. Tanno, M.; Sakamoto, J.; Miura, T.; Shimamoto, K.; Horio, Y. Nucleocytoplasmic shuttling of the NAD⁺-dependent histone deacetylase SIRT1. *J. Biol. Chem.* 2007, 282, 6823–6832.
8. Morigi, M.; Perico, L.; Benigni, A. Sirtuins in Renal Health and Disease. *J. Am. Soc. Nephrol.* 2018, 29, 1799–1809.

9. Rodgers, J.T.; Lerin, C.; Haas, W.; Gygi, S.P.; Spiegelman, B.M.; Puigserver, P. Nutrient control of glucose homeostasis through a complex of PGC-1 α and SIRT1. *Nature* 2005, 434, 113–118.
10. Kume, S.; Haneda, M.; Kanasaki, K.; Sugimoto, T.; Araki, S.; Isono, M.; Isshiki, K.; Uzu, T.; Kashiwagi, A.; Koya, D. Silent information regulator 2 (SIRT1) attenuates oxidative stress-induced mesangial cell apoptosis via p53 deacetylation. *Free Radic. Biol. Med.* 2006, 40, 2175–2182.
11. Khader, A.; Yang, W.L.; Kuncewitch, M.; Jacob, A.; Prince, J.M.; Asirvatham, J.R.; Nicastro, J.; Coppa, G.F.; Wang, P. Sirtuin 1 activation stimulates mitochondrial biogenesis and attenuates renal injury after ischemia-reperfusion. *Transplantation* 2014, 98, 148–156.
12. Kume, S.; Uzu, T.; Horiike, K.; Chin-Kanasaki, M.; Isshiki, K.; Araki, S.; Sugimoto, T.; Haneda, M.; Kashiwagi, A.; Koya, D. Calorie restriction enhances cell adaptation to hypoxia through Sirt1-dependent mitochondrial autophagy in mouse aged kidney. *J. Clin. Investig.* 2010, 120, 1043–1055.
13. Moynihan, K.A.; Grimm, A.A.; Plueger, M.M.; Bernal-Mizrachi, E.; Ford, E.; Cras-Meneur, C.; Permutt, M.A.; Imai, S. Increased dosage of mammalian Sir2 in pancreatic beta cells enhances glucose-stimulated insulin secretion in mice. *Cell Metab.* 2005, 2, 105–117.
14. Chen, D.; Bruno, J.; Easlon, E.; Lin, S.J.; Cheng, H.L.; Alt, F.W.; Guarente, L. Tissue-specific regulation of SIRT1 by calorie restriction. *Genes Dev.* 2008, 22, 1753–1757.
15. Xie, J.; Zhang, X.; Zhang, L. Negative regulation of inflammation by SIRT1. *Pharmacol. Res.* 2013, 67, 60–67.
16. Vaquero, A.; Scher, M.; Lee, D.; Erdjument-Bromage, H.; Tempst, P.; Reinberg, D. Human SirT1 interacts with histone H1 and promotes formation of facultative heterochromatin. *Mol. Cell* 2004, 16, 93–105.
17. Vaziri, H.; Dessain, S.K.; Ng Eaton, E.; Imai, S.I.; Frye, R.A.; Pandita, T.K.; Guarente, L.; Weinberg, R.A. hSIR2(SIRT1) functions as an NAD-dependent p53 deacetylase. *Cell* 2001, 107, 149–159.
18. Yeung, F.; Hoberg, J.E.; Ramsey, C.S.; Keller, M.D.; Jones, D.R.; Frye, R.A.; Mayo, M.W. Modulation of NF-kappaB-dependent transcription and cell survival by the SIRT1 deacetylase. *EMBO J.* 2004, 23, 2369–2380.
19. Bernier, M.; Paul, R.K.; Martin-Montalvo, A.; Scheibye-Knudsen, M.; Song, S.; He, H.J.; Armour, S.M.; Hubbard, B.P.; Bohr, V.A.; Wang, L.; et al. Negative regulation of STAT3 protein-mediated cellular respiration by SIRT1 protein. *J. Biol. Chem.* 2011, 286, 19270–19279.
20. Burnett, C.; Valentini, S.; Cabreiro, F.; Goss, M.; Somogyvari, M.; Piper, M.D.; Hoddinott, M.; Sutphin, G.L.; Leko, V.; McElwee, J.J.; et al. Absence of effects of Sir2 overexpression on lifespan in *C. elegans* and *Drosophila*. *Nature* 2011, 477, 482–485.

21. Lim, J.H.; Lee, Y.M.; Chun, Y.S.; Chen, J.; Kim, J.E.; Park, J.W. Sirtuin 1 modulates cellular responses to hypoxia by deacetylating hypoxia-inducible factor 1 α . *Mol. Cell* 2010, 38, 864–878.
22. Laemmle, A.; Lechleiter, A.; Roh, V.; Schwarz, C.; Portmann, S.; Furer, C.; Keogh, A.; Tschan, M.P.; Candinas, D.; Vorburger, S.A.; et al. Inhibition of SIRT1 impairs the accumulation and transcriptional activity of HIF-1 α protein under hypoxic conditions. *PLoS ONE* 2012, 7, e33433.
23. Yacoub, R.; Lee, K.; He, J.C. The Role of SIRT1 in Diabetic Kidney Disease. *Front. Endocrinol. (Lausanne)* 2014, 5, 166.
24. Scarpulla, R.C. Metabolic control of mitochondrial biogenesis through the PGC-1 family regulatory network. *Biochim. Biophys. Acta* 2011, 1813, 1269–1278.
25. Price, N.L.; Gomes, A.P.; Ling, A.J.; Duarte, F.V.; Martin-Montalvo, A.; North, B.J.; Agarwal, B.; Ye, L.; Ramadori, G.; Teodoro, J.S.; et al. SIRT1 is required for AMPK activation and the beneficial effects of resveratrol on mitochondrial function. *Cell Metab.* 2012, 15, 675–690.
26. Cetrullo, S.; D’Adamo, S.; Tantini, B.; Borzi, R.M.; Flamigni, F. mTOR, AMPK, and Sirt1: Key Players in Metabolic Stress Management. *Crit. Rev. Eukaryot. Gene Expr.* 2015, 25, 59–75.
27. Zhou, Y.; Song, T.; Peng, J.; Zhou, Z.; Wei, H.; Zhou, R.; Jiang, S.; Peng, J. SIRT1 suppresses adipogenesis by activating Wnt/ β -catenin signaling in vivo and in vitro. *Oncotarget* 2016, 7, 77707–77720.
28. North, B.J.; Marshall, B.L.; Borra, M.T.; Denu, J.M.; Verdin, E. The human Sir2 ortholog, SIRT2, is an NAD⁺-dependent tubulin deacetylase. *Mol. Cell* 2003, 11, 437–444.
29. Jing, H.; Lin, H. Sirtuins in epigenetic regulation. *Chem. Rev.* 2015, 115, 2350–2375.
30. Vaquero, A.; Scher, M.B.; Lee, D.H.; Sutton, A.; Cheng, H.L.; Alt, F.W.; Serrano, L.; Sternglanz, R.; Reinberg, D. SirT2 is a histone deacetylase with preference for histone H4 Lys 16 during mitosis. *Genes Dev.* 2006, 20, 1256–1261.
31. Das, C.; Lucia, M.S.; Hansen, K.C.; Tyler, J.K. CBP/p300-mediated acetylation of histone H3 on lysine 56. *Nature* 2009, 459, 113–117.
32. Serrano, L.; Martinez-Redondo, P.; Marazuela-Duque, A.; Vazquez, B.N.; Dooley, S.J.; Voigt, P.; Beck, D.B.; Kane-Goldsmith, N.; Tong, Q.; Rabanal, R.M., et al. The tumor suppressor SirT2 regulates cell cycle progression and genome stability by modulating the mitotic deposition of H4K20 methylation. *Genes Dev.* 2013, 27, 639–653.
33. Zhang, M.; Pan, Y.; Dorfman, R.G.; Yin, Y.; Zhou, Q.; Huang, S.; Liu, J.; Zhao, S. Sirtinol promotes PEPCK1 degradation and inhibits gluconeogenesis by inhibiting deacetylase SIRT2. *Sci. Rep.* 2017, 7, 7.
34. Jung, Y.J.; Lee, A.S.; Nguyen-Thanh, T.; Kim, D.; Kang, K.P.; Lee, S.; Park, S.K.; Kim, W. SIRT2 Regulates LPS-Induced Renal Tubular CXCL2 and CCL2 Expression. *J. Am. Soc. Nephrol.* 2015,

- 26, 1549–1560.
35. Jing, E.; Gesta, S.; Kahn, C.R. SIRT2 regulates adipocyte differentiation through FoxO1 acetylation/deacetylation. *Cell Metab.* 2007, 6, 105–114.
 36. Wang, F.; Nguyen, M.; Qin, F.X.; Tong, Q. SIRT2 deacetylates FOXO3a in response to oxidative stress and caloric restriction. *Aging Cell* 2007, 6, 505–514.
 37. Ramakrishnan, G.; Davaakhuu, G.; Kaplun, L.; Chung, W.C.; Rana, A.; Atfi, A.; Miele, L.; Tzivion, G. Sirt2 deacetylase is a novel AKT binding partner critical for AKT activation by insulin. *J. Biol. Chem.* 2014, 289, 6054–6066.
 38. Scher, M.B.; Vaquero, A.; Reinberg, D. SirT3 is a nuclear NAD⁺-dependent histone deacetylase that translocates to the mitochondria upon cellular stress. *Genes Dev.* 2007, 21, 920–928.
 39. Hirschey, M.D.; Shimazu, T.; Capra, J.A.; Pollard, K.S.; Verdin, E. SIRT1 and SIRT3 deacetylate homologous substrates: AceCS1,2 and HMGCS1,2. *Aging (Albany NY)* 2011, 3, 635–642.
 40. Hebert, A.S.; Dittenhafer-Reed, K.E.; Yu, W.; Bailey, D.J.; Selen, E.S.; Boersma, M.D.; Carson, J.J.; Tonelli, M.; Balloon, A.J.; Higbee, A.J.; et al. Calorie restriction and SIRT3 trigger global reprogramming of the mitochondrial protein acetylome. *Mol. Cell* 2013, 49, 186–199.
 41. Qiu, X.; Brown, K.; Hirschey, M.D.; Verdin, E.; Chen, D. Calorie restriction reduces oxidative stress by SIRT3-mediated SOD2 activation. *Cell Metab.* 2010, 12, 662–667.
 42. Chen, C.J.; Fu, Y.C.; Yu, W.; Wang, W. SIRT3 protects cardiomyocytes from oxidative stress-mediated cell death by activating NF-kappaB. *Biochem. Biophys. Res. Commun.* 2013, 430, 798–803.
 43. Pillai, V.B.; Sundaresan, N.R.; Kim, G.; Gupta, M.; Rajamohan, S.B.; Pillai, J.B.; Samant, S.; Ravindra, P.V.; Isbatan, A.; Gupta, M.P. Exogenous NAD blocks cardiac hypertrophic response via activation of the SIRT3-LKB1-AMP-activated kinase pathway. *J. Biol. Chem.* 2010, 285, 3133–3144.
 44. Shi, T.; Wang, F.; Stieren, E.; Tong, Q. SIRT3, a mitochondrial sirtuin deacetylase, regulates mitochondrial function and thermogenesis in brown adipocytes. *J. Biol. Chem.* 2005, 280, 13560–13567.
 45. Sundaresan, N.R.; Gupta, M.; Kim, G.; Rajamohan, S.B.; Isbatan, A.; Gupta, M.P. Sirt3 blocks the cardiac hypertrophic response by augmenting Foxo3a-dependent antioxidant defense mechanisms in mice. *J. Clin. Investig.* 2009, 119, 2758–2771.
 46. Rardin, M.J.; Newman, J.C.; Held, J.M.; Cusack, M.P.; Sorensen, D.J.; Li, B.; Schilling, B.; Mooney, S.D.; Kahn, C.R.; Verdin, E.; et al. Label-free quantitative proteomics of the lysine acetylome in mitochondria identifies substrates of SIRT3 in metabolic pathways. *Proc. Natl. Acad. Sci. USA* 2013, 110, 6601–6606.

47. Koyama, T.; Kume, S.; Koya, D.; Araki, S.; Isshiki, K.; Chin-Kanasaki, M.; Sugimoto, T.; Haneda, M.; Sugaya, T.; Kashiwagi, A.; et al. SIRT3 attenuates palmitate-induced ROS production and inflammation in proximal tubular cells. *Free Radic. Biol. Med.* 2011, 51, 1258–1267.
48. Haigis, M.C.; Mostoslavsky, R.; Haigis, K.M.; Fahie, K.; Christodoulou, D.C.; Murphy, A.J.; Valenzuela, D.M.; Yancopoulos, G.D.; Karow, M.; Blander, G.; et al. SIRT4 inhibits glutamate dehydrogenase and opposes the effects of calorie restriction in pancreatic beta cells. *Cell* 2006, 126, 941–954.
49. Anderson, K.A.; Huynh, F.K.; Fisher-Wellman, K.; Stuart, J.D.; Peterson, B.S.; Douros, J.D.; Wagner, G.R.; Thompson, J.W.; Madsen, A.S.; Green, M.F.; et al. SIRT4 Is a Lysine Deacylase that Controls Leucine Metabolism and Insulin Secretion. *Cell Metab.* 2017, 25, 838–855.e815.
50. Laurent, G.; German, N.J.; Saha, A.K.; de Boer, V.C.; Davies, M.; Koves, T.R.; Dephoure, N.; Fischer, F.; Boanca, G.; Vaitheesvaran, B.; et al. SIRT4 coordinates the balance between lipid synthesis and catabolism by repressing malonyl CoA decarboxylase. *Mol. Cell* 2013, 50, 686–698.
51. Laurent, G.; de Boer, V.C.; Finley, L.W.; Sweeney, M.; Lu, H.; Schug, T.T.; Cen, Y.; Jeong, S.M.; Li, X.; Sauve, A.A.; et al. SIRT4 represses peroxisome proliferator-activated receptor α activity to suppress hepatic fat oxidation. *Mol. Cell Biol.* 2013, 33, 4552–4561.
52. Ho, L.; Titus, A.S.; Banerjee, K.K.; George, S.; Lin, W.; Deota, S.; Saha, A.K.; Nakamura, K.; Gut, P.; Verdin, E.; et al. SIRT4 regulates ATP homeostasis and mediates a retrograde signaling via AMPK. *Aging (Albany NY)* 2013, 5, 835–849.
53. Jeong, S.M.; Xiao, C.; Finley, L.W.; Lahusen, T.; Souza, A.L.; Pierce, K.; Li, Y.H.; Wang, X.; Laurent, G.; German, N.J.; et al. SIRT4 has tumor-suppressive activity and regulates the cellular metabolic response to DNA damage by inhibiting mitochondrial glutamine metabolism. *Cancer Cell* 2013, 23, 450–463.
54. Hershberger, K.A.; Martin, A.S.; Hirschey, M.D. Role of NAD(+) and mitochondrial sirtuins in cardiac and renal diseases. *Nat. Rev. Nephrol.* 2017, 13, 213–225.
55. Schlicker, C.; Gertz, M.; Papatheodorou, P.; Kachholz, B.; Becker, C.F.; Steegborn, C. Substrates and regulation mechanisms for the human mitochondrial sirtuins Sirt3 and Sirt5. *J. Mol. Biol.* 2008, 382, 790–801.
56. Park, J.; Chen, Y.; Tishkoff, D.X.; Peng, C.; Tan, M.; Dai, L.; Xie, Z.; Zhang, Y.; Zwaans, B.M.; Skinner, M.E.; et al. SIRT5-mediated lysine desuccinylation impacts diverse metabolic pathways. *Mol. Cell* 2013, 50, 919–930.
57. Chen, X.F.; Tian, M.X.; Sun, R.Q.; Zhang, M.L.; Zhou, L.S.; Jin, L.; Chen, L.L.; Zhou, W.J.; Duan, K.L.; Chen, Y.J.; et al. SIRT5 inhibits peroxisomal ACOX1 to prevent oxidative damage and is downregulated in liver cancer. *EMBO Rep.* 2018, 19, e45124.

58. de Moura, M.B.; Uppala, R.; Zhang, Y.; Van Houten, B.; Goetzman, E.S. Overexpression of mitochondrial sirtuins alters glycolysis and mitochondrial function in HEK293 cells. *PLoS ONE* 2014, 9, e106028.
59. Du, J.; Zhou, Y.; Su, X.; Yu, J.J.; Khan, S.; Jiang, H.; Kim, J.; Woo, J.; Kim, J.H.; Choi, B.H.; et al. Sirt5 is a NAD-dependent protein lysine demalonylase and desuccinylase. *Science* 2011, 334, 806–809.
60. Kumar, S.; Lombard, D.B. Functions of the sirtuin deacylase SIRT5 in normal physiology and pathobiology. *Crit. Rev. Biochem. Mol. Biol.* 2018, 53, 311–334.
61. Rardin, M.J.; He, W.; Nishida, Y.; Newman, J.C.; Carrico, C.; Danielson, S.R.; Guo, A.; Gut, P.; Sahu, A.K.; Li, B.; et al. SIRT5 regulates the mitochondrial lysine succinylome and metabolic networks. *Cell Metab.* 2013, 18, 920–933.
62. Tan, M.; Peng, C.; Anderson, K.A.; Chhoy, P.; Xie, Z.; Dai, L.; Park, J.; Chen, Y.; Huang, H.; Zhang, Y.; et al. Lysine glutarylation is a protein posttranslational modification regulated by SIRT5. *Cell Metab.* 2014, 19, 605–617.
63. Nishida, Y.; Rardin, M.J.; Carrico, C.; He, W.; Sahu, A.K.; Gut, P.; Najjar, R.; Fitch, M.; Hellerstein, M.; Gibson, B.W.; et al. SIRT5 Regulates both Cytosolic and Mitochondrial Protein Malonylation with Glycolysis as a Major Target. *Mol. Cell* 2015, 59, 321–332.
64. Yu, J.; Sadhukhan, S.; Noriega, L.G.; Moullan, N.; He, B.; Weiss, R.S.; Lin, H.; Schoonjans, K.; Auwerx, J. Metabolic characterization of a Sirt5 deficient mouse model. *Sci. Rep.* 2013, 3, 2806.
65. Chiba, T.; Peasley, K.D.; Cargill, K.R.; Maringer, K.V.; Bharathi, S.S.; Mukherjee, E.; Zhang, Y.; Holtz, A.; Basisty, N.; Yagobian, S.D.; et al. Sirtuin 5 Regulates Proximal Tubule Fatty Acid Oxidation to Protect against AKI. *J. Am. Soc. Nephrol.* 2019, 30, 2384–2398.
66. Sosnowska, B.; Mazidi, M.; Penson, P.; Gluba-Brzozka, A.; Rysz, J.; Banach, M. The sirtuin family members SIRT1, SIRT3 and SIRT6: Their role in vascular biology and atherogenesis. *Atherosclerosis* 2017, 265, 275–282.
67. Liszt, G.; Ford, E.; Kurtev, M.; Guarente, L. Mouse Sir2 homolog SIRT6 is a nuclear ADP-ribosyltransferase. *J. Biol. Chem.* 2005, 280, 21313–21320.
68. Masri, S.; Rigor, P.; Cervantes, M.; Ceglia, N.; Sebastian, C.; Xiao, C.; Roqueta-Rivera, M.; Deng, C.; Osborne, T.F.; Mostoslavsky, R.; et al. Partitioning circadian transcription by SIRT6 leads to segregated control of cellular metabolism. *Cell* 2014, 158, 659–672.
69. Lerrer, B.; Gertler, A.A.; Cohen, H.Y. The complex role of SIRT6 in carcinogenesis. *Carcinogenesis* 2016, 37, 108–118.
70. Zhong, L.; D’Urso, A.; Toiber, D.; Sebastian, C.; Henry, R.E.; Vadysirisack, D.D.; Guimaraes, A.; Marinelli, B.; Wikstrom, J.D.; Nir, T.; et al. The histone deacetylase Sirt6 regulates glucose

homeostasis via Hif1 α . *Cell* 2010, 140, 280–293.

71. Sundaresan, N.R.; Vasudevan, P.; Zhong, L.; Kim, G.; Samant, S.; Parekh, V.; Pillai, V.B.; Ravindra, P.V.; Gupta, M.; Jeevanandam, V.; et al. The sirtuin SIRT6 blocks IGF-Akt signaling and development of cardiac hypertrophy by targeting c-Jun. *Nat. Med.* 2012, 18, 1643–1650.
72. Huang, W.; Liu, H.; Zhu, S.; Woodson, M.; Liu, R.; Tilton, R.G.; Miller, J.D.; Zhang, W. Sirt6 deficiency results in progression of glomerular injury in the kidney. *Aging (Albany NY)* 2017, 9, 1069–1083.
73. Liu, M.; Liang, K.; Zhen, J.; Zhou, M.; Wang, X.; Wang, Z.; Wei, X.; Zhang, Y.; Sun, Y.; Zhou, Z.; et al. Sirt6 deficiency exacerbates podocyte injury and proteinuria through targeting Notch signaling. *Nat. Commun.* 2017, 8, 413.
74. Cai, J.; Liu, Z.; Huang, X.; Shu, S.; Hu, X.; Zheng, M.; Tang, C.; Liu, Y.; Chen, G.; Sun, L.; et al. The deacetylase sirtuin 6 protects against kidney fibrosis by epigenetically blocking β -catenin target gene expression. *Kidney Int.* 2020, 97, 106–118.
75. Ford, E.; Voit, R.; Liszt, G.; Magin, C.; Grummt, I.; Guarente, L. Mammalian Sir2 homolog SIRT7 is an activator of RNA polymerase I transcription. *Genes Dev.* 2006, 20, 1075–1080.
76. Kiran, S.; Anwar, T.; Kiran, M.; Ramakrishna, G. Sirtuin 7 in cell proliferation, stress and disease: Rise of the Seventh Sirtuin! *Cell. Signal.* 2015, 27, 673–682.
77. Vazquez, B.N.; Thackray, J.K.; Serrano, L. Sirtuins and DNA damage repair: SIRT7 comes to play. *Nucleus* 2017, 8, 107–115.
78. Li, L.; Shi, L.; Yang, S.; Yan, R.; Zhang, D.; Yang, J.; He, L.; Li, W.; Yi, X.; Sun, L.; et al. SIRT7 is a histone desuccinylase that functionally links to chromatin compaction and genome stability. *Nat. Commun.* 2016, 7, 12235.
79. Miyasato, Y.; Yoshizawa, T.; Sato, Y.; Nakagawa, T.; Miyasato, Y.; Kakizoe, Y.; Kuwabara, T.; Adachi, M.; Ianni, A.; Braun, T.; et al. Sirtuin 7 Deficiency Ameliorates Cisplatin-induced Acute Kidney Injury Through Regulation of the Inflammatory Response. *Sci. Rep.* 2018, 8, 5927.
80. Wakino, S.; Hasegawa, K.; Itoh, H. Sirtuin and metabolic kidney disease. *Kidney Int.* 2015, 88, 691–698.
81. Martin, J.E.; Sheaff, M.T. Renal ageing. *J. Pathol.* 2007, 211, 198–205.
82. Lim, J.H.; Kim, E.N.; Kim, M.Y.; Chung, S.; Shin, S.J.; Kim, H.W.; Yang, C.W.; Kim, Y.S.; Chang, Y.S.; Park, C.W.; et al. Age-associated molecular changes in the kidney in aged mice. *Oxidative Med. Cell. Longev.* 2012, 2012, 171383.
83. Ryu, D.R.; Yu, M.R.; Kong, K.H.; Kim, H.; Kwon, S.H.; Jeon, J.S.; Han, D.C.; Noh, H. Sirt1-hypoxia-inducible factor-1 α interaction is a key mediator of tubulointerstitial damage in the aged kidney. *Aging Cell* 2019, 18, e12904.

84. Hsu, S.C.; Huang, S.M.; Chen, A.; Sun, C.Y.; Lin, S.H.; Chen, J.S.; Liu, S.T.; Hsu, Y.J. Resveratrol increases anti-aging Klotho gene expression via the activating transcription factor 3/c-Jun complex-mediated signaling pathway. *Int. J. Biochem. Cell Biol.* 2014, 53, 361–371.
85. Chuang, P.Y.; Cai, W.; Li, X.; Fang, L.; Xu, J.; Yacoub, R.; He, J.C.; Lee, K. Reduction in podocyte SIRT1 accelerates kidney injury in aging mice. *Am. J. Physiol. Renal Physiol.* 2017, 313, F621–f628.
86. Ning, Y.C.; Cai, G.Y.; Zhuo, L.; Gao, J.J.; Dong, D.; Cui, S.; Feng, Z.; Shi, S.Z.; Bai, X.Y.; Sun, X.F.; et al. Short-term calorie restriction protects against renal senescence of aged rats by increasing autophagic activity and reducing oxidative damage. *Mech. Ageing Dev.* 2013, 134, 570–579.
87. Kim, E.N.; Lim, J.H.; Kim, M.Y.; Ban, T.H.; Jang, I.A.; Yoon, H.E.; Park, C.W.; Chang, Y.S.; Choi, B.S. Resveratrol, an Nrf2 activator, ameliorates aging-related progressive renal injury. *Aging (Albany NY)* 2018, 10, 83–99.
88. Benigni, A.; Corna, D.; Zoja, C.; Sonzogno, A.; Latini, R.; Salio, M.; Conti, S.; Rottoli, D.; Longaretti, L.; Cassis, P.; et al. Disruption of the Ang II type 1 receptor promotes longevity in mice. *J. Clin. Investig.* 2009, 119, 524–530.
89. Uneda, K.; Wakui, H.; Maeda, A.; Azushima, K.; Kobayashi, R.; Haku, S.; Ohki, K.; Haruhara, K.; Kinguchi, S.; Matsuda, M.; et al. Angiotensin II Type 1 Receptor-Associated Protein Regulates Kidney Aging and Lifespan Independent of Angiotensin. *J. Am. Heart Assoc.* 2017, 6, e006120
90. Sundaresan, N.R.; Bindu, S.; Pillai, V.B.; Samant, S.; Pan, Y.; Huang, J.Y.; Gupta, M.; Nagalingam, R.S.; Wolfgeher, D.; Verdin, E.; et al. SIRT3 Blocks Aging-Associated Tissue Fibrosis in Mice by Deacetylating and Activating Glycogen Synthase Kinase 3 β . *Mol. Cell Biol.* 2015, 36, 678–692.
91. Zhang, N.; Li, Z.; Mu, W.; Li, L.; Liang, Y.; Lu, M.; Wang, Z.; Qiu, Y.; Wang, Z. Calorie restriction-induced SIRT6 activation delays aging by suppressing NF- κ B signaling. *Cell Cycle* 2016, 15, 1009–1018.
92. Anson, R.M.; Guo, Z.; de Cabo, R.; Iyun, T.; Rios, M.; Hagepanos, A.; Ingram, D.K.; Lane, M.A.; Mattson, M.P. Intermittent fasting dissociates beneficial effects of dietary restriction on glucose metabolism and neuronal resistance to injury from calorie intake. *Proc. Natl. Acad. Sci. USA* 2003, 100, 6216–6220.
93. Svendsen, P.F.; Jensen, F.K.; Holst, J.J.; Haugaard, S.B.; Nilas, L.; Madsbad, S. The effect of a very low calorie diet on insulin sensitivity, beta cell function, insulin clearance, incretin hormone secretion, androgen levels and body composition in obese young women. *Scand. J. Clin. Lab. Investig.* 2012, 72, 410–419.

94. Friedman, A.N.; Chambers, M.; Kamendulis, L.M.; Temmerman, J. Short-term changes after a weight reduction intervention in advanced diabetic nephropathy. *Clin. J. Am. Soc. Nephrol.* 2013, 8, 1892–1898.
95. Tikoo, K.; Tripathi, D.N.; Kabra, D.G.; Sharma, V.; Gaikwad, A.B. Intermittent fasting prevents the progression of type I diabetic nephropathy in rats and changes the expression of Sir2 and p53. *FEBS Lett.* 2007, 581, 1071–1078.
96. Kitada, M.; Takeda, A.; Nagai, T.; Ito, H.; Kanasaki, K.; Koya, D. Dietary restriction ameliorates diabetic nephropathy through anti-inflammatory effects and regulation of the autophagy via restoration of Sirt1 in diabetic Wistar fatty (fa/fa) rats: A model of type 2 diabetes. *Exp. Diabetes Res.* 2011, 2011, 908185.
97. Wang, X.L.; Wu, L.Y.; Zhao, L.; Sun, L.N.; Liu, H.Y.; Liu, G.; Guan, G.J. SIRT1 activator ameliorates the renal tubular injury induced by hyperglycemia in vivo and in vitro via inhibiting apoptosis. *Biomed. Pharmacother.* 2016, 83, 41–50.
98. Kitada, M.; Kume, S.; Imaizumi, N.; Koya, D. Resveratrol improves oxidative stress and protects against diabetic nephropathy through normalization of Mn-SOD dysfunction in AMPK/SIRT1-independent pathway. *Diabetes* 2011, 60, 634–643.
99. Kim, M.Y.; Lim, J.H.; Youn, H.H.; Hong, Y.A.; Yang, K.S.; Park, H.S.; Chung, S.; Ko, S.H.; Shin, S.J.; Choi, B.S.; et al. Resveratrol prevents renal lipotoxicity and inhibits mesangial cell glucotoxicity in a manner dependent on the AMPK-SIRT1-PGC-1 α axis in db/db mice. *Diabetologia* 2013, 56, 204–217.
100. Park, H.S.; Lim, J.H.; Kim, M.Y.; Kim, Y.; Hong, Y.A.; Choi, S.R.; Chung, S.; Kim, H.W.; Choi, B.S.; Kim, Y.S.; et al. Resveratrol increases AdipoR1 and AdipoR2 expression in type 2 diabetic nephropathy. *J. Transl. Med.* 2016, 14, 176.
101. Huang, K.; Huang, J.; Xie, X.; Wang, S.; Chen, C.; Shen, X.; Liu, P.; Huang, H. Sirt1 resists advanced glycation end products-induced expressions of fibronectin and TGF- β 1 by activating the Nrf2/ARE pathway in glomerular mesangial cells. *Free Radic. Biol. Med.* 2013, 65, 528–540.
102. Wu, L.; Zhang, Y.; Ma, X.; Zhang, N.; Qin, G. The effect of resveratrol on FoxO1 expression in kidneys of diabetic nephropathy rats. *Mol. Biol. Rep.* 2012, 39, 9085–9093.
103. Shao, Y.; Lv, C.; Wu, C.; Zhou, Y.; Wang, Q. Mir-217 promotes inflammation and fibrosis in high glucose cultured rat glomerular mesangial cells via Sirt1/HIF-1 α signaling pathway. *Diabetes Metab. Res. Rev.* 2016, 32, 534–543.
104. Liu, R.; Zhong, Y.; Li, X.; Chen, H.; Jim, B.; Zhou, M.M.; Chuang, P.Y.; He, J.C. Role of transcription factor acetylation in diabetic kidney disease. *Diabetes* 2014, 63, 2440–2453.
105. Wen, D.; Huang, X.; Zhang, M.; Zhang, L.; Chen, J.; Gu, Y.; Hao, C.M. Resveratrol attenuates diabetic nephropathy via modulating angiogenesis. *PLoS ONE* 2013, 8, e82336.

106. Hong, Q.; Zhang, L.; Das, B.; Li, Z.; Liu, B.; Cai, G.; Chen, X.; Chuang, P.Y.; He, J.C.; Lee, K. Increased podocyte Sirtuin-1 function attenuates diabetic kidney injury. *Kidney Int.* 2018, 93, 1330–1343.
107. Hasegawa, K.; Wakino, S.; Simic, P.; Sakamaki, Y.; Minakuchi, H.; Fujimura, K.; Hosoya, K.; Komatsu, M.; Kaneko, Y.; Kanda, T.; et al. Renal tubular Sirt1 attenuates diabetic albuminuria by epigenetically suppressing Claudin-1 overexpression in podocytes. *Nat. Med.* 2013, 19, 1496–1504.
108. Umino, H.; Hasegawa, K.; Minakuchi, H.; Muraoka, H.; Kawaguchi, T.; Kanda, T.; Tokuyama, H.; Wakino, S.; Itoh, H. High Basolateral Glucose Increases Sodium-Glucose Cotransporter 2 and Reduces Sirtuin-1 in Renal Tubules through Glucose Transporter-2 Detection. *Sci. Rep.* 2018, 8, 6791.
109. Hawley, S.A.; Ford, R.J.; Smith, B.K.; Gowans, G.J.; Mancini, S.J.; Pitt, R.D.; Day, E.A.; Salt, I.P.; Steinberg, G.R.; Hardie, D.G. The Na⁺/Glucose Cotransporter Inhibitor Canagliflozin Activates AMPK by Inhibiting Mitochondrial Function and Increasing Cellular AMP Levels. *Diabetes* 2016, 65, 2784–2794.
110. Packer, M. Interplay of adenosine monophosphate-activated protein kinase/sirtuin-1 activation and sodium influx inhibition mediates the renal benefits of sodium-glucose co-transporter-2 inhibitors in type 2 diabetes: A novel conceptual framework. *Diabetes Obes. Metab.* 2020, 22, 734–742.
111. Jiao, X.; Li, Y.; Zhang, T.; Liu, M.; Chi, Y. Role of Sirtuin3 in high glucose-induced apoptosis in renal tubular epithelial cells. *Biochem. Biophys. Res. Commun.* 2016, 480, 387–393.
112. Srivastava, S.P.; Li, J.; Kitada, M.; Fujita, H.; Yamada, Y.; Goodwin, J.E.; Kanasaki, K.; Koya, D. SIRT3 deficiency leads to induction of abnormal glycolysis in diabetic kidney with fibrosis. *Cell Death Dis.* 2018, 9, 997.
113. Shi, J.X.; Wang, Q.J.; Li, H.; Huang, Q. SIRT4 overexpression protects against diabetic nephropathy by inhibiting podocyte apoptosis. *Exp. Ther. Med.* 2017, 13, 342–348.
114. Ji, L.; Chen, Y.; Wang, H.; Zhang, W.; He, L.; Wu, J.; Liu, Y. Overexpression of Sirt6 promotes M2 macrophage transformation, alleviating renal injury in diabetic nephropathy. *Int. J. Oncol.* 2019, 55, 103–115.
115. Muraoka, H.; Hasegawa, K.; Sakamaki, Y.; Minakuchi, H.; Kawaguchi, T.; Yasuda, I.; Kanda, T.; Tokuyama, H.; Wakino, S.; Itoh, H. Role of Nampt-Sirt6 Axis in Renal Proximal Tubules in Extracellular Matrix Deposition in Diabetic Nephropathy. *Cell Rep.* 2019, 27, 199–212.e195.
116. Humphreys, B.D. Mechanisms of Renal Fibrosis. *Annu. Rev. Physiol.* 2018, 80, 309–326.
117. He, W.; Wang, Y.; Zhang, M.Z.; You, L.; Davis, L.S.; Fan, H.; Yang, H.C.; Fogo, A.B.; Zent, R.; Harris, R.C.; et al. Sirt1 activation protects the mouse renal medulla from oxidative injury. *J. Clin.*

- Investig. 2010, 120, 1056–1068.
118. Li, J.; Qu, X.; Ricardo, S.D.; Bertram, J.F.; Nikolic-Paterson, D.J. Resveratrol inhibits renal fibrosis in the obstructed kidney: Potential role in deacetylation of Smad3. *Am. J. Pathol.* 2010, 177, 1065–1071.
 119. Huang, X.Z.; Wen, D.; Zhang, M.; Xie, Q.; Ma, L.; Guan, Y.; Ren, Y.; Chen, J.; Hao, C.M. Sirt1 activation ameliorates renal fibrosis by inhibiting the TGF- β /Smad3 pathway. *J. Cell Biochem.* 2014, 115, 996–1005.
 120. Xia, Y.; Deng, J.; Zhou, Q.; Shao, X.; Yang, X.; Sha, M.; Zou, H. Expression and significance of Sirt1 in renal allografts at the early stage of chronic renal allograft dysfunction. *Transpl. Immunol.* 2018, 48, 18–25.
 121. Koh, E.S.; Kim, S.; Kim, M.; Hong, Y.A.; Shin, S.J.; Park, C.W.; Chang, Y.S.; Chung, S.; Kim, H.S. D-Pinitol alleviates cyclosporine A-induced renal tubulointerstitial fibrosis via activating Sirt1 and Nrf2 antioxidant pathways. *Int. J. Mol. Med.* 2018, 41, 1826–1834.
 122. Vasko, R.; Xavier, S.; Chen, J.; Lin, C.H.; Ratliff, B.; Rabadi, M.; Maizel, J.; Tanokuchi, R.; Zhang, F.; Cao, J.; et al. Endothelial sirtuin 1 deficiency perpetrates nephrosclerosis through downregulation of matrix metalloproteinase-14: Relevance to fibrosis of vascular senescence. *J. Am. Soc. Nephrol.* 2014, 25, 276–291.
 123. Kida, Y.; Zullo, J.A.; Goligorsky, M.S. Endothelial sirtuin 1 inactivation enhances capillary rarefaction and fibrosis following kidney injury through Notch activation. *Biochem. Biophys. Res. Commun.* 2016, 478, 1074–1079.
 124. Kalluri, R.; Weinberg, R.A. The basics of epithelial-mesenchymal transition. *J. Clin. Investig.* 2009, 119, 1420–1428.
 125. Xiao, Z.; Chen, C.; Meng, T.; Zhang, W.; Zhou, Q. Resveratrol attenuates renal injury and fibrosis by inhibiting transforming growth factor- β pathway on matrix metalloproteinase 7. *Exp. Biol. Med.* (Maywood) 2016, 241, 140–146.
 126. Ponnusamy, M.; Zhou, X.; Yan, Y.; Tang, J.; Tolbert, E.; Zhao, T.C.; Gong, R.; Zhuang, S. Blocking sirtuin 1 and 2 inhibits renal interstitial fibroblast activation and attenuates renal interstitial fibrosis in obstructive nephropathy. *J. Pharmacol. Exp. Ther.* 2014, 350, 243–256.
 127. He, F.F.; You, R.Y.; Ye, C.; Lei, C.T.; Tang, H.; Su, H.; Zhang, C. Inhibition of SIRT2 Alleviates Fibroblast Activation and Renal Tubulointerstitial Fibrosis via MDM2. *Cell Physiol. Biochem.* 2018, 46, 451–460.
 128. Lin, J.R.; Zheng, Y.J.; Zhang, Z.B.; Shen, W.L.; Li, X.D.; Wei, T.; Ruan, C.C.; Chen, X.H.; Zhu, D.L.; Gao, P.J. Suppression of Endothelial-to-Mesenchymal Transition by SIRT (Sirtuin) 3 Alleviated the Development of Hypertensive Renal Injury. *Hypertension* 2018, 72, 350–360.

129. Cruz-Solbes, A.S.; Youker, K. Epithelial to Mesenchymal Transition (EMT) and Endothelial to Mesenchymal Transition (EndMT): Role and Implications in Kidney Fibrosis. *Results Probl. Cell Differ.* 2017, 60, 345–372.
 130. Quan, Y.; Park, W.; Jin, J.; Kim, W.; Park, S.K.; Kang, K.P. Sirtuin 3 Activation by Honokiol Decreases Unilateral Ureteral Obstruction-Induced Renal Inflammation and Fibrosis via Regulation of Mitochondrial Dynamics and the Renal NF- κ B-TGF- β 1/Smad Signaling Pathway. *Int. J. Mol. Sci.* 2020, 21, 402.
-

Retrieved from <https://encyclopedia.pub/entry/history/show/4976>