

Global Distribution of Geogenic High-Arsenic Groundwater

Subjects: **Water Resources**

Contributor: Jiju Guo , Wengeng Cao , Guohui Lang , Qifa Sun , Tian Nan , Xiangzhi Li , Yu Ren , Zeyan Li

Groundwater constitutes a vital source of freshwater, accounting for roughly 95% of the total available freshwater resources on Earth. It is utilized not only for daily water needs but also for agricultural irrigation, industrial purposes, ecological recharge, and power generation.

high-arsenic groundwater

worldwide scale

in situ remediation of arsenic

human health risk assessment

1. Introduction

Groundwater constitutes a vital source of freshwater, accounting for roughly 95% of the total available freshwater resources on Earth ^[1]. It is utilized not only for daily water needs but also for agricultural irrigation, industrial purposes, ecological recharge, and power generation ^[2]. Therefore, groundwater holds significant value as a resource and plays a critical role in the environment. The degradation of groundwater quality represents a significant issue within the context of global environmental and climate change today. Since the Industrial Revolution, there has been widespread concern over the deterioration of groundwater quality ^[3]. Among the various groundwater quality issues, the release of high concentrations of heavy metals has had a significant impact on groundwater quality, and serious consideration must be given to its potential risks and hazards to human health. In particular, As is considered by the United States Agency for Toxic Substances and Disease Registry (ATSDR) to be the pollutant that poses the highest potential risk to human health due to the release from natural sources and the resulting high geogenic concentrations in groundwater ^[4]. The sources of As in groundwater primarily include natural origins such as geological formations, volcanic activity, and hydrothermal processes, as well as anthropogenic activities including mining, coal combustion, and petroleum extraction ^[5]. The majority of global health issues caused by As are linked to the consumption of water with high As concentrations. Due to the wide range of negative effects of high As concentrations on human health, the World Health Organization (WHO), the United States, and the European Union (EU) have lowered the Maximum Contaminant Level (MCL) of As in drinking water from 50 µg/L to 10 µg/L as a safe limit for As concentration in drinking water ^{[6][7]}. High-As groundwater is defined as groundwater with As concentrations above the WHO drinking water standard. The enrichment of high-As groundwater is primarily influenced by a combination of natural sources and hydrogeochemical conditions, with the majority of natural high-As groundwater primarily being a result of geological arsenic contamination ^[5]. Despite the established risks, many countries, such as Bangladesh, Nepal, Pakistan, Mexico, and Argentina, continue to adopt the 50 µg/L standard for arsenic concentration in their national drinking

water guidelines, due to a lack of professional expertise, economic considerations, and the low-level arsenic detection technology [8].

Human exposure to As occurs through direct and indirect pathways. Direct exposure involves drinking water with a high As concentration, contact with skin, and inhalation of gasses with a high As concentration. Indirect exposure mainly occurs through the food chain; this includes eating crops, vegetables, and fruits cultivated in As-contaminated soil or irrigated with As-rich groundwater, as well as consuming meat products from animals raised in such environments. Prolonged exposure to As, regardless of the route, can result in serious health disorders affecting the skin, blood vessels, and nervous system. Extended periods of high As exposure also notably increase the risk of developing cancers in organs like the lungs, liver, kidneys, and skin [9][10] (Figure 1).




Figure 1. Different pathways of arsenic exposure in groundwater and effects on humans (MMA-Monomethylarsenite; DMA-Dimethylarsenite).

Environmental As exists in groundwater in both organic and inorganic forms, with varying levels of toxicity associated with different forms. The three primary forms of inorganic arsenic are as follows: pentavalent arsenate [As(V)], trivalent arsenite [As(III)], and metallic arsenic. Arsenic in organic form often occurs as various organic arsenic compounds such as Monomethylarsenite (MMA) and Dimethylarsenite (DMA) [11]. Among these, inorganic arsenic is more toxic to humans, and the toxicity significantly differs between the oxidation states of As(III) and As(V). The toxicity of As(III) is more than 60 times higher than that of As(V) and 70 times higher than that of methylated arsenic [12]. The heightened toxicity of As(III) is partially because of its reactivity towards biologically relevant molecules [13]. The methylated arsenic forms, including MMA and DMA, exhibit moderate toxicity, while other organic forms, such as arsenobetaine (AsB) and arsenocholine (AsC), are generally considered non-toxic [12]. In aqueous solutions, As(III) and As(V) primarily exist as oxyanions due to the high charge and small ion radius of As^{3+} and As^{5+} . The presence and dispersion of distinct arsenic compounds within hydrological systems are markedly influenced by both the redox potential and pH levels prevailing in aquatic environments [14]. Under circumstances characterized by moderate-to-high redox potentials, As tends to stabilize into the As(V) form (H_3AsO_4 , H_2AsO_4^- , HAsO_4^{2-} , or AsO_4^{3-}). Conversely, in environments featuring predominantly acidic or weakly alkaline reducing conditions, and lower redox potentials, As(III) tends to be prevalent as the uncharged H_3AsO_3 molecule [15].

2. Global Distribution of Geogenic High-Arsenic Groundwater

High-As groundwater is widespread worldwide. According to statistics, 107 countries are affected by high-As groundwater, with the highest number in Asia (32) and Europe (31), followed by Africa (20), North America (11), South America (9), and Australia (4) [16]. The most affected countries are Bangladesh, India, Pakistan, China, Nepal [7], Laos [17], Cambodia [18], Myanmar [19], Vietnam [20], and the United States. The world map (**Figure 2**) displays the global distribution of geogenic high-As groundwater, predominantly found in inland basins and river deltas in South Asian, East Asian, and South American countries. Major countries are shown in **Table 1**. Generally, the river-marine sedimentary shallow (Holocene) aquifers in the river deltas are the main areas where high-As groundwater occurs naturally, and it occurs mainly under reducing aquifer conditions [21].



Figure 2. Main countries worldwide affected by geogenic high-arsenic groundwater ($\geq 10 \mu\text{g/L}$) (taking the maximum arsenic concentration), see **Table 1** for specific data.

Globally, the problem of geogenic high-As groundwater is particularly prominent in South and Southeast Asia, especially in Bangladesh and India [22]. In Bangladesh, 61 areas have been identified as having high-As groundwater. The potential population at risk is approximately 20 million people [23]. According to the National Drinking Water Survey of Bangladesh, around 8% of the water samples had As levels exceeding the Bangladesh standard of $50 \mu\text{g/L}$, while around 18% of the samples were above the WHO guideline of $10 \mu\text{g/L}$ [24]. The concentration of As in groundwater is higher in Bangladesh compared to other countries, and some tube wells even contain As concentrations as high as $4730 \mu\text{g/L}$ [25][26]. In India, high-As groundwater has already affected twenty states and four union territories, and about 100 million people are under threat from the toxicity of high-As groundwater [16][27][28][29]. The impact of high-As groundwater in India is concentrated on the Ganges–Yarlung Tsangpo Plain, seen on the neo-alluvial (Holocene) floodplains of the rivers in the Himalayas [30][31]. Approximately 50–60 million individuals in Pakistan consume high-As groundwater ($>50 \mu\text{g/L}$) in vulnerable areas [32]. A meta-analysis of groundwater affected by As in Pakistan showed that 73% of these groundwater samples contained

arsenic above 10 µg/L [33]. China is also one of the world's most representative areas of high-As groundwater, with more than 20 provinces/autonomous regions having high-As groundwater problems. These high-As groundwater provinces are mainly located in the fluvial/alluvial-lacustrine plains and basins (Yinchuan Plain, Hetao Plain, Guide Basin, Hohhot Basin, Junggar Basin, Datong Basin, etc.) located in arid/semi-arid regions and alluvial plains/basins and river deltas in humid/semi-humid regions (Yangtze River, Yellow River Delta, Pearl River Delta, Delta, Huaihe River, Alluvial Plain, Yellow River, Yuncheng Basin, Taiyuan Basin, Songnen Plain, etc.) [34][35][36][37]. The population affected by high-As groundwater contamination in China was estimated to be about 19.6 million according to a statistical risk assessment model developed by Rodríguez-Lado et al. [38].

Table 1. The occurrence of high-As groundwater reported by major countries in the world.

Country	Study Area	Max As conc. (µg/L)	Samples	Environmental Condition and/or Enrichment Mechanism	References
Afghanistan	Ghazni and maidan Wardak provinces	990	746	The weathering and leaching action	[39]
Argentina	Santiago del Estero Province	14,969	40	Volcanic ash sedimentary environment; agricultural irrigation	[40]
	La Pampa	5300	44	The geological factors; weathering of volcanic ash and loess; oxidizing condition	[41]
Australia	Stuarts Point coastal	85	140	Desorption of As from Al-hydroxides and As-enriched Fe-oxyhydroxides; high concentrations of HCO_3^- and PO_4^-	[42]
Bangladesh	Noakhali	4730	52,202	Eroded by flood plain rivers	[25]
Bolivia		364	24	The alteration of volcanic rocks; evaporation and redox reactions	[43]
Botswana	Botswana	116	20	Delta; evaporation concentration; weakly alkaline environment; pH 6.29–8.60	[44]
Brazil		2980		Anthropogenic; volcanic activity and weathering of rocks	[43]
Burkina Faso		1630	45	Zones of gold mineralization in volcano-sedimentary rocks	[45]
China	Datong Basin	1932	1022	The weak alkaline reductive environment; high HCO_3^-	[46]

Country	Study Area	Max As conc. (µg/L)	Samples	Environmental Condition and/or Enrichment Mechanism	References
				concentration; water–rock interactions	
	Hetao Basin	572	63	The reducing conditions; the dissolved organic; the competitive effects of other anions	[47]
	Jiangnan Basin	2330	34	The high HCO_3^- concentrations; microorganisms and exogenous substances; the seasonal variation; strongly reducing environment; reducing environment	[48]
	Taiwan (Lanyang and Chianan Plain)	1010		Alluvial plain; high DOC; strong reducing conditions	[49]
	Tarim Basin	91.2	233	Reducing environment; the dissolved organic; reductive dissolution release;	[50]
	Yinchuan	177	92	Agricultural irrigation; the reductive dissolution of Fe oxides; the high PO_4^- concentrations	[51]
	Pearl River Delta	161	18	Reductive environment; the high NH_4^+ concentrations; high concentrations of NH_4^+ and organic matter	[52]
Cambodian		1610	207	Holocene alluvial sediments; reducing environment	[53]
Costa Rica	Northern Costa Rica	29,100	35	Associated with the volcanic rock	[43]
Czech Republic	Mokrsko	1690	62	pH > 9	[54]
Ecuador		969	67	In hot springs	[43]
Ethiopia	Southwestern Ethiopia	184.5	44	pH < 7	[55]
Ghana		1760	357	Spillages of the mines; pH 4.8–6.99	[56]
Hungary	Southern Hungary	260	73	At a depth of 0.8–2.4 km and containing CH_4	[57]

Country	Study Area	Max As conc. (µg/L)	Samples	Environmental Condition and/or Enrichment Mechanism	References
India	Bhair	1466	1365	Ganga Plain; Holocene newer alluvium and the Pleistocene older alluvium	[58]
	Shahpur block, Bhojpur district, Bihar state	1805	4704	Ganges plain	[28]
	Punjab	3192	4780	Alluvial aquifers	[58]
Iran	Kurdistan Some villages	1500	27	Mining and sedimentary environment	[59]
	East Azarbaijan-Tabriz Plain	2000	18	Hydrogeological and environmental reducing conditions	
	Ardabil-A city	5834	163	Interaction of hydrothermal fluids with the rocks and geogenic source-geological structure	
	Mazandar an-Haraz River	110	20	Geogenic source and mining	
	Tabas South Khorasan	53	29	Weathering	
	Razavi Khorasan Chelpu Kashmar	606	12	Geogenic Origin sedimentary environment	
	Isfahan Mutehgold mining district	1061	17	Weathering and mining	
Japan		38	136	Reducing environment and factory blowdown	[26]
Korea	Geumsan County	113	150	Oxidation reaction of sulfide minerals in metasedimentary rocks and desorption process under high pH conditions	[60]
Nigeria	Warri-Port Harcourt, Ogun State, Kaduna	750	20	Alluvial sediments, reducing environment, slightly acidic	[16]
Pakistan	Kasur, Shhiwal, Bahawalpur, and Rahim Yar Khan	3090	395	Irrigation and factory sewage	[61]

Country	Study Area	Max As conc. (µg/L)	Samples	Environmental Condition and/or Enrichment Mechanism	References
	Lahore municipality	85	41	Topsoil and extensive irrigation of unconfined aquifers, reductive dissolution	[32]
	Mailsi	812	44	Human activity	[49]
Paraguay		120	37	Human activity and volcanic ash deposition environment	[43]
Lao PDR	Vientiane	24.4	3	Reducing environment	[17]
	Borikhamxay	30	7	Reducing environment	
	Champasack	25.6	27	Reducing environment	
	Attapeu	31.6	10	Reducing environment	
Myanmar	Ayeyarwady	630	55	Reductive dissolution of Fe oxyhydroxides	[49]
Mexico	La Laguna Region	5000	29	Adsorption or coprecipitation on iron oxides, clay-mineral surfaces, and organic carbon	[57] [49]
	Zacatecas [67]	75.4	182	Geological origin, water–rock interaction	
Nepal	Nawalparasi	2620	18,000	Seasons and climate change, water–rock interaction	[62]
Pakistan	Larkana Sindh, [68]	318	58	pH 6.8–8.1	[63]
	Punjab	655	141	pH 7.0–9.3	[64]
Spain	Duero Cenozoic Basin	613	514	pH 5.87–1.58 [69]	[16]
Thailand	Suphan Buri	5000	21	pH 5.20–5.90; Eh 250–370 mV [43]	[70]
USA	San Joaquin Valley, California	148.5	4983 [71]	Arid and semi-arid basins; alluvial, fluvial, and lacustrine deposits; pH > 7.8; reducing conditions	[65]
	Lahontan Valley, in Churchill County, Nevada [54]	4100	59	Lacustrine sediments	[66]

In Africa, high-As groundwater has been found in only a few areas across the continent, primarily in the western and southern regions, more due to insufficient research rather than a shortage of problems [67]. Twenty countries in Africa have recorded high concentrations of arsenic in groundwater, including Botswana, Burkina Faso, Ethiopia,

Country ^[67]	Study Area	Max As conc. (µg/L)	Samples	Environmental Condition and/or Enrichment Mechanism ^[51]	References ^[49]
Vietnam	Mekong Delta	850	109	pH 7.22–8.63	

References

- Lall, U.; Josset, L.; Russo, T. A Snapshot of the World's Groundwater Challenges. *Annu. Rev. Env. Resour.* 2020, 45, 171–194.
- Grogan, D.S.; Wisser, D.; Prusevich, A.; Lammers, R.B.; Froliking, S. The Use and Re-Use of Unsustainable Groundwater for Irrigation: A Global Budget. *Environ. Res. Lett.* 2017, 12, 034017.
- Burri, N.M.; Weatherl, R.; Moeck, C.; Schirmer, M. A Review of Threats to Groundwater Quality in the Anthropocene. *Sci. Total Environ.* 2019, 684, 136–154.
- ATSDR. The ATSDR 2019 Substance Priority List, Agency for Toxic Substances and Disease Registry. Available online: <https://www.atsdr.cdc.gov/spl/index.html> (accessed on 5 November 2023).
- Fendorf, S.; Michael, H.A.; van Geen, A. Spatial and Temporal Variations of Groundwater Arsenic in South and Southeast Asia. *Science* 2010, 328, 1123–1127.
- Gorchev, H.G.; Ozolins, G. WHO Guidelines for Drinking-Water Quality. *WHO Chron.* 1984, 38, 104–108.
- Thakur, J.K.; Thakur, R.K.; Ramanathan, A.L.; Kumar, M.; Singh, S.K. Arsenic Contamination of Groundwater in Nepal—An Overview. *Water* 2011, 3, 1–20.
- Koby, M.; Soltani, R.D.C.; Omwene, P.I.; Khataee, A. A Review on Decontamination of Arsenic-Contained Water by Electrocoagulation: Reactor Configurations and Operating Cost along with Removal Mechanisms. *Environ. Technol. Innov.* 2020, 17, 100519.
- Rahaman, M.S.; Rahaman, M.M.; Mise, N.; Sikder, M.T.; Ichihara, G.; Uddin, M.K.; Kurasaki, M.; Ichihara, S. Environmental Arsenic Exposure and Its Contribution to Human Diseases, Toxicity Mechanism and Management. *Environ. Pollut.* 2021, 289, 117940.
- Yadav, M.K.; Saidulu, D.; Gupta, A.K.; Ghosal, P.S.; Mukherjee, A. Status and Management of Arsenic Pollution in Groundwater: A Comprehensive Appraisal of Recent Global Scenario, Human Health Impacts, Sustainable Field-Scale Treatment Technologies. *J. Environ. Chem. Eng.* 2021, 9, 105203.
- Hung, D.Q.; Nekrassova, O.; Compton, R.G. Analytical Methods for Inorganic Arsenic in Water: A Review. *Talanta* 2004, 64, 269–277.
- Kumaresan, M.; Riyazuddin, P. Overview of Speciation Chemistry of Arsenic. *Curr. Sci.* 2001, 80, 837–846.

13. Kalman, J.; Smith, B.D.; Bury, N.R.; Rainbow, P.S. Biodynamic Modelling of the Bioaccumulation of Trace Metals (Ag, As and Zn) by an Infaunal Estuarine Invertebrate, the Clam *Scrobicularia Plana*. *Aquat. Toxicol.* 2014, 154, 121–130.
14. Ansari, M.A.; Saravana Kumar, U.; Noble, J.; Akhtar, N.; Akhtar, M.A.; Deodhar, A. Isotope Hydrology Tools in the Assessment of Arsenic Contamination in Groundwater: An Overview. *Chemosphere* 2023, 340, 139898.
15. Mandal, B.K.; Suzuki, K.T. Arsenic Round the World: A Review. *Talanta* 2002, 58, 201–235.
16. Shaji, E.; Santosh, M.; Sarath, K.V.; Prakash, P.; Deepchand, V.; Divya, B.V. Arsenic Contamination of Groundwater: A Global Synopsis with Focus on the Indian Peninsula. *Geosci. Front.* 2021, 12, 101079.
17. Cho, K.H.; Sthiannopkao, S.; Pachepsky, Y.A.; Kim, K.-W.; Kim, J.H. Prediction of Contamination Potential of Groundwater Arsenic in Cambodia, Laos, and Thailand Using Artificial Neural Network. *Water Res.* 2011, 45, 5535–5544.
18. Buschmann, J.; Berg, M.; Stengel, C.; Sampson, M.L. Arsenic and Manganese Contamination of Drinking Water Resources in Cambodia: Coincidence of Risk Areas with Low Relief Topography. *Environ. Sci. Technol.* 2007, 41, 2146–2152.
19. Van Geen, A.; Ahmed, E.B.; Pitcher, L.; Mey, J.L.; Ahsan, H.; Graziano, J.H.; Ahmed, K.M. Comparison of Two Blanket Surveys of Arsenic in Tubewells Conducted 12years Apart in a 25km² Area of Bangladesh. *Sci. Total Environ.* 2014, 488–489, 484–492.
20. Stopelli, E.; Duyen, V.T.; Mai, T.T.; Trang, P.T.K.; Viet, P.H.; Lightfoot, A.; Kipfer, R.; Schneider, M.; Eiche, E.; Kontny, A.; et al. Spatial and Temporal Evolution of Groundwater Arsenic Contamination in the Red River Delta, Vietnam: Interplay of Mobilisation and Retardation Processes. *Sci. Total Environ.* 2020, 717, 137143.
21. Winkel, L.; Berg, M.; Amini, M.; Hug, S.J.; Annette Johnson, C. Predicting Groundwater Arsenic Contamination in Southeast Asia from Surface Parameters. *Nat. Geosci.* 2008, 1, 536–542.
22. Brammer, H.; Ravenscroft, P. Arsenic in Groundwater: A Threat to Sustainable Agriculture in South and South-East Asia. *Environ. Int.* 2009, 35, 647–654.
23. Ganguli, S.; Rifat, M.A.H.; Das, D.; Islam, S.; Islam, M.N. Groundwater Pollution in Bangladesh: A Review. *Grassroots J. Nat. Resour.* 2021, 04, 115–145.
24. Bangladesh Bureau of Statistics. Bangladesh National Drinking Water Quality Survey of 2009; Bangladesh Bureau of Statistics: Dhaka, Bangladesh, 2011.
25. Chakraborti, D.; Rahman, M.M.; Das, B.; Murrill, M.; Dey, S.; Chandra Mukherjee, S.; Dhar, R.K.; Biswas, B.K.; Chowdhury, U.K.; Roy, S.; et al. Status of Groundwater Arsenic Contamination in Bangladesh: A 14-Year Study Report. *Water Res.* 2010, 44, 5789–5802.

26. Tashdedul, H.M.; Reyes, N.J.D.G.; Jeon, M.; Kim, L.-H. Current Status and Technologies for Treating Groundwater Arsenic Pollution in Bangladesh. *J. Wetl. Res.* 2022, 24, 142–154.
27. Chakraborti, D.; Rahman, M.M.; Chatterjee, A.; Das, D.; Das, B.; Nayak, B.; Pal, A.; Chowdhury, U.K.; Ahmed, S.; Biswas, B.K.; et al. Fate of over 480 Million Inhabitants Living in Arsenic and Fluoride Endemic Indian Districts: Magnitude, Health, Socio-Economic Effects and Mitigation Approaches. *J. Trace Elem. Med. Biol.* 2016, 38, 33–45.
28. Chakraborti, D.; Rahman, M.M.; Ahamed, S.; Dutta, R.N.; Pati, S.; Mukherjee, S.C. Arsenic Contamination of Groundwater and Its Induced Health Effects in Shahpur Block, Bhojpur District, Bihar State, India: Risk Evaluation. *Environ. Sci. Pollut. R.* 2016, 23, 9492–9504.
29. Bindal, S.; Singh, C.K. Predicting Groundwater Arsenic Contamination: Regions at Risk in Highest Populated State of India. *Water Res.* 2019, 159, 65–76.
30. Chakraborti, D.; Ghorai, S.; Das, B.; Pal, A.; Nayak, B.; Shah, B. Arsenic Exposure through Groundwater to the Rural and Urban Population in the Allahabad-Kanpur Track in the Upper Ganga Plain. *J. Environ. Monit. JEM* 2009, 11, 1455–1459.
31. Mukherjee, A.; Verma, S.; Gupta, S.; Henke, K.R.; Bhattacharya, P. Influence of Tectonics, Sedimentation and Aqueous Flow Cycles on the Origin of Global Groundwater Arsenic: Paradigms from Three Continents. *J. Hydrol.* 2014, 518, 284–299.
32. Podgorski, J.E.; Eqani, S.A.M.A.S.; Khanam, T.; Ullah, R.; Shen, H.; Berg, M. Extensive Arsenic Contamination in High-pH Unconfined Aquifers in the Indus Valley. *Sci. Adv.* 2017, 3, e1700935.
33. Shahid, M. A Meta-Analysis of the Distribution, Sources and Health Risks of Arsenic-Contaminated Groundwater in Pakistan. *Environ. Pollut.* 2018, 242, 307–319.
34. Cao, W.G.; Zhang, Z.; Guo, H.M.; Fu, Y.; Gao, Z.P.; Nan, T.; Ren, Y.; Li, Z.Y. Spatial Distribution and Controlling Mechanisms of High Fluoride Groundwater in the Coastal Plain of Bohai Rim, North China. *J. Hydrol.* 2023, 617, 128952.
35. Wen, D.G.; Zhang, F.C.; Zhang, E.Y.; Wang, C.; Han, S.B.; Zheng, Y. Arsenic, Fluoride and Iodine in Groundwater of China. *J. Geochem. Explor.* 2013, 135, 1–21.
36. He, X.D.; Li, P.Y.; Ji, Y.J.; Wang, Y.H.; Su, Z.M.; Elumalai, V. Groundwater Arsenic and Fluoride and Associated Arsenicosis and Fluorosis in China: Occurrence, Distribution and Management. *Expo. Health* 2020, 12, 355–368.
37. Guo, H.; Wen, D.; Liu, Z.; Jia, Y.; Guo, Q. A Review of High Arsenic Groundwater in Mainland and Taiwan, China: Distribution, Characteristics and Geochemical Processes. *Appl. Geochem.* 2014, 41, 196–217.
38. Rodríguez-Lado, L.; Sun, G.F.; Berg, M.; Zhang, Q.; Xue, H.B.; Zheng, Q.M.; Johnson, C.A. Groundwater Arsenic Contamination Throughout China. *Science* 2013, 341, 866–868.

39. Saffi, M.H.; Eqrar, M. Arsenic Contamination of Groundwater in Ghazni and Maidan Wardak Provinces: Afghanistan. In *Arsenic Research and Global Sustainability: Proceedings of the Sixth International Congress on Arsenic in the Environment (As2016)*, Stockholm, Sweden, 19–23 June 2016; CRC Press: Boca Raton, FL, USA, 2016; pp. 41–42. ISBN 978-1-138-02941-5.
40. Bhattacharya, P.; Claesson, M.; Bundschuh, J.; Sracek, O.; Fagerberg, J.; Jacks, G.; Martin, R.A.; Storniolo, A. del R.; Thir, J.M. Distribution and Mobility of Arsenic in the Río Dulce Alluvial Aquifers in Santiago Del Estero Province, Argentina. *Sci. Total Environ.* 2006, 358, 97–120.
41. Aullón Alcaine, A.; Schulz, C.; Bundschuh, J.; Jacks, G.; Thunvik, R.; Gustafsson, J.-P.; Mörtz, C.-M.; Sracek, O.; Ahmad, A.; Bhattacharya, P. Hydrogeochemical Controls on the Mobility of Arsenic, Fluoride and Other Geogenic Co-Contaminants in the Shallow Aquifers of Northeastern La Pampa Province in Argentina. *Sci. Total Environ.* 2020, 715, 136671.
42. Smith, J.V.S.; Jankowski, J.; Sammut, J. Vertical Distribution of As(III) and As(V) in a Coastal Sandy Aquifer: Factors Controlling the Concentration and Speciation of Arsenic in the Stuarts Point Groundwater System, Northern New South Wales, Australia. *Appl. Geochem.* 2003, 18, 1479–1496.
43. Bundschuh, J.; Armienta, M.A.; Morales-Simfors, N.; Alam, M.A.; López, D.L.; Delgado Quezada, V.; Dietrich, S.; Schneider, J.; Tapia, J.; Sracek, O.; et al. Arsenic in Latin America: New Findings on Source, Mobilization and Mobility in Human Environments in 20 Countries Based on Decadal Research 2010-2020. *Crit. Rev. Env. Sci. Tec.* 2021, 51, 1727–1865.
44. Huntsman-Mapila, P.; Mapila, T.; Letshwenyo, M.; Wolski, P.; Hemond, C. Characterization of Arsenic Occurrence in the Water and Sediments of the Okavango Delta, NW Botswana. *Appl. Geochem.* 2006, 21, 1376–1391.
45. Smedley, P.L.; Knudsen, J.; Maiga, D. Arsenic in Groundwater from Mineralised Proterozoic Basement Rocks of Burkina Faso. *Appl. Geochem.* 2007, 22, 1074–1092.
46. He, X.D.; Li, P.Y.; Wu, J.H.; Wei, M.J.; Ren, X.F.; Wang, D. Poor Groundwater Quality and High Potential Health Risks in the Datong Basin, Northern China: Research from Published Data. *Environ. Geochem. Health* 2021, 43, 791–812.
47. Guo, H.; Yang, S.; Tang, X.; Li, Y.; Shen, Z. Groundwater Geochemistry and Its Implications for Arsenic Mobilization in Shallow Aquifers of the Hetao Basin, Inner Mongolia. *Sci. Total Environ.* 2008, 393, 131–144.
48. Wang, Z.; Guo, H.M.; Liu, H.Y.; Zhang, W.M. Source, Migration, Distribution, Toxicological Effects and Remediation Technologies of Arsenic in Groundwater in China. *China Geol.* 2023, 6, 476–493.
49. Wang, Y.X.; Li, J.X.; Ma, T.; Xie, X.J.; Deng, Y.M.; Gan, Y.Q. Genesis of Geogenic Contaminated Groundwater: As, F and I. *Crit. Rev. Env. Sci. Tec.* 2021, 51, 2895–2933.

50. Sun, Y.; Zhou, J.L.; Yang, F.Y.; Ji, Y.Y.; Zeng, Y.Y. Distribution and Co-Enrichment Genesis of Arsenic, Fluorine and Iodine in Groundwater of the Oasis Belt in the Southern Margin of Tarim Basin. *Earth Sci. Front.* 2022, 29, 99–114.
51. Guo, Q.; Guo, H.M.; Yang, Y.C.; Han, S.B.; Zhang, F.C. Hydrogeochemical Contrasts between Low and High Arsenic Groundwater and Its Implications for Arsenic Mobilization in Shallow Aquifers of the Northern Yinchuan Basin, P.R. China. *J. Hydrol.* 2014, 518, 464–476.
52. Wang, Y.; Jiao, J.J.; Cherry, J.A. Occurrence and Geochemical Behavior of Arsenic in a Coastal Aquifer–Aquitard System of the Pearl River Delta, China. *Sci. Total Environ.* 2012, 427–428, 286–297.
53. Berg, M.; Stengel, C.; Trang, P.; Hungviet, P.; Sampson, M.; Leng, M.; Samreth, S.; Fredericks, D. Magnitude of Arsenic Pollution in the Mekong and Red River Deltas—Cambodia and Vietnam. *Sci. Total Environ.* 2007, 372, 413–425.
54. Litter, M.I.; Ingallinella, A.M.; Olmos, V.; Savio, M.; Difeo, G.; Botto, L.; Farfán Torres, E.M.; Taylor, S.; Frangie, S.; Herkovits, J.; et al. Arsenic in Argentina: Occurrence, Human Health, Legislation and Determination. *Sci. Total Environ.* 2019, 676, 756–766.
55. Dilpazeer, F.; Munir, M.; Baloch, M.Y.J.; Shafiq, I.; Iqbal, J.; Saeed, M.; Abbas, M.M.; Shafique, S.; Aziz, K.H.H.; Mustafa, A.; et al. A Comprehensive Review of the Latest Advancements in Controlling Arsenic Contaminants in Groundwater. *Water* 2023, 15, 478.
56. Kusimi, J.M.; Kusimi, B.A. The Hydrochemistry of Water Resources in Selected Mining Communities in Tarkwa. *J. Geochem. Explor.* 2012, 112, 252–261.
57. Rowland, H.A.L.; Omoregie, E.O.; Millot, R.; Jimenez, C.; Mertens, J.; Baciú, C.; Hug, S.J.; Berg, M. Geochemistry and Arsenic Behaviour in Groundwater Resources of the Pannonian Basin (Hungary and Romania). *Appl. Geochem.* 2011, 26, 1–17.
58. Dhillon, A.K. Arsenic Contamination of India's Groundwater: A Review and Critical Analysis. In *Arsenic Water Resources Contamination*; Springer: Cham, Switzerland, 2020; pp. 177–205.
59. Hamidian, A.H.; Razeghi, N.; Zhang, Y.; Yang, M. Spatial Distribution of Arsenic in Groundwater of Iran, a Review. *J. Geochem. Explor.* 2019, 201, 88–98.
60. Jadhav, S.V.; Bringas, E.; Yadav, G.D.; Rathod, V.K.; Ortiz, I.; Marathe, K.V. Arsenic and Fluoride Contaminated Groundwaters: A Review of Current Technologies for Contaminants Removal. *J. Environ. Manage.* 2015, 162, 306–325.
61. Tsuji, J.S.; Chang, E.T.; Gentry, P.R.; Clewell, H.J.; Boffetta, P.; Cohen, S.M. Dose-Response for Assessing the Cancer Risk of Inorganic Arsenic in Drinking Water: The Scientific Basis for Use of a Threshold Approach. *Crit. Rev. Toxicol.* 2019, 49, 36–84.

62. Ali, W.; Mushtaq, N.; Javed, T.; Zhang, H.; Ali, K.; Rasool, A.; Farooqi, A. Vertical Mixing with Return Irrigation Water the Cause of Arsenic Enrichment in Groundwater of District Larkana Sindh, Pakistan. *Environ. Pollut.* 2019, 245, 77–88.
63. Mushtaq, N.; Masood, N.; Khattak, J.A.; Hussain, I.; Khan, Q.; Farooqi, A. Health Risk Assessment and Source Identification of Groundwater Arsenic Contamination Using Agglomerative Hierarchical Cluster Analysis in Selected Sites from Upper Eastern Parts of Punjab Province, Pakistan. *Hum. Ecol. Risk Assess. Int. J.* 2021, 27, 999–1018.
64. Gómez, J.J.; Lillo, J.; Sahún, B. Naturally Occurring Arsenic in Groundwater and Identification of the Geochemical Sources in the Duero Cenozoic Basin, Spain. *Environ. Geol.* 2006, 50, 1151–1170.
65. Haugen, E.A.; Jurgens, B.C.; Arroyo-Lopez, J.A.; Bennett, G.L. Groundwater Development Leads to Decreasing Arsenic Concentrations in the San Joaquin Valley, California. *Sci. Total Environ.* 2021, 771, 145223.
66. Walkera, M.; Seiler, R.L.; Meinert, M. Effectiveness of Household Reverse-Osmosis Systems in a Western U.S. Region with High Arsenic in Groundwater. *Sci. Total Environ.* 2008, 389, 245–252.
67. Medunić, G.; Fiket, Ž.; Ivanić, M. Arsenic Contamination Status in Europe, Australia, and Other Parts of the World. In *Arsenic in Drinking Water and Food*; Srivastava, S., Ed.; Springer: Singapore, 2020; pp. 183–233. ISBN 9789811385872.
68. Daniele, L. Distribution of Arsenic and Other Minor Trace Elements in the Groundwater of Ischia Island (Southern Italy). *Environ. Geol.* 2004, 46, 96–103.
69. Sorg, T.J.; Chen, A.S.C.; Wang, L. Arsenic Species in Drinking Water Wells in the USA with High Arsenic Concentrations. *Water Res.* 2014, 48, 156–169.
70. McClintock, T.R.; Chen, Y.; Bundschuh, J.; Oliver, J.T.; Navoni, J.; Olmos, V.; Lepori, E.V.; Ahsan, H.; Parvez, F. Arsenic Exposure in Latin America: Biomarkers, Risk Assessments and Related Health Effects. *Sci. Total Environ.* 2012, 429, 76–91.
71. Ortega-Guerrero, A. Evaporative Concentration of Arsenic in Groundwater: Health and Environmental Implications, La Laguna Region, Mexico. *Environ. Geochem. Health* 2017, 39, 987–1003.
72. Morales-Arredondo, J.I.; Esteller-Alberich, M.V.; Armienta Hernández, M.A.; Martínez-Florentino, T.A.K. Characterizing the Hydrogeochemistry of Two Low-Temperature Thermal Systems in Central Mexico. *J. Geochem. Explor.* 2018, 185, 93–104.
73. Smedley, P.L.; Nicolli, H.B.; Macdonald, D.M.J.; Barros, A.J.; Tullio, J.O. Hydrogeochemistry of Arsenic and Other Inorganic Constituents in Groundwaters from La Pampa, Argentina. *Appl. Geochem.* 2002, 17, 259–284.

Retrieved from <https://encyclopedia.pub/entry/history/show/124480>