

Hepatic Veno-Occlusive Disease with Immunodeficiency

Subjects: Genetics & Heredity

Contributor: Camila Xu

Hepatic veno-occlusive disease with immunodeficiency (also called VODI) is a hereditary disorder of the liver and immune system. Its signs and symptoms appear after the first few months of life. Hepatic veno-occlusive disease is a condition that blocks (occludes) small veins in the liver, disrupting blood flow in this organ. This condition can lead to enlargement of the liver (hepatomegaly), a buildup of scar tissue (hepatic fibrosis), and liver failure.

Keywords: genetic conditions

1. Introduction

Children with VODI are prone to recurrent infections caused by certain bacteria, viruses, and fungi. The organisms that cause infection in people with this disorder are described as opportunistic because they ordinarily do not cause illness in healthy people. These infections are usually serious and may be life-threatening. In most people with VODI, infections occur before hepatic veno-occlusive disease becomes evident.

Many people with VODI live only into childhood, although some affected individuals have lived to early adulthood.

2. Frequency

VODI appears to be a rare disorder; approximately 20 affected families have been reported worldwide. Most people diagnosed with the condition have been of Lebanese ancestry. However, the disorder has also been identified in several individuals with other backgrounds in the United States and Italy.

3. Causes

VODI results from mutations in the *SP110* gene. This gene provides instructions for making a protein called SP110 nuclear body protein, which is involved in the normal function of the immune system. This protein likely helps regulate the activity of genes needed for the body's immune response to foreign invaders (such as viruses and bacteria).

Mutations in the *SP110* gene prevent cells from making functional SP110 nuclear body protein, which impairs the immune system's ability to fight off infections. It is unclear how a lack of this protein affects blood flow in the liver.

3.1. The gene associated with Hepatic veno-occlusive disease with immunodeficiency

- SP110

4. Inheritance

This condition is inherited in an autosomal recessive pattern, which means both copies of the gene in each cell have mutations. The parents of an individual with an autosomal recessive condition each carry one copy of the mutated gene, but they typically do not show signs and symptoms of the condition.

5. Other Names for This Condition

- familial veno-occlusive disease with immunodeficiency
- hepatic venoocclusive disease with immunodeficiency

- veno-occlusive disease and immunodeficiency syndrome
 - VODI
-

References

1. Cliffe ST, Wong M, Taylor PJ, Ruga E, Wilcken B, Lindeman R, Buckley MF, Roscioli T. The first prenatal diagnosis for veno-occlusive disease and immunodeficiency syndrome, an autosomal recessive condition associated with mutations in SP110. *Prenat Diagn.* 2007 Jul;27(7):674-6.
 2. Roscioli T, Cliffe ST, Bloch DB, Bell CG, Mullan G, Taylor PJ, Sarris M, Wang J, Donald JA, Kirk EP, Ziegler JB, Salzer U, McDonald GB, Wong M, Lindeman R, Buckley MF. Mutations in the gene encoding the PML nuclear body protein Sp 110 are associated with immunodeficiency and hepatic veno-occlusive disease. *Nat Genet.* 2006 Jun;38(6):620-2.
 3. Roscioli T, Ziegler JB, Buckley M, Wong M. Hepatic Veno-Occlusive Disease with Immunodeficiency. 2007 Feb 21 [updated 2017 Jan 12]. In: Adam MP, Ardinger HH, Pagon RA, Wallace SE, Bean LJH, Stephens K, Amemiya A, editors. *GeneReviews®* [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2020. Available from <http://www.ncbi.nlm.nih.gov/books/NBK1271/>
-

Retrieved from <https://encyclopedia.pub/entry/history/show/11535>