

# MEGDEL Syndrome

Subjects: Genetics

Submitted by:  Rita

Xu

(This entry belongs to Entry Collection "["MedlinePlus"](#)")

## Definition

MEGDEL syndrome is an inherited disorder that affects multiple body systems. It is named for several of its features: 3-methylglutaconic aciduria (MEG), deafness (D), encephalopathy (E), and Leigh-like disease (L).

---

## 1. Introduction

MEGDEL syndrome is characterized by abnormally high levels of an acid, called 3-methylglutaconic acid, in the urine (3-methylglutaconic aciduria). MEGDEL syndrome is one of a group of metabolic disorders that can be diagnosed by presence of this feature. People with MEGDEL syndrome also have high urine levels of another acid called 3-methylglutaric acid.

In infancy, individuals with MEGDEL syndrome develop hearing loss caused by changes in the inner ear (sensorineural deafness); the hearing problems gradually worsen over time.

Another feature of MEGDEL syndrome is brain dysfunction (encephalopathy). In infancy, encephalopathy leads to difficulty feeding, an inability to grow and gain weight at the expected rate (failure to thrive), and weak muscle tone (hypotonia). Infants with MEGDEL syndrome later develop involuntary muscle tensing (dystonia) and muscle stiffness (spasticity), which worsen over time. Because of these brain and muscle problems, affected babies have delayed development of mental and movement abilities (psychomotor delay), or they may lose skills they already developed. Individuals with MEGDEL syndrome have intellectual disability and never learn to speak.

People with MEGDEL syndrome have changes in the brain that resemble those in another condition called Leigh syndrome. These changes, which can be seen with medical imaging, are referred to as Leigh-like disease.

Other features that occur commonly in MEGDEL syndrome include low blood sugar (hypoglycemia) in affected newborns; liver problems (hepatopathy) in infancy, which can be serious but improve by early childhood; and episodes of abnormally high amounts of lactic acid in the blood (lactic acidosis).

The life expectancy of individuals with MEGDEL syndrome is unknown. Because of the severe health problems caused by the disorder, some affected individuals do not survive past infancy.

## 2. Frequency

MEGDEL syndrome is a rare disorder; its prevalence is unknown. At least 40 affected individuals have been mentioned in the medical literature.

## 3. Causes

MEGDEL syndrome is caused by mutations in the *SERAC1* gene. The function of the protein produced from this gene is not completely understood, although research suggests that it is involved in altering (remodeling) certain fats called phospholipids, particularly a phospholipid known as phosphatidylglycerol. Another phospholipid called cardiolipin is made from phosphatidylglycerol. Cardiolipin is a component of the membrane that surrounds cellular structures called mitochondria, which convert the energy from food into a form that cells can use, and is important for the proper functioning of these structures.

*SERAC1* gene mutations involved in MEGDEL syndrome lead to little or no *SERAC1* protein function. As a

result, phosphatidylglycerol remodeling is impaired, which likely alters the composition of cardiolipin. Researchers speculate that the abnormal cardiolipin affects mitochondrial function, reducing cellular energy production and leading to the neurological and hearing problems characteristic of MEGDEL syndrome. It is unclear how *SERAC1* gene mutations lead to abnormal release of 3-methylglutaconic acid in the urine, although it is thought to be related to mitochondrial dysfunction.

### 3.1. The Gene Associated with MEGDEL Syndrome

SERAC1

## 4. Inheritance

This condition is inherited in an autosomal recessive pattern, which means both copies of the gene in each cell have mutations. The parents of an individual with an autosomal recessive condition each carry one copy of the mutated gene, but they typically do not show signs and symptoms of the condition.

## 5. Other Names for This Condition

- 3-methylglutaconic aciduria type IV with sensorineural deafness, encephalopathy, and Leigh-like syndrome
- 3-methylglutaconic aciduria with deafness, encephalopathy, and Leigh-like syndrome
- MEGDHEL syndrome
- SERAC1 defect

## References

1. Sarig O, Goldsher D, Nousbeck J, Fuchs-Telem D, Cohen-Katsenelson K, Iancu TC, Manov I, Saada A, Sprecher E, Mandel H. Infantile mitochondrial hepatopathy is a cardinal feature of MEGDEL syndrome (3-methylglutaconic aciduria type IV with sensorineural deafness, encephalopathy and Leigh-like syndrome) caused by novel mutations in *SERAC1*. *Am J Med Genet A*. 2013 Sep;161A(9):2204-15. doi:10.1002/ajmg.a.36059.
2. Tort F, García-Silva MT, Ferrer-Cortès X, Navarro-Sastre A, García-Villoria J, Coll MJ, Vidal E, Jiménez-Almazán J, Dopazo J, Briones P, Elpeleg O, Ribes A. Exome sequencing identifies a new mutation in *SERAC1* in a patient with 3-methylglutaconic aciduria. *Mol Genet Metab*. 2013 Sep-Oct;110(1-2):73-7. doi:10.1016/j.ymgme.2013.04.021.
3. Wortmann S, Rodenburg RJ, Huizing M, Loupatty FJ, de Koning T, Kluijtmans LA, Engelke U, Wevers R, Smeitink JA, Morava E. Association of 3-methylglutaconic aciduria with sensori-neural deafness, encephalopathy, and Leigh-like syndrome (MEGDEL association) in four patients with a disorder of the oxidative phosphorylation. *Mol Genet Metab*. 2006 May;88(1):47-52.
4. Wortmann SB, de Brouwer APM, Wevers RA, Morava E. *SERAC1* Deficiency. 2014 Apr 17 [updated 2020 Jul 23]. In: Adam MP, Ardinger HH, Pagon RA, Wallace SE, Bean LJH, Stephens K, Amemiya A, editors. *GeneReviews*® [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2020. Available from <http://www.ncbi.nlm.nih.gov/books/NBK195853/>
5. Wortmann SB, Duran M, Anikster Y, Barth PG, Sperl W, Zschocke J, Morava E, Wevers RA. Inborn errors of metabolism with 3-methylglutaconic aciduria as discriminative feature: proper classification and nomenclature. *J Inher Metab Dis*. 2013 Nov;36(6):923-8. doi: 10.1007/s10545-012-9580-0. Review.
6. Wortmann SB, Vaz FM, Gardeitchik T, Vissers LE, Renkema GH, Schuurs-Hoeijmakers JH, Kulik W, Lammens M, Christin C, Kluijtmans LA, Rodenburg RJ, Nijtmans LG, Grünwald A, Klein C, Gerhold JM, Kozicz T, van Hasselt PM, Harakalova M, Kloosterman W, Barić I, Pronicka E, Ucar SK, Naess K, Singhal KK, Krumina Z, Gilissen C, van Bokhoven H, Veltman JA, Smeitink JA, Lefeber DJ, Spelbrink JN, Wevers RA, Morava E, de Brouwer AP. Mutations in the phospholipid remodeling gene *SERAC1* impair mitochondrial function and intracellular cholesterol trafficking and cause dystonia and deafness. *Nat Genet*. 2012 Jun;44(7):797-802. doi: 10.1038/ng.2325.

## Keywords

genetic conditions

Retrieved from <https://encyclopedia.pub/4563>