

Ion Channels in Ovarian Cancer

Subjects: **Oncology**

Contributor: Concetta Altamura

Ovarian cancer (OC) is the deadliest gynecologic cancer, due to late diagnosis, development of platinum resistance, and inadequate alternative therapy. It has been demonstrated that membrane ion channels play important roles in cancer processes, including cell proliferation, apoptosis, motility, and invasion. Here, we review the contribution of ion channels in the development and progression of OC, evaluating their potential in clinical management. Increased expression of voltage-gated and epithelial sodium channels has been detected in OC cells and tissues and shown to be involved in cancer proliferation and invasion. Potassium and calcium channels have been found to play a critical role in the control of cell cycle and in the resistance to apoptosis, promoting tumor growth and recurrence. Overexpression of chloride and transient receptor potential channels was found both *in vitro* and *in vivo*, supporting their contribution to OC.

Ion channels

Ovarian cancer

Drug resistance

chemoresistance

1. Sodium Channels

Two major classes of sodium channels are expressed in mammals: the voltage-gated sodium channels (VGSC) and the epithelial sodium channels (eNaC). Nine isoforms (Nav1.1–Nav1.9) of VGSC are expressed throughout the body in various excitable cell types, while epithelial sodium channels are found mainly in the skin, lung, and kidney [1]. Altered expression of these channels has been observed in various tumors, especially in breast cancer, colon cancer, prostate cancer, and small cell lung cancer [2][3][4][5][6].

The first direct hints that VGSCs may be important for ovarian carcinogenesis came in 2010, when Gao and collaborators investigated the functional expression of these channels in OC and the possible correlation between VGSC activity and carcinogenic processes [7]. Gene expression analysis demonstrated that mRNA levels of Nav1.1, Nav1.3, Nav1.4, and Nav1.5 are significantly higher in OC cells than those in normal ovarian tissue, suggesting a role of VGSC in ovarian tumorigenesis. Furthermore, Nav1.2, Nav1.4, Nav1.5, and Nav1.7 mRNA levels are greatly increased in the highly metastatic OC cell lines compared to the low-metastatic cells, correlating the enhanced expression of VGSC with the cellular invasion process. Additional experiments revealed that the Nav1.5 protein is the main VGSC involved in OC and that increased Nav1.5 expression was associated with high histological grade and metastasis in OC [7].

To date, the molecular pathway by which sodium channels contribute to cancer progression and invasiveness is not fully understood yet, but there are some hypotheses. The VGSCs expressed in cancer cells, including Nav1.5, carry a small persistent Na^+ current that depolarizes the resting potential and induces cytosolic Na^+ accumulation

(Figure 1) [8][9]. Elevated intracellular Na^+ leads to Ca^{2+} entry through the reverse mode activity of the sodium-calcium exchanger (NCX1) [10], thus triggering activation of Src kinases that promote cancer cell invasion and metastatic progression (Figure 1) [11]. The NCX1-mediated increase in Na^+ efflux may activate the isoform 1 of the sodium-proton exchanger NHE1 [10] consequently leading to intracellular alkalinisation and extracellular acidification. These conditions stimulate tumor cell metabolism, promote the formation and the activity of actin-rich invasive protrusions, the invadopodia, and activates acidic-pH-dependent proteases such as the zinc-dependent matrix metalloproteases MMP-2 and -9 [12] and/or the acidic cysteine proteases [13], leading to an increase in proteolysis-dependent cell invasion (Figure 1) [14]. In line with this, inhibition of NHE1 activity with selective inhibitors, such as Cariporide and Zoniporide reduces cancer cell proliferation and invasive capability in different tumor types, including OC (Figure 1) [15][16]. Moreover, it has been shown that both in CHO cells and in the human OC cells A2780 NHE1 and NCX1 are physically and functionally linked such that an increase in sodium concentration caused by NHE1 can further drive NCX1 to generate a coordinate sodium efflux in exchange for calcium entry [10][17]. Thus, the resulting calcium increase inside the cell can promote tumorigenesis and tumor progression [18] (Figure 1).




Figure 1. Ion channels expression in ovarian cancer (OC). Schematic representation of the main ion channels and transporters involved in OC development and progression. Their dysregulated role in cell proliferation, migration and metastasis has been studied by modulating their activity with selective ion channel inhibitors. See text for further details.

Because the functional activity of VGSC is required to promote OC cell invasiveness, they constitute a promising target for the OC drug discovery. It was demonstrated that the treatment of OC cell lines (SKOV-3 and Caov-3) with the selective VGSC blocker tetrodotoxin (TTX) reduces the migration and invasion by 50–60%, without any significant effect on proliferation [7]. However, this natural agent is unequivocally toxic for mammals.

Recently, several population-based studies reported that polyunsaturated fatty acids, especially the omega 3 eicosapentaenoic acid (EPA), have a protective effect against OC [19][20]. Treatment of OC cells with EPA is associated with suppression of growth and reduction in invasivity [21]. It appeared that EPA exerts these beneficial effects through inhibition of Nav1.5 channels in a dose-dependent manner [22]. Additionally, various VGSC blockers belonging to class 1B antiarrhythmic drugs, antiepileptic drugs, and local anesthetics have been studied in *in vitro* and *in vivo* cancer models [23]. For instance, treatment of OC cells with the antiepileptic topiramate reduced OC cell proliferation and migration, suggesting a potential anticancer effect [24]. As regards the local anesthetics, retrospective studies have revealed that the use of these drugs for regional anesthesia during perioperative period reduced the recurrence rate of patients with breast cancer [25]. In keeping with this, *in vitro* experiments demonstrated that bupivacaine and lidocaine inhibited OC cells proliferation and metastasis by blocking Nav1.5, confirming their role as potential anticancer agents [26][27]. The multi-effects of lidocaine were also confirmed *in vivo*, by using murine syngeneic OC (ID8) model. The molecular mechanism by which lidocaine hampers OC malignancy was recently studied by Liu and collaborators, who demonstrated that lidocaine binding to Nav1.5 blocked the activation of FAK/Paxillin signaling pathway, inhibiting proliferation and metastasis, and decreased cisplatin resistance of OC cells [48]. However, it should be noted that these compounds might elicit their anticancer effects also through mechanisms other than VGSC inhibition. Indeed, Xuan et al. demonstrated that bupivacaine showed antiproliferative effects via the direct activation of intrinsic and extrinsic apoptotic pathways [26].

2. Potassium Channels

Potassium channels (K^+ channels) are the largest and most varied group of ion channels, being expressed in both excitable and non-excitable cells. Based on their structure and function, these channels can be classified into four main families: Voltage-gated K^+ channels (Kv); Ca^{2+} – or Na^+ – activated K^+ channels (KCa, KNa); inwardly rectifying K^+ channels (Kir); and two-pore domain K^+ channels (K2P) [28].

Using a variety of K^+ channel inhibitors on the OC cell line A2780, Zhanping et al. demonstrated that Kv channels are the main K^+ channels affecting OC cell proliferation and cell cycle progression [29]. Among these, Kv10.1 (Eag) and Kv11.1 (hERG) are significantly upregulated in OC cell lines and tissue, contributing to the hyperpolarization of the cell membrane [30]. Such hyperpolarization may increase Ca^{2+} influx into the cell through transport systems (ion channels and transporters) by increasing the driving force for Ca^{2+} [31][32]. Increased intracellular Ca^{2+} may then facilitate the opening of calcium-dependent K^+ channels (KCa), the expression of which is stimulated by mitogenic factors, thereby maintaining hyperpolarization and driving force for a sustained Ca^{2+} entry [33][34]. This condition also maintains the driving force for Na^+ dependent nutrient transport and intracellular acidification through NHE1 [32], driving the cells into a more potentially proliferative state. In addition, we can hypothesize that, in tumor cells over-expressing voltage-gated Ca^{2+} channels (especially T-type channels) and Na^+ channels, the depolarized

resting membrane potential of non-excitatory cells may force these channels to enter an inactivated state. Thus, membrane hyperpolarization at rest may remove channel inactivation and allows Ca^{2+} or Na^+ influx at any further little depolarizing oscillation. Finally, an alternative or complementary hypothesis is that opening of certain K^+ channels, independently of cell hyperpolarization, may contribute to cell volume regulation, which is critical during cell cycle progression [32].

In this scenario, Eag and HERG channels have been studied as possible molecular targets for anticancer drug design. Indeed, a variety of blockers was shown to inhibit OC cell viability through various pathways. Kv channel inhibition disrupts cell cycle, increasing the proportion of cells in G0/G1 phase with respect to S/G2/M phase (Figure 1). More recently, knockdown of HERG reduced proliferation, migration, and invasive ability of SKOV-3 cells, confirming HERG involvement in the pathogenesis of OC [35][34]. For example, the use of Eag blockers, such as imipramine and clofilium, increased the proportion of SKOV-3 cells undergoing early apoptosis, whereas treatment with the HERG channel specific blockers, E-4031 and ergtoxin, affected the cell cycle, resulting in the block of OC cells in the S phase (Figure 1) [36]. In addition, the natural alkaloid berberine (BBR) induced a strong cytotoxic effect against OC cells through inhibition of hERG channels [35]. Additional studies would be necessary to define the mechanisms by which BBR induces hERG1 channel blocking and by which hERG1 inhibition mediates phenotypic alterations in OC.

An important problem with the use of HERG blockers is the risk of side effects, specifically regarding ventricular arrhythmia and a slowed cardiac repolarization. Maybe the use of state-dependent blockers targeting open Kv11.1 channels in cancer cells, while sparing inactivated cardiac Kv11.1 channels might be helpful [37]. Another strategy to generate highly selective inhibitors of potassium channels is represented by the use of monoclonal antibodies. The first monoclonal antibody against Eag, mAb56, was shown to inhibit OC cell growth in vitro [38], thereby encouraging further studies.

Among the KCa family, the maxi-conductance BK channel KCa1.1 (KCNMA1 gene) plays an important role in the progression of OC cancer, contributing to cell proliferation and migration [39]. The KCa1.1 channel is over-expressed in several cancers including glioma [40] and hormone-sensitive tumors like breast, prostate and ovarian cancer [41][42]. Its role in OC has been studied by using selective BK channel blockers, such as iberiotoxin and tetrandrine [43]. Tetrandrine inhibits BK channel currents in OC cells and OC cell proliferation by inducing G1 phase arrest and accelerating apoptosis (Figure 1). Inhibition of BK channels also reduced the expression of the heat shock proteins hsp90 and hsp70, which are molecular chaperones involved in the folding and maturation of key regulatory proteins, including p53 [44]. Thus, the decrease of hsp expression may favor the expression of p53, p21, and Bax proteins, thereby triggering apoptosis (Figure 1) [43].

On the other hand, the opening of BK channels by NS1619 inhibits proliferation of the OC cell line A2780 in a time- and dosage-dependent manner. The efflux of K^+ from the cell results in cell shrinkage and the release of cytochrome c and activation of caspase, inducing apoptosis [45]. Thus, activity of BK channels finely tunes OC cell survival (Figure 1).

Recent studies revealed an involvement of KCa2.3 (*KCNN3* gene) and KCa3.1 (*KCNN4* gene) in OC pathogenesis [46][47]. *KCNN3* mRNA expression was significantly lower in OC tissues compared with normal controls, suggesting its negative contribution in OC progression [46].

Blockage of the KCa3.1 or its down-regulation in SKOV-3 human OC cells prevented OC migration as a consequence of a loss of interaction between KCa3.1 and the purinergic receptor P2Y2 [47].

An emerging class of K⁺ channels involved in OC is represented by the ATP-sensitive K⁺ channel Kir6.2/SUR2. Such channel results from the combination of the inward rectifier K⁺ channel Kir6.2 (encoded by *KCNJ11*) and the sulfonylurea receptor SUR2 (encoded by *ABCC9*) that is essential for the modulation of channel gating in response to the binding of nucleotides or drugs [48]. Recently, it was reported that Kir6.2 and SUR2 genes are downregulated in OC compared to healthy tissues and that high expression of the SUR2 gene is associated with improved overall survival in all OC patients [49]. The stimulation of the Kir6.2/SUR2 channel activity with the pharmacological activator minoxidil caused a reduction of OC cells proliferation and tumor growth in an OC xenograft model, producing mitochondrial disruption and severe DNA damage [49]. Thus, repurposing of regulatory agency approved K⁺ channel activators could pave the way for novel therapeutic approach in OC, able to improve OC patients' benefits.

Regarding the K2P channel family, the main members involved in OC are the TWIK-related acid-sensitive potassium channel 3 (TASK-3 encoded by *KCNK9*) and the TWIK- related K⁺ channel type 1 and 2 (TREK-1 encoded by *KCNK2* and TREK-2 encoded by *KCNK10*). Many studies have established *KCNK9* as a proto-oncogene, whose amplification promotes cell proliferation [50]. In keeping with this, an artificial point mutation (G95E) in *KCNK9* abolished TASK-3 channel activity and oncogenic activities, including proliferation, resistance to apoptosis, and tumor cell growth [51]. Pharmacological inhibition of TASK-3 by methanandamide and zinc reduced cell proliferation and apoptosis in the SKOV-3 and OVCAR-3 cell lines (Figure 1). Conversely, increased immunopositivity of TASK-3 in OC patients was associated with clinically significant increased survival, in contrast with the definition of TASK-3 as an oncogene. Therefore, as for BK channels, TASK-3 channels might have a dual role depending on the level of expression/activity, suggesting that results of K⁺ channel modulation may be quite unpredictable [52].

Interestingly, TREK-1-blocking agents, such as curcumin, reduced OC cells proliferation and increased late apoptosis processes, suggesting that these channels could represent a new therapeutic option for OC that requires further investigation (Figure 1) [38].

As for TREK channels, TREK-1 overexpression was found in OC cells and tissues and was associated to a significant increase in cancer cell proliferation. A variety of natural compounds has been studied for their anticancer properties, including curcumin. This polyphenol extracted from the turmeric plants has gained attention for its antioxidant, anti-inflammatory and antimicrobial activities. Curcumin exhibits antioxidant effect through free-radical-scavenging activity and has been involved in many signaling pathways and molecular targets (growth factors, inflammatory cytokines, and apoptotic proteins) that influence cancer growth [53]. Interestingly, it has been

demonstrated that curcumin is able to block TREK-1 expression, inducing late apoptosis processes and determining a reduction of OC cells proliferation (Figure 1) [54]. Thus, these results pinpoint TREK-1 channel as a potential target for new therapeutic option for OC.

3. Chloride Channels

While most studies have focused on the role of potassium channels up to now, chloride channels have recently gained more attention in carcinogenesis [55]. Like other ion channels, chloride channels and downstream signaling are involved in tumor growth and aggressiveness through the regulation of cell motility, cell cycle progression or apoptosis resistance [56].

Mammalian chloride channels include five classes classified according to their regulation: voltage-gated chloride channels (CLCs), cystic fibrosis transmembrane conductance regulator (CFTR), volume-regulated chloride channels (VRAC), calcium-activated chloride channels (CaCCs), and ligand-gated chloride channels (GABA (y-aminobutyric acid) and glycine-activated) [57]. Additionally, chloride intracellular channel (CLIC) proteins are the most recent Cl- channels to be discovered and are classified separately from other Cl- channels [58].

Chloride channels are involved in many different cellular functions that include the regulation of the cell excitability, ion homeostasis, trans-epithelial fluid transport, pH level, and cell volume regulation. The latter is particularly relevant for cancer cell migration and infiltration [57]. These channels also contribute to the regulation of the cell cycle, probably as key players in the progression from G1 to S phase [59].

The heterogeneity of chloride channel activity suggests a multitask contribution to OC tumorigenesis. The study of the effects of various inhibitors of CLC and VRAC channels on the OC A2780 cell line demonstrated that treatment with niflumic acid (NFA), tamoxifen and 5-nitro-2-(3-phenylpropylamino)-benzoate (NPPB) inhibited cell proliferation and suppressed cell adhesion and invasion (Figure 1) [60][61]. Further in vitro studies also revealed that CIC-3 silencing by antisense oligonucleotides decreased protein levels of MMP-2, MMP-9, and VEGF in culture medium of SKOV3 cells, confirming CIC-3 involvement in proliferation, invasion, and migration of OC cells (Figure 1) [61].

CFTR is a cyclic AMP-dependent chloride ion-conducting channel expressed mainly in epithelial cells, including the female reproductive tract [62]. Xu et al. investigated for the first time the relationship between CFTR and OC, analyzing CFTR expression in human epithelial ovary cells in normal conditions and in cancer through immunohistochemical staining [63][64]. Their results showed that CFTR expression was significantly increased in OC compared with both benign tumor and normal ovarian tissue and that high expression levels were correlated with poor prognosis, advanced histological grades, and an increased CA125 biomarker level. The role of CFTR was also investigated in vitro in SKOV3 and A2780 cells and in vivo. The knockdown of CFTR suppressed the malignant behavior of OC cells, including cell proliferation, colony formation, migration, invasion, and adhesion (Figure 1). Furthermore, inhibition of CFTR inhibition in vivo suppressed xenograft tumor development, pinpointing CFTR as a potential biomarker in OC [65].

Ca²⁺-activated chloride channels are known to be involved in the regulation of carcinogenesis and are believed to be important emerging drug targets in cancer [66][67]. In particular, a member of the Ca²⁺-activated chloride channels, TMEM16A, also known as ANO1, was proposed to contribute to tumor growth and invasion in head and neck squamous cell carcinomas, breast cancer, hepatocellular carcinoma, lung cancer, gastric cancer, and prostate cancer [68][69][70][71][72][73]. TMEM16A is upregulated also in human epithelial OC cells and peripheral blood mononuclear cells (PBMCs) from patients with OC and its inhibition suppresses growth and invasiveness of OC cells and xenograft tumors in mice (Figure 1) [74]. It is worth noting that TMEM16A gene expression is upregulated in PBMCs from patients with serous OC, but reduced in patients after surgical removal of tumor. All these results suggest that the change of TMEM16A gene expression in PBMCs might be a biomarker for diagnosis or monitoring the progression of OC [74]. However, this hypothesis merits further clinical validation. Since its identification as a CaCC, several regulatory mechanisms of TMEM16A chloride channel have been identified. Among these, it was shown that TMEM16A ion conduction could be affected by lipid environment [75][76]. In cancer cells, lipid distribution in microdomains at the plasma membrane is critical for cancer cell proliferation and migration [77]. Recently, Ye and collaborators revealed that TMEM16A-related chloride currents are involved in the distribution and clustering of phosphatidylinositol 4,5-bisphosphate (PIP2) at the membrane [78]. The cleavage of PIP2 by phospholipase C (PLC) promotes the formation of IP₃ that, in turn, activates IP₃R and mobilizes Ca²⁺ from intracellular stores. Thus, disturbing TMEM function or expression may abrogate the formation of such macro-complexes and inhibit the associated downstream signaling pathways [76].

Recent studies have focused attention on TMEM16A channel regulation by the calcium-activated chloride channel regulator 1 (CLCA1) (Figure 1). A growing body of evidence suggests that CaCC-1 may contribute to tumorigenesis in colorectal cancer, pancreatic cancer, and OC [144,149]. Comparative proteomics revealed an increased expression of CaCC-1 in the OC OV-90 cell line and in cell models of tumor aggregate formation (TOV-112D and ES-2). The knockdown of CaCC-1 with the chloride channel blocker NFA or with siRNA reduced the ability of cancer cells to form multicellular aggregates, which is a critical process in metastasis development (Figure 1) [79]. These findings highlight the importance of CaCC-1 in OC progression, and further investigation is warranted to delineate the role of CaCC-1 in the pathogenesis of OC.

An emerging class of chloride channels involved in cancer is represented by the intracellular chloride channels (CLICs) [80]. There are six members in the CLIC family (CLIC1-6), of which CLIC-1, CLIC-3, CLIC-4, and CLIC-5 are the most likely involved in OC. Since CLIC1 and CLIC4 were the first CLIC proteins to be cloned and functionally studied [81], they remain so far the most characterized within the family. These proteins were initially identified as blood biomarkers of OC because human CLIC1 and CLIC4 were found into the blood of xenografted tumor-bearing mice and were significantly elevated in sera of EOC patients [81][82][83][84]. Furthermore, in vitro experiments demonstrated that CLIC1 and CLIC4 knockdown slowed the proliferation and migration of OC cells, indicating a potential role in tumor progression (Figure 1) [85][86]. Singha and collaborators showed that elevated CLIC4 expression, but not CLIC1 expression, was a negative indicator of patient survival, suggesting its potential as a diagnostic and prognostic marker for OC and as a target for therapy [86]. These results have been recently supported by an in silico analysis that correlated the expression of CLIC proteins with cancer related patient mortality. The data showed that low or high CLIC1 expression in the OCs does not influence patient survival, but

that high CLIC3 and CLIC4 expression correlates with poor patient survival in all the six cancers analyzed, including OC [87]. It is worth noting that, although CLIC proteins can form functional chloride channel in lipid bilayer, their actual function in live cells remains elusive [88]. Thus, if their involvement in cancer is documented, there is no available data showing that they act effectively as chloride channels.

4. Calcium Channels

Calcium is an essential signal transduction element involved in many different cell processes such as proliferation, differentiation, growth, cell death, and apoptosis [89]. Calcium fluxes through cytoplasmic and intracellular Ca^{2+} channels are required for cell cycle progression from G1/S phase through mitosis. Conversely, depletion of intracellular calcium blocks the cell cycle in the G0/G1 and S phases [90]. Thus, the remodeling of Ca^{2+} homeostasis is thought of as a critical hallmark of cancer cells. In OC, the calcium channels involved in calcium signaling pathways promoting cancer behaviors are mostly voltage-gated calcium channels (VGCC), non-voltage activated calcium channels (belonging to the TRP superfamily, see Section 6), and intracellular calcium channels such as the IP₃R and Orai families. Among VGCC, T-type Ca^{2+} channels resulted the most studied for its altered expression in several cancers, including OC. Expression of T-type Ca^{2+} channel is greatly increased in OC A2780 and H08910 cell lines compared with their benign counterparts [91][92]. Treatment of these cells with T-type Ca^{2+} channel blockers, such as mibepradil or 3,4-dihydroquinazoline derivatives (e.g., KYS05090), disturbs cell-cycle progression, slows proliferation, and enhances cell death (Figure 1). The mechanism for the induction of those effects involves caspase dependent apoptosis accompanied by a reduction of the antiapoptotic protein, survivin [93]. Survivin is a direct downstream target of the PI3K/AKT pathway: the activation of this pathway induces the inactivation of FOXO-containing transcription repressor complexes and increases expression of survivin. Inhibition of T-type Ca^{2+} channels decreases AKT phosphorylation and modifies FOXO expression, which culminates in reduced survivin expression (Figure 1) [93]. Although the direct molecular connection between T-type channels and the PI3K/AKT/FOXO pathway is still unknown, inhibitors of T-type channels or survivin expression could be developed. Aberrant expression of VGCC in OC was also implicated in OC cell migration. The Ca^{2+} influx mediated by L-type calcium channels was responsible for the activation of the MAPK signaling cascade and ERK signaling pathway, which subsequently led to an increase of OC cells migration [94]. Growing evidence also revealed a tight cross-talk between L-type Ca^{2+} channel and lipid metabolism in OC progression [95]. High level of lysophosphatidic acid (LPA) and arachidonic acid (AA)-derived eicosanoids were found in ascites of OC patients and in vivo studies referred to them as triggers of the Ca^{2+} influx in the OC cells inducing proliferation, migration or drug resistance [96][97]. The ability of LPA to increase cellular Ca^{2+} entry could be due to a variety of signaling pathways. It mainly acts via GPCRs, activating Gq protein and increasing the cytosolic Ca^{2+} concentration through the classic phospholipase C (PLC)-dependent pathway. Recent studies also hypothesized that LPA-induced Ca^{2+} mobilization may be due to a directly modulation by LPA of L-type Ca^{2+} channels and Ca^{2+} activated K⁺ channels [95]. (Indeed, treatment of SKOV-3 OC cells with the L-type Ca^{2+} channel blocker nifedipine significantly blocked LPA-induced cell migration and adhesion in OC cancer [98].

In addition, AA may exert a critical role in regulating cytosolic Ca²⁺ concentration and consequently cell migration and invasion. It was demonstrated that AA directly inhibits TRPC3 (See Section 6) in mammary MCF-7 cells, attenuating breast cancer cell proliferation and migration [99]. However, the link between lipid mediators and Ca²⁺ channels needs to be further investigated in OC in order to suggest novel therapeutic strategies for OC treatment.

Calcium cytosolic concentration is also regulated by intracellular channels allowing Ca²⁺ movement between subcellular compartments. Over the last years, a critical role of the IP3Rs and Orai1-STIM1 in OC progression has become increasingly apparent in numerous studies. IP3Rs are Ca²⁺ channels, mainly located in the endoplasmic reticulum (ER), responsible for Ca²⁺ release from ER into the cytosol after activation by IP3 and modulation by calcium [100]. The close proximity of the ER to mitochondria, lysosomes, and nucleus, has allowed IP3Rs to emerge as critical regulators of cell fate. Three IP3R homologous isoforms, IP3R1s, IP3R2s, and IP3R3s, have been identified as key sites for regulation by pro- and anti-apoptotic factors. The flow of Ca²⁺ specifically from IP3Rs caused mitochondrial permeability transition, thereby releasing several pro-apoptotic factors into the cytoplasm, which activates the apoptotic cascade. The role of IP3R1 isoform in apoptosis induction was widely described in a variety of cancer cells. In OC, it was demonstrated that natural compounds like sulforaphane (SFN) [101] or hormones like melatonin [102] may act as potential chemotherapeutic agents in OC cells lines, partially upregulating IP3R1. In keeping with this, in vivo experiments showed that SFN treatment was able to reduce OC volume in athymic nude mice with OC-induced by A2780 cells injection. When SFN was co-administered with IP3R blocker, tumor volume was higher compared to sole SFN treatment, but still lower than in untreated mice. These results supported the idea that IP3R1 was involved in pro-apoptotic processes, taking part to the tumor suppressor effect in OC [101]. However, the expression profile of IP3R1 remains uncertain in cancer progression because other results showed contrasting actions of this channel in tumor progression and drug resistance. For instance, an enhanced activity of IP3R1 in prostate cancer promotes cell survival and resistance to hormonal deprivation therapy [103]. More recently, it was reported that IP3R1 plays a different role in MDA-MB-231 breast cancer cells compared to JIMT1 breast cancer cells, DLD1 colorectal and A2780 OC cells [104]. Treatment of these cell lines with the VGCC blocker nifedipine induced an increase of cytosolic Ca²⁺ levels in A2780 and DLD1 cells, whereas no changes have been seen in MDA-MB-231 and JIMT1 breast cancer cells. The high cytoplasm Ca²⁺ concentration in A2780 and DLD1 cells was mainly due to the upregulation of Cav1.3 and IP3R1, which allows linking to the activation of apoptotic process. Indeed, IP3R1 silencing determined an increase in proliferation and migration in DLD1 cells, whereas a decrease of these processes have been seen when IP3R1 was silenced in MDA-MB-231. These results support different functions of the IP3R1 in the invasiveness of various cancer cells in response to nifedipine treatment [105]. In stark contrast to IP3R1, it was demonstrated that IP3R3 has anti-apoptotic and proliferative function in several cancers, including OC [105]. IP3R3 expression was upregulated in glioblastoma, colorectal carcinoma, and breast cancer, and was correlated to an enhanced capacity of cancer cell migration and invasion [105]. Indeed, the inhibition or downregulation of IP3R3 also abrogated proliferative and invasive phenotype and extended survival rate [106][107].

The Ca²⁺ release from intracellular stores determined the activation of Store Operated Ca²⁺ Entry (SOCE), a process by which extracellular Ca²⁺ entries into the cytoplasm to refill the internal Ca²⁺ stores. This event is controlled at a molecular level by the canonical TRP channels, the Ca²⁺ release-activated calcium channel protein

1 (ORAI1) and the ER Ca²⁺ sensors STIM1 (stromal interaction molecule 1) and STIM2 [108]. A number of studies have shown that the overexpression of these channels is positively correlated to cell proliferation, migration, and invasion in several cancers, including glioblastoma, cervical cancer, and breast cancer [109][110]. Recently, it was found a tight connection between Orai1-STIM1 expression and placental growth factor (PIGF) in OC cells. Exposure of OC cells to PIGF triggered a significant increase of Orai1 as well as STIM1 transcript and protein levels, enhancing SOCE and the expression of Hypoxia Inducible Factor HIF1 α , which may support proliferation and survival of metastatic tumor cells [111]. Thus, targeting all these pathways may offer new therapeutic options for the treatment of OC.

5. TRP Channels

TRP channels are calcium-permeable and non-selective ion channels grouped into six subfamilies, including TRPC ("C" for canonical), TRPV ("V" for vanilloid), TRPM ("M" for melastatin), TRPA ("A" for ankyrin), TRPP ("P" for polycystic), and TRPML ("ML" for mucolipin) [112]. TRP subfamilies could be involved in calcium influx and downstream pathways, regulating cell survival, proliferation, and apoptosis. Upregulation or downregulation of TRP channels is linked to various phases of tumor progression, and strongly correlates with clinical parameters, such as overall survival and progression-free survival.

Immunohistochemistry of healthy and OC tissue revealed that TRPV1 and TRPV6 channels were markedly increased in OC tissues compared to healthy tissues [113][114]. TRPV6 mRNA and protein are also highly expressed in all OC subtypes, both in early and in late stage, indicating TRPV6 mRNA levels as a promising biomarker for the presence of OC [115]. In prostate and breast cancers, elevated TRPV6 and the resulting sustained elevation of intracellular Ca²⁺ promote the activation of Nuclear Factor of Activated T-cell transcription factor (NFAT), which activates genes involved in cell proliferation and migration [116][117]. Thus, reduction of TRPV6 activity by decreasing its expression or by pharmacological intervention may represent a way to develop new therapeutic molecular interventions in cancer. Indeed, targeting TRPV6 with specific channel blockers (such as soricidin and its derivates SOR-C13 and SOR-C27) reduces growth of OC SKOV-3 tumor xenografts in mice, further supporting this channel as a viable target (Figure 1) [115]. Moreover, the peptide inhibitor of TRPV6, SOR-C13, completed a phase 1 clinical trial [118], resulting as safe and well tolerated in patients with advanced epithelial OC, without evidence of hematological, neurological or other significant toxicities.

Evidences on the role of TRPC channels in cancer are expanded with their detection in a number of different tumors including breast, ovary, colon, thyroid, and pancreas [119][120][121][122][123]. Several spliced variants of TRPC1, TRPC3, TRPC4, and TRPC6 were found in OC [124]. TRPC3 channels resulted overexpressed both in human OC tissue and in OC cells, suggesting a role in OC progression. Accordingly, knockdown of these channels reduced the proliferation of OC cells and suppressed tumor formation in mice after OC cells injection, suggesting a functional role for TRP-mediated Ca²⁺ entry in cell proliferation and invasion [125]. Recently, it was also demonstrated that TRPC3 expression is upregulated in OC cells after estrogen stimulation, promoting proliferation and migration of cells through intracellular calcium signaling pathways [126].

Lastly, TRPM7 is highly expressed in OC cells and tissues and its expression level is significantly associated with poor prognosis in patients with OC [127]. Silencing of TRPM7 with siRNA inhibited OC cells proliferation, migration, and invasion, through direct regulation of calcium influx that, in turn, modulates many signaling transduction pathways including Akt, Src, and p38 (Figure 1) [128][129].

6. Ion Channels and Drug Resistance

The high mortality rate in OC is related to late diagnosis in advanced clinical stages, metastasis in the peritoneal cavity, and resistance to chemotherapy [130]. It is estimated that 75% of patients who are highly responsive to initial therapy, relapse within 2 years because of acquired resistance [131]. For instance, resistance to platinum agents remains a significant obstacle to overcome. Various mechanisms causing cell drug resistance have been described *in vitro* and *in vivo*.

Ion channels have been shown to play key roles in chemosensitivity and several ion channels have been shown to modify sensitivity to neoplastic drugs by modulating apoptosis pathways or the efficiency of drug permeation due to electrochemical gradients. The abnormal expression of antiapoptotic proteins is often associated with OC aggressiveness and resistance to cytotoxic therapy [132], suggesting that intervention on this pathway may be an effective approach to increase OC response to therapy. In this respect, inhibition of T-type Ca^{2+} channels with the antagonist mibepradil or siRNA induces proapoptotic responses through reduced expression of the antiapoptotic gene *survivin* [94]. Thus, the inhibition of *survivin* expression enhances the ability of platinum agents to increase DNA double strand breaks and induce apoptosis in OC [133]. Indeed, combining carboplatin with mibepradil synergistically increases apoptosis and inhibits growth of platinum-resistant OC cells *in vitro*. More importantly, *in vivo* pretreatment with mibepradil in carboplatin-based therapy reduced xenograft tumor burden in mice model of platinum-resistant OC (Figure 2) [94], supporting the idea that T-type Ca^{2+} channel blockers could be useful to improve patients' response to platinum agents. Few studies focused on the roles of Orai1 and STIM1 in OC drug resistance. One work from Schmidt and collaborators showed that these channels are highly expressed in the A2780 cisplatin-resistant OC cells compared to A2780 non-resistant cells, leading to elevated SOCE [134]. Pharmacological inhibition of Orai1 with 2-aminoethoxydiphenyl borate (2-APB) increased cisplatin-induced apoptosis of A2780 cisplatin-resistant OC cells and abrogated the differences in cisplatin sensitivity between them and A2780 non-resistant cells. Taken together, these data suggest that enhanced SOCE could contribute to drug resistance in OC cells and might become potential targets in OC treatment [133]. The role of TRPC1 channels in platinum resistance has also been investigated. A marked decrease in TRPC1 mRNA levels was measured in cisplatin and carboplatin-resistant OC cells compared with their sensitive counterparts [135], suggesting a TRPC1 contribution to drug resistance (Figure 2). A mechanism that contributes to cisplatin resistance in OC is autophagy. A tight interaction was found between TRPC1 and two proteins involved in ERK-mediated autophagy (PIK3C3 and SPARCL1) [136][137], confirming the role of TRPC1 in the regulation of autophagy and in drug resistance in OC. More recently, it was demonstrated that Nav1.5 was involved in cisplatin resistance in OC. *In vitro* experiments showed that lidocaine enhanced OC cells sensitivity to cisplatin by blocking Nav1.5 and FAK/Paxillin signaling pathway. The OC cells treated with cisplatin and lidocaine combination expressed higher levels of apoptotic

proteins such as caspase-3 and 8, and lower level of the antiapoptotic protein Bcl-2 [27]. Thus, targeting these channels maybe a promising strategy for overcoming cisplatin resistance.

Figure 2. Ion channels roles in OC drug resistance. Table on the left summarizes the main ion channels likely involved in OC drug resistance, focusing on their expression level on OC cells and tissues. Table on the right lists the effects of ion channel blockers on chemosensitivity to platinum-based agents.

Drug resistance phenomenon in OC is reportedly attributed to the existence of cancer stem cells (CSC), which establish a reservoir of cells with the exclusive capacity of self-renew. These cells are generally resistant to conventional anticancer treatments [138]. Thus, the discovery of drugs that selectively target CSCs may help preventing cancer recurrence, improving the survival rate of OC patients. Indeed, Lee and collaborators screened an FDA-approved compound library of ~1000 compounds on OC stem cells (CSCs) and found that four calcium channel blockers (manidipine, lacidipine, benidipine, and Iomerizine) had anticancer effects against ovarian CSC by reducing stemness and inducing apoptosis in CSCs (Figure 2) [139]. These results suggest that calcium channels may represent a selective target for new therapeutic approaches in the prevention of OC recurrence.

The importance of K⁺ influx in cisplatin resistance in OC was showed for the first time by Sharp et al., who demonstrated that the K⁺ ionophore amphotericin B could sensitize cells to cisplatin and other platinum-containing agents [140]. Reduction of intracellular K⁺ concentration by amphotericin B or the Na-K-Cl cotransporter inhibitor bumetanide boost cisplatin-induced apoptosis [141][142]. These early results suggested that there might be a tight correlation between the expression levels of K⁺ channels and the sensitivity to cytotoxic drugs. Accordingly, hEag1 channel silencing increased the chemosensitivity of SKOV3 cell lines to cisplatin, promoting apoptosis through the NF-κB pathway (Figure 2) [143]. The decreased hEag1 expression found in OC tissue of patients treated with cisplatin-based adjuvant chemotherapy was correlated with a favorable prognosis and predicted higher sensitivity to cisplatin treatment.

The KCNN3 gene encoding the calcium-activated K⁺ channel KCa2.3 was predicted to be one of 1298 genes contributing to drug resistance in OC [144]. Using immunohistochemistry and big data analyses, Liu and collaborators found a significantly lower expression of KCNN3 in drug-resistant OC tissues, confirming an involvement of KCa2.3 channels in drug resistance (Figure 2) [33][46]. It was further hypothesized that KCa2.3 and TRPC1 may act in concert in drug resistance through inhibition of drug-induced cell death [145]. This promising combination strategy warrants further study. In addition, the decreased KCNMA1 expression in OC cell lines was associated with minor sensitivity to platinum-based chemotherapy (Figure 2) [146].

Regarding chloride channels, OC patients with lower CLIC1 levels showed increased sensitivity to cisplatin treatment [84], likely because CLIC1 participates in the regulation of anti-apoptosis signaling pathways in response to chemotherapy (Figure 2).

Altogether, these results shed new light on how ion channels contribute to chemotherapy resistance, providing a potential therapeutic alternative for OC patients who frequently succumb to platinum resistant disease.

7. Concluding remarks and perspectives

A variety of ion channels are remarkably expressed in human OC cells and tissues, where they are involved in the development of various malignant behaviors, including proliferation, migration, invasion, and apoptosis resistance. For some of these channels, a correlation of their expression levels with prognosis or resistance to cytotoxic drugs has also been reported, supporting a putative role as predictive markers for combination chemotherapy regimens. In line with this, the specific inhibition of some ion channels increases the cytotoxicity induced by different chemotherapeutic drugs, thus opposing tumor resistance. Accumulating data have revealed that the aberrant expression of ion channels in OC greatly contribute to the main hallmarks of OC process, from initial proliferation to metastasis to chemoresistance. The T-type Ca^{2+} channels, for example, are overexpressed in ovarian cancer where they are involved in tumor growth and its chemoresistance to platinum-based drugs [46]. Indeed, blocking T-type Ca^{2+} channels with mibepradil during administration of carboplatin sensitizes platinum-resistant ovarian tumors to the anticancer drug-mediated toxicity [114]. Most studies reported in this review describe the implication of one channel in the regulation of OC development and progression. However, it is likely that channels may work in tandem or in more complex network to promote specific cell function. Studies about complexes of ion channels are emerging to explain their involvement in the control of these processes. For example, voltage-gated Na^+ channels can be associated with NCX and NHE1 cotransports to stimulate tumor cell metabolism and promote cancer cell invasion and metastatic progression [16]. In colorectal cancer cells, the simultaneous activation of KCa3.1 and inhibition of Kv11.1 with riluzole synergistically reduces tumor proliferation while increasing apoptosis and overcoming cisplatin resistance [114]. Thus, the identification of the complete set of ion channels involved in OC and their interrelationships might allow to set up new promising therapeutic approaches, pointing for instance to a combination of different channel inhibitors or employing the drugs together with chemotherapeutics. Furthermore, the use of selective modulators directed on transmembrane ion channels allows reaching easily the target without entering the cells. This means that these drugs are not subjected to efflux pumps, bypassing this major chemoresistance mechanism. Membrane localization is also an invaluable opportunity for the use of selective monoclonal antibodies [115].

Importantly, many of the available pharmacological modulators of ion channels are already successfully used for the treatment of various non-oncological diseases, which may facilitate their repurposing in OC treatment. The use of clinically approved drugs could bring many benefits including known safety profile and well understood pharmacokinetics in humans, as well as saving time and money in drug development.

However, there is a risk that these modulators may have off-target effects on normal physiological function, because many of the channels identified in OC cells are also expressed in healthy normal cells. In this regard, it would be important to focus on and selectively target those ion channels having aberrant overexpression in the tumor tissues compared to the normal counterpart tissue or showing cancer-specific isoforms/splice variants. Antibodies or small molecules that target tumor-specific isoforms may represent a way to increase the sensitivity

and specificity of ion channel-targeted therapies in cancer. An example is the highly selective monoclonal antibody against the oncogenic Kv 10.1 potassium channel which has been demonstrated to inhibit tumor growth both in xenograft models of human melanoma cells and in xenografted human pancreatic cancer [116]. Antibodies or fragments against ion channels might be also useful for cancer diagnosis[117]. However, the monoclonal antibody-based approach is limited by the continuous dynamics of the tumor environment and by the size-dependent tissue penetration and further investigation is needed to overcome these limitations. In addition, it is worth to note that the activity and expression of ion channels may vary in a cell cycle-dependent manner and between tumor cell subtypes (e.g. cancer stem cells versus parenchymal cells), limiting the efficacy of ion channel modulators.

The study of ion channels expression in OC has been steadily/increasingly gaining interest with time, also because of their involvement in cancer drug resistance. Several benefits could be obtained through a combined strategy of ion channel-targeting drugs and chemotherapy, improving pharmacological response in chemoresistant OC patients. Further studies are warranted in animal models and in humans, in order to develop new specific therapeutic approaches.

References

1. Hernandez, C.M.; Richards, J.R. Physiology, Sodium Channels; StatPearls Publishing: Treasure Island, FL, USA, 2020.
2. Djamgoz, M.B.A.; Fraser, S.P.; Brackenbury, W.J. In Vivo Evidence for Voltage-Gated Sodium Channel Expression in Carcinomas and Potentiation of Metastasis. *Cancers* **2019**, *11*, 1675.
3. Nelson, M.; Yang, M.; Millican-Slater, R.; Brackenbury, W. Nav1.5 regulates breast tumor growth and metastatic dissemination in vivo. *Oncotarget* **2015**, *6*, 32914–32929.
4. House, C.; Vaske, C.; Schwartz, A.; Obias, V.; Frank, B.; Luu, T.; Sarvazyan, N.; Irby, R.; Strausberg, R.L.; Hales, T.G.; et al. Voltage gated Na⁺ channel SCN5A is a key regulator of a gene transcriptional network that controls colon cancer invasion. *Cancer Res.* **2010**, *70*, 6957–6967.
5. Suy, S.; Hansen, T.; Auto, H.D.; Kallakury, B.V.S.; Dailey, V.; Danner, M.; Macarthur, L.; Zhang, Y.; Miessau, M.J.; Collins, S.P.; et al. Expression of Voltage-gated sodium channel Na1.8 in human prostate cancer is associated with high histological grade. *J. Clin. Exp. Oncol.* **2012**, *1*, 1–13.
6. He, M.; Liu, S.; Gallolu Kankamalage, S.; Borromeo, M.D.; Girard, L.; Gazdar, A.F.; Minna, J.D.; Johnson, J.E.; Cobb, M.H. The Epithelial Sodium Channel (αENaC) Is a Downstream Therapeutic Target of ASCL1 in Pulmonary Neuroendocrine Tumors. *Transl. Oncol.* **2018**, *11*, 292–299.
7. Gao, R.; Shen, Y.; Cai, J.; Lei, M. Expression of voltage-gated sodium channel subunit in human ovarian cancer. *Oncol. Rep.* **2010**, *23*, 1293–1299.

8. Gillet, L.; Roger, S.; Besson, P.; Lecaille, F.; Gore, J.; Bougnoux, P.; Lalmanach, G.; Le Guennec, J.Y. Voltage-gated sodium channel activity promotes cysteine cathepsin-dependent invasiveness and colony growth of human cancer cells. *J. Biol. Chem.* 2009, 284, 8680–8691.

9. Yang, M.; Kozminski, D.J.; Wold, L.A.; Modak, R.; Calhoun, J.D.; Isom, L.L.; Brackenbury, W.J. Therapeutic potential for phenytoin: Targeting Na(v)1.5 sodium channels to reduce migration and invasion in metastatic breast cancer. *Breast Cancer Res. Treat.* 2012, 134, 603–615.

10. Liskova, V.; Hudecova, S.; Lencesova, L.; Iuliano, F.; Sirova, M.; Ondrias, K.; Pastorekova, S.; Krizanova, S. Type 1 Sodium Calcium Exchanger Forms a Complex with Carbonic Anhydrase IX and Via Reverse Mode Activity Contributes to pH Control in Hypoxic Tumors. *Cancers* 2019, 11, 1139.

11. Besson, P.; Driffort, V.; Bon, É.; Gradek, F.; Chevalier, S.; Roger, S. How do voltage-gated sodium channels enhance migration and invasiveness in cancer cells? *Biochim. Biophys. Acta* 2015, 1848, 2493 2501.

12. Greco, M.R.; Antelmi, E.; Busco, G.; Guerra, L.; Rubino, R.; Casavola, V.; Reshkin, S.J.; Cardone, R.A. Protease activity at invadopodial focal digestive areas is dependent on NHE1-driven acidic pH. *Oncol. Rep.* 2014, 31, 940–946.

13. Matarrese, P.; Ascione, B.; Ciarlo, L.; Vona, R.; Leonetti, C.; Scarsella, M.; Mileo, A.M.; Catricalà, C.; Paggi, M.G.; Malorni, W. Cathepsin B inhibition interferes with metastatic potential of human melanoma: An in vitro and in vivo study. *Mol. Cancer* 2010, 9, 207.

14. Brisson, L.; Driffort, V.; Benoist, L.; Poet, M.; Counillon, L.; Antelmi, E.; Rubino, R.; Besson, P.; Labbal, F.; Chevalier, S.; et al. NaV1.5 Na⁺ channels allosterically regulate the NHE-1 exchanger and promote the activity of breast cancer cell invadopodia. *J. Cell Sci.* 2013, 126, 4835–4842.

15. Harguindeguy, S.; Arranz, J.L.; Polo Orozco, J.D.; Rauch, C.; Fais, S.; Cardone, R.A.; Reshkin, S.J. Cariporide and other new and powerful NHE1 inhibitors as potentially selective anticancer drugs —An integral molecular/biochemical/metabolic/clinical approach after one hundred years of cancer research. *J. Transl. Med.* 2013, 11, 282.

16. Sanhueza, C.; Araos, J.; Naranjo, L.; Toledo, F.; Beltrán, A.R.; Ramírez, M.A.; Gutiérrez, J.; Pardo, F.; Leiva, A.; Sobrevia, L. Sodium/proton exchanger isoform 1 regulates intracellular pH and cell proliferation in human ovarian cancer. *Biochim. Biophys. Acta Mol. Basis Dis.* 2017, 1863, 81–91.

17. Yi, Y.H.; Ho, P.Y.; Chen, T.W.; Lin, W.J.; Gukassyan, V.; Tsai, T.H.; Wang, D.W.; Lew, T.S.; Tang, C.Y.; Lo, S.J.; et al. Membrane targeting and coupling of NHE1-integrinαIIbβ3-NCX1 by lipid rafts following integrin-ligand interactions trigger Ca²⁺ oscillations. *J. Biol. Chem.* 2009, 284, 3855–3864.

18. Xu, J.; Ji, B.; Wen, G.; Yang, Y.; Jin, H.; Liu, X.; Xie, R.; Song, W.; Song, P.; Dong, H.; et al. Na+/H⁺ exchanger 1, Na+/Ca²⁺ exchanger 1 and calmodulin complex regulates interleukin 6-mediated cellular behavior of human hepatocellular carcinoma. *Carcinogenesis*. 2016, 37, 290–300.

19. Bosetti, C.; Negri, E.; Franceschi, S.; Pelucchi, C.; Talamini, R.; Montella, M.; Conti, E.; La Vecchia, C. Diet and ovarian cancer risk: A case-control study in Italy. *Int. J. Cancer* 2001, 93, 911–915.

20. Merritt, M.A.; Cramer, D.W.; Missmer, S.A.; Vitonis, A.F.; Titus, L.J.; Terry, K.L. Dietary fat intake and risk of epithelial ovarian cancer by tumor histology. *Br. J. Cancer* 2014, 110, 1392–1401.

21. Sharma, A.; Belna, J.; Logan, J.; Espat, J.; Hurteau, J.A. The effects of ω-3 fatty acids on growth regulation of epithelial ovarian cancer cell lines. *Gynecol. Oncol.* 2005, 99, 58–64.

22. Liu, J.; Liu, D.; Liu, J.; Zhao, C.; Yao, S.; Hong, L. Blocking the Nav1.5 channel using eicosapentaenoic acid reduces migration and proliferation of ovarian cancer cells. *Int. J. Oncol.* 2018, 53, 855–865.

23. Martin, F.; Ufodiama, C.; Watt, I.; Bland, M.; Brackenbury, W.J. Therapeutic value of voltage gated sodium channel inhibitors in breast, colorectal and prostate cancer: A systematic review. *Front. Pharmacol.* 2015, 6, 273.

24. Xu, G.; Fang, Z.; Clark, L.; Sun, W.; Yin, Y.; Zhang, R.; Sullivan, S.; Tran, A.; Kong, W.; Wang, J.; et al. Topiramate exhibits anti-tumorigenic and metastatic effects in ovarian cancer cells. *Am. J. Transl. Res.* 2018, 10, 1663–1676.

25. Exadaktylos, A.K.; Buggy, D.J.; Moriarty, D.C.; Mascha, E.; Sessler, D.I. Can anesthetic technique for primary breast cancer surgery affect recurrence or metastasis? *Anesthesiology* 2006, 105, 660–664.

26. Xuan, W.; Zhao, H.; Hankin, J.; Chen, L.; Yao, S.; Ma, D. Local anesthetic bupivacaine induced ovarian and prostate cancer apoptotic cell death and underlying mechanisms in vitro. *Sci. Rep.* 2016, 6, 26277.

27. Liu, C.; Yu, M.; Li, Y.; Wang, H.; Xu, C.; Zhang, X.; Li, M.; Guo, H.; Ma, D.; Guo, X. Lidocaine inhibits the metastatic potential of ovarian cancer by blocking Na V 1.5-mediated EMT and FAK/Paxillin signaling pathway. *Cancer Med.* 2020, doi:10.1002/cam4.3621.

28. Kuang, Q.; Purhonen, P.; Hebert, H. Structure of potassium channels. *Cell Mol. Life Sci.* 2015, 72, 3677–3693.

29. Zhanping, W.; Xiaoyu, P.; Na, C.; Shenglan, W.; Bo, W. Voltage-gated K⁺ channels are associated with cell proliferation and cell cycle of ovarian cancer cell. *Gynecol. Oncol.* 2007, 104, 455–460.

30. Asher, V.; Khan, R.; Warren, A.; Shaw, R.; Schalkwyk, G.V.; Bali, A.; Sowter, H.M. The Eag potassium channel as a new prognostic marker in ovarian cancer. *Diagn. Pathol.* 2010, 5, 78.

31. Wonderlin, W.M.; Strobl, J.S. Potassium channels, proliferation and G1 progression. *J. Membr. Biol.* 1996, 154, 91–107.

32. Pardo, L.A. Voltage-gated potassium channels in cell proliferation. *Physiology* 2004, 19, 285–292.

33. Iamshanova, O.; Fiorio Pla, A.; Prevarskaya, N. Molecular mechanisms of tumour invasion: Regulation by calcium signals. *J. Physiol.* 2017, 595, 3063–3075.

34. He, S.; Moutaoufik, M.; Islam, S.; Persad, A.; Wu, A.; Aly, K.A.; Fonge, H.; Babu, M.; Cayabyab, F.S. HERG channel and cancer: A mechanistic review of carcinogenic processes and therapeutic potential. *Biochim. Biophys. Acta Rev. Cancer* 2020, 1873, 188355.

35. Zhi, D.; Zhou, K.; Yu, D.; Fan, X.; Zhang, J.; Li, X.; Dong, M. hERG1 is involved in the pathophysiological process and inhibited by berberine in SKOV3 cells. *Oncol. Lett.* 2019, 17, 5653–5661.

36. Asher, V.; Warren, A.; Shaw, R.; Sowter, H.; Bali, A.; Khan, R. The role of Eag and HERG channels in cell proliferation and apoptotic cell death in SK-OV-3 ovarian cancer cell line. *Cancer Cell Int.* 2011, 11, 6.

37. Arcangeli, A.; Crociani, O.; Lastraioli, E.; Masi, A.; Pilozzi, S.; Becchetti, A. Targeting ion channels in cancer: A novel frontier in antineoplastic therapy. *Curr. Med. Chem.* 2009, 16, 66–93.

38. Gómez-Varela, D.; Zwick-Wallasch, E.; Knötgen, H.; Sánchez, A.; Hettmann, T.; Ossipov, D.; Weseloh, R.; Contreras-Jurado, C.; Rothe, M.; Stühmer, W.; et al. Monoclonal antibody blockade of the human Eag1 potassium channel function exerts antitumor activity. *Cancer Res.* 2007, 67, 7343–7349.

39. Chelintseva, E.; Djamgoz, M.B.A. Mesenchymal stem cell differentiation: Control by calcium-activated potassium channels. *J. Cell Physiol.* 2018, 233, 3755–3768.

40. Liu, X.; Chang, Y.; Reinhart, P.H.; Sontheimer, H.; Chang, Y. Cloning and characterization of glioma BK, a novel BK channel isoform highly expressed in human glioma cells. *J. Neurosci.* 2002, 22, 1840–1849.

41. Bloch, M.; Ousingsawat, J.; Simon, R.; Schraml, P.; Gasser, T.C.; Mihatsch, M.J.; Kunzelmann, K.; Bubendorf, L. KCNMA1 gene amplification promotes tumor cell proliferation in human prostate cancer. *Oncogene* 2007, 26, 2525–2534.

42. Oeggerli, M.; Tian, Y.; Ruiz, C.; Wijker, B.; Sauter, G.; Obermann, E.; Güth, U.; Zlobec, I.; Sausbier, M.; Kunzelmann, K.; et al. Role of KCNMA1 in breast cancer. *PLoS ONE* 2012, 7, e41664.

43. Han, X.; Wang, F.; Yao, W.; Xing, H.; Weng, D.; Song, X.; Chen, G.; Xi, L.; Zhu, T.; Zhou, J.; et al. Heat shock proteins and p53 play a critical role in K⁺ channel-mediated tumor cell proliferation and apoptosis. *Apoptosis* 2007, 12, 1837–1846.

44. Zylacz, M.; King, F.W.; Wawrzynow, A. Hsp70 interactions with the p53 tumour suppressor protein. *Embo J.* 2001, 20, 4634–4638.

45. Han, X.; Xi, L.; Wang, H.; Huang, X.; Ma, X.; Han, Z.; Wu, P.; Ma, X.; Lu, Y.; Wang, G.; et al. The potassium ion channel opener NS1619 inhibits proliferation and induces apoptosis in A2780 ovarian cancer cells. *Biochem. Biophys. Res. Commun.* 2008, 375, 205–209.

46. Liu, X.; Wei, L.; Zhao, B.; Cai, X.; Dong, C.; Yin, F. Low expression of KCNN3 may affect drug resistance in ovarian cancer. *Mol. Med. Rep.* 2018, 18, 1377–1386.

47. Robles-Martinez, L.; Garay, E.; Martel-Gallegos, M.G.; Cisneros-Mejorado, A.; Pérez-Montiel, D.; Lara, A.; Arellano, R. Kca3.1 Activation Via P2y2 Purinergic Receptors Promotes Human Ovarian Cancer Cell (Skov-3) Migration. *Sci. Rep.* 2017, 7, 4340.

48. Tinker, A.; Aziz, Q.; Li, Y.; Specterman, M. ATP-Sensitive Potassium Channels and Their Physiological and Pathophysiological Roles. *Compr. Physiol.* 2018, 8, 1463–1511.

49. Fukushiro-Lopes, D.; Hegel, A.D.; Russo, A.; Senyuk, V.; Liotta, M.; Beeson, G.C.; Beeson, C.C.; Burdette, J.; Potkul, R.K.; Gentile, S. Repurposing Kir6/SUR2 Channel Activator Minoxidil to Arrests Growth of Gynecologic Cancers. *Front. Pharmacol.* 2020, 11, 577.

50. Mu, D.; Chen, L.; Zhang, X.; See, L.H.; Koch, C.M.; Yen, C.; Tong, J.J.; Spiegel, L.; Nguyen, K.C.Q.; Servoss, A.; et al. Genomic amplification and oncogenic properties of the KCNK9 potassium channel gene. *Cancer Cell*. 2003, 3, 297–302.

51. Pei, L.; Wiser, O.; Slavin, A.; Mu, D.; Powers, S.; Jan, L.Y.; Hoey, T. Oncogenic potential of TASK3 (KCNK9) depends on K⁺ channel function. *Proc. Natl Acad Sci USA* 2003, 100, 7803–7807.

52. Innamaa, A.; Jackson, L.; Asher, V.; Van Shalkwyk, G.; Warren, A.; Hay, D.; Bali, A.; Sowter, H.; Khan, R. Expression and prognostic significance of the oncogenic K2P potassium channel KCNK9 (TASK-3) in ovarian carcinoma. *Anticancer Res.* 2013, 33, 1401–1408.

53. Giordano, A.; Tommonaro, G. Curcumin and Cancer. *Nutrients* 2019, 11, 2376.

54. Innamaa, A.; Jackson, L.; Asher, V.; Van Schalkwyk, G.; Warren, A.; Keightley, A.; Hay, D.; Bali, A.; Sowter, H.; Khan, R. Expression and effects of modulation of the K2P potassium channels TREK-1 (KCNK2) and TREK-2 (KCNK10) in the normal human ovary and epithelial ovarian cancer. *Clin. Transl. Oncol.* 2013, 15, 910–918.

55. Cuddapah, V.A.; Sontheimer, H. Ion channels and transporters (corrected) in cancer. 2. Ion channels and the control of cancer cell migration. *Am. J. Physiol. Cell Physiol.* 2011, 301, C541–C549.

56. Prevarskaya, N.; Skryma, R.; Shuba, Y. Ion channels and the hallmarks of cancer. *Trends Mol. Med.* 2010, 16, 107–121.

57. Verkman, A.; Galietta, L. Chloride channels as drug targets. *Nat. Rev. Drug Discov.* 2009, 8, 153–171.

58. Ashley, R.H. Challenging accepted ion channel biology: p64 and the CLIC family of putative intracellular anion channel proteins (Review). *Mol. Membr. Biol.* 2003, 20, 1–11.

59. Ye, D.; Luo, H.; Lai, Z.; Zou, L.; Zhu, L.; Mao, J.; Jacob, T.; Ye, W.; Wang, L.; Chenc, L. CIC-3 Chloride Channel Proteins Regulate the Cell Cycle by Up-regulating cyclin D1-CDK4/6 through Suppressing p21/p27 Expression in Nasopharyngeal Carcinoma Cells. *Sci. Rep.* 2016, 6, 30276.

60. Li, M.; Wang, Q.; Lin, W.; Wang, B. Regulation of ovarian cancer cell adhesion and invasion by chloride channels. *Int. J. Gynecol. Cancer* 2009, 19, 526–530.

61. Li, M.; Wu, D.B.; Wang, J.; Chen, L. CLC-3 Cl-channel-mediated invasion and migration of human ovarian cancer cells. *Eur. J. Gynaecol. Oncol.* 2016, 37, 689–695.

62. Tizzano, E.F.; Silver, M.M.; Chitayat, D.; Benichou, J.C.; Buchwald, M. Differential cellular expression of cystic fibrosis transmembrane regulator in human reproductive tissues. Clues for the infertility in patients with cystic fibrosis. *Am. J. Pathol.* 1994, 144, 906–914.

63. Li, M.; Wang, B.; Lin, W. Cl-channel blockers inhibit cell proliferation and arrest the cell cycle of human ovarian cancer cells. *Eur. J. Gynaecol. Oncol.* 2008, 29, 267–271.

64. Xu, J.; Yong, M.; Li, J.; Dong, X.; Yu, T.; Fu, X.; Hu, L. High level of CFTR expression is associated with tumor aggression and knockdown of CFTR suppresses proliferation of ovarian cancer in vitro and in vivo. *Oncol. Rep.* 2015, 33, 2227–2234.

65. Xu, J.; Lin, L.; Yong, M.; Dong, X.; Yu, T.; Hu, L. Adenovirus mediated overexpression of cystic fibrosis transmembrane conductance regulator enhances invasiveness and motility of serous ovarian cancer cells. *Mol. Med. Rep.* 2016, 13, 265–272.

66. Lang, F.; Stournaras, C. Ion channels in cancer: Future perspectives and clinical potential. *Philos. Trans. R Soc. Lond. B Biol. Sci.* 2014, 369, 20130108.

67. Stock, C.; Schwab, A. Ion channels and transporters in metastasis. *Biochim. Biophys. Acta* 2015, 1848, 2638–2646.

68. Liu, W.; Lu, M.; Liu, B.; Huang, Y.; Wang, K.W. Inhibition of Ca(2+)-activated Cl(–) channel ANO1/TMEM16A expression suppresses tumor growth and invasiveness in human prostate carcinoma. *Cancer Lett.* 2012, 326, 41–51.

69. Britschgi, A.; Bill, A.; Brinkhaus, H.; Rothwell, C.; Clay, I.; Duss, S.; Rebhan, M.; Raman, P.; Guy, C.T.; Wetzel, K. Calcium activated chloride channel ANO1 promotes breast cancer progression by activating EGFR and CAMK signaling. *Proc. Natl. Acad. Sci. USA* 2013, 110, 1026–1034.

70. Bill, A.; Gutierrez, A.; Kulkarni, S.; Kemp, C.; Bonenfant, D.; Voshol, H.; Duvvuri, U.; Gaither, A. ANO1/ TMEM16A interacts with EGFR and correlates with sensitivity to EGFR-targeting therapy in head and neck cancer. *Oncotarget* 2015, 6, 9173–9188.

71. Jia, L.; Liu, W.; Guan, L.; Lu, M.; Wang, K.W. Inhibition of calcium activated Chloride Channel ANO1/TMEM16A suppresses tumor growth and invasion in human lung cancer. *PLoS ONE* 2015, 10, e0136584.

72. Liu, F.; Cao, Q.H.; Lu, D.J.; Luo, B.; Lu, X.F.; Luo, R.C.; Wang, X.C. TMEM16A overexpression contributes to tumor invasion and poor prognosis of human gastric cancer through TGF β signaling. *Oncotarget* 2015, 6, 11585–11599.

73. Deng, L.; Yang, J.; Chen, H.; Ma, B.; Pan, K.; Su, C.; Xu, F.; Zhang, J. Knockdown of TMEM16A suppressed MAPK and inhibited cell proliferation and migration in hepatocellular carcinoma. *Oncotargets Ther.* 2016, 9, 325–333.

74. Liu, Z.; Zhang, S.; Hou, F.; Zhang, C.; Gao, J.; Wang, K. Inhibition of Ca^{2+} -activated chloride channel ANO1 suppresses ovarian cancer through inactivating PI3K/Akt signaling. *Int. J. Cancer.* 2019, 144, 2215–2226.

75. Schreiber, R.; Ousingsawat, J.; Wanitchakool, P.; Sirianant, L.; Benedetto, R.; Reiss, K.; Kunzelmann, K. Regulation of TMEM16A/ANO1 and TMEM16F/ANO6 ion currents and phospholipid scrambling by Ca^{2+} and plasma membrane lipid. *J. Physiol.* 2018, 596, 217–229.

76. Crottès, D.; Yeh Jan, L. The multifaceted role of TMEM16A in cancer. *Cell Calcium* 2019, 82, 102050.

77. Gueguinou, M.; Gambade, A.; Felix, R.; Chantome, A.; Fourbon, Y.; Bougnoux, P.; Weber, G.; Potier-Cartereau, M.; Vandier, C. Lipid rafts, $\text{KCa}/\text{ClCa}/\text{Ca}^{2+}$ channel complexes and EGFR signaling: Novel targets to reduce tumor development by lipids? *Biochim. Biophys. Acta* 2015, 1848, 2603–2620.

78. Ye, W.; Han, T.W.; Nassar, L.M.; Zubia, M.; Jan, Y.N.; Jan, L.Y. Phosphatidylinositol-(4, 5)-bisphosphate regulates calcium gating of small-conductance cation channel TMEM16F. *Proc. Natl. Acad. Sci. USA* 2018, 115, E1667–E1674.

79. Musrap, N.; Tuccitto, A.; Karagiannis, G.S.; Saraon, P.; Batruch, I.; Diamandis, E.P. Comparative proteomics of ovarian cancer aggregate formation reveals an increased expression of calcium activated chloride channel regulator 1 (ClCa1). *J. Biol. Chem.* 2015, 290, 17218–17227.

80. Peretti, M.; Angelini, M.; Savalli, N.; Florio, T.; Yuspa, S.H.; Mazzanti, M. Chloride channels in cancer: Focus on chloride intracellular channel 1 and 4 (CLIC1 AND CLIC4) proteins in tumor development and as novel therapeutic targets. *Biochim. Biophys. Acta* 2014, 1848, 2523–2531.

81. Valenzuela, S.M.; Martin, D.K.; Por, S.B.; Robbins, J.M.; Warton, K.; Bootcov, M.R.; Schofield, P.R.; Campbell, T.J.; Breit, S.N. Molecular cloning and expression of a chloride ion channel of cell

nuclei. *J. Biol. Chem.* 1997, 272, 12575–12582.

82. Tang, H.Y.; Beer, L.A.; Chang-Wong, T.; Hammond, R.; Gimotty, P.; Coukos, G.; Speicher, D.W. A xenograft mouse model coupled with in-depth plasma proteome analysis facilitates identification of novel serum biomarkers for human ovarian cancer. *J. Proteome Res.* 2012, 11, 678–691.

83. Tang, H.Y.; Beer, L.A.; Tanyi, J.L.; Zhang, R.; Liu, Q.; Speicher, D.W. Protein isoform-specific validation defines multiple chloride intracellular channel and tropomyosin isoforms as serological biomarkers of ovarian cancer. *J. Proteom.* 2013, 89, 165–178.

84. Yu, W.; Cui, R.; Qu, H.; Liu, C.; Deng, H.; Zhang, Z. Expression and prognostic value of CLIC1 in epithelial ovarian cancer. *Exp. Ther. Med.* 2018, 15, 4943–4949.

85. Qu, H.; Chen, Y.; Cao, G.; Liu, C.; Xu, J.; Deng, H.; Zhang, Z. Identification and validation of differentially expressed proteins in epithelial ovarian cancers using quantitative proteomics. *Oncotarget* 2016, 7, 83187 83199.

86. Singha, B.; Harper, S.L.; Goldman, A.R.; Bitler, B.G.; Aird, K.M.; Borowsky, M.; Cadungog, M.G.; Liu, Q.; Zhang, R.; Jean, S.; et al. CLIC1 and CLIC4 complement CA125 as a diagnostic biomarker panel for all subtypes of epithelial ovarian cancer. *Sci. Rep.* 2018, 8, 14725.

87. Hernandez-Fernaud, J.R.; Ruengeler, E.; Casazza, A.; Neilson, L.J.; Pulleine, E.; Santi, A.; Ismail, S.; Lilla, S.; Dhayade, S.; MacPherson, I.R.; et al. Secreted CLIC3 drives cancer progression through its glutathione-dependent oxidoreductase activity. *Nat. Commun.* 2017, 8, 14206.

88. Argenzio, E.; Moolenaar, W.H. Emerging biological roles of Cl⁻intracellular channel proteins. *J. Cell Sci.* 2016, 129, 4165–4174.

89. Berridge, M.J.; Bootman, M.D.; Roderick, H.L. Calcium signalling: Dynamics, homeostasis and remodelling. *Nat. Rev. Mol. Cell Biol.* 2003, 4, 517–529.

90. Clapham, D.E. Calcium signaling. *Cell* 1995, 80, 259.

91. Li, W.; Zhang, S.L.; Wang, N.; Zhang, B.B.; Li, M. Blockade of T-type Ca(2+) channels inhibits human ovarian cancer cell proliferation. *Cancer Invest.* 2011, 29, 339–346.

92. Jang, S.J.; Choi, H.W.; Choi, D.L.; Cho, S.; Rim, H.K.; Choi, H.E.; Kim, K.S.; Huang, M.; Rhim, H.; Lee, K.T.; et al. In vitro cytotoxicity on human ovarian cancer cells by T-type calcium channel blockers. *Bioorg. Med. Chem. Lett.* 2013, 23, 6656–6662.

93. Dziegielewska, B.; Casarez, E.V.; Yang, W.Z.; Gray, L.S.; Dziegielewski, J.; Slack-Davis, J.K. T-type Ca²⁺ channel inhibition sensitizes ovarian cancer to carboplatin. *Mol. Cancer Ther.* 2016, 15, 460–470.

94. Mertens-Walker, I.; Bolitho, C.; Baxter, R.C.; Marsh, D.J. Gonadotropin-induced ovarian cancer cell migration and proliferation require extracellular signal-regulated kinase 1/2 activation regulated by calcium and protein kinase C δ . *Endocr. Relat. Cancer* 2010, 17, 335–349.

95. Kouba, S.; Ouldamer, L.; Garcia, C.; Fontaine, D.; Chantome, A.; Vandier, C.; Goupille, C.; Potier-Cartereau, M. Lipid metabolism and Calcium signaling in epithelial ovarian cancer. *Cell Calcium* 2019, 81, 38–50.

96. Xu, Y.; Gaudette, D.C.; Boynton, J.D.; Frankel, A.; Fang, X.J.; Sharma, A.; Hurteau, J.; Casey, G.; Goodbody, A.; Mellors, A. et al., Characterization of an ovarian cancer activating factor in ascites from ovarian cancer patients. *Clin. Cancer Res.* 1995, 1, 1223–1232.

97. Yu, X.; Zhang, Y.; Chen, H. LPA receptor 1 mediates LPA-induced ovarian cancer metastasis: An in vitro and in vivo study. *BMC Cancer* 2016, 16, 846.

98. Kim, E.K.; Ha, J.M.; Kim, Y.W.; Jin, S.Y.; Ha, H.K.; Bae, S.S. Inhibitory role of polyunsaturated fatty acids on lysophosphatidic acid-induced cancer cell migration and adhesion. *FEBS Lett.* 2014, 588, 2971–2977.

99. Zhang, H.; Zhou, L.; Shi, W.; Song, N.; Yu, K.; Gu, Y. A mechanism underlying the effects of polyunsaturated fatty acids on breast cancer. *Int. J. Mol. Med.* 2012, 30, 487–494.

100. Lencesova, L.; Krizanova, O. IP(3) receptors, stress and apoptosis. *Gen. Physiol. Biophys.* 2012, 31, 119–130.

101. Hudecova, S.; Markova, J.; Simko, V.; Csaderova, L.; Stracina, T.; Sirova, M.; Fojtu, M.; Svastova, E.; Gronesova, P.; Pastorek, M. Sulforaphane-induced apoptosis involves the type 1 IP3 receptor. *Oncotarget* 2016, 7, 61403–61418.

102. Chovancova, B.; Hudecova, S.; Lencesova, L.; Babula, P.; Rezuchova, I.; Penesova, A.; Grman, M.; Moravcik, R.; Zeman, M.; Krizanova, O. Melatonin-induced changes in cytosolic calcium might be responsible for apoptosis induction in tumour cells. *Cell Physiol. Biochem.* 2017, 44, 763–777.

103. Shibao, K.; Fiedler, M.J.; Nagata, J.; Minagawa, N.; Hirata, K.; Nakayama, Y.; Iwakiri, Y.; Nathanson, M.H.; Yamaguchi, K. The type III inositol 1, 4, 5-trisphosphate receptor is associated with aggressiveness of colorectal carcinoma. *Cell Calcium* 2010, 48, 315–323.

104. Chovancova, B.; Liskova, V.; Miklikova, S.; Hudecova, S.; Babula, P.; Penesova, A.; Sevcikova, A.; Durinikova, E.; Novakova, M.; Matuskova, M. et al. Calcium signaling affects migration and proliferation differently in individual cancer cells due to nifedipine treatment. *Biochem. Pharmacol.* 2020, 171, 113695.

105. Rezuchova, I.; Hudecova, S.; Soltysova, A.; Matuskova, M.; Durinikova, E.; Chovancova, B.; Zuzcak, M.; Cihova, M.; Burikova, M.; Penesova, A.; et al. Type 3 inositol 1,4,5-trisphosphate receptor has antiapoptotic and proliferative role in cancer cells. *Cell Death Dis.* 2019, 10, 186.

106. Mound, A.; Vautrin-Glabik, A.; Foulon, A.; Botia, B.; Hague, F.; Parys, J.B.; Ouadid-Ahidouch, H.; Rodat-Despoix, L. Downregulation of type 3 inositol (1, 4, 5)-trisphosphate receptor decreases breast cancer cell migration through an oscillatory Ca2+ signal. *Oncotarget* 2017, 8, 72324.

107. Kang, S.S.; Han, K.S.; Ku, B.M.; Lee, Y.K.; Hong, J.; Shin, H.Y.; Almonte, A.G.; Woo, D.H.; Brat, D.J.; Hwang, E.M. et al. Caffeine-mediated inhibition of calcium release channel inositol 1, 4, 5-trisphosphatereceptor subtype 3 blocks glioblastoma invasion and extends survival. *Cancer Res.* 2010, 70, 1173–1183.

108. Lewis, R.S. Store-operated calcium channels: New perspectives on mechanism and function. *Cold Spring Harb. Perspect. Biol.* 2011, 3, a003970.

109. Yang, S.; Zhang, J.J.; Huang, X.Y. Orai1 and STIM1 are critical for breast tumor cell migration and metastasis. *Cancer Cell* 2009, 15, 124–134.

110. Chen, Y.F.; Chiu, W.T.; Chen, Y.T.; Lin, P.Y.; Huang, H.J.; Chou, C.Y.; Chang, H.C.; Tang, M.J.; Shen, M.R. Calcium store sensor stromal-interaction molecule 1-dependent signaling plays an important role in cervicalcancer growth, migration, and angiogenesis. *Proc. Natl. Acad. Sci. USA* 2011, 108, 15225–15230.

111. Abdelazeem, K.N.M.; Droppova, B.; Sukkar, B.; Al-Maghout, T.; Pelzl, L.; Zacharopoulou, N.; Hassan Ali Hassan, N.; Abdel-Fattah, K.I.; Stournaras, C.; Lang, F. Upregulation of Orai1 and STIM1 expression as well as store-operated Ca²⁺ entry in ovary carcinoma cells by placental growth factor. *Biochem. Biophys. Res. Commun.* 2019, 512, 467–472.

112. Pedersen, S.F.; Owsianik, G.; Nilius, B. TRP channels: An overview. *Cell Calcium* 2005, 38, 233–252.

113. Zhuang, L.; Peng, J.B.; Tou, L.; Takanaga, H.; Adam, R.M.; Hediger, M.A.; Freeman, M.R. Calcium-selective ion channel, CaT1, is apically localized in gastrointestinal tract epithelia and is aberrantly expressed in human malignancies. *Lab. Invest.* 2002, 82, 1755–1764.

114. Han, H.; Chay, D.B.; Nam, S.; Cho, H.; Chung, J.Y.; Kim, J.H. Prognostic Significance of Transient Receptor Potential Vanilloid Type 1 (TRPV1) and Phosphatase and Tension Homolog (PTEN) in Epithelial Ovarian Cancer. *Cancer Genom. Proteom.* 2020, 17, 309–319.

115. Xue, H.; Wang, Y.; MacCormack, T.J.; Lutes, T.; Rice, C.; Davey, M.; Dugourd, D.; Ilenchuk, T.T.; Stewart, J.M. Inhibition of Transient Receptor Potential Vanilloid 6 channel, elevated in human ovarian cancers, reduces tumour growth in a xenograft model. *J. Cancer* 2018, 9, 3196–3207.

116. Lehen'kyi, V.; Flourakis, M.; Skryma, R.; Prevarskaia, N. TRPV6 channel controls prostate cancer cell proliferation via Ca(2+)/NFAT-dependent pathways. *Oncogene* 2007, 26, 7380–7385.

117. Quang, C.T.; Leboucher, S.; Passaro, D.; Fuhrmann, L.; Nourieh, M.; Vincent-Salomon, A.; Ghysdael, J. The calcineurin/NFAT pathway is activated in diagnostic breast cancer cases and is essential to survival and metastasis of mammary cancer cells. *Cell Death Dis.* 2015, 6, e1658.

118. Fu, S.; Hirte, H.; Welch, S.; Ilenchuk, T.T.; Lutes, T.; Rice, C.; Fields, N.; Nemet, A.; Dugourd, D.; Piha-Paul, S.; et al. First-in-human phase I study of SOR-C13, a TRPV6 calcium channel inhibitor, in patients with advanced solid tumors. *Invest. New Drugs* 2017, 35, 324–333.

119. Faouzi, M.; Hague, F.; Geerts, D.; Ay, A.S.; Potier-Cartereau, M.; Ahidouch, A.; Ouadid-Ahidouch, H. Functional cooperation between KCa3.1 and TRPC1 channels in human breast cancer: Role in cell proliferation and patient prognosis. *Oncotarget* 2016, 7, 36419–36435.

120. Chen, J.; Luan, Y.; Yu, R.; Zhang, Z.; Zhang, J.; Wang, W. Transient receptor potential (TRP) channels, promising potential diagnostic and therapeutic tools for cancer. *Biosci. Trends* 2014, 8, 1–10.

121. Villalobos, C.; Hernandez-Morales, M.; Gutierrez, L.G.; Nunez, L. TRPC1 and ORAI1 channels in colon cancer. *Cell Calcium* 2019, 81, 59–66.

122. Asghar, M.Y.; Magnusson, M.; Kemppainen, K.; Sukumaran, P.; Lof, C.; Pulli, I.; Kalhori, V.; Tornquist, K. Transient Receptor Potential Canonical 1 (TRPC1) Channels as Regulators of Sphingolipid and VEGF Receptor Expression: Implication for thyroid cancer cell migration and proliferation. *J. Biol. Chem.* 2015, 290, 16116–16131.

123. Dong, H.; Shim, K.N.; Li, J.M.; Estrema, C.; Ornelas, T.A.; Nguyen, F.; Liu, S.; Ramamoorthy, S.L.; Ho, S.; Carethers, J.M.; et al. Molecular mechanisms underlying Ca²⁺–mediated motility of human pancreatic duct cells. *Am. J. Physiol. Cell Physiol.* 2010, 299, C1493–C1503.

124. Zeng, B.; Yuan, C.; Yang, X.; Atkin, S.L.; Xu, S.Z. TRPC Channels and their splice variants are essential for promoting human ovarian cancer cell proliferation and tumorigenesis. *Curr. Cancer Drug Targets* 2013, 13, 103–116.

125. Yang, S.L.; Cao, Q.; Zhou, K.C.; Feng, Y.J.; Wang, Y.Z. Transient receptor potential channel C3 contributes to the progression of human ovarian cancer. *Oncogene* 2009, 28, 1320–1328.

126. Li, S.; Jiang, K.; Li, J.; Hao, X.; Chu, W.; Luo, C.; Zhu, Y.; Xie, R.; Chen, B. Estrogen enhances the proliferation and migration of ovarian cancer cells by activating transient receptor potential channel C3. *J. Ovarian Res.* 2020, 13, 20.

127. Wang, J.; Liao, Q.J.; Zhang, Y.; Zhou, H.; Luo, C.H.; Tang, J.; Wang, Y.; Tang, Y.; Zhao, M.; Zhao, X.H.; et al. TRPM7 is required for ovarian cancer cell growth, migration and invasion. *Biochem. Biophys. Res. Commun.* 2014, 454, 547–553.

128. Wei, C.; Wang, X.; Chen, M.; Ouyang, K.; Song, L.S.; Cheng, H. Calcium flickers steer cell migration. *Nature* 2009, 457, 901–905.

129. Yang, D.; Kim, J. Emerging role of transient receptor potential (TRP) channels in cancer progression. *BMB Rep.* 2020, 53, 125–132.

130. Ottevanger, P.B. Ovarian cancer stem cells more questions than answers. *Semin. Cancer Biol.* 2017, 44, 67–71.

131. French, B.; Gasch, C.; O’Leary, J.J.; Gallagher, M.F. Developing ovarian cancer stem cell models: Laying the pipeline from discovery to clinical intervention. *Mol. Cancer* 2014, 13, 262.

132. Yang, X.; Zheng, F.; Xing, H.; Gao, Q.; Wei, W.; Lu, Y.; Wang, S.; Zhou, J.; Hu, W.; Ma, D. Resistance to chemotherapy-induced apoptosis via decreased caspase-3 activity and overexpression of antiapoptotic proteins in ovarian cancer. *J. Cancer Res. Clin. Oncol.* 2004, 130, 423–428.

133. Schmidt, S.; Liu, G.; Liu, G.; Yang, W.; Honisch, S.; Pantelakos, S.; Stournaras, C.; Höning, A.; Lang, F. Enhanced Orai1 and STIM1 expression as well as store operated Ca²⁺ entry in therapy resistant ovary carcinoma cells. *Oncotarget* 2014, 5, 4799–4810.

134. Mir, R.; Stanzani, E.; Martinez-Soler, F.; Villanueva, A.; Vidal, A.; Condom, E.; Ponce, J.; Gil, J.; Tortosa, A.; Giménez-Bonafé, P. YM155 sensitizes ovarian cancer cells to cisplatin inducing apoptosis and tumor regression. *Gynecol. Oncol.* 2014, 132, 211–220.

135. Liu, X.; Zou, J.; Su, J.; Lu, Y.; Zhang, J.; Li, L.; Yin, F. Downregulation of transient receptor potential cation channel, subfamily C, member 1 contributes to drug resistance and high histological grade in ovarian cancer. *Int. J. Oncol.* 2016, 48, 243–252.

136. Jaber, N.; Dou, Z.; Lin, R.Z.; Zhang, J.; Zong, W.X. Mammalian PIK3C3/VPS34: The key to autophagic processing in liver and heart. *Autophagy* 2012, 8, 707–708.

137. Yin, F.; Liu, X.; Li, D.; Wang, Q.; Zhang, W.; Li, L. Bioinformatic analysis of chemokine (C-C motif) ligand 21 and SPARC-like protein 1 revealing their associations with drug resistance in ovarian cancer. *Int. J. Oncol.* 2013, 42, 1305–1316.

138. Dean, M.; Fojo, T.; Bates, S. Tumour stem cells and drug resistance. *Nat. Rev. Cancer* 2005, 5, 275–284.

139. Lee, H.; Kim, J.W.; Kim, D.K.; Choi, D.K.; Lee, S.; Yu, J.H.; Kwon, O.B.; Lee, J.; Lee, D.S.; Kim, J.H.; et al. Calcium Channels as Novel Therapeutic Targets for Ovarian Cancer Stem Cells. *Int. J. Mol. Sci.* 2020, 21, 2327.

140. Sharp, S.Y.; Mistry, P.; Valenti, M.R.; Bryant, A.P.; Kelland, L.R. Selective potentiation of platinum drug cytotoxicity in cisplatin-sensitive and -resistant human ovarian carcinoma cell lines by amphotericin B. *Cancer Chemother. Pharmacol.* 1994, 35, 137–143.

141. Marklund, L.; Henriksson, R.; Grankvist, K. Cisplatin-induced apoptosis of mesothelioma cells is affected by potassium ion flux modulator amphotericin B and bumetanide. *Int. J. Cancer* 2001, 93, 577–583.

142. Marklund, L.; Andersson, B.; Behnam-Motlagh, P.; Sandstrom, P.E.; Henriksson, R.; Grankvist, K. Cellular potassium ion deprivation enhances apoptosis induced by cisplatin. *Basic Clin. Pharmacol. Toxicol.* 2004, 94, 245–251.

143. Hui, C.; Lan, Z.; Yue-li, L.; Li-lin, H. Knockdown of Eag1 Expression by RNA Interference Increases Chemosensitivity to Cisplatin in Ovarian Cancer Cells. *Reprod. Sci.* 2015, 22, 1618–1626.

144. Lloyd, K.L.; Cree, I.A.; Savage, R.S. Prediction of resistance to chemotherapy in ovarian cancer: A systematic review. *BMC Cancer* 2015, 15, 117.

145. Gueguinou, M.; Harnois, T.; Crottes, D.; Uguen, A.; Deliot, N.; Gambade, A.; Chantôme, A.; Haelters, J.P.; Jaffrè, P.A.; Jourdan, M.L.; et al. SK3/TRPC1/Orai1 complex regulates SOCE dependent colon cancer cell migration: A novel opportunity to modulate anti EGFR mAb action by the alkyl lipid Ohmeline. *Oncotarget* 2016, 7, 36168–36184.

146. Samuel, P.; Pink, R.C.; Caley, D.P.; Currie, J.M.; Brooks, S.A.; Carter, D.R. Over-expression of miR-31 or loss of KCNMA1 leads to increased cisplatin resistance in ovarian cancer cells. *Tumour Biol.* 2016, 37, 2565–2573.

Retrieved from <https://encyclopedia.pub/entry/history/show/17795>