

GPCRs in Insects

Subjects: [Agriculture, Dairy & Animal Science](#)

Contributor: Nannan Liu , Ting Li , Yifan Wang , Shikai Liu

G-protein coupled receptors (GPCRs) play important roles in cell biology and insects' physiological processes, toxicological response and the development of insecticide resistance. New information on genome sequences, proteomic and transcriptome analysis and expression patterns of GPCRs in organs such as the central nervous system in different organisms has shown the importance of these signaling regulatory GPCRs and their impact on vital cell functions. Our growing understanding of the role played by GPCRs at the cellular, genome, transcriptome and tissue levels is now being utilized to develop new targets that will sidestep many of the problems currently hindering human disease control and insect pest management.

G-protein coupled receptor regulation pathway

GPCR physiological functions

tissue specific expression

genome sequences analysis

phylogenetic tree

insect physiology

insecticide resistance

1. Introduction

G-protein-coupled receptors (GPCRs), which are proteins that share a seven α -helical transmembrane structure, govern a number of biological and physiological processes in both the vertebrate and invertebrate kingdoms. The main function of GPCRs is to transduce extracellular and environmental signals and regulate intracellular second messengers through coupling to heterotrimeric G-proteins and their downstream effectors ^[1]. GPCRs are known to be involved in recognizing extracellular messengers, transducing signals to the cytosol, and mediating the cellular responses necessary for the normal physiological functions of organisms ^{[2][3][4][5][6][7]}. GPCR binding to a wide variety of molecules (ligands), including hormones, neurotransmitters, ions, photons, odorants, neuropeptides and other stimuli through coupling with G proteins and arrestins ^[1], results in interactions with G proteins that, in turn, activate the downstream effectors of GPCR pathways, including the adenylate cyclase/cyclic AMP (AC/cAMP), phosphatidylinositol/diacylglycerol/protein kinase C (PI/DAG/PKC) and voltage gated calcium channel (Ca²⁺ channels) pathways, ultimately playing an indispensable role in the organism's growth, development, reproduction and other physiological activities ^{[6][7][8][9][10]}. These critical functions mean that a better understanding of the role played by GPCRs in cell physiology and biochemistry is crucial for efforts to develop new molecular-level human disease therapies ^{[10][11][12][13][14][15][16]}.

In just a few short years, GPCR research in insects has progressed from the initial GPCR gene identification to comprehensive bioinformatics analyses, from single GPCR gene analysis to whole genome sequences of GPCRs and explorations of their pathways, and from traditional transcriptional analysis of the gene expression to gene

functional characterization of the GPCR genes in insect physiology and cellular biology. The incredible progress being made in related fields includes a wide range of complementary technologies, including bioinformatics and quantitative expression analyses, with functional studies using RNA interferon revealing potential biological functions that significantly impact insect physiology [17][18], including reproduction [19][20][21][22][23][24], regulating growth and development [21][25][26][27][28][29][30][31][32][33][34][35], the stress response [36][37][38][39][40][41][42][43][44][45][46][47], feeding [21][35][48][49][50][51][52][53][54][55][56], general behaviors [20][57][58][59][60][61][62] and other physiological processes [39][63][64][65][66][67][68][69][70]. In addition, the overexpression of GPCRs in insecticide resistance has been reported in both mosquitoes [42][43][44], and house flies [71][72]. The functions of GPCR regulatory pathways and GPCR downstream intracellular cascades have been explored in the development of insecticide resistance in *Culex quinquefasciatus* [42][43][44], providing a good understanding of the complex molecular processes that govern the development of insecticide resistance through the GPCR signally transduction pathways. The outcomes of these physiological and toxicological studies not only provide us with a clear global picture that is enabling us to develop a clearer understanding of the highly complex mechanisms, genes and pathways involved in these insect physiological and toxicological processes, but are also providing a strong foundation that will allow us to develop tightly GPCR targeted new insecticides and/or environmentally sound insecticides for better vector control that circumvent the problems associated with resistance, representing a highly practical application of scientific research in this area. The findings reported in the most recent studies of GPCR functions in insect are opening up promising new avenues that will undoubtedly revolutionize future research on insect pest management.

2. Whole Genome Sequencing and Transcriptome Analysis—Sequence Comparison and GPCR Characterization in Insects

Recent developments in high-throughput sequencing (HTS) technologies have created powerful analytic tools that enable researchers to study the complex gene interactions in individual organisms and the diverse relationships between organisms. Whole-genome analysis and transcriptome gene expression have revolutionized efforts to identify and annotate insect GPCRs in many different insect species, with comparative genomics of insect populations being used to identify novel targets for insect pest control. Given the growing problem of the development of resistance to the insecticides that are conventionally used to control insects, especially those that are medically, agriculturally and economically important [73], there is an urgent need to shift from traditional chemical pesticides towards more targeted gene-centric approaches. Indeed, genome resequencing analysis followed by functional characterization has opened up many opportunities for discovering new targets, such as GPCRs, for new insecticides that not only control insect pests effectively but also overcome the problem of insecticide resistance. This section reviews GPCRs and their possible functions for several insect species at the whole genome level.

2.1. Classification Systems Used in Characterizing GPCRs

In general, two classification systems are used to characterize GPCRs in organisms, namely the A-F system [74][75] and GRAFS [76]. The A-F system classifies GPCRs primarily in terms of their sequences and functional similarities using six classes, labelled A, B, C, D, E and F. Class A, known as the “rhodopsin-like family”, is the largest group of

GPCRs; Class B is the “secretin receptor family”; Class C contains the metabotropic glutamate family; Class D refers to the fungal mating pheromone receptors; Class E includes all the cAMP receptors; Class F are the frizzled/smoothened receptors [77]. The GRAFS system is based on the phylogenetic tree of the human GPCR sequences, with GPCRs being classified into five families: glutamate (G), rhodopsin (R), adhesion (A), frizzled/taste 2 (F) and secretin (S) [76]. In insects, the classification of GPCRs is based on the A-F system. Work on insect GPCRs was revolutionized by the publication of first insect genome sequence of *Drosophila melanogaster*, an important model organism [78]. To date, more than 100 insect genomes have been sequenced and published, including *Anopheles gambiae* [79], *Aedes aegypti* [80], *Cx. quinquefasciatus* [81], *Musca domestica* [82] and some other species [83]. Sequencing and annotating these insect genomes provide a strong foundation for future research and new insights that enrich our understanding of the potential functions of GPCRs in insect physiology and molecular biology.

As an important model species in the insect kingdom, *D. melanogaster* has 200 GPCR genes, over 70 of which belong to Class A (the rhodopsin-like family), 20 to Class B (the secretin-like family), 5 to Class C (the metabotropic glutamate-like family) and 5 to Class F (the frizzled/smoothened family); the remainder have not yet been clearly classified [84][85]. A homology-based bioinformatics analysis conducted on the genome of the mosquito *An. gambiae* revealed 276 GPCR genes, of which over 80 belong to Class A (the rhodopsin-like family), 20 to Class B (the secretin-like family), 8 to Class C (the metabotropic glutamate-like family), 8 belong to Class F (frizzled/smoothened) and the remainder were other GPCRs [86]. Several GPCRs were explored via reannotation in the mosquito *Ae. aegypti*, coding over 135 GPCR genes, with 89 belonging to Class A (the rhodopsin-like family), 24 to Class B (the secretin-like family), 8 to Class C (the metabotropic glutamate-like family), 11 belonging to Class F (frizzled/smoothened family) and the rest not been clearly classified [80]. Around 90 GPCRs from the silkworm *Bombyx mori* have also been annotated by bioinformatics analysis, with 69 being classified as Class A (the rhodopsin-like family), 8 as Class B (the secretin-like family), 9 as Class C (the metabotropic glutamate-like family), 3 as Class F (the frizzled/smoothened family) and others [87]. For *Apis mellifera*, around 50 genes have been coded for GPCR, of which 31 belong to Class A (the rhodopsin-like family) and 4 to Class B (the secretin-like family) [17]. For *Cx. quinquefasciatus*, around 115 GPCR genes have been identified, with 52 belonging to Class A (the rhodopsin-like family), 4 to Class B (the secretin-like family) and the rest not been classified [42]. For *Musca domestica*, around 94 genes have been identified as GPCR genes, 55 of which are classified as Class A (the rhodopsin-like family), 27 are classified as Class B (the secretin-like family), 4 are classified as Class C (the metabotropic glutamate-like family) and the rest are still unclear [72] (Table 1, Figure 1).




Figure 1. Sequence comparison of GPCRs in insects. The phylogenetic tree includes 64 *Ae. aegypti* GPCRs, 98 *D. melanogaster* GPCRs, 14 *Aedes albopictus* GPCRs, 26 *An. Gambiae* GPCRs, 9 *Cx. Quinquefasciatus* GPCRs and 7 *M. domestica* GPCRs. Different colored circles represent different species. Except for the genes listed for *D. melanogaster*, which are taken from Flybase (<https://flybase.org>, accessed on 7 May 2021), all the genes are from Vectorbase (<https://vectorbase.org>, accessed on 7 May 2021). Sequence alignment was conducted in MEGAX and the phylogenetic tree was developed using a neighbor-joining method by MEGAX with 2000 bootstrap replicates. The evolutionary distances were computed using the p-distance method. Different classes of GPCRs are represented by colored branches—green for the Class A (the rhodopsin-like family), yellow for the Class B (the secretin receptor family), purple for the Class C (the metabotropic glutamate family) and blue for the Class F (the frizzled and smoothened receptor family). The major types of GPCRs in each of classes are labeled.

Table 1. The information of GPCR genes in insect species.

Insect Species	Total Number of Genes	Class A Gene Number	Class B Gene Number	Class C Gene Number	Class F Gene Number	Source of Genome Information	Reference
<i>D. melanogaster</i>	~200	>70	~20	~5	~5	https://flybase.org/ (accessed on 7 May 2021)	[84][85]
<i>An. gambiae</i>	~276	81	21	8	~8	https://www.ncbi.nlm.nih.gov/genome/46?genome_assembly_id=22679 (accessed on 7 May 2021)	[86]
<i>Ae. aegypti</i>	135	89	24	8	11	https://www.ncbi.nlm.nih.gov/genome/44?genome_assembly_id=322291 (accessed on 7 May 2021)	[80]
<i>B. mori</i>	~90	~70	~7	~8	~4	https://www.ncbi.nlm.nih.gov/genome/76?genome_assembly_id=1491718 (accessed on 7 May 2021)	[87]
<i>A. mellifera</i>	~50	~31	~4	Not clear	Not clear	https://www.ncbi.nlm.nih.gov/genome/48?genome_assembly_id=403979 (accessed on 7 May 2021)	[17]
<i>L. cuprina</i>	197	73	18	9	Not clear	https://www.ncbi.nlm.nih.gov/genome/12732?genome_assembly_id=358015 (accessed on 7 May 2021)	[88]
<i>Cx. quinquefasciatus</i>	115	52	4	Not clear	Not clear	https://www.ncbi.nlm.nih.gov/genome/393?genome_assembly_id=1502880 (accessed on 7 May 2021)	[42]
<i>M. domestica</i>	94	55	27	4	Not clear	https://www.ncbi.nlm.nih.gov/genome/14461?genome_assembly_id=44793 (accessed on 7 May 2021)	[72]

ies have
rized by
typical genomic sequencing, RNA-seq and the assembly method. The subsequent genome sequence and gene/protein identification and annotation revealed 197 GPCRs, including 73 to be in Class A (the rhodopsin-like family), 18 in Class B (the secretin-like family) and 9 in Class C (the metabotropic glutamate receptor family), along with some unclassified proteins [88] (Figure 1, Table 1). In recent work, a transcriptome study on GPCRs in the stick insect, *Carausius morosus*, conducted via RNA-seq and de novo RNA assembly revealed 430 putative GPCR genes [89]; a transcriptome analysis predicted over 300 transcripts for coding GPCRs in fire ants [90] and 65 genes for putative neuropeptide GPCRs were identified in the spider mite [91].

Generally, Class A GPCRs are the predominant class of GPCRs in insects, comprising their neuropeptide and protein hormone receptors, muscarinic acetylcholine receptors, dopamine receptors, 5-hydroxytryptamine receptors, tyramine receptors, opsin receptors, vasopressin receptors and orphan receptors, among others [92][93]. These perform a variety of have different functions [94][95][96], including metamorphosis (e.g., dopamine receptors) [29][34][35][97][98], feeding modulation (e.g., allatostatin receptors) [99][100], behavioral changes (e.g., sex peptide receptors and SIF amide neuropeptide receptors) [101][102][103][104], metabolism regulation (e.g., CCHamide-2 receptors and adipokinetic hormone receptor) [51][53][62][105][106] and visual photoreceptors (e.g., rhodopsin proteins) [107], among many others [92][95][108][109][110]. The Class B GPCRs, which include diuretic hormone receptors, methuselah-type receptors and others [111][112], mainly focus on the insect's water balance (e.g., corticotropin-releasing factor receptors) [113], body temperature modulation (e.g., diuretic hormone receptors) [114], stress

response (e.g., methuselah GPCRs) [115][116] and other functions [117]. Their Class C receptors are mainly composed of metabotropic GABA receptors, which serve as neurotransmitters receptors and are involved in signal transportation [118]. The frizzled receptors and smoothed-type receptors, which are mainly Class F GPCRs, focus on Wnt signaling [119][120][121] and the hedgehog signaling pathway, respectively [122][123] (Figure 1). Both the Wnt signaling pathway and the hedgehog pathway are important for insect development [124][125].

2.2. Receptors of GPCR Involved in Insect Physiology or Insecticide Resistance That Are Potential Targets for Insecticide Development

Several GPCRs have been reported to be involved in different functions in insect physiology as described below (Table 2) and could thus be potential targets for new insecticides. The majority are Class A GPCRs, including the dopamine receptor, which mainly regulates sexual activity [60][126], morphogenesis [29][34][35][97], mushroom and locomotor activity [127] and ethanol-induced sedation [68]. Neuropeptide and hormone receptors such as the adipokinetic hormone receptor (AKHR) can receive signals from the adipokinetic hormone and regulate lipid mobilization [51][53][62][106], while the allatostatin receptor (AstR) regulates the juvenile hormone synthesis [26][128], the diapause hormone receptor (DHR) is involved in insect development [39][41][45] and the neuropeptide receptors focus on the regulation of insect feeding behavior [50][129] and ecdysone synthesis [33]. Orphan receptor, like BNGR in *B. mori* regulates the insect's food intake and growth [130], and DLGR2 in *D. melanogaster* regulates the insect's bursicon bioactivity [25]. Tyramine receptors are responsible for the development of antiparasitic compounds [131], while calcitonin receptors regulate hindgut contraction and primary urine secretion [66] and the 5-HT (5-hydroxytryptamine) receptors and muscarinic acetylcholine receptors are important for the insect's neural processes [132][133][134]. Rhodopsin receptors perform vital functions in both the insect's reproduction system [22] and its vision [107]. Class B GPCRs such as methuselah receptors are also involved in insect longevity [135], and oxidative stress resistance [136] and diuretic hormone receptors regulate the body temperature and homeostasis [114]; GPCRs in the Class C family, such as metabotropic GABA receptors, are important for the central nervous system (CNS) [118].

Table 2. The functions of GPCR receptors in insect physiology.

Receptor Group	Receptor Name	Classes	Species	Function	Reference
5-HT receptors	Trica5-HT7 R	Class A	<i>Tribolium castaneum</i>	Insect's neural processes	[134]
Adipokinetic hormone receptor	Akh receptor	Class A	<i>Bactrocera dorsalis</i>	Lipid mobilization	[62]
Adipokinetic hormone receptor	Akh receptor	Class A	<i>D. melanogaster</i>	Lipid mobilization	[51]
Adipokinetic hormone receptor	Akh receptor	Class A	<i>Nilaparvata lugens</i>	Lipid mobilization	[53]
Allatostatin	AstAR1	Class	<i>D. melanogaster</i>	Metamorphosis	[98]

Receptor Group	Receptor Name	Classes	Species	Function	Reference
receptor		A			
Allatostatin receptor	DAR-1/DAR-2	Class A	<i>D. melanogaster</i>	Feeding modulation	[99]
Allatostatin receptor	Dippu-AstR	Class A	<i>Diptera punctata</i>	Juvenile hormone synthesis	[128]
Arginine vasopressin-like receptor	AVPL receptor	Class A	<i>T. castaneum</i>	Diuretic signaling pathway	[92]
Calcitonin receptors	GPCRCAL1	Class A	<i>Ae. aegypti</i>	primary urine secretion	[66]
CCHa2 receptor	CCHa2-R	Class A	<i>D. melanogaster</i>	Insulin production	[105]
Diapause hormone receptor	DH-R	Class A	<i>Ae. aegypti</i>	Development	[39]
Diapause hormone receptor	Bommo-DHR	Class A	<i>B. mori</i>	Development	[45]
Diapause hormone receptor	HzDHR	Class A	<i>Helicoverpa zea</i>	Development	[41]
Dopamine receptor	Dop1R2, DmDopEcR	Class A	<i>D. melanogaster</i>	Morphogenesis	[34][97]
Dopamine receptor	DopEcR	Class A	<i>D. melanogaster</i>	Mushroom and locomotor activity	[127]
Dopamine receptor	DopEcR	Class A	<i>D. melanogaster</i>	Ethanol-induced sedation	[68]
Dopamine receptor	AipsDopEcR	Class A	<i>Agrotis ipsilon</i>	Sexual activity regulation	[60][126]
Dopamine receptor	DopEcR	Class A	<i>Helicoverpa armigera</i>	Morphogenesis	[35]
Dopamine receptor	D2R	Class A	<i>T. castaneum</i>	Morphogenesis	[29]
Leucokinin receptor	LKr	Class A	<i>D. melanogaster</i>	Feeding modulation	[100][103]
Myosuppressin receptors	CG8985/CG13803	Class A	<i>D. melanogaster</i>	visceral muscles inhibition	[108]

Receptor Group	Receptor Name	Classes	Species	Function	Reference
Neuropeptide receptors	GPCR-B2	Class A	<i>B. mori</i>	Ecdysone synthesis	[33]
Neuropeptide receptors	Schgr-sNPFR	Class A	<i>Schistocerca gregaria</i>	Feeding behavior	[50]
Neuropeptide Drosulfakinin receptor	CCKLR-17D1	Class A	<i>D. melanogaster</i>	Fighting behavior	[104]
Orphan receptor	DLGR2	Class A	<i>D. melanogaster</i>	Bursicon bioactivity	[25]
Orphan receptor	BNGR-A4 receptor	Class A	<i>B. mori</i>	Food intake and growth	[130]
Rhodopsin receptors	Rh2	Class A	<i>T. castaneum</i>	Reproduction	[22]
Sex peptide receptor	SPR	Class A	<i>D. melanogaster</i>	Reproductive behavior	[101]
SIFamide receptor	SIFaR	Class A	<i>D. melanogaster</i>	Reproductive behavior	[102]
Tyramine receptor	TAR1	Class A	<i>Rhipicephalus (Boophilus) microplus</i>	Development of antiparasitic	[131]
Corticotropin releasing factor receptor	CG12370	Class B	<i>D. melanogaster</i>	Water balance	[113]
Diuretic hormone receptors	DH31R	Class B	<i>D. melanogaster</i>	temperature regulation and homeostasis	[114]
Methuselah receptor	mth	Class B	<i>D. melanogaster</i>	Oxidative stress resistance	[136]
Methuselah receptor	Ldmth1	Class B	<i>Lymantria dispar</i>	Insect longevity	[135]
Metabotropic GABA receptors	D-GABABR1, R2 and R3	Class C	<i>D. melanogaster</i>	Central nervous system	[118]

development. Discovering and identifying the GPCR gene profiles in different tissues will help broaden our understanding of the functions of various types of tissue and gene functions, and the biological mechanisms involved [137]. The aberrant expression of tissue specific expressed GPCRs may also be implicated in various abnormal functions of the insects, and hence important for the control of insect pests. This section reviews recent discoveries related to the GPCR genes present in insect nervous, digestive and reproductive systems, and in others, such as fat bodies, muscles and salivary glands (Figure 2).

3.1. Brain Tissue and Central Nervous System

Several GPCRs have been identified as being predominately expressed in the brain and CNS, corresponding to neuropeptide expression in diverse insect species. A neuropeptide bona fide natalisin receptor found to be highly expressed in the CNS of *D. melanogaster*, *T. castaneum* and *B. mori* regulates sexual activity and fecundity in insects [20]. A mutant *methuselah* gene expressed in the brain of *Drosophila*, a model insect often used to investigate gene functions, plays a critical function in oxidative stress resistance [136]; a dopamine/ecdyseroid receptor that is a head-specific expression gene is also overexpressed in *Drosophila* adults and embryos [97]. Another study also found the dopamine/ecdyseroid receptor (*DopEcR*) to be highly expressed in *Drosophila* nervous system and brain tissues, and its mutant has been implicated in the insect's resistance to alcohol [68]. A *Drosophila* *sex/myo-inhibiting* peptide receptor expressed in the CNS is highly conserved in insects [19], while an octopamine receptor *DmOcta2R* transcript present at much higher levels in *Drosophila* males than in females is mainly expressed in the head of adults [138] and a *Drosophila* neuropeptide *capa* GPCR expressed in larvae central nerve system is responsible for sensitivity to desiccation stress [37]. Of two IGRs (*LGR3/LGR4*) characterized in diverse *Drosophila* tissues, *Lgr3* transcripts were predominately expressed in the CNS, while *Lgr4* transcripts were highly expressed in the CNS of the wandering larval stage. In adults, the *Lgr4* transcripts showed significantly high expression in the male thoracic-abdominal ganglion and brain tissues [139].

The silkworm, *B. mori*, which is classed as a beneficial insect, is another model species frequently used in gene functional studies. A neuropeptide GPCR A4 receptor gene (*BomNPFR*) amplified from its brain tissue was predicted to be involved in food intake and growth processes [130]. Three GPCR receptors responding to neuron ion transport peptides (ITPs) were identified in silkworm larvae using tissue specific expression, revealing that *bngr-A2* was mainly expressed in the brain, with no expression in the CNS, and *bngr-A34* was not present in nervous tissues [21]. A neuropeptide allatotropin receptor was significantly overexpressed in the corpora cardiaca and brain of *B. mori*, indicating the potential involvement of juvenile hormone (JH) biosynthesis processes [140]. In honeybees, an important beneficial insect that plays a critical role in pollination, an octopamine receptor was identified in the honeybee brain [63]. A tyramine receptor was later identified that was highly expressed in nurse bees and forager brain neuropils [141].

The red import fire ant, *Solenopsis invicta*, is an invasive urban insect species whose gene expression and functions have been widely studied over the last couple of decades. Ten GPCR gene expressions have exhibited significant differences in the brain tissue of workers, alate virgin queens, and mated queens [90]. One neuropeptide F-Like receptor was isolated from *S. invicta* and its differential expression levels in the brain suggest its potential function to be feeding regulation for mated queens [48]. In the medically important kissing bug, *Rhodnius prolixus*, an important Chagas disease vector, GPCRs have been found to be involved in multiple physiological pathways. Three variants of a pyrokinin-1 receptor gene characterized in different tissues of fifth instar larvae of *R. prolixus* revealed two variants that were mainly expressed in the CNS [142]. A corazonin receptor was found to be overexpressed in the brain and involved in *R. prolixus* heartbeat control [143] and a serotonin type 2b receptor transcript has also been shown to be upregulated in the CNS of *R. prolixus* [144]. To predict the potential function of

a kinin receptor in *R. prolixus* blood post-feeding, a kinin receptor was characterized in various tissues and found to be overexpressed in the CNS of late instar larvae [56]. In mosquitoes, which are responsible for transmitting a number of human and animal diseases, GPCRs have been identified as being involved in mosquito blood feeding and insecticide resistance. The overexpression of an allatotropin GPCR receptor (*AeATr*) gene was characterized in the nervous system and corpora alata-corpora cardiac complex of *Ae. aegypti*. Blood feeding depressed the transcript level of *AeATr*, and was associated with JH biosynthesis in mosquitoes [32]. A rhodopsin-like GPCR overexpressed in the head of adult mosquitoes was found to play a critical role in the development of permethrin insecticide resistance in the mosquito, *Cx. Quinquefasciatus* [43]. Tissue specific studies conducted on this GPCR gene revealed its significantly high expression in the insect's brain tissue along with a G-protein alpha subunit, two adenylyl cyclase and one protein kinase A gene, playing important roles in the GPCR-leading intracellular pathway in the insect's neuron system and regulating its insecticide resistance [44]. A corazonin neuropeptide receptor (*MdCrz*) has been found to be overexpressed in the larval CNS of the house fly, another medically important pest, including in its ventral nerve cord, the protocerebral DL neurons of the brain lobes and its *vCrz* neurons. Although it was overexpressed in the heads of both male and female houseflies, there was no expression in other body tissues, indicating that *MdCrz* is a brain-specific expression gene [145].

The red flour beetle, *T. castaneum*, is a stored product pest that causes problem for communities around the world. The spatial expression of a 5-HT₇-type serotonin receptor identified in *T. castaneum* showed the highest expression levels to be in the head of both male and female adults, predominantly expressed in the brain but accompanied by high levels of expression in the optic lobes, predicting the functional importance of this receptor in neural processes [134]. A D₂-like dopamine receptor gene predominantly expressed in the head and CNS of *T. castaneum* adults [146] and an inotocin receptor was found to be overexpressed in the head of the insect's early larval stage [147]. In the desert locust, *S. gregaria*, an agricultural pest species, two novel octopamine receptors were characterized in adults, with *SgOctaR* being overexpressed in the CNS, including the brain, optic lobes, subesophageal ganglion and thoracic ganglions, while *SgOctβR* was overexpressed in the CNS. Interestingly, these receptors were highly expressed in the long-term gregarious locusts but not in the solitary locusts, suggesting the receptors' function in locust behavior [148]. GPCRs have also been widely studied in other agricultural pest species. The GPCR *NIA42* was found to be highly expressed in the brain and abdominal integuments of the adult brown planthopper, *N. lugens*, where it was linked to the neuropeptide NI-elevenin that is involved in the regulation of planthopper melanization [70][149]. A novel octopamine receptor gene expression has been characterized in diverse tissues of the fifth-instar larva rice stem borer, *Chilo suppressalis*, showing a high level of expression in the nerve cord [150]. A large screen of GPCRs that was characterized in *C. suppressalis* identified 51 putative GPCR genes. The expression of these genes was examined in tissues of the insect's CNS, fat body, gut and hemocytes, with most of the receptor genes being highly expressed in the CNS [151]. A serotonin receptor (*Pr5-HT₈*) was identified as being highly expressed in the nerve cord of the larva small white butterfly, *Pieris rapae*. 5-hydroxytryptamine (5-HT) has been identified as a neurotransmitter that plays critical roles in the regulation of a number of physiological processes, which is consistent with the expression of *Pr5-HT₈* in the butterfly's CNS [133]. An allatotropin receptor has also been found to be overexpressed in the brain, thoracic ganglion and abdomen ganglion of the bollworm, *H. armigera* [152].

3.2. Digestion and Reproduction Systems

In mosquito species, GPCRs are known to be expressed in the digestion system and involved in blood-feeding behavior. A leucokinin receptor has diverse functions, responding to multiple kinins in the mosquito, *Ae. aegypti*, and is expressed in the mosquito's hindgut and Malpighian tubules [153]. Piermarini's group also identified several GPCRs and metabolic genes upregulated or downregulated in the Malpighian tubules of blood-taken *Aedes albopictus* [154]. An allatotropin GPCR receptor (*AeATr*) gene has also been found to be overexpressed in the ovary of adult mosquitoes [32]. In another blood-feeding insect, *R. prolixus*, a serotonin type 2b receptor transcript was overexpressed in the Malpighian tubules, salivary glands and guts [144]. A kinin receptor has also been found to be highly overexpressed in the gut system of late instar larvae of *R. prolixus* [56]. Two variants of a pyrokinin-1 receptor gene were mainly expressed in the male testes and prothoracic glands of fifth instar larvae of *R. prolixus* [142]. In *Drosophila*, several GPCRs were identified as overexpressed in the digestion and reproduction systems. These *Drosophila Lgr4* transcripts were highly expressed in the gut system of the wandering larval stage, and significantly highly expressed in the male midgut and crop [139]. Additionally, a sex/myo-inhibiting peptide receptor was highly expressed in the male *Drosophila* reproduction organs [19] and an octopamine receptor gene was highly expressed in the Malpighian tubules, with lesser amounts found in the midgut and hemocytes of the fifth-instar larva rice stem borer, *C. suppressalis* [150]. Several GPCRs in *C. suppressalis* were overexpressed in the gut system [151]. A neuropeptide F-Like receptor detected in the gut and reproduction systems of *S. invicta* suggests its potential function was involved in the feeding regulation of mated queens [48]. One ITP gene identified in *B. mori*, *bngR-A2*, was found to be mainly expressed in the reproduction system [21]. A serotonin receptor (*Pr5-HT₈*) is known to be highly expressed in the Malpighian tubules, fat body and midgut of larvae of the small white butterfly, *P. rapae* [133].

3.3. Other Insect Organs

In addition to the GPCRs identified in the CNS, digestive and reproductive systems in insect species, many GPCRs have also been characterized in other organs. In *Drosophila* adults, *Lgr3* transcripts were found to be highly expressed in the female fat body and uterus, and the male salivary glands [139]. An adipokinetic hormone receptor was predominately expressed in the fat body of the oriental fruit fly, *B. dorsalis*, and was also involved in its triacylglycerol mobilization and sexual behavior [62]. Forty six putative GPCR transcripts have been isolated from the foreleg tissues of the cattle tick, *Phipicephalus australis*, providing valuable information for GPCR studies of signal transduction, host preference and mating behavior in insects [155]. An allatotropin receptor has been identified that is overexpressed in the male bumblebee accessory glands, predicting its potential involvement in JH biosynthesis [156] and in *B. mori* a sex peptide receptor has been found to be highly expressed in the prothoracic gland, predicting the critical roles of receptor in regulating ecdysteroidogenesis [28]. A novel octopamine receptor, *SgOct β R*, was also overexpressed in the flight muscles of *S. gregaria*, suggesting its involvement in locust gregarious behavior [148], while a GPCR receptor, *NIA42*, was significantly highly expressed in the integument and salivary gland of the brown planthopper indicating its function in the insect's melanization [149].




Figure 2. Tissue specific expression of GPCR genes in insect species. Insect GPCRs are mainly expressed in the brain and central nervous system, highlighted in light blue; those in the digestive system are highlighted in gray; those in the reproductive system are highlighted in light purple; those in other organs are highlighted in orange. Twenty-six GPCRs that are highly expressed in the brain and central nervous system in 13 insect species are involved in sexual activity and fecundity, oxidative stress resistance, desiccation stress, food intake and growth, juvenile hormone biosynthesis, feeding regulation, heartbeat control, permethrin insecticide resistance, behavior and cuticle melanization.

However, while the precise functions of the GPCRs in specific tissues are still in need of further characterization, the impact of the GPCRs on insect behaviors can now be predicted to some extent based on the functions of those specific tissues in insects. The importance of GPCRs in insects can be partially appreciated by considering their distribution, location and function within the cells of specific tissues. In humans, if any of the fundamental processes governed by GPCRs in specific tissues are dysfunctional, this will lead to acute or chronic diseases [157] [158] [159] [160]. The physical location and expression of GPCRs may provide a direct mechanism for the transduction of extracellular messages into tissue responses and functions. Consequently, the identification of the GPCRs in brains and CNS of insects could shed new light on the important roles they play in neuronal firing, the regulation of ion transportation across cell membranes and the modulation of neuron membrane potentials in other species.

References

- Hilger, D.; Masureel, M.; Kobilka, B.K. Structure and dynamics of GPCR signaling complexes. *Nat. Struct. Mol. Biol.* 2018, 25, 4–12.
- Bockaert, J.; Pin, J.P. Molecular tinkering of G protein-coupled receptors: An evolutionary success. *EMBO J.* 1999, 18, 1723–1729.

3. Maudsley, S.; Martin, B.; Luttrell, L.M. The Origins of Diversity and Specificity in G Protein-Coupled Receptor Signaling. *J. Pharmacol. Exp. Ther.* 2005, 314, 485–494.
4. Lagerström, M.C.; Schiöth, H.B. Structural diversity of G protein-coupled receptors and significance for drug discovery. *Nat. Rev. Drug Discov.* 2008, 7, 339–357.
5. Spehr, M.; Munger, S.D. Olfactory receptors: G protein-coupled receptors and beyond. *J. Neurochem.* 2009, 109, 1570–1583.
6. Millar, R.P.; Newton, C.L. The Year in G Protein-Coupled Receptor Research. *Mol. Endocrinol.* 2010, 24, 261–274.
7. Eglen, R.M.; Reisine, T. GPCRs Revisited: New Insights Lead to Novel Drugs. *Pharmaceuticals* 2011, 4, 244–272.
8. Gether, U. Uncovering Molecular Mechanisms Involved in Activation of G Protein-Coupled Receptors. *Endocr. Rev.* 2000, 21, 90–113.
9. Goldsmith, Z.G.; Dhanasekaran, D.N. G Protein regulation of MAPK networks. *Oncogene* 2007, 26, 3122–3142.
10. Gerald, W.Z. Calcium channel signaling complexes with receptors and channels. *Curr. Mol. Pharmacol.* 2015, 8, 8–11.
11. Flower, D.R. Modelling G-protein-coupled receptors for drug design. *Biochim. Biophys. Acta (BBA)-Rev. Biomembr.* 1999, 1422, 207–234.
12. Hopkins, A.L.; Groom, C.R. The druggable genome. *Nat. Rev. Drug Discov.* 2002, 1, 727–730.
13. Robas, N.; O'Reilly, M.; Katugampola, S.; Fidock, M. Maximizing serendipity: Strategies for identifying ligands for orphan G-protein-coupled receptors. *Curr. Opin. Pharmacol.* 2003, 3, 121–126.
14. Jacoby, E.; Bouhelal, R.; Gerspacher, M.; Seuwen, K. The 7 TM G-Protein-Coupled Receptor Target Family. *ChemMedChem* 2006, 1, 760–782.
15. Hauser, A.S.; Attwood, M.M.; Rask-Andersen, M.; Schiöth, H.B.; Gloriam, D.E. Trends in GPCR drug discovery: New agents, targets and indications. *Nat. Rev. Drug Discov.* 2017, 16, 829–842.
16. Congreve, M.; de Graaf, C.; Swain, N.A.; Tate, C.G. Impact of GPCR Structures on Drug Discovery. *Cell* 2020, 181, 81–91.
17. Hauser, F.; Cazzamali, G.; Williamson, M.; Blenau, W.; Grimmelhuijzen, C.J. A review of neurohormone GPCRs present in the fruitfly *Drosophila melanogaster* and the honey bee *Apis mellifera*. *Prog. Neurobiol.* 2006, 80, 1–19.
18. Hauser, F.; Cazzamali, G.; Williamson, M.; Park, Y.; Li, B.; Tanaka, Y.; Predel, R.; Neupert, S.; Schachtner, J.; Verleyen, P.; et al. A genome-wide inventory of neurohormone GPCRs in the red

- flour beetle *Tribolium castaneum*. *Front. Neuroendocr.* 2008, 29, 142–165.
19. Poels, J.; Van Loy, T.; Vandersmissen, H.P.; Van Hiel, B.; Van Soest, S.; Nachman, R.J.; Broeck, J.V. Myoinhibiting peptides are the ancestral ligands of the promiscuous *Drosophila* sex peptide receptor. *Cell. Mol. Life Sci.* 2010, 67, 3511–3522.
 20. Jiang, H.; Lkhagva, A.; Park, Y.; Kim, Y.-J.; Daubnerová, I.; Chae, H.-S.; Šimo, L.; Jung, S.-H.; Yoon, Y.-K.; Lee, N.-R.; et al. Natalisin, a tachykinin-like signaling system, regulates sexual activity and fecundity in insects. *Proc. Natl. Acad. Sci. USA* 2013, 110, E3526–E3534.
 21. Nagai, C.; Mabashi-Asazuma, H.; Nagasawa, H.; Nagata, S. Identification and Characterization of Receptors for Ion Transport Peptide (ITP) and ITP-like (ITPL) in the Silkworm *Bombyx mori*. *J. Biol. Chem.* 2014, 289, 32166–32177.
 22. Bai, H.; Palli, S.R. Identification of G protein-coupled receptors required for vitellogenin uptake into the oocytes of the red flour beetle, *Tribolium castaneum*. *Sci. Rep.* 2016, 6, 27648.
 23. Jing, Y.-P.; An, H.; Zhang, S.; Wang, N.; Zhou, S. Protein kinase C mediates juvenile hormone–dependent phosphorylation of Na⁺/K⁺-ATPase to induce ovarian follicular patency for yolk protein uptake. *J. Biol. Chem.* 2018, 293, 20112–20122.
 24. Marciniak, P.; Urbański, A.; Lubawy, J.; Szymczak, M.; Pacholska-Bogalska, J.; Chowański, S.; Kuczer, M.; Rosiński, G. Short neuropeptide F signaling regulates functioning of male reproductive system in *Tenebrio molitor* beetle. *J. Comp. Physiol. B* 2020, 190, 521–534.
 25. Mendive, F.M.; Van Loy, T.; Claeysen, S.; Poels, J.; Williamson, M.; Hauser, F.; Grimmelikhuijzen, C.J.; Vassart, G.; Broeck, J.V. *Drosophila* molting neurohormone bursicon is a heterodimer and the natural agonist of the orphan receptor DLGR2. *FEBS Lett.* 2005, 579, 2171–2176.
 26. Kim, Y.-J.; Zitnan, D.; Cho, K.-H.; Schooley, D.A.; Mizoguchi, A.; Adams, M.E. Central peptidergic ensembles associated with organization of an innate behavior. *Proc. Natl. Acad. Sci. USA* 2006, 103, 14211–14216.
 27. Žitňan, D.; Kim, Y.-J.; Žitňanová, I.; Roller, L.; Adams, M. Complex steroid–peptide–receptor cascade controls insect ecdysis. *Gen. Comp. Endocrinol.* 2007, 153, 88–96.
 28. Yamanaka, N.; Hua, Y.-J.; Roller, L.; Spalovská-Valachová, I.; Mizoguchi, A.; Kataoka, H.; Tanaka, Y. *Bombyx* prothoracicostatic peptides activate the sex peptide receptor to regulate ecdysteroid biosynthesis. *Proc. Natl. Acad. Sci. USA* 2010, 107, 2060–2065.
 29. Bai, H.; Zhu, F.; Shah, K.; Palli, S.R. Large-scale RNAi screen of G protein-coupled receptors involved in larval growth, molting and metamorphosis in the red flour beetle. *BMC Genom.* 2011, 12, 388.
 30. Li, B.; Beeman, R.W.; Park, Y. Functions of duplicated genes encoding CCAP receptors in the red flour beetle, *Tribolium castaneum*. *J. Insect Physiol.* 2011, 57, 1190–1197.

31. Li, K.; Jia, Q.; Li, S. Juvenile hormone signaling—A mini review. *Insect Sci.* 2019, 26, 600–606.
32. Nouzova, M.; Brockhoff, A.; Mayoral, J.G.; Goodwin, M.; Meyerhof, W.; Noriega, F.G. Functional characterization of an allatotropin receptor expressed in the corpora allata of Mosquitoes. *Peptides* 2012, 34, 201–208.
33. Iga, M.; Nakaoka, T.; Suzuki, Y.; Kataoka, H. Pigment Dispersing Factor Regulates Ecdysone Biosynthesis via Bombyx Neuropeptide G Protein Coupled Receptor-B2 in the Prothoracic Glands of Bombyx mori. *PLoS ONE* 2014, 9, e103239.
34. Regna, K.; Kurshan, P.T.; Harwood, B.N.; Jenkins, A.M.; Lai, C.-Q.; Muskavitch, M.A.; Kopin, A.S.; Draper, I. A critical role for the Drosophila dopamine D1-like receptor Dop1R2 at the onset of metamorphosis. *BMC Dev. Biol.* 2016, 16, 15.
35. Kang, X.-L.; Zhang, J.-Y.; Wang, D.; Zhao, Y.-M.; Han, X.-L.; Wang, J.-X.; Zhao, X.-F. The steroid hormone 20-hydroxyecdysone binds to dopamine receptor to repress lepidopteran insect feeding and promote pupation. *PLoS Genet.* 2019, 15, e1008331.
36. Homma, T.; Watanabe, K.; Tsurumaru, S.; Kataoka, H.; Imai, K.; Kamba, M.; Niimi, T.; Yamashita, O.; Yaginuma, T. G protein-coupled receptor for diapause hormone, an inducer of Bombyx embryonic diapause. *Biochem. Biophys. Res. Commun.* 2006, 344, 386–393.
37. Terhzaz, S.; Cabrero, P.; Robben, J.H.; Radford, J.C.; Hudson, B.D.; Milligan, G.; Dow, J.A.T.; Davies, S.-A. Mechanism and Function of Drosophila capa GPCR: A Desiccation Stress-Responsive Receptor with Functional Homology to Human NeuromedinU Receptor. *PLoS ONE* 2012, 7, e29897.
38. Bryon, A.; Wybouw, N.; Dermauw, W.; Tirry, L.; Van Leeuwen, T. Genome wide gene-expression analysis of facultative reproductive diapause in the two-spotted spider mite Tetranychus urticae. *BMC Genom.* 2013, 14, 1–20.
39. Choi, M.-Y.; Estep, A.; Sanscrainte, N.; Becnel, J.; Meer, R.K.V. Identification and expression of PBAN/diapause hormone and GPCRs from Aedes aegypti. *Mol. Cell. Endocrinol.* 2013, 375, 113–120.
40. Devambez, I.; Agha, M.A.; Mitri, C.; Bockaert, J.; Parmentier, M.-L.; Marion-Poll, F.; Grau, Y.; Soustelle, L. Gαo Is Required for L-Canavanine Detection in Drosophila. *PLoS ONE* 2013, 8, e63484.
41. Jiang, H.; Wei, Z.; Nachman, R.J.; Park, Y. Molecular cloning and functional characterization of the diapause hormone receptor in the corn earworm Helicoverpa zea. *Peptides* 2014, 53, 243–249.
42. Li, T.; Liu, L.; Zhang, L.; Liu, N. Role of G-protein-coupled Receptor-related Genes in Insecticide Resistance of the Mosquito, Culex quinquefasciatus. *Sci. Rep.* 2015, 4, 6474.

43. Li, T.; Cao, C.; Yang, T.; Zhang, L.; He, L.; Xi, Z.; Bian, G.; Liu, N. A G-protein-coupled receptor regulation pathway in cytochrome P450-mediated permethrin-resistance in Mosquitoes, *Culex quinquefasciatus*. *Sci. Rep.* 2015, 5, 17772.
44. Li, T.; Liu, N. Regulation of P450-mediated permethrin resistance in *Culex quinquefasciatus* by the GPCR/G α s/AC/cAMP/PKA signaling cascade. *Biochem. Biophys. Rep.* 2017, 12, 12–19.
45. Shen, Z.; Jiang, X.; Yan, L.; Chen, Y.; Wang, W.; Shi, Y.; Shi, L.; Liu, D.; Zhou, N. Structural basis for the interaction of diapause hormone with its receptor in the silkworm, *Bombyx mori*. *FASEB J.* 2018, 32, 1338–1353.
46. Chen, C.-H.; Di, Y.-Q.; Shen, Q.-Y.; Wang, J.-X.; Zhao, X.-F. The steroid hormone 20-hydroxyecdysone induces phosphorylation and aggregation of stromal interacting molecule 1 for store-operated calcium entry. *J. Biol. Chem.* 2019, 294, 14922–14936.
47. Petruccelli, E.; Lark, A.; Mrkvicka, J.A.; Kitamoto, T. Significance of DopEcR, a G-protein coupled dopamine/ecdysteroid receptor, in physiological and behavioral response to stressors. *J. Neurogenet.* 2020, 34, 55–68.
48. Chen, M.-E.; Pietrantonio, P.V. The short neuropeptide F-like receptor from the red imported fire ant, *Solenopsis invicta* Buren (Hymenoptera: Formicidae). *Arch. Insect Biochem. Physiol.* 2006, 61, 195–208.
49. Kersch, C.N.; Pietrantonio, P.V. Mosquito *Aedes aegypti* (L.) leucokinin receptor is critical for in vivo fluid excretion post blood feeding. *FEBS Lett.* 2011, 585, 3507–3512.
50. Dillen, S.; Zels, S.; Verlinden, H.; Spit, J.; Van Wielendaele, P.; Broeck, J.V. Functional Characterization of the Short Neuropeptide F Receptor in the Desert Locust, *Schistocerca gregaria*. *PLoS ONE* 2013, 8, e53604.
51. Baumbach, J.; Xu, Y.; Hehlert, P.; Kühnlein, R.P. G α q, Gy1 and Plc21C Control *Drosophila* Body Fat Storage. *J. Genet. Genom.* 2014, 41, 283–292.
52. Lin, F.; Hossain, M.A.; Post, S.; Karashchuk, G.; Tatar, M.; De Meyts, P.; Wade, J.D. Total Solid-Phase Synthesis of Biologically Active *Drosophila* Insulin-Like Peptide 2 (DILP2). *Aust. J. Chem.* 2017, 70, 208–212.
53. Lu, K.; Zhang, X.; Chen, X.; Li, Y.; Li, W.; Cheng, Y.; Zhou, J.; You, K.; Zhou, Q. Adipokinetic Hormone Receptor Mediates Lipid Mobilization to Regulate Starvation Resistance in the Brown Planthopper, *Nilaparvata lugens*. *Front. Physiol.* 2018, 9, 1730.
54. Marchal, E.; Schellens, S.; Monjon, E.; Bruyninckx, E.; Marco, H.G.; Gäde, G.; Broeck, J.V.; Verlinden, H. Analysis of Peptide Ligand Specificity of Different Insect Adipokinetic Hormone Receptors. *Int. J. Mol. Sci.* 2018, 19, 542.

55. Yang, H.; Huang, J.; Liu, Y.; Li, J.; Luo, S.; Wu, J. Prediction of the post-translational modifications of adipokinetic hormone receptors from solitary to eusocial bees. *Sociobiology* 2018, 65, 271–279.
56. Sangha, V.; Lange, A.B.; Orchard, I. Identification and cloning of the kinin receptor in the Chagas disease vector, *Rhodnius prolixus*. *Gen. Comp. Endocrinol.* 2020, 289, 113380.
57. Bainton, R.J.; Tsai, L.T.-Y.; Schwabe, T.; DeSalvo, M.; Gaul, U.; Heberlein, U. *moody* Encodes Two GPCRs that Regulate Cocaine Behaviors and Blood-Brain Barrier Permeability in *Drosophila*. *Cell* 2005, 123, 145–156.
58. Hyun, S.; Lee, Y.; Hong, S.-T.; Bang, S.; Paik, D.; Kang, J.; Shin, J.; Lee, J.; Jeon, K.; Hwang, S.; et al. *Drosophila* GPCR *Han* Is a Receptor for the Circadian Clock Neuropeptide PDF. *Neuron* 2005, 48, 267–278.
59. Thamm, M.; Balfanz, S.; Scheiner, R.; Baumann, A.; Blenau, W. Characterization of the 5-HT_{1A} receptor of the honeybee (*Apis mellifera*) and involvement of serotonin in phototactic behavior. *Cell. Mol. Life Sci.* 2010, 67, 2467–2479.
60. Abrieux, A.; Debernard, S.; Maria, A.; Gaertner, C.; Anton, S.; Gadenne, C.; Duportets, L. Involvement of the G-Protein-Coupled Dopamine/Ecdysteroid Receptor *DopEcR* in the Behavioral Response to Sex Pheromone in an Insect. *PLoS ONE* 2013, 8, e72785.
61. Kwon, H.; Agha, M.A.; Smith, R.C.; Nachman, R.J.; Marion-Poll, F.; Pietrantonio, P.V. Leucokinin mimetic elicits aversive behavior in Mosquito *Aedes aegypti* (L.) and inhibits the sugar taste neuron. *Proc. Natl. Acad. Sci. USA* 2016, 113, 6880–6885.
62. Hou, Q.-L.; Chen, E.-H.; Jiang, H.-B.; Wei, D.-D.; Gui, S.-H.; Wang, J.-J.; Smagghe, G. Adipokinetic hormone receptor gene identification and its role in triacylglycerol mobilization and sexual behavior in the oriental fruit fly (*Bactrocera dorsalis*). *Insect Biochem. Mol. Biol.* 2017, 90, 1–13.
63. Grohmann, L.; Blenau, W.; Erber, J.; Ebert, P.R.; Strünker, T.; Baumann, A. Molecular and functional characterization of an octopamine receptor from honeybee (*Apis mellifera*) brain. *J. Neurochem.* 2003, 86, 725–735.
64. Johnson, E.C.; Bohn, L.M.; Taghert, P.H. *Drosophila* CG8422 encodes a functional diuretic hormone receptor. *J. Exp. Biol.* 2004, 207, 743–748.
65. Kwon, H.; Lu, H.-L.; Longnecker, M.T.; Pietrantonio, P.V. Role in Diuresis of a Calcitonin Receptor (GPCRCAL1) Expressed in a Distal-Proximal Gradient in Renal Organs of the Mosquito *Aedes aegypti* (L.). *PLoS ONE* 2012, 7, e50374.
66. Kwon, H.; Pietrantonio, P.V. Calcitonin receptor 1 (*AedaeGPCRCAL1*) hindgut expression and direct role in myotropic action in females of the Mosquito *Aedes aegypti* (L.). *Insect Biochem. Mol. Biol.* 2013, 43, 588–593.

67. Lee, D.; Broeck, J.V.; Lange, A.B. Identification and Expression of the CCAP Receptor in the Chagas' Disease Vector, *Rhodnius prolixus*, and Its Involvement in Cardiac Control. *PLoS ONE* 2013, 8, e68897.
68. Petruccelli, E.; Li, Q.; Rao, Y.; Kitamoto, T. The Unique Dopamine/Ecdysteroid Receptor Modulates Ethanol-Induced Sedation in *Drosophila*. *J. Neurosci.* 2016, 36, 4647–4657.
69. Pietrantonio, P.V.; Xiong, C.; Nachman, R.J.; Shen, Y. G protein-coupled receptors in arthropod vectors: Omics and pharmacological approaches to elucidate ligand-receptor interactions and novel organismal functions. *Curr. Opin. Insect Sci.* 2018, 29, 12–20.
70. Uchiyama, H.; Maehara, S.; Ohta, H.; Seki, T.; Tanaka, Y. Elevenin regulates the body color through a G protein-coupled receptor NIA42 in the brown planthopper *Nilaparvata lugens*. *Gen. Comp. Endocrinol.* 2018, 258, 33–38.
71. Li, M.; Reid, W.R.; Zhang, L.; Scott, J.G.; Gao, X.; Kristensen, M.; Liu, N. A whole transcriptomal linkage analysis of gene co-regulation in insecticide resistant house flies, *Musca domestica*. *BMC Genom.* 2013, 14, 803.
72. Ma, Z.; Zhang, Y.; You, C.; Zeng, X.; Gao, X. The role of G protein-coupled receptor-related genes in cytochrome P450-mediated resistance of the house fly, *Musca domestica* (Diptera: Muscidae), to imidacloprid. *Insect Mol. Biol.* 2019, 29, 92–103.
73. Liu, N. Insecticide Resistance in Mosquitoes: Impact, Mechanisms, and Research Directions. *Annu. Rev. Entomol.* 2015, 60, 537–559.
74. Attwood, T.K.; Findlay, J.B.C. Fingerprinting G-protein-coupled receptors. *Protein Eng. Des. Sel.* 1994, 7, 195–203.
75. Kolakowski, L.F., Jr. GCRDb: A G-protein-coupled receptor database. *Recept. Channels* 1994, 2, 1–7.
76. Schiöth, H.B.; Fredriksson, R. The GRAFS classification system of G-protein coupled receptors in comparative perspective. *Gen. Comp. Endocrinol.* 2005, 142, 94–101.
77. Hu, G.-M.; Mai, T.-L.; Chen, C.-M. Visualizing the GPCR Network: Classification and Evolution. *Sci. Rep.* 2017, 7, 1–15.
78. Adams, M.D.; Celniker, S.E.; Holt, R.A.; Evans, C.A.; Gocayne, J.D.; Amanatides, P.G.; Scherer, S.E.; Li, P.W.; Hoskins, R.A.; Galle, R.F.; et al. The Genome Sequence of *Drosophila melanogaster*. *Science* 2000, 287, 2185–2195.
79. Holt, R.A.; Subramanian, G.M.; Halpern, A.; Sutton, G.G.; Charlab, R.; Nusskern, D.R.; Wincker, P.; Clark, A.G.; Ribeiro, J.C.; Wides, R.; et al. The Genome Sequence of the Malaria Mosquito *Anopheles gambiae*. *Science* 2002, 298, 129–149.

80. Nene, V.; Wortman, J.R.; Lawson, D.; Haas, B.; Kodira, C.; Tu, Z.; Loftus, B.; Xi, Z.; Megy, K.; Grabherr, M.; et al. Genome Sequence of *Aedes aegypti*, a Major Arbovirus Vector. *Science* 2007, 316, 1718–1723.
81. Arensburger, P.; Megy, K.; Waterhouse, R.M.; Abrudan, J.; Amedeo, P.; Antelo, B.; Bartholomay, L.; Bidwell, S.; Caler, E.; Camara, F.; et al. Sequencing of *Culex quinquefasciatus* Establishes a Platform for Mosquito Comparative Genomics. *Science* 2010, 330, 86–88.
82. Scott, J.G.; Warren, W.C.; Beukeboom, L.W.; Bopp, D.; Clark, A.G.; Giers, S.D.; Hediger, M.; Jones, A.K.; Kasai, S.; A Leichter, C.; et al. Genome of the house fly, *Musca domestica* L., a global vector of diseases with adaptations to a septic environment. *Genome Biol.* 2014, 15, 1–17.
83. Li, F.; Zhao, X.; Li, M.; He, K.; Huang, C.; Zhou, Y.; Li, Z.; Walters, J.R. Insect genomes: Progress and challenges. *Insect Mol. Biol.* 2019, 28, 739–758.
84. Brody, T.; Cravchik, A. *Drosophila melanogaster* G Protein–Coupled Receptors. *J. Cell Biol.* 2000, 150, F83–F88.
85. Hanlon, C.D.; Andrew, D.J. Outside-in signaling—A brief review of GPCR signaling with a focus on the *Drosophila* GPCR family. *J. Cell Sci.* 2015, 128, 3533–3542.
86. Hill, C.A.; Fox, A.N.; Pitts, R.J.; Kent, L.B.; Tan, P.L.; Chrystal, M.A.; Cravchik, A.; Collins, F.H.; Robertson, H.M.; Zwiebel, L.J. G Protein-Coupled Receptors in *Anopheles gambiae*. *Science* 2002, 298, 176–178.
87. Fan, Y.; Sun, P.; Wang, Y.; He, X.; Deng, X.; Chen, X.; Zhang, G.; Chen, X.; Zhou, N. The G protein-coupled receptors in the silkworm, *Bombyx mori*. *Insect Biochem. Mol. Biol.* 2010, 40, 581–591.
88. Anstead, C.A.; Korhonen, P.K.; Young, N.D.; Hall, R.S.; Jex, A.R.; Murali, S.C.; Hughes, D.S.; Lee, S.F.; Perry, T.; Stroehlein, A.J.; et al. *Lucilia cuprina* genome unlocks parasitic fly biology to underpin future interventions. *Nat. Commun.* 2015, 6, 7344.
89. Şahbaz, B.D. Prediction and expression analysis of G protein-coupled receptors in the laboratory stick insect, *Carausius morosus*. *Turk. J. Boil.* 2019, 43, 77–88.
90. Calkins, T.L.; Tamborindeguy, C.; Pietrantonio, P.V. GPCR annotation, G proteins, and transcriptomics of fire ant (*Solenopsis invicta*) queen and worker brain: An improved view of signaling in an invasive superorganism. *Gen. Comp. Endocrinol.* 2019, 278, 89–103.
91. Veenstra, J.A.; Rombauts, S.; Grbić, M. In silico cloning of genes encoding neuropeptides, neurohormones and their putative G-protein coupled receptors in a spider mite. *Insect Biochem. Mol. Biol.* 2012, 42, 277–295.
92. Aikins, M.J.; Schooley, D.A.; Begum, K.; Detheux, M.; Beeman, R.W.; Park, Y. Vasopressin-like peptide and its receptor function in an indirect diuretic signaling pathway in the red flour beetle.

- Insect Biochem. Mol. Biol. 2008, 38, 740–748.
93. Caers, J.; Verlinden, H.; Zels, S.; Vandersmissen, H.P.; Vuerinckx, K.; Schoofs, L. More than two decades of research on insect neuropeptide GPCRs: An overview. *Front. Endocrinol.* 2012, 3, 151.
 94. Garczynski, S.F.; Brown, M.R.; Shen, P.; Murray, T.F.; Crim, J.W. Characterization of a functional neuropeptide F receptor from *Drosophila melanogaster*. *Peptides* 2002, 23, 773–780.
 95. Vogel, K.J.; Brown, M.R.; Strand, M.R. Phylogenetic Investigation of Peptide Hormone and Growth Factor Receptors in Five Dipteran Genomes. *Front. Endocrinol.* 2013, 4, 193.
 96. Xia, R.; Li, M.; Wu, Y.; Qi, Y.; Ye, G.; Huang, J. A new family of insect muscarinic acetylcholine receptors. *Insect Mol. Biol.* 2016, 25, 362–369.
 97. Srivastava, D.P.; Yu, E.J.; Kennedy, K.; Chatwin, H.; Reale, V.; Hamon, M.; Smith, T.; Evans, P.D. Rapid, Nongenomic Responses to Ecdysteroids and Catecholamines Mediated by a Novel *Drosophila* G-Protein-Coupled Receptor. *J. Neurosci.* 2005, 25, 6145–6155.
 98. Deveci, D.; Martin, F.A.; Leopold, P.; Romero, N.M. AstA Signaling Functions as an Evolutionary Conserved Mechanism Timing Juvenile to Adult Transition. *Curr. Biol.* 2019, 29, 813–822.
 99. Chen, J.; Reiher, W.; Hermann-Luibl, C.; Sellami, A.; Cognigni, P.; Kondo, S.; Helfrich-Förster, C.; Veenstra, J.A.; Wegener, C. Allatostatin A Signalling in *Drosophila* Regulates Feeding and Sleep and Is Modulated by PDF. *PLoS Genet.* 2016, 12, e1006346.
 100. Zandawala, M.; Yurgel, M.E.; Texada, M.J.; Liao, S.; Rewitz, K.F.; Keene, A.C.; Nässel, D.R. Modulation of *Drosophila* post-feeding physiology and behavior by the neuropeptide leucokinin. *PLoS Genet.* 2018, 14, e1007767.
 101. Yapici, N.; Kim, Y.-J.; Ribeiro, C.; Dickson, B.J. A receptor that mediates the post-mating switch in *Drosophila* reproductive behaviour. *Nat. Cell Biol.* 2007, 451, 33–37.
 102. Sellami, A.; Veenstra, J.A. SIFamide acts on fruitless neurons to modulate sexual behavior in *Drosophila melanogaster*. *Peptides* 2015, 74, 50–56.
 103. Yurgel, M.E.; Kakad, P.; Zandawala, M.; Nässel, D.R.; Godenschwege, T.A.; Keene, A.C. A single pair of leucokinin neurons are modulated by feeding state and regulate sleep–metabolism interactions. *PLoS Biol.* 2019, 17, e2006409.
 104. Wu, F.; Deng, B.; Xiao, N.; Wang, T.; Li, Y.; Wang, R.; Shi, K.; Luo, D.-G.; Rao, Y.; Zhou, C. A neuropeptide regulates fighting behavior in *Drosophila melanogaster*. *eLife* 2020, 9.
 105. Sano, H.; Nakamura, A.; Texada, M.J.; Truman, J.W.; Ishimoto, H.; Kamikouchi, A.; Nibu, Y.; Kume, K.; Ida, T.; Kojima, M. The Nutrient-Responsive Hormone CCHamide-2 Controls Growth by Regulating Insulin-like Peptides in the Brain of *Drosophila melanogaster*. *PLoS Genet.* 2015, 11, e1005209.

106. Iyison, N.B.; Sinmaz, M.G.; Sahbaz, B.D.; Shahraki, A.; Aksoydan, B.; Durdagi, S. In silico characterization of adipokinetic hormone receptor and screening for pesticide candidates against stick insect, *Carausius morosus*. *J. Mol. Graph. Model.* 2020, 101, 107720.
107. Posnien, N.; Hopfen, C.; Hilbrant, M.; Ramos-Womack, M.; Murat, S.; Schönauer, A.; Herbert, S.L.; Nunes, M.D.S.; Arif, S.; Breuker, C.J.; et al. Evolution of Eye Morphology and Rhodopsin Expression in the *Drosophila melanogaster* Species Subgroup. *PLoS ONE* 2012, 7, e37346.
108. Egerod, K.; Reynisson, E.; Hauser, F.; Cazzamali, G.; Williamson, M.; Grimmelikhuijzen, C.J.P. Molecular cloning and functional expression of the first two specific insect myosuppressin receptors. *Proc. Natl. Acad. Sci. USA* 2003, 100, 9808–9813.
109. Hauser, F.; Williamson, M.; Cazzamali, G.; Grimmelikhuijzen, C.J.P. Identifying neuropeptide and protein hormone receptors in *Drosophila melanogaster* by exploiting genomic data. *Briefings Funct. Genom. Proteom.* 2006, 4, 321–330.
110. Bayliss, A.; Roselli, G.; Evans, P.D. A comparison of the signalling properties of two tyramine receptors from *Drosophila*. *J. Neurochem.* 2013, 125, 37–48.
111. Li, C.; Chen, M.; Sang, M.; Liu, X.; Wu, W.; Li, B. Comparative genomic analysis and evolution of family-B G protein-coupled receptors from six model insect species. *Gene* 2013, 519, 1–12.
112. De Mendoza, A.; Jones, J.W.; Friedrich, M. Methuselah/Methuselah-like G protein-coupled receptors constitute an ancient metazoan gene family. *Sci. Rep.* 2016, 6, 21801.
113. Hector, C.E.; Bretz, C.A.; Zhao, Y.; Johnson, E.C. Functional differences between two CRF-related diuretic hormone receptors in *Drosophila*. *J. Exp. Biol.* 2009, 212, 3142–3147.
114. Goda, T.; Doi, M.; Umezaki, Y.; Murai, I.; Shimatani, H.; Chu, M.L.; Nguyen, V.H.; Okamura, H.; Hamada, F.N. Calcitonin receptors are ancient modulators for rhythms of preferential temperature in insects and body temperature in mammals. *Genes Dev.* 2018, 32, 140–155.
115. Lin, Y.-J.; Seroude, L.; Benzer, S. Extended Life-Span and Stress Resistance in the *Drosophila* Mutant methuselah. *Science* 1998, 282, 943–946.
116. Cvejic, S.; Zhu, Z.; Felice, S.J.; Berman, Y.; Huang, X.-Y. The endogenous ligand Stunted of the GPCR Methuselah extends lifespan in *Drosophila*. *Nat. Cell Biol.* 2004, 6, 540–546.
117. Harmor, A.J. Family-B G-protein-coupled receptors. *Genome Biol.* 2001, 2, 1–3013.
118. Mezler, M.; Müller, T.; Raming, K. Cloning and functional expression of GABA_B receptors from *Drosophila*. *Eur. J. Neurosci.* 2001, 13, 477–486.
119. Adler, P.N.; Vinson, C.; Park, W.J.; Conover, S.; Klein, L. Molecular structure of frizzled, a *Drosophila* tissue polarity gene. *Genetics* 1990, 126, 401–416.

120. I Strutt, D. Asymmetric Localization of Frizzled and the Establishment of Cell Polarity in the *Drosophila* Wing. *Mol. Cell* 2001, 7, 367–375.
121. Povelones, M.; Nusse, R. The role of the cysteine-rich domain of Frizzled in Wingless-Armadillo signaling. *EMBO J.* 2005, 24, 3493–3503.
122. Heuvel, M.V.D.; Ingham, P.W. *smoothed* encodes a receptor-like serpentine protein required for hedgehog signalling. *Nat. Cell Biol.* 1996, 382, 547–551.
123. Fan, J.; Liu, Y.; Jia, J. Hh-induced *Smoothed* conformational switch is mediated by differential phosphorylation at its C-terminal tail in a dose- and position-dependent manner. *Dev. Biol.* 2012, 366, 172–184.
124. Martin, B.L.; Kimelman, D. Wnt Signaling and the Evolution of Embryonic Posterior Development. *Curr. Biol.* 2009, 19, R215–R219.
125. Villarreal, C.M.; Darakananda, K.; Wang, V.R.; Jayaprakash, P.M.; Suzuki, Y. Hedgehog signaling regulates imaginal cell differentiation in a basally branching holometabolous insect. *Dev. Biol.* 2015, 404, 125–135.
126. Abrieux, A.; Duportets, L.; Debernard, S.; Gadenne, C.; Anton, S. The GPCR membrane receptor, DopEcR, mediates the actions of both dopamine and ecdysone to control sex pheromone perception in an insect. *Front. Behav. Neurosci.* 2014, 8, 312.
127. Lark, A.; Kitamoto, T.; Martin, J.-R. Modulation of neuronal activity in the *Drosophila* mushroom body by DopEcR, a unique dual receptor for ecdysone and dopamine. *Biochim. Biophys. Acta (BBA)-Bioenerg.* 2017, 1864, 1578–1588.
128. Lungchukiet, P.; Zhang, J.; Tobe, S.S.; Bendena, W.G. Quantification of allatostatin receptor mRNA levels in the cockroach, *Diploptera punctata*, using real-time PCR. *J. Insect Physiol.* 2008, 54, 981–987.
129. Ida, T.; Takahashi, T.; Tominaga, H.; Sato, T.; Kume, K.; Ozaki, M.; Hiraguchi, T.; Maeda, T.; Shiotani, H.; Terajima, S.; et al. Identification of the novel bioactive peptides dRYamide-1 and dRYamide-2, ligands for a neuropeptide Y-like receptor in *Drosophila*. *Biochem. Biophys. Res. Commun.* 2011, 410, 872–877.
130. Deng, X.; Yang, H.; He, X.; Liao, Y.; Zheng, C.; Zhou, Q.; Zhu, C.; Zhang, G.; Gao, J.; Zhou, N. Activation of *Bombyx* neuropeptide G protein-coupled receptor A4 via a Gai-dependent signaling pathway by direct interaction with neuropeptide F from silkworm, *Bombyx mori*. *Insect Biochem. Mol. Biol.* 2014, 45, 77–88.
131. Gross, A.D.; Temeyer, K.B.; Day, T.A.; de León, A.A.P.; Kimber, M.J.; Coats, J.R. Pharmacological characterization of a tyramine receptor from the southern cattle tick, *Rhipicephalus (Boophilus) microplus*. *Insect Biochem. Mol. Biol.* 2015, 63, 47–53.

132. Collin, C.; Hauser, F.; Krogh-Meyer, P.; Hansen, K.K.; De Valdivia, E.G.; Williamson, M.; Grimmeliikhuijzen, C.J. Identification of the *Drosophila* and *Tribolium* receptors for the recently discovered insect RYamide neuropeptides. *Biochem. Biophys. Res. Commun.* 2011, 412, 578–583.
133. Qi, Y.-X.; Xia, R.-Y.; Wu, Y.-S.; Stanley, D.; Huang, J.; Ye, G.-Y. Larvae of the small white butterfly, *Pieris rapae*, express a novel serotonin receptor. *J. Neurochem.* 2014, 131, 767–777.
134. Vleugels, R.; Lenaerts, C.; Broeck, J.V.; Verlinden, H. Signalling properties and pharmacology of a 5-HT7-type serotonin receptor from *Tribolium castaneum*. *Insect Mol. Biol.* 2013, 23, 230–243.
135. Cao, C.; Sun, L.; Du, H.; Moural, T.W.; Bai, H.; Liu, P.; Zhu, F. Physiological functions of a methuselah-like G protein coupled receptor in *Lymantria dispar* Linnaeus. *Pestic. Biochem. Physiol.* 2019, 160, 1–10.
136. Pandey, A.; Khatoon, R.; Saini, S.; Vimal, D.; Patel, D.K.; Narayan, G.; Chowdhuri, D.K. Efficacy of methuselah gene mutation toward tolerance of dichlorvos exposure in *Drosophila melanogaster*. *Free. Radic. Biol. Med.* 2015, 83, 54–65.
137. Tobin, A.B.; Butcher, A.J.; Kong, K.C. Location, location, location...site-specific GPCR phosphorylation offers a mechanism for cell-type-specific signalling. *Trends Pharmacol. Sci.* 2008, 29, 413–420.
138. Qi, Y.-X.; Xu, G.; Gu, G.-X.; Mao, F.; Ye, G.-Y.; Liu, W.; Huang, J. A new *Drosophila* octopamine receptor responds to serotonin. *Insect Biochem. Mol. Biol.* 2017, 90, 61–70.
139. Van Hiel, M.B.; Vandersmissen, H.P.; Proost, P.; Broeck, J.V. Cloning, constitutive activity and expression profiling of two receptors related to relaxin receptors in *Drosophila melanogaster*. *Peptides* 2015, 68, 83–90.
140. Yamanaka, N.; Yamamoto, S.; Žitňan, D.; Watanabe, K.; Kawada, T.; Satake, H.; Kaneko, Y.; Hiruma, K.; Tanaka, Y.; Shinoda, T.; et al. Neuropeptide Receptor Transcriptome Reveals Unidentified Neuroendocrine Pathways. *PLoS ONE* 2008, 3, e3048.
141. Reim, T.; Balfanz, S.; Baumann, A.; Blenau, W.; Thamm, M.; Scheiner, R. AmTAR2: Functional characterization of a honeybee tyramine receptor stimulating adenylyl cyclase activity. *Insect Biochem. Mol. Biol.* 2017, 80, 91–100.
142. Paluzzi, J.-P.; O'Donnell, M.J. Identification, spatial expression analysis and functional characterization of a pyrokinin-1 receptor in the Chagas' disease vector, *Rhodnius prolixus*. *Mol. Cell. Endocrinol.* 2012, 363, 36–45.
143. Hamoudi, Z.; Lange, A.B.; Orchard, I. Identification and Characterization of the Corazonin Receptor and Possible Physiological Roles of the Corazonin-Signaling Pathway in *Rhodnius prolixus*. *Front. Neurosci.* 2016, 10, 357.

144. Paluzzi, J.-P.V.; Ebhatt, G.; Wang, C.-H.J.; Ezandawala, M.; Lange, A.B.; Eorchar, I. Identification, functional characterization, and pharmacological profile of a serotonin type-2b receptor in the medically important insect, *Rhodnius prolixus*. *Front. Neurosci.* 2015, 9, 175.
145. Sha, K.; Conner, W.C.; Choi, D.Y.; Park, J.H. Characterization, expression, and evolutionary aspects of Corazonin neuropeptide and its receptor from the House Fly, *Musca domestica* (Diptera: Muscidae). *Gene* 2012, 497, 191–199.
146. Verlinden, H.; Vleugels, R.; Verdonck, R.; Urlacher, E.; Broeck, J.V.; Mercer, A. Pharmacological and signalling properties of a D2-like dopamine receptor (Dop3) in *Tribolium castaneum*. *Insect Biochem. Mol. Biol.* 2015, 56, 9–20.
147. Stafflinger, E.; Hansen, K.K.; Hauser, F.; Schneider, M.; Cazzamali, G.; Williamson, M.; Grimmelhuijzen, C.J.P. Cloning and identification of an oxytocin/vasopressin-like receptor and its ligand from insects. *Proc. Natl. Acad. Sci. USA* 2008, 105, 3262–3267.
148. Verlinden, H.; Vleugels, R.; Marchal, E.; Badisco, L.; Tobback, J.; Pflüger, H.-J.; Blenau, W.; Broeck, J.V. The cloning, phylogenetic relationship and distribution pattern of two new putative GPCR-type octopamine receptors in the desert locust (*Schistocerca gregaria*). *J. Insect Physiol.* 2010, 56, 868–875.
149. Wang, S.; Wang, W.; Ma, Q.; Shen, Z.; Zhang, M.; Zhou, N.; Zhang, C. Elevenin signaling modulates body color through the tyrosine-mediated cuticle melanism pathway. *FASEB J.* 2019, 33, 9731–9741.
150. Wu, S.-F.; Xu, G.; Qi, Y.-X.; Xia, R.-Y.; Huang, J.; Ye, G.-Y. Two splicing variants of a novel family of octopamine receptors with different signaling properties. *J. Neurochem.* 2014, 129, 37–47.
151. Xu, G.; Gu, G.-X.; Teng, Z.-W.; Wu, S.-F.; Huang, J.; Song, Q.-S.; Ye, G.-Y.; Fang, Q. Identification and expression profiles of neuropeptides and their G protein-coupled receptors in the rice stem borer *Chilo suppressalis*. *Sci. Rep.* 2016, 6, 28976.
152. Zhang, F.; Wang, J.; Thakur, K.; Hu, F.; Zhang, J.-G.; Jiang, X.-F.; An, S.-H.; Jiang, H.; Jiang, L.; Wei, Z.-J. Isolation functional characterization of allatotropin receptor from the cotton bollworm, *Helicoverpa armigera*. *Peptides* 2019, 122, 169874.
153. Pietrantonio, P.V.; Jagge, C.; Taneja-Bageshwar, S.; Nachman, R.J.; Barhoumi, R. The Mosquito *Aedes aegypti* (L.) leucokinin receptor is a multiligand receptor for the three *Aedes* kinins. *Insect Mol. Biol.* 2005, 14, 55–67.
154. Esquivel, C.J.; Cassone, B.J.; Piermarini, P.M. A de novo transcriptome of the Malpighian tubules in non-blood-fed and blood-fed Asian tiger Mosquitoes *Aedes albopictus*: Insights into diuresis, detoxification, and blood meal processing. *PeerJ* 2016, 4, e1784.
155. Munoz, S.; Guerrero, F.D.; Kellogg, A.; Heekin, A.M.; Leung, M.-Y. Bioinformatic prediction of G protein-coupled receptor encoding sequences from the transcriptome of the foreleg, including the

- Haller's organ, of the cattle tick, *Rhipicephalus australis*. PLoS ONE 2017, 12, e0172326.
156. Verlinden, H.; Lismont, E.; Bil, M.; Urlacher, E.; Mercer, A.; Broeck, J.V.; Huybrechts, R. Characterisation of a functional allatotropin receptor in the bumblebee, *Bombus terrestris* (Hymenoptera, Apidae). *Gen. Comp. Endocrinol.* 2013, 193, 193–200.
 157. Kawakami, N.; Miyoshi, K.; Horio, S.; Fukui, H. β 2-Adrenergic Receptor-Mediated Histamine H1 Receptor Down-Regulation: Another Possible Advantage of β 2 Agonists in Asthmatic Therapy. *J. Pharmacol. Sci.* 2004, 94, 449–458.
 158. Drake, M.T.; Shenoy, S.K.; Lefkowitz, R.J. Trafficking of G Protein–Coupled Receptors. *Circ. Res.* 2006, 99, 570–582.
 159. Chen, J.-F.; Sonsalla, P.K.; Pedata, F.; Melani, A.; Domenici, M.R.; Popoli, P.; Geiger, J.; Lopes, L.V.; de Mendonça, A. Adenosine A2A receptors and brain injury: Broad spectrum of neuroprotection, multifaceted actions and “fine tuning” modulation. *Prog. Neurobiol.* 2007, 83, 310–331.
 160. Duan, W.; Gui, L.; Zhou, Z.; Liu, Y.; Tian, H.; Chen, J.-F.; Zheng, J. Adenosine A2A receptor deficiency exacerbates white matter lesions and cognitive deficits induced by chronic cerebral hypoperfusion in mice. *J. Neurol. Sci.* 2009, 285, 39–45.

Retrieved from <https://encyclopedia.pub/entry/history/show/24606>