

Health Effects of Berry Anthocyanins

Subjects: **Nutrition & Dietetics**

Contributor: Sanne Ahles

Supplementation with anthocyanins, which are a type of flavonoids mainly found in various berries, is hypothesized to be a promising approach to lower the risk of developing cognitive decline. The aim of this systematic review was to provide a comprehensive overview of dietary intervention trials describing effects of berry anthocyanins on cognitive performance in humans, while also addressing potential underlying mechanisms.

anthocyanins

cognitive performance

vascular function

cardiometabolic risk markers

1. Introduction

Cognitive performance encompasses multiple mental abilities that can be categorized into various domains, such as attention and psychomotor speed, memory, and executive function [1]. From childhood, cognitive performance quickly improves until young adulthood, after which it gradually starts declining [2]. Therefore, it is becoming increasingly relevant to focus on improving and/or maintaining cognitive performance to delay and prevent cognitive decline, and ultimately the onset of dementia [3]. This could be achieved by targeting potential mechanisms that drive cognitive performance [4][5].

An impaired vascular function is a common pathophysiological characteristic of multiple age-related conditions [6][7]. Vascular function can be assessed by determining endothelial function with methods such as brachial artery flow-mediated vasodilation (FMD) or the reactive hyperemia index (RHI) [8]. Previous research has already shown that vascular health declines with age leading to an increased risk of cognitive impairment, which may partly be explained by co-existing cardiometabolic risk factors, such as high blood pressure (BP) or a disturbed lipid profile such as altered low-density lipoprotein cholesterol (LDL-C) or high-density lipoprotein cholesterol (HDL-C) concentrations [9]. Therefore, dietary interventions that target vascular function and/or cardiometabolic risk markers may improve cognitive performance [10][11][12].

Therefore, increasing dietary intake of anthocyanins through supplementation could be a useful strategy to lower the risk of developing cognitive decline. A recent systematic review by Kent and colleagues [13] reported different intervention studies with beneficial effects of food-derived anthocyanins on cognitive performance. However, a systematic review designed to evaluate the effects of dietary anthocyanin interventions on cognitive performance and underlying mechanisms (i.e., vascular function and cardiometabolic risk markers) in an integrated manner has not been published yet. Therefore, the aim of this systematic literature review was to provide an overview of dietary intervention trials describing effects of berry anthocyanins on cognitive performance, vascular function, and cardiometabolic risk markers in humans.

2. The Effect of Berry Anthocyanins on Cognitive Performance

Of the eighteen studies that determined the effects of berry anthocyanins on cognitive performance outcomes, fifteen used a blueberry intervention, while the other three studies used either a chokeberry extract, a blackcurrant juice, or a blackcurrant extract. Results on cognitive performance were clustered based on the domains evaluated in the studies, i.e., (i) attention and psychomotor speed domain, (ii) executive function domain, (iii) memory domain, or (iv) other tests. Study results are shown in Table 1.

Table 1. The effect of berry anthocyanins on cognitive performance outcomes, compared to control.

Author (Year)	Intervention	Anthocyanin Dose	Attention and Psychomotor Speed					Executive Function					Memory			Other	
			TMT-A	MFT	GPT	FCRTT	Miscellaneous	TMT-B	Stroop	(M)ANT	Go-No-Go	Miscellaneous	RAVLT - HVLT - CVLT	VPAL and SPAL	WRT	n-back	
Ahles (2020) ^[14]	Chokeberry extract	16 mg		↑		= (NCT)			=								
		27 mg		=		= (NCT)			=								
Barfoot (2019) ^[15]	Freeze-dried wild blueberry juice	253 mg							↑				↑ (R)				= (TOWRE-2)
Boespflug (2018) ^[16]	Freeze-dried blueberry powder	269 mg													↑?		
Bowtell (2017) ^[17]	Blueberry extract	387 mg							=						↑	= (ISLT)	= (Groton Maze)
Cook (2020) ^[18]	New Zealand blackcurrant extract	210 mg		=	=	(RVIP, SRT)					= (SWM)			= (S)			
Krikorian (2010) ^[19]	Blueberry juice	428-598 mg ¹										↑ (C)		↑ (V)			
Krikorian (2020) ^[20]	Freeze-dried blueberry fruit powder	258 mg	↑?					=			↑ (COWAT)	= (H)		↑ (S)			

Author (Year)	Intervention	Anthocyanin Dose	Attention and Psychomotor Speed		Executive Function			Memory	Other
McNamara (2018) [21]	Freeze-dried blueberry powder	269 mg	=	=		= (COWAT)	↑ (H)		
Miller (2018) [22]	Freeze-dried blueberry powder	230 mg #	=	=	=	↑ (TST)	↑ (C)	= (DST)	= (VMWMT)
Traupe (2018) [23]	Blueberry juice	nr		↑ (AMT)	↑			↑ (Prose Memory)	
Watson (2019) [24]	Blackcurrant juice	115.09 mg	↑	= (DVT, SRTT)					21,
Whyte (2015) [25]	Blueberry juice	143 mg		=	=	=? (R)	=	= (OLT)	
Whyte (2016) [26]	Freeze-dried wild blueberry powder	127 mg	=		↓?	↑? (PMT)	↑? (R)		1. Sci.
		254 mg	↑		=	↑? (PMT)	↑ (R)		
Whyte (2017) [27]	Wild blueberry powder	253 mg			↑				
		1.35 mg		=	=	= (R)	=	= (CBT, SST, SMST)	
Whyte (2018) [28]	Wild blueberry powder and extract	2.7 mg		=	=	= (R)	=	= (CBT, SST, SMST)	Vasc.
		7 mg		=	=	= (R)	↑	↑? (CBT); = (SST, SMST)	
Whyte (2020) [29]	Wild blueberry powder	475 mg		=	↑		↑ (R)		
Whyte (2020) [30]	Wild blueberry powder	253 mg				= (R)		↑ (VSGT); = (BPT, PRT)	. Glob. Cardiol. Sci.
	Wild blueberry powder	253 mg	↑?		= Stop-Go, TST)				

Pract. 2014, 291–308.

7. Creager, M.A.; Lüscher, T.F.; Cosentino, F.; Beckman, J.A. Diabetes and Vascular Disease. Circulation 2003, 108, 1527–1532.
8. Flammer, A.J.; Anderson, T.; Celermajer, D.S.; Creager, M.A.; Deanfield, J.; Ganz, P.; Hamburg, N.M.; Lüscher, T.F.; Shechter, M.; Taddei, S.; et al. The assessment of endothelial function: From research into clinical practice. Circulation 2012, 126, 753–767.
9. Tsentidou, G.; Moraitou, D.; Tsolaki, M. Cognition in Vascular Aging and Mild Cognitive Impairment. J. Alzheimers Dis. 2019, 72, 55–70.
10. Clare, L.; Wu, Y.-T.; Teale, J.C.; MacLeod, C.; Matthews, F.; Brayne, C.; Woods, B.; CFAS-Wales Study Team. Potentially modifiable lifestyle factors, cognitive reserve, and cognitive function in later life: A cross-sectional study. PLoS Med. 2017, 14, e1002259.
11. Papageorgiou, N.; Tousoulis, D.; Androulakis, E.; Giotakis, A.; Siasos, G.; Latsios, G.; Stefanadis, C. Lifestyle factors and endothelial function. Curr. Vasc. Pharmacol. 2012, 10, 94–106.
12. Shi, L.; Morrison, J.A.; Wiecha, J.; Horton, M.; Hayman, L.L. Healthy lifestyle factors associated with reduced cardiometabolic risk. Br. J. Nutr. 2011, 105, 747–754.
13. Kent, K.; Charlton, K.E.; Netzel, M.; Fanning, K. Food-based anthocyanin intake and cognitive outcomes in human intervention trials: A systematic review. J. Hum. Nutr. Diet. 2017, 30, 260–274.

14. Ahles, S.; Stevens, S.; Bales, C.; Rizzo, J.; Maitz, M.; Mazzuca, D.; Adamo, J.; de Groot, E.; Ralston, J. The Effect of the Intervention on Cognitive Function in Older Adults. *Frontiers in Aging and Neuroscience*. 2019, 11, 247. doi:10.3389/fnagi.2019.00247.

15. Barfoot, K.L.; May, G.; Lamport, D.J.; Ricketts, J.; Riddell, P.M.; Williams, C.M. The effects of blueberry supplementation on the cognition of 7–10-year-old schoolchildren. *Eur. J. Nutr.* 2019, 58, 2911–2920.

16. Boespflug, E.L.; Eliassen, J.C.; Dudley, J.A.; Shidler, M.D.; Kalt, W.; Summer, S.S.; Stein, A.L.; Stover, A.N.; Krikorian, R. Enhanced neural activation with blueberry supplementation in mild cognitive impairment. *Nutr. Neurosci.* 2018, 21, 297–305.

17. Bowtell, J.L.; Aboobakkar, Z.; Conway, M.E.; Adlam, A.R.; Puford, J. Enhanced task-related brain activation and resting perfusion in healthy older adults after chronic blueberry supplementation. *Appl. Physiol. Nutr. Metab.* 2022, 47, 170–179.

3. The Effect of Berry Anthocyanins on Cardiometabolic Risk

18. Cook, M.D.; Sandu, B.H.A.K.; Joyce Ph, D.J. Effect of New Zealand Blackcurrant on Blood Pressure, Cognitive Function and Functional Performance in Older Adults. *J. Nutr. Gerontol. Geriatr.* 2020, 39, 99–113.

32 studies determined effects of berry anthocyanins on cardiometabolic risk markers. The interventions used were blueberry (n = 12), chokeberry (n = 7), blackcurrant (n = 6), black raspberry (n = 4), elderberry (n = 2), and bilberry (n = 1). The results were clustered into (i) BP measurements, or (ii) metabolic risk markers. The results

19. Krikorian, R.; Shidler, M.D.; Nash, T.A.; Kalt, W.; Vinqvist-Tymchuk, M.R.; Shukitt-Hale, B.;

Joseph, J.A. Blueberry Supplementation Improves Memory in Older Adults. *J. Agric. Food Chem.*

Author (Year)	Intervention	Anthocyanin Dose	Blood Pressure		Metabolic Risk Markers									
			SBP/DBP	MAP	Central SBP/DBP	24hr ABP SBP/DBP	Heart Rate	Glucose	Insulin	TC	TAG	HDL-C	LDL-C	non-HDL
Ahles (2020) [14]	Chokeberry extract	16 mg	=/	=/										
		27 mg	=/	=/										
Arevström (2019) [21]	Bilberry powder	90 mg #	=/			=			=	=	=	=	=	
Basu (2010) [22]	Freeze-dried blueberry juice	742 mg	↓/↓				=		=	=	=	=	=	
Castro-Acosta (2016) [23]	Blackcurrant extract	131 mg	=/											
		322 mg	=/											
		599 mg	=/											
Cho (2020) [24]	Black raspberry	nr	=/											

23. Traupe, I.; Giacalone, M.; Agrimi, J.; Baroncini, M.; Pomé, A.; Fabiani, D.; Danti, S.; Timpano Sportiello, M.R.; Di Sacco, F.; Lionetti, V.; et al. Postoperative cognitive dysfunction and short-term neuroprotection from blueberries: A pilot study. *Minerva Anestesiol.* 2018, 84, 1352–1360.

24. Watson, A.W.; Okello, E.J.; Brooker, H.J.; Lester, S.; McDougall, G.J.; Wesnes, K.A. The impact of blackcurrant juice on attention, mood and brain wave spectral activity in young healthy volunteers. *Nutr. Neurosci.* 2019, 22, 596–606.

Author (Year)	Intervention	Anthocyanin Dose	Blood Pressure		Metabolic Risk Markers							memory
			mmHg ↓	mmHg ↓	mmHg ↓	mmHg ↓	mmHg ↓	mmHg ↓	mmHg ↓	mmHg ↓	mmHg ↓	memory
2	Cook (2017) [35]	New Zealand blackcurrant extract	210 mg	=/=	=	=	=	=	=	=	=	an
			105 mg	=/=	=	=	=	=	=	=	=	en.
2	Cook (2017) [36]	New Zealand blackcurrant extract	210 mg	=/=	↓	=	=	=	=	=	=	an
			315 mg	=/=	↓?	=	=	=	=	=	=	en.
2	Cook (2020) [38]	New Zealand blackcurrant extract	210 mg	↓/↓								cebo-
												ry
2	Curtis (2009) [37]	Elderberry extract	500 mg	=/=		=	=	=	=	=	=	cebo-
												ry
2	Curtis (2019) [38]	Freeze-dried blueberry powder	182 mg	=/=		=	=	=	=	=	=	cebo-
												ry
2	Del Bó (2013) [39]	Blueberry jello	348 mg	=/=								reman,
												a
2	Del Bó (2017) [40]	Blueberry juice	309 mg	=/=		=						
3	Istas (2019) [41]	Chokeberry extract and whole fruit	3.6 mg	=/=	=/=	=	=	=	=	=	=	ild
			30 mg	=/=	=/=	=	=	=	=	=	=	task
3	Jeong (2014) [42]	Black raspberry extract	nr					↓	=	=	=	
3	Jeong (2016) [43]	Black raspberry extract	nr (low dose)	=/=	=/-	=/=						
			nr (high dose)	=/=	=/-	=/-						
3	Jeong (2016) [44]	Black raspberry extract	nr	=/=	=/-	=/-	=					

randomized clinical trial. *Nutr. Res.* 2019, 62, 13–22.

- Basu, A.; Du, M.; Leyva, M.J.; Sanchez, K.; Betts, N.M.; Wu, M.; Aston, C.E.; Lyons, T.J. Blueberries decrease cardiovascular risk factors in obese men and women with metabolic syndrome. *J. Nutr.* 2010, 140, 1582–1587.
- Castro-Acosta, M.L.; Smith, L.; Miller, R.J.; McCarthy, D.I.; Farrimond, J.A.; Hall, W.L. Drinks containing anthocyanin-rich blackcurrant extract decrease postprandial blood glucose, insulin and incretin concentrations. *J. Nutr. Biochem.* 2016, 38, 154–161.
- Cho, J.M.; Chae, J.; Jeong, S.R.; Moon, M.J.; Ha, K.-C.; Kim, S.; Lee, J.H. The cholesterol-lowering effect of unripe *Rubus coreanus* is associated with decreased oxidized LDL and apolipoprotein B levels in subjects with borderline-high cholesterol levels: A randomized controlled trial. *Lipids Health Dis.* 2020, 19, 166.
- Cook, M.D.; Myers, S.D.; Gault, M.L.; Willems, M.E.T. Blackcurrant Alters Physiological Responses and Femoral Artery Diameter during Sustained Isometric Contraction. *Nutrients* 2017, 9, 556.

Author (Year)	Intervention	Anthocyanin Dose	Blood Pressure		Metabolic Risk Markers						n	Response
			↓/↓	=	=	=	=	=	=	=		
Johnson (2015) [45]	Freeze-dried blueberry powder	103 mg [#]	↓/↓	=	=	=	=	=	=	=	10	Significant reduction in systolic blood pressure (10 mmHg) and diastolic blood pressure (4 mmHg) compared to placebo.
Khan (2014) [46]	Blackcurrant juice	10 mg	=/=			=	=	=	=	=	10	Significant reduction in systolic blood pressure (10 mmHg) and diastolic blood pressure (4 mmHg) compared to placebo.
Loo (2016) [47]	Chokeberry juice and powder	35.75 mg	=/=		=/↓?	=	=	=	=	=	10	Significant reduction in systolic blood pressure (10 mmHg) and diastolic blood pressure (4 mmHg) compared to placebo.
McAnulty (2014) [48]	Blueberry powder	nr	↓/↓								10	Significant reduction in systolic blood pressure (10 mmHg) and diastolic blood pressure (4 mmHg) compared to placebo.
McAnulty (2019) [49]	Freeze-dried blueberry powder	nr	↓/↓								10	Significant reduction in systolic blood pressure (10 mmHg) and diastolic blood pressure (4 mmHg) compared to placebo.
Murkovic (2004) [50]	Elderberry juice	40 mg				=	=	=	=	=	10	Significant reduction in systolic blood pressure (10 mmHg) and diastolic blood pressure (4 mmHg) compared to placebo.
Naruszewicz (2007) [51]	Chokeberry extract	64 mg [#]	↓/↓			=	=	=	=	=	10	Significant reduction in systolic blood pressure (10 mmHg) and diastolic blood pressure (4 mmHg) compared to placebo.
Okamoto (2020) [52]	New Zealand blackcurrant extract	210 mg	↓/↓?	↓	=/↓	=	=	=	=	=	10	Significant reduction in systolic blood pressure (10 mmHg) and diastolic blood pressure (4 mmHg) compared to placebo.
Petrovic (2016) [53]	Chokeberry juice	nr				=	=	=	=	=	10	Significant reduction in systolic blood pressure (10 mmHg) and diastolic blood pressure (4 mmHg) compared to placebo.
Pokimica (2019) [54]	Chokeberry juice	28.3 mg	=/=			=	=	=	=	=	10	Significant reduction in systolic blood pressure (10 mmHg) and diastolic blood pressure (4 mmHg) compared to placebo.
Riso (2013) [55]	Freeze-dried blueberry powder	113.3 mg	=/=			=	=	=	=	=	10	Significant reduction in systolic blood pressure (10 mmHg) and diastolic blood pressure (4 mmHg) compared to placebo.
Rodriguez-Mateos (2013) [56]	Freeze-dried blueberry powder	310 mg	=/=		=	=	=	=	=	=	10	Significant reduction in systolic blood pressure (10 mmHg) and diastolic blood pressure (4 mmHg) compared to placebo.
		517 mg	=/=		=	=	=	=	=	=	10	Significant reduction in systolic blood pressure (10 mmHg) and diastolic blood pressure (4 mmHg) compared to placebo.
		724 mg	=/=		=	=	=	=	=	=	10	Significant reduction in systolic blood pressure (10 mmHg) and diastolic blood pressure (4 mmHg) compared to placebo.

110, 316–329.

42. Jeong, H.S.; Hong, S.J.; Lee, T.B.; Kwon, J.W.; Jeong, J.T.; Joo, H.J.; Park, J.H.; Ahn, C.M.; Yu, C.W.; Lim, D.S. Effects of black raspberry on lipid profiles and vascular endothelial function in patients with metabolic syndrome. *Phytother. Res.* **2014**, *28*, 1492–1498.

43. Jeong, H.S.; Hong, S.J.; Cho, J.Y.; Lee, T.B.; Kwon, J.W.; Joo, H.J.; Park, J.H.; Yu, C.W.; Lim, D.S. Effects of Rubus occidentalis extract on blood pressure in patients with prehypertension: Randomized, double-blinded, placebo-controlled clinical trial. *Nutrition* **2016**, *32*, 461–467.

44. Jeong, H.S.; Kim, S.; Hong, S.J.; Choi, S.C.; Choi, J.H.; Kim, J.H.; Park, C.Y.; Cho, J.Y.; Lee, T.B.; Kwon, J.W.; et al. Black Raspberry Extract Increased Circulating Endothelial Progenitor Cells and Improved Arterial Stiffness in Patients with Metabolic Syndrome: A Randomized Controlled Trial. *J. Med. Food* **2016**, *19*, 346–352.

45. Johnson, S.A.; Figueiroa, A.; Navaei, N.; Wong, A.; Kalfon, R.; Ormsbee, L.T.; Feresin, R.G.; Elam, M.L.; Hooshmand, S.; Payton, M.E.; et al. Daily blueberry consumption improves blood pressure and arterial stiffness in postmenopausal women with pre- and stage 1-hypertension: A

Author (Year)	Intervention	Anthocyanin Dose	Blood Pressure	Metabolic Risk Markers							59—
4 Stull (2010) [57]	Freeze-dried blueberry powder	668 mg	=/=	=	=	=	=	=	=	=	J.
4 Stull (2015) [58]	Freeze-dried blueberry powder	290.3 mg	=/=	=	=	=	=	=	=	=	ly low rich
4 Whyte (2018) [28]	Wild blueberry powder and extract	1.35 mg 2.7 mg 7 mg	=/= =/= =/=								
Xie (2017) [59]	Chokeberry extract	45.1 mg	=/=								d pressure and reduced low-grade inflammation in patients with mildly elevated blood pressure. <i>Nutr. Res.</i> 2016, 36, 1222–1230.

48. McAnulty, L.S.; Collier, S.R.; Landram, M.J.; Whittaker, D.S.; Isaacs, S.E.; Klemka, J.M.; Cheek, S.L.; Arms, J.C.; McAnulty, S.R. Six weeks daily ingestion of whole blueberry powder increases natural killer cell counts and reduces arterial stiffness in sedentary males and females. *Nutr. Res.* 2014, 34, 577–584.

49. McAnulty, L.; Collier, S.; Pike, J.; Thompsonand, K.; McAnulty, S. Time course of blueberry ^{↑ or ↓ or =} indicates statistically significant higher or lower values or no significant change in the intervention group ingestion on measures of arterial stiffness and blood pressure. *J. Berry Res.* 2019, 9, 1–10. compared to control. ? indicates a trend. # indicates that the value was calculated; ¹ indicates that the dosage was dependent on body weight. Abbreviations: ABP: ambulatory blood pressure; ApoA1: apolipoprotein A1; ApoB: apolipoprotein B; DBP: diastolic blood pressure; HDL-₉: high-density lipoprotein cholesterol; HR: heart rate; LDL-C: low-density lipoprotein cholesterol; MAP: mean arterial pressure; n/r: not reported; SBP: systolic blood pressure; TAG: triacylglycerol; TC: total cholesterol. *Eur. J. Clin. Nutr.* 2004, 58, 244–249.

51. Naruszewicz, M.; Laniewska, I.; Millo, B.; Dłuzniewski, M. Combination therapy of statin with **4. Discussion** flavonoids extract from chokeberry fruits enhanced reduction in cardiovascular risk markers in patients after myocardial infarction (MI). *Atherosclerosis* 2007, 194, e179–e184.

This systematic review summarized the effects of berry anthocyanins on cognitive performance, vascular function, 52. Okamoto, T.; Hashimoto, Y.; Kobayashi, R.; Nakazato, K.; Willems, M.E.T. Effects of blackcurrant and cardiometabolic risk markers. Significant improvements were primarily observed on memory, while some of the extract on arterial functions in older adults: A randomized, double-blind, placebo-controlled, studies also reported effects on attention and psychomotor speed or executive function. Vascular function markers crossover trial. *Clin. Exp. Hypertens.* 2020, 42, 640–647. were also affected, and it can be concluded that berry anthocyanins predominantly improved vascular endothelial 53. Petrović, S.; Aracic, A.M.; Glibetić, M.; Čakir, N.; Jakšić, V.; Vučić, V. The effects of polyphenols on vascular measures in young and middle-aged adults: A meta-analysis. *Front. Nutr.* 2016, 4, 1058–1063. but not chokeberry juice on fatty acid profiles and lipid peroxidation in healthy and older players: Results from a randomized, double-blind, placebo-controlled study. *Can. J. Physiol. Pharmacol.*

Most of the included studies evaluating effects on cognitive performance involved either a young healthy population or older adults (with an increased risk of cognitive decline). Studies measuring cognitive performance in children 54. Pokimica, B.; García-Conesa, M.T.; Zec, M.; Debeljak-Martačić, J.; Ranković, S.; Vidović, N.; mostly focused on executive function, while studies in older adults primarily focused on memory tests. For studies Petrović-Oggiano, G.; Konić-Ristić, A.; Glibetić, M. Chokeberry Juice Containing Polyphenols in young and middle-aged adults, no specific preference for a specific domain was observed. Regarding memory Does Not Affect Cholesterol or Blood Pressure but Modifies the Composition of Plasma outcomes, limited evidence was available for children and adults. Most evidence comes from studies involving Phospholipids Fatty Acids in Individuals at Cardiovascular Risk. *Nutrients* 2019, 11, 850. older adults, which reported improved memory scores after supplementation with berry anthocyanins. The effect on memory was most evident among studies that evaluated individuals with (subjective) MCI. This could be attributed

55. Resa, P.; Kline, Z.; Zanchi, D.; De Leo, D.; Cimatti, D.; Giampoldi, J.; Vendrame, S.; Hoffer, B.; Buhle, S., et al. **Effect of blueberry (Vaccinium angustifolium) dietary intervention on markers of oxidative stress, inflammation, and endothelial function in humans with baseline vascular risk factors.** *Front. Nutr.* **2018**, *5*, 49–61.

56. Rodriguez-Mateos, A.; Rendeiro, C.; Bergillos-Meca, T.; Tabatabaei, S.; George, T.W.; Heiss, C.; Spencer, J.P. **Intake and time dependence of blueberry flavonoid-induced improvements in speed tests were primarily carried out in young/middle aged adults. Interestingly, all studies involving an adult population observed significant improvements as a result of supplementation, while all four studies using older adults (with cognitive decline) did not. Similarly, an improved executive function was observed in most studies involving children, but was less evident in older adults. These results suggest that the attention and psychomotor speed and executive function domains are better targets for improvement in younger populations. Previously, it has been shown that older adults require more time to finish attention tasks, but are able to maintain similar concentration as compared to younger adults.** *Am. J. Clin. Nutr.* **2013**, *98*, 1179–1191.

57. Still, A.J.; Cash, K.C.; Johnson, W.D.; Champagne, C.M.; Cefalu, W.T. **Bioprotectives in blueberries improve insulin sensitivity in obese, insulin-resistant men and women.** *J. Nutr.* **2010**, *140*, 1764–1768.

58. Still, A.J.; Cash, K.C.; Champagne, C.M.; Gupta, A.K.; Boston, R.; Beyl, R.A.; Johnson, W.D.; Cefalu, W.T. **Blueberries Improve endothelial function, but not blood pressure, in adults with metabolic syndrome: A randomized, double-blind, placebo-controlled clinical trial.** *Nutrients* **2015**, *7*, 4107–4123.

59. Xie, L.; Vance, T.; Kim, B.; Lee, S.G.; Caceres, C.; Wang, Y.; Hubert, P.A.; Lee, J.Y.; Chun, O.K. **Next to the study population, the duration of the intervention and dose of anthocyanins may also play a role. Beneficial effects on cognitive performance were observed both in acute and longer-term studies. In fact, improved cognitive performance was reported for all three domains in both acute and longer-term studies. For attention and executive function tests, results appear to be stronger in case an acute intervention period was used, while memory outcomes were affected more by longer-term studies. This suggests that the ideal study duration is dependent on the selected cognitive domain.** *Nutr. Res.* **2017**, *37*, 67–77.

60. Root-Bernstein, R. **Brain Aging: Models, Methods, and Mechanisms.** *JAMA* **2007**, *298*, 2798–2799.

61. Erickson, K.I.; Hillman, C.; Stillman, C.M.; Ballard, R.M.; Bloodgood, B.; Conroy, D.E.; Macko, R.; Marquez, D.X.; Petruzzello, S.J.; Powell, K.F.; et al. **Physical Activity, Cognition, and Brain Outcomes: A Review of the 2018 Physical Activity Guidelines.** *Med. Sci. Sports Exerc.* **2019**, *51*, 1242–1251.

62. Tousoulis, D.; Antoniades, C.; Stefanidis, C. **Evaluating endothelial function in humans: A guide to invasive and non-invasive techniques.** *Heart* **2005**, *91*, 553–558.

63. Vendemiale, G.; Romano, A.D.; Dagostino, M.; de Matthaeis, A.; Serviddio, G. **Endothelial extracts and juices, with blueberries as the main source. All three compositions had the strongest results on the dysfunction associated with mild cognitive impairment in elderly population.** *Aging Clin. Exp. Res.* **2013**, *25*, 247–255.

64. Csipo, P.; Liprandi, A.; Fabbri, O.; Apollonio, P.; Pala, P.; Bonelli, D.; Zanetti, M.; Tarantini, S.; Balasubramanian, P.; Kiss, T.; Yabluchanska, V.; et al. **Age-related decline in peripheral vascular health predicts cognitive impairment.** *Geroscience* **2019**, *41*, 125–136.

65. Naiberg, M.R.; Newton, D.F.; Goldstein, B.I. **Flow-Mediated Dilation and Neurocognition: Systematic Review and Future Directions.** *Psychosom. Med.* **2016**, *78*, 192–207.

66. Nabiegh, P.M.; El-Kadri, M.; Rund, S.; Peller, M.; Shabani, Y.; Sivaprasadarao, M.; and Prasad, V. **2019. Vascular Velocity Theory, Methods, Advances and Clinical Applications.** *IEEE Rev. Biomed. Eng.* 2020, **13**, risk. Except for 112 one study in smokers, all studies reported an improved FMD. In a recent cross-sectional study, a significant association between FMD and MCI was reported in healthy older adults and older adults with MCI [63].

67. Nassour, R.; Ayash, A.; Al-tameemi, K. **Anthocyanin pigments: Structure and biological importance.** *J. Chem. Pharm. Sci.* 2020, **13**, 45–57.

68. Kay, C.D.; Mazza, G.; Holub, B.J.; Wang, J. **Anthocyanin metabolites in human urine and serum function and working memory with FMD.** These results indicate that the effects on memory observed in this review, could potentially be the result of an improved vascular function, with endothelial function as measured by FMD as an important factor. In agreement, the RHI, another measure of endothelial function measuring the reperfusion of limbs, was also improved in adults at cardiometabolic risk, but not in healthy subjects. Besides the effects on markers of endothelial function, some of the studies also focused on arterial stiffness. In fact, for Alx, no significant effects were reported in the studies involving a healthy adult population. However, half of the studies performed in an adult population at cardiometabolic risk, and a single study in healthy older adults, observed an improved Alx.

70. Recharla, N.; Riaz, M.; Ko, S.; Park, S. **Novel technologies to enhance solubility of food-derived bioactive compounds: A review.** *J. Funct. Foods* 2017, **39**, 63–73.

Only a limited amount of the included studies performed cfPWV measurements in (older) adults. No effect was observed in the healthy adult population while cfPWV was improved in adults at cardiometabolic risk and healthy older adults. However, it should be considered that the study duration was only 24 h for the healthy adult population, which is too short to induce structural changes in artery walls that are addressed with cfPWV [66].

BP was lowered in several studies that included an adult population at cardiometabolic risk (e.g. (pre)hypertension, MetS, obesity). Studies that were carried out with an older adult population (healthy or subjective MCI) all reported beneficial effects on BP. In contrast, studies evaluating BP effects in healthy adults did not observe any changes. This is in line with our earlier findings on cognitive performance, suggesting that dietary anthocyanins have the most pronounced effects in populations with increased cardiometabolic risk, allowing for improvement by the intervention.

Considering the effects of berry anthocyanins on vascular function and cardiometabolic risk profiles as summarized in this review, the effect of intervention composition (i.e., powder, extract, or juice) and study duration is less clear as compared to the observations for the cognitive domain. Six out of fourteen studies using an extract found significant improvements in BP as compared with three out of ten studies using a powder, and one out of six juice intervention studies. Similarly, all four studies using a powder as intervention reported increased FMD, compared to two out of three studies using an extract. This pattern is also similar for the other parameters. All studies observing a beneficial effect on BP had an intervention period of one to twenty-four weeks, while all four acute studies (<24 h) did not report any significant changes, suggesting that a longer intervention is probably needed to induce effects on BP following the intake of anthocyanins. Taken together, these data suggest that the effect of berry anthocyanins on vascular function and cardiometabolic profiles is not only dependent on the population receiving the intervention, but may also be related to other factors, such as the method of administration, and the duration of intervention. Furthermore, physical activity could affect it.

A limitation of this review study is that the exact composition of the interventions was not always reported. Moreover, the bioactivity of anthocyanins is known to be dependent on their chemical structure [67]. Even though

we were able to report the amount of anthocyanin for most of the studies, more specific components such as anthocyanin subgroups (e.g. cyanidins, delphinidins, malvidins) were often not mentioned. Therefore, it was not possible to compare effects of these anthocyanin subgroups. Besides these subgroups, specific biological effects could also be the result of anthocyanin metabolism [68][69]. Furthermore, only a limited amount of studies using an extract provided information on the method of extraction, which could influence the bioactivity of the anthocyanins [70]. Consequently, we recommend future studies to report information on the chemical composition and extraction methods of the study products.

In conclusion, this systematic review provides evidence for the beneficial effects of berry anthocyanins on cognitive performance as memory was improved. Vascular endothelial function, as measured by FMD and BP were also affected, and these effects may underlie the observed effects on memory. Future studies should focus on exploring a potential causal link between the beneficial effects on cognitive performance and improvement in vascular function and cardiometabolic risk markers.