

# The Hikikomori Phenomenon

Subjects: Sociology

Contributor: Vincenzo Esposito, Felice Addeo, Valentina D'Auria, Francesca Romana Lenzi

The Hikikomori phenomenon can be classified with the classification of “social pathology”: the Hikikomori phenomenon, and its spread in society, appear to be a real danger to the sustainability and resilience of the very society in which it occurs. This is because the social isolation of an individual, especially if young and non-independent, impacts the community of reference in human, economic and psychological terms.

Keywords: Hikikomori ; mixed method ; sustainability

---

## 1. Introduction

Hikikomori is a Japanese term formed from two words “Hiku” (to pull) and “Komuru” (to withdraw), which means “to stand aside, to withdraw” <sup>[1][2]</sup>. This term is first used with scientific value, to refer to both the syndrome and the sufferer, by psychiatrist Tamaki Saito in 1998 in his work “Endless Adolescence”. In his book, the author takes a radical stance by pointing to Hikikomori as a new form of isolation and a new psychopathology that cannot be superimposed on any other known ones.

In 2003, the Japanese government, through the Minister of Health, designated Hikikomori syndrome as new psychopathology and identified its characteristics: (1) continued isolation for 6 or more months; (2) motivations for isolation, dictated by the desire to escape the pressures of social fulfillment; and (3) distrust of relationships, the individual who decides to isolate himself develops a strong distrust of interpersonal relationships, society and its dynamics.

In the first research (medical), Hikikomori was confused with depression or schizophrenia and treated with psychotropic drugs: most often with poor and inconclusive results <sup>[2][3][4]</sup>. After identifying Hikikomori as a new psychopathology and creating ad hoc therapies for its treatment <sup>[1][4]</sup>, today there is a tendency to confuse this syndrome with two other disorders, Internet Addiction and Gaming Disorder. While in these cases the isolation is due to an addiction to the Internet or video games, in the case of a Hikikomori the isolation is due to social malaise <sup>[2][5]</sup>. According to Ricci and Pierdominici, only 30 percent of individuals who self-isolate are offline and it is estimated that only one-tenth of the total number of individuals make heavy use of the Internet <sup>[6][7]</sup>.

Hikikomori syndrome has been classified in the 2019 version of the DSM as a cultural syndrome: this type of label is given only when a certain pathology affects only a state or a single population. In the 2022 updated version of the DSM, Hikikomori syndrome is included in the appendix of psychopathologies ready to become an integral and fixed part of the volume.

## 2. The Hikikomori Phenomenon

In the 1970s and 1980s in Japan, for the first time, several studies identified symptoms of “withdrawal neurosis” <sup>[2][8]</sup> in situations where patients decided to withdraw from their everyday contexts, such as school or work. Early cases of withdrawal (occurring mostly among young people) were often confused with various existing psychopathologies, such as schizophrenia or depression. Due to the increasing attention paid to these symptoms by various scholars, more in-depth studies of the disorder began to be conducted, eventually recognizing it as a real syndrome different from all others already known <sup>[5][8][9]</sup>.

In the 1990s, these symptoms were traced to the syndrome that is now known and identified as Hikikomori <sup>[2][3][5][10][11]</sup>. Since voluntary isolation is also a prodromal symptom of schizophrenia, in early diagnoses, Hikikomori syndrome was treated with the drug and nursing therapies provided for schizophrenic patients <sup>[2][5]</sup>. However, in Hikikomori, all of the symptomatology that greatly alters the perception of reality present in schizophrenics is absent. Far from being considered persons incapable of understanding, those who become Hikikomori first and foremost choose it; that is, they decide it with lucidity <sup>[12][13]</sup>.

According to some authors [8][10][13][14], Hikikomori can be classified as a true state of anomie, a term coined by Durkheim [15] that means “absence of norms”. However, their loss of normative references is not related to the general context, but only to those aspects related to social dynamics, personal success and one's public image. It would thus be a form of narrow anomie, as other norms, such as moral ones, are maintained.

One of the first general characteristics that emerge from early studies on Hikikomori syndrome is the slow course of the pathology, which does not manifest itself suddenly and blindly, but as a long process. Crepaldi [16] tried to identify the stages that a Hikikomori goes through before total isolation. What can be inferred is that the isolation resulting from Hikikomori syndrome is not a choice dictated by a bad personal moment. It is a reasoned choice that the person matures over time and in full clarity. In the first stage, the person begins to feel the urge to isolate himself or herself, finds relief in isolation and feels uncomfortable with other people; in the middle stage, the person begins to consciously process the impulse to isolate, quits school or work, reverses the sleep-wake cycle [3][17] and begins to spend most of their time in their own home [16].

Among the main reasons why young people become Hikikomori is the pressure for social achievement and the race for perfectionism [2][5]. In young people, the onset of Hikikomori disorder often coincides with the end of school age (age 19) and the beginning of adulthood, when important decisions about one's future must be made. At this delicate stage, those who have been termed “emerging adults” [13][18], ages 18–29, must build their future and identity. In high-income societies, young people are often under constant stress caused by the competitive climate at both school and work. In this regard, other reasons related to the sociocultural context of young people in self-isolation have been highlighted: frequent incidents of bullying suffered and bad relationships with peers [10][19]; the pressure exerted by schools and inattention of educational institutions toward young people who are Hikikomori or prone to become Hikikomori [4][8][20]; and an overly protective and comforting family context [2][4][5][20].

Japan is considered the home of the Hikikomori phenomenon [21][22], and to date it remains the country with the largest number of isolated children. Data dating back to 2011, reported in a study by Takasu et al. [17], showed that about 200,000 people in Japan can be classified as Hikikomori (out of a population of about 120 million). One of the reasons that could be attributed to the phenomenon is certain Japanese cultural traits. From a cultural point of view, in Japan great importance is placed on “sekentei” (世間体), which means “appearance in the eyes of others” and reflects social expectations that the individual cannot disappoint. Therefore, dependence and conformity in Japanese culture are part of cultural values that can guide and condition each individual's being in society. According to several authors [4][6][13][20][21][23], the boys who are most likely to become Hikikomori are those who prefer to spend a lot of time alone, have an economically well-off family, perform well in school, and are often bullied. Along with these “personal” factors, there are some contextual factors that facilitate withdrawal; school and/or university pressure, fear of facing new realities (many Hikikomori decide to isolate themselves at the end of a school career, ready to start a career path), and finally Japanese work environment, which is very competitive.

In 2014, Fansten et al. [24] hypothesized the existence of different blocking modes in the socialization process of Hikikomori. Their study proposed the construction of a typology to define the different types of Hikikomori, later revised by Crepaldi [16]. Currently, from the typological classifications developed by these scholars, four types of Hikikomori can be distinguished:

- **Alternative:** Those who belong to this type decide to isolate themselves because they do not accept the social dynamics of modern society. Individuals categorized in this way isolate themselves to avoid “regular” adolescence and rebel against society. Is possible that in this case, the boy before reclusion is preceded by a severe existential depression;
- **Reactionary:** Those in this group experience an many times familiar, context, already difficult one that has contributed to exacerbating an already pre-existing. Imprisonment is a reaction to their already difficult situation and is linked to an especially traumatic events;
- **Resigning:** This type of withdrawal concerns those individuals who cannot withstand the pressure arising from others' expectations. The subjects manage to alleviate their malaise by isolating themselves by abandoning social competition, whether in school and work settings;
- **Chrysalis:** The person in this typology seeks an escape from the responsibilities of future adult life in isolation. He feels that he does not have the skills to deal with the future, which is seen as a source of anxiety; therefore, every thought about it is avoided through the mechanism of avoidance. The Hikikomori is like freezing time and adopting strategies,

unconsciously or consciously, aimed at flattening one's life and trying to freeze the present. There is an alteration in the mechanisms of nourishment, eating meals in a way that is fast and irregularly.

On the other hand, Li and Wong <sup>[23]</sup> proposed a classification of the Hikikomori syndrome about whether or not the person is affected by other diseases. In this regard, the distinction between primary Hikikomori and secondary Hikikomori has been proposed. In the former case, the syndrome is not related to any pre-existing psychopathology of the person. Secondary Hikikomori, on the other hand, is when the confinement is a direct consequence of an already present problem, such as anxiety disorders or obsessive-compulsive disorders.

Over the past decade, studies on Hikikomori syndrome have spread outside Japan and around the world, allowing for a deeper understanding of a condition that is as silent as it is extensive. Major studies include those conducted in France <sup>[24]</sup>; in Italy <sup>[16]</sup>; Canada <sup>[25]</sup>; Australia and South Korea <sup>[26]</sup>; Spain <sup>[27]</sup>; Belgium <sup>[28]</sup>;

Although studies have spread to most states in industrialized society, there is still a gap that has not been closed. Although Hikikomori disorder was initially treated with depression and anxiety, this is fortunately no longer the case today. However, there is still a lack of clear diagnosis to identify ad hoc treatment for this type of patient <sup>[10]</sup>.

Some recent studies have pointed out the strong relationship between Hikikomori and the use of technological devices, especially the Internet. The development and spread of the Internet have influenced the behavior of people living ordinary lives and that of Hikikomori. While early case studies showed no link between Hikikomori and technology use habits <sup>[5]</sup>, more recent studies <sup>[23][26][29][30]</sup> show the opposite: Hikikomori spend most of their day online, mostly playing online games, developing in many cases what is now recognized as Internet Gaming Disorder (IGD) and Internet Addiction (IA). The relationship between these disorders and the pathology of Hikikomori continues to intensify, as evidenced by studies such as that of Tateno et al. <sup>[31]</sup>, which show an increase in the amount of time spent online. In recent years, the intensified coexistence of these disorders has led to confusion, resulting in the overlap between them, to the point that the pathology of Hikikomori is confused with both Internet Gaming Disorder and Internet Addiction, and conversely <sup>[26][30][32]</sup>. For Hikikomori, the Internet is an escape route since using it they can create new relationships and use that Web-mediated sociality as a substitute for the sociality lost by isolating themselves at home <sup>[31][33][34]</sup>.

---

## References

1. Koyama, N.; Miyake, Y.; Kawakami, N.; Tsuchiya, M.; Tachimori, H.; Takeshima, T. The World Mental Health Japan Survey Group. In Lifetime Prevalence, Psychiatric Comorbidity and Demographic Correlates of Hikikomori in a Community Population in Japan; Elsevier: Dublin, Ireland, 2008; p. 176.
2. Tajan, T. Hikikomori: The Japanese Cabinet Office's 2016 Survey of Acute Social Withdrawal. *Asia-Pac. J.* 2017, 15, 5017.
3. Kato, T.; Kanba, S.; Teo, A. Hikikomori experience in Japan and International relevance. *World Psychiatry* 2018, 17, 105–106.
4. Teo, A.; Chen, J.; Kubo, H.; Katsuki, R.; Sato-Kasai, M.; Shimokawa, N.; Hayakawa, K.; Umene-Nakano, W.; Aikens, J.E.; Kanba, S.; et al. Development and validation of the 25-item Hikikomori Questionnaire (HQ-25). *Psychiatry Clin. Neurosci.* 2018, 72, 780–788.
5. Kondo, N.; Sakai, M.; Kuroda, Y.; Kiyota, Y.; Kitabata, Y.; Kurosawa, M. General condition of Hikikomori (prolonged social withdrawal) in Japan: Pscyhiatric diagnosis and outcome in mental health welfare centres. *Int. J. Soc. Pscyhiatric* 2011, 59, 79–86.
6. Saito, T. *Adolescence without End*; Minnesota University Press: Minneapolis, MI, USA, 1988.
7. Ricci, C. *Hikikomori: Adolescenti in Volontaria Reclusione*; Franco Angeli: Milan, Italy, 2008.
8. Kato, T.; Hashimoto, R.; Hayakawa, K.; Kubo, H.; Watabe, M.; Teo, A.; Kanba, S. Multidimensional anatomy of modern tipe depression in Japan: A proposal for a different diagnostic approach to depression beyond the DSM-5. *Psychiatry Clin. Neurosci.* 2016, 70, 7–23.
9. Pierdominici, C. Intervista a Tamaki Saito sul fenomeno Hikikomori. *Psychomedia* 2008.
10. Teo, A.; Gaw, R. Hikikomori, a Japanese Culture-Bound Syndrome of Social Withdrawal? A Proposal for DSM-5. *J. Nerv. Ment. Dis.* 2010, 198, 444–449.
11. Suwa, M.; Suzuki, K. The phenomenon of “hikikomori” (social withdrawal) and the socio-cultural situation in Japan today. *J. Psychopathol.* 2013, 19, 191–198.

12. Kato, T.A.; Suzuki, Y.; Horie, K.; Teo, A.R.; Sakamoto, S. One month version of Hikikomori Questionnaire-25 (HQ-25M): Development and initial validation. *Psychiatry Clin. Neurosci.* 2022.
13. Lin, P.K.; Koh, A.H.; Liew, K. The relationship between Hikikomori risk factors and social withdrawal tendencies among emerging adults—An exploratory study of Hikikomori in Singapore. *Front. Psychiatry* 2022, 13, 1065304.
14. Yong, R.K.F.; Kaneko, Y. Hikikomori, a Phenomenon of Social Withdrawal and Isolation in Young Adults Marked by an Anomic Response to Coping Difficulties: A Qualitative Study Exploring Individual Experiences from First- and Second-Person Perspectives. *Open J. Prev. Med.* 2016, 06, 1–20.
15. Durkheim, É. De La Définition Des Phénomènes Religieux. In *L'Année Sociologique (1896/1897–1924/1925)*; Presses Universitaires de France: Paris, France, 1897; Volume 2, pp. 1–28. Available online: <http://www.jstor.org/stable/27880719> (accessed on 9 February 2023).
16. Crepaldi, M.H. *I Giovani Che non Escono di Casa, Psiche e Dintorni*; Alpes Italia: Roma, Italy, 2019.
17. Takasu, N.N.; Toichi, M.; Nakamura, W. Importance of regular lifestyle with daytime bright light exposure on circadian rhythm sleep–wake disorders in pervasive developmental disorders. *Jpn. Dent. Sci. Rev.* 2011, 47, 141–149.
18. Arnett, J.J.; Žukauskienė, R.; Sugimura, K. The new life stage of emerging adulthood at ages 18–29 years: Implications for mental health. *Lancet Psychiatry* 2014, 1, 569–576.
19. Sakamoto, N.; Martin, R.G.; Kumano, H.; Kuboki, T.; Al-Adawi, S. Hikikomori, is it a culture-reactive or culture-bound syndrome? Nidotherapy and a clinical vignette from Oman. *J. Psychiatry Med.* 2005, 35, 191–198.
20. Litvintsev, D. Social and Home Reclusion: A Review of Foreign Research. *Ideas Ideals* 2022, 14, 374–384.
21. Miller, A.; Toivonen, T. To Discipline or Accommodate? On the Rehabilitation of Japanese ‘Problem Youth’. *Asia-Pac. J. Jpn. Focus* 2010, 7, 3368.
22. Ogino, T. Managing Categorization and Social Withdrawal in Japan: Rehabilitation Process in a Private Support Group for Hikikomori. *Jpn. J. Sociol.* 2004, 13, 120–133.
23. Wong, P.W.C.; Li, T.M.H.; Chan, M.K.Y.; Law, F.Y.W.; Chau, M.; Cheng, C.; Fu, K.; Bacon-Shone, J.; Yip, P.S. The prevalence and correlates of severe social withdrawal (Hikikomori) in Hong Kong: A cross-sectional telephone-based survey study. *Int. J. Soc. Psychiatry* 2015, 61, 330–342.
24. Fansten, M. *Hikikomori, Ces Adolescents en Retrait*; Colin, A., Ed.; Elliot Edizioni: Paris, France, 2014.
25. Chong, S.; Chan, K. A case study of a chinese ‘hikikomorian’ in canada—Theorizing the process of hikikomorization. *Ефектология Теорія И Практика* 2012, 13, 99–114.
26. Lee, Y.S.; Lee, J.Y.; Choi, T.Y.; Choi, J.T. Home visitation program for detecting, evaluating and treating socially withdrawn youth in Korea. *Psychiatry Clin.* 2013, 67, 193–202.
27. Malagón-Amor, A.; Córcoles-Martínez, D.; Martín-López, L.M.; Pérez-Solà, V. Hikikomori in Spain: A descriptive Study. *Int. J. Psychiatry* 2015, 61, 475–483.
28. Vanlath, J.; Bart, S.; Koen, L.; Stijn, V.P.; Molly, W.S.; Steven, A. Why do the lonely stay lonely? Chronically lonely adolescent. Attribution and emotions in situations of social inclusion and exclusion. *J. Personal. Soc. Psychol.* 2015, 109, 932.
29. Estip, E.; Ethibault, A.; Chatel, A.E.; Kisely, S. Internet Addiction, Hikikomori Syndrome, and the Prodromal Phase of Psychosis. *Front. Psychiatry* 2016, 7, 6.
30. Roza, T.H.; Noronha, L.T.; Makrakis, M.A.; Spritzer, D.T.; Gadelha, A.; Kessler FH, P.; Passos, I.C. Gaming Disorder and Problematic Use of Social Media. In *Digital Mental Health: A Practitioner's Guide*; Springer International Publishing: Berlin/Heidelberg, Germany, 2023; pp. 237–253.
31. Kato, T.A.; Shinfuku, N.; Tateno, M. Internet society, internet addiction, and pathological social withdrawal: The chicken and egg dilemma for internet addiction and hikikomori. *Curr. Opin. Psychiatry* 2020, 33, 264–270.
32. Teo, A.R.; Fetters, M.D.; Stufflebam, K.; Tateno, M.; Balhara, Y.; Choi, T.Y.; Kanba, S.; Mathews, C.A.; Kato, T.A. Identification of the hikikomori syndrome of social withdrawal: Psychosocial features and treatment preferences in four countries. *Int. J. Soc. Psychiatry* 2014, 61, 64–72.
33. Unterreiner, A. Hikikomori: Une expérience de confinement, Natacha Vellut, Claude Martin, Cristina Figueiredo et Maïa Fansten (dir.), 2021, Rennes, Presses de l'École des hautes études en santé publique (EHESP), 192 pages. *Rev. Des Polit. Soc. Fam.* 2022, 1, 127–129.
34. Di Renzo, M.; D'Oria, P. Ritirata ma non troppo Il travagliato percorso terapeutico di una adolescente. *Franco Angeli. Psicobiettivo* 2022, XMLII, 109–117.

