

Gigantea in Flower Development

Subjects: Genetics & Heredity

Contributor: Julia Weiss

ANTEA (GI) is a gene involved in multiple biological functions, which have been analysed and are partially conserved in a series of mono- and dicotyledonous plant species. The identified biological functions include control over the circadian rhythm, light signalling, cold tolerance, hormone signalling and photoperiodic flowering. The latter function is a central role of GI, as it involves a multitude of pathways, both dependent and independent of the gene CONSTANS(CO), as well as on the basis of interaction with miRNA. The complexity of the gene function of GI increases due to the existence of paralogs showing changes in genome structure as well as incidences of sub- and neofunctionalization. We present an updated report of the biological function of GI, integrating late insights into its role in floral initiation, flower development and volatile flower production.

Keywords: Gene ontology ; molecular function ; cellular localisation ; biological function ; circadian clock ; flowering time ; flower development ; floral scent

1. Introduction

GIGANTEA (GI) is a plant-specific nuclear protein, identified for the first time in *Arabidopsis thaliana* as a late flowering mutant^[1]. Although six decades have passed since its discovery, its precise molecular function has not been completely elucidated. Only at the end of the XXth century was it possible to obtain information describing its chromosomal organization^[2]. The mapping identified the genomic locus on chromosome 1, consisting of 14 exons and encoding for a protein of 1,173 amino acids^{[2][3]}. The *GI* gene, which appeared early in land plants, is present in a single copy in most plants, such as *Arabidopsis* or rice^[4], while in *Solanaceae* it is found in two or three copies^[5]. Evolutionary phylogenetic analysis of the gene has shown that *GI* in *Petunia* and in general in *Solanaceae*, is grouped separately from the clade of *Brassicaceae*, *Rosaceae* and *Fabaceae*. This indicates an evolutionary departure that appears to be specific to plant families. Further gene duplications have been found in the subclades of tomato, *Nicotiana benthamiana* and *Petunia inflata* ^[5].

Recent studies carried out in *Petunia*^[6] support the hypothesis that the structural evolution of the main circadian clock genes such as *PhGI*, occurred through changes in the number of paralogues via gene duplications. Further changes have occurred in the gene structure and in the coding region, including both deletions and substitutions, depending on the species and paralog.

This review gives an overview of the molecular functions, cellular localisation and biological functions of GI paralogs, described in model plants as well as crops of agronomic importance.

2. The Cellular Localisation of Gigantea

The cellular localisation of *GI* in *Arabidopsis* was identified by constitutively expressing *GI-GFP* both transiently in protoplasts and transgenic plants. Fluorescent microscopy analysis revealed that GI protein is predominantly present in the nucleus of some cell types, forming nuclear bodies^{[7][8]}. Importantly, GI proteins lack a nuclear localisation signal. To better understand the nature of these formations, specific sub-nuclear marker-proteins of different compartments were used, directed specifically to nucleoli, spliceosomes, heterochromatin beams and Cajal bodies. GI was not localized in any of the aforementioned nuclear compartments, showing that GI does not play any role in the processes of protein degradation, pre-mRNA splicing and biogenesis of rRNA and snRNP. The evening complex gene *EARLY FLOWERING 4* (*ELF4*) was found to sequester GI from the nucleoplasm, where it binds to the *CO* promoter, to discrete nuclear bodies [8]. This process is induced under long days in *Arabidopsis*. *ELF4* protein synthesis oscillates during the day and consequently does the formation of GI nuclear bodies, peaking at night, and it is proposed that GI can then be released in the morning, avoiding the necessity for de novo GI synthesis^[8].

GI is also localized in the cytosol where it stabilizes the F-box protein ZEITLUPE (ZTL) through heterodimerization. Under blue light, GI binds to the LOV (light, oxygen or voltage sensing) domain of ZTL, thus ensuring a robust and accurate oscillation of its target proteins^[9]. It was proposed that sequestration of GI by ZTL to the cytosol, regulates the nuclear pool of GI and controls thereby the distribution of GI between nucleus and cytosol as well as its protection from degradation^[10]. Degradation in the nucleus coincides with the high accumulation of *CONSTITUTIVE PHOTOMORPHOGENIC 1* (COP1) and clock-associated protein ELF3. COP1 is an E3 ubiquitin-ligase, and *ELF3* allows COP1 to interact with GI leading to GI degradation under short day conditions. Thus, the destabilisation of GI through COP1 and ELF3 plays an important role in the modulation of circadian rhythms and regulation of flowering transition in *Arabidopsis*^[11].

3. The Molecular Functions of Gigantea

3.1. Gigantea Coordinates Light Signalling, Protein Degradation and Transcription of the Circadian Clock

The most recent clock model, based on *Arabidopsis*, describes the endogenous clock as an intricate system of negative autoregulatory feedback loops interacting with each other via transcriptional and post-translational activation and repression^[12] (Figure 1). Two MYB domain transcription factors, *CIRCADIAN CLOCK ASSOCIATED 1* (*CCA1*) and *LATE ELONGATED HYPOCOTYL* (*LHY*) form the central loop together with *PSEUDO-RESPONSE REGULATOR 1* (*PRR1*), also known as *TIMING OF CAB EXPRESSION 1* (*TOC1*). The morning loop is formed by *PSEUDO-RESPONSE REGULATORS* (*PRRs*) 9,7 and 5^[13]. The evening loop is composed of *EARLY FLOWERING 3 and 4* (*ELF3 and ELF4*) and *LUX ARHYTHMO* (*LUX*), together with *ZEITLUPE* (*ZTL*)^[14]. *ELF* genes and *ZTL* control light input signals to the clock and the ability of plants to distinguish between different day lengths^{[2][9][15]}. GI interacts with several of these clock genes, ensuring expression peaks, period length and amplitude of the different clock genes at specific times of the day. In *Arabidopsis*, *GI* transcript levels are controlled by the circadian clock with peak transcript levels at 8–10 h after dawn^[2]. In *Petunia*, a lack of robust circadian rhythmicity under continuous darkness for *PhGI1* hints to the necessity of correct light signalling for oscillation^[16].

One molecular function within the circadian oscillator complex consists of the binding of GI protein to ZEITLUPE (ZTL), a protein codified by a gene belonging to the evening complex. ZTL receives light inputs through its LIGHT, OXYGEN, VOLTAGE (LOV) domain, sensing blue-light, but it also has an F-box, targeting protein for degradation. ZTL degrades the central clock protein *TOC1*^[14]. GI is crucial for stabilizing and maintaining the oscillations of ZTL blue-light photoreceptor through direct protein protein interaction. In this connection, GI interacts with the blue light-sensitive domain of ZTL, stabilising it post-translationally under blue light^{[9][17]}. In *Arabidopsis*, the expression of *ZTL* is uniform during 24 hours, while protein levels oscillate^[18]. In contrast, transcription of *PhCHL*, the *Petunia* ortholog of *ZTL*, is characterized by a significant rise at midday^{[16][19]}. There are two paralogs of *GI* in *Petunia x hybrida*, *PhGI1* and *PhGI2*. The Silencing of *PhGI1* causes a significant prolongation of the rhythmic period of expression of *PhCHL*^[16]. This suggests that coordination of *PhCHL* by *PhGI* does not only occur at the post-translational level, but that *PhGI* also coordinates the timing of expression of *PhCHL* during the day. However, it is not clear yet if this is a direct effect or happens as a result of modified levels of proteins involved in transcription such as *PhTOC1*.

According to the theory of the repression feedback circuits, the morning elements *LHY* and *CCA1* heterodimerize and repress the expression of *TOC1* and the evening complex members as *GI*, *LUX*, *ELF3*, *ELF4*, *PRR9*, *PRR7* and *PRR5*, which in turn, when expressed, repress the expression of *CCA1* and *LHY*. In the evening, *TOC1* represses all of the previously expressed components. The molecular interaction between GI and ZTL on one side and ZTL and *TOC1* on the other side, predicts changes in the expression pattern of clock genes affected by *TOC1* protein levels. Indeed, experiments carried out by Fowler et al. (1999) in *Arabidopsis* demonstrated that under both long days (LDs) and short days (SDs), a mutation in the *GI* locus affects the *CCA1* and *LHY* gene expression. On the other hand, over-expression or mutations of *CCA1* and *LHY* disrupt the *GI* expression. In agreement with these results, the double mutant of *LHY* and *CCA1* shows an overabundance of the transcription of *GI* [7]. These results led to the conclusion that *GI* operates in a feedback loop for adjusting and maintaining the length of the clock period. The involvement of *GI* in circadian clock control was confirmed in rice, where *OsGI* controls amplitude and circadian rhythm phases of global gene expression under natural field conditions^[4].




Figure 1. Schematic representation of the clock model indicating interactions by autoregulatory feedback loops via transcriptional and post-translational activation and repression. The clock components are depicted according to their peak-time expression from left to right. Black bars indicate repression. Green arrowheads lines indicate activation of transcription. Fuchsia arrowheads lines indicate proteasomal degradation. The red curve represents the expression of GI during the day. TOC1 is protected during the day from ZTL-induced proteasome degradation thanks to the direct protein-protein interaction of GI with ZTL. During the evening, ELF3 promotes the proteasome degradation of GI through its interaction with COP1, thus promoting the ZTL-dependent TOC degradation. Boxes indicate sets of genes acting in coordination. The PRR box: PRR 5,7,9—these proteins are sequentially expressed during the day, and all repress LHY and CCA1; GI-ZTL box—protein-protein interaction leads to ZTL maturation; ELF3-ELF4-LUX complex box—protein complex peaking at dusk.

3.2. Flowering Time and Photoperiod Related Molecular Functions

In Arabidopsis, photoperiod, causing the transition from the vegetative phase to flowering, is regulated by the diurnal expression of *CONSTANS* (CO)^{[20][21]}. It has been shown that during long days (LD), light stabilises CO protein. In Arabidopsis, the expression of CO coinciding with the period of light leads to the activation of *FLOWERING LOCUS T* (FT) gene. On the other hand, under short day (SD) conditions, the peak expression of CO occurs after sunset because the CO protein is not sufficiently stabilized by light^[22]. The transcription of CO is repressed during sunrise, due to the activity of the *CYCLING DOF FACTOR 1* (CDF1) transcriptional repressor bound to the CO promoter.

GIGANTEA plays a key role in this flowering regulation pathway. Figure 2 illustrates the involvement of GI in flowering time control in Arabidopsis as well as in other model plants. Although several paralogs of GI exist in different species, i.e., maize, thus far, the existing models on flowering time control represent the function of only one of the paralogs, and further research may elucidate further functions of GI in other plants. In Arabidopsis, an enzymatic complex is formed thanks to a direct protein-protein interaction between GI and FLAVIN-BINDING, KELCH REPEAT, F BOX protein 1 (FKF1), through its FKF1 LOV (Light, Oxygen or Voltage) domain^[23]. This complex mediates the degradation of *CYCLING DOF FACTOR 1* (CDF1), a main CO repressor. This complex is strictly dependent on light, being the expression of GI under the control of the circadian clock. Therefore, towards the middle of the day, when the accumulation of GI along with FKF1 reaches the peak, the DOF-degradation complex is repressed, leading to an increase in the transcription of CO and, therefore, to the transcription of *FLOWERING LOCUS T* (FT)^{[23][24]}. This does not happen in short days in Arabidopsis, as the GI accumulation peak occurs about three hours earlier than that of FKF1. This prevents the formation of the DOF-degradation complex and the consequently reduced abundance of the CO transcript.

Another light-dependent control of flowering through GIGANTEA involves *SPINDLY* (SPY). GI inhibits SPY in a light dependant manner. At the same time, SPY represses the expression of CO and FT, as observed in *spy-4* plants, where the late-flowering phenotype of *gi-2* plants was abolished due to a partial suppression of abundance reduction in CO and FT RNA^[25]. Furthermore, SPY also affects flowering by inhibiting GA signaling, required for flower promotion^[26].

Additionally, *GI* can regulate the expression of *FT* independently of *CO*. It inhibits the *FT* transcriptional repressors such as *SHORT VEGETATIVE PHASE* (*SVP*), *TEMPRANILLO 1* (*TEM1*) and *TEMPRANILLO 2* (*TEM2*). *GI* alters their stability or neutralises their repressor effect by blocking their access to the *FT* promoter region. Likewise, *GI* binds their specific target regions on the *FT* promoter^[27], thus affecting the abundance of the *FT* transcripts.

GI regulates *FT* expression and photoperiodic flowering, independent of *CO*, via its interaction with a microRNA. The *miRNA172* inhibits the expression of the main transcriptional repressors of *FT*, *TARGET OF EAT 1* (*TOE1*) and *APETALA 2* (*AP2*). It was demonstrated that *miRNA172* processing is positively regulated in the presence of *GI* protein through an unclear molecular interaction^[28]. Growth of plants under natural conditions, i.e., with photo and thermoperiod, shows that it has a major impact in the coordination of *FT*, suggesting that *GI* may have a role in temperature and light coordination^[29].

Many *GIGANTEA* orthologs have been described in the last decades, in a wide range of plant species, from gymnosperms to mono y dicotyledon angiosperms. Many of these have similar expression patterns to those of *Arabidopsis GI*, suggesting broad conservation of the photoperiodic flowering regulation mechanisms (Figure 2)^{[30][31][32][33][34][35][36]}. Indeed, the *GI* ortholog of soybean, *GmGla*, controls flowering time by inducing the expression of the soybean florigen gene ortholog *GmFT2a*^[37]. Among three haplotypes (*H1*, *H2*, *H3*) of *GmGla*, *H1* rescues the late-flowering phenotype of *gi-2* in *Arabidopsis*, while *H2* and *H3* delay flowering in transgenic *Arabidopsis* with a wild type background. This diversification was proposed to be related to flowering time adaptation during soybean domestication^[38]. Similar to *Arabidopsis*, *GmGla* also positively regulates *gma-miR172a*, which in turn represses *Glyma03g33470*, leading to upregulation of *FT*, *AP2* and *LFY* and early flowering^[39].

Wheat is a long-day (LD) plant and one *Gla* ortholog *TaGI1*, has been described. It shares 63% identity with *AtGI* and a superior homology to *GI* from grasses such as rice and barley^[30]. Overexpression of *TaGI1* alters flowering time in wheat, resulting in early flowering both under LD and SD^[30]. In barley, an upregulation of *HvGI* observed in *HvELF3* mutants results in an early flowering phenotype^[40].

Conservation of *GI* function upon flowering time is also observed in short-day (SD) plants. The comparison of the sequences between *Arabidopsis GI* and its ortholog in *Oryza sativa GI* (*OsGI*) reveals 67% of identity as well as conservation of its nuclear localisation^{[41][42]}. In rice, a quantitative SD plant, over-expression of the *GIGANTEA* gene (*OsGI*) under LD and SD leads to a delay in flowering time with a more pronounced effect under SD [43]. Rice overexpressing *OsGI* and under LD condition, show increased expression of the *CO* ortholog *Hd1*, while the expression of *Hd3a*, the *FT* ortholog, was reduced. Thus, a reverse regulation of *Hd3a* by *Hd1* occurs in rice under LD conditions. Opposite to *Arabidopsis*, *OsGI* acts as a flowering time suppressor^{[43][44][45]}. However, under SD conditions *Hd1* promotes heading coinciding with increased transcript levels of *Hd3a*. This suggests that the regulatory network of the key flowering genes is conserved among rice and *Arabidopsis*^[44]. Likewise, in the SD and C4 plant maize, which has two circadian regulated *GI* paralogs, *Zmgi1* mutants flower earlier than non-mutant plants in LD photoperiods, which is the opposite to the late-flowering *gi* mutants from *Arabidopsis*, but not in SD photoperiods [36]. The flowering time control mechanism also involves *FT*-like floral activator gene *Centroradialis8* (*ZCN8*) and the *CONSTANS*-like flowering regulatory gene *Constans* of *Zea mays1* (*CONZ1*), which are both upregulated in the *Zmgi1* mutant, confirming the existence of the conserved network of the three key regulatory genes *Zmgi1*, *CONZ1*, *ZCN8* in maize, but gene interaction is photoperiod dependent and differs whether the species is a LD or SD plant.

Petunia x hybrida has two *GI* paralogs, *PhGI1* and *PhGI2*. Downregulation of *PhGI1* by interference RNA does not induce late flowering, but a flowering time effect of *PhGI2* cannot be ruled out^[46]. In the perennial poplar, three *GI-like* genes *PagGla*, *PagGlb* and *PagGlc* were identified and overexpressed in *Arabidopsis* wild type ecotype Columbia-0 (*Col-0*), leading to early flowering^[46]. Recent work in white lupin has identified *LaGI* as a major QTL involved in flowering time^[47]. As the single copy gene *MpGI* coordinates phase transition into reproductive development in the basal angiosperm *Marchantia polymorpha*^[48]. Indeed, loss of *MpGI* causes complete loss of long-day dependent phase transition. We can conclude that that *GI* plays a key role in flowering time in a variety of species.




Figure 2. Models of GIGANTEA dependent flowering time regulation in (a) Arabidopsis, (b) soybean, (c) rice (d) and maize under long day condition. Regulation of flowering through GI differs between LD and SD plants. In contrast to LD plant Arabidopsis, *OsGI* inhibits flowering in rice, an SD plant, because the *FT* ortholog *Hd3a* is repressed by *Hd1*, the *CO* ortholog.

3.3. Light Signalling Related Molecular Functions

Photoreceptors, such as phytochromes, phototropins, cryptochromes and UV-light photoreceptors, control light-induced plant development by the integration of light cues from the environment, such as quality, intensity and duration. Specific wavelength inputs are transformed into physiological signals, a process called photomorphogenesis^{[49][50]}. Cryptochromes and phototropins absorb mainly the blue spectrum (B, $\lambda = 400\text{--}499\text{ nm}$) and are involved in the regulation of flowering time, inhibition of hypocotyl growth and phototropism^[51]. Five members of phytochromes (Phy) exist in Arabidopsis, from PhyA to PhyE^{[42][52]}. They act as red (R, $\lambda = 660\text{ nm}$; PhyB-E) and far-red (FR, $\lambda = 730\text{ nm}$; PhyA) photoreceptors. PhyA mediates two distinct photobiological responses: The very-low-fluence response (VLFR) and the high-irradiance response (HIR)^[49]. Arabidopsis *gi*-mutants, grown under saturated R light condition, have shown an elongated hypocotyl compared to the wild type and little or no change in responsiveness to continuous FR light^[42], indicating that *GI* appears to be a positive regulator of *PhyB* signalling during seedling de-etiolation^[25]. Considering also that neither the genes nor the abundance of PhyA-B proteins are affected in *gi*-mutants, it appears that *GI* works downstream of PhyA-B, after their migration to the nucleus in response to light, where *GI* is constitutively localised. Additional data regarding *gi*-mutants grown in FR light condition revealed low VLFR levels, deficient cotyledon unfolding and a low seed germination index. These phenotypes were rescued through *GI* over-expression, demonstrating that *GI* plays a role in the PhyA signalling^[53]. The *gi*-mutant seedlings also exhibited a long hypocotyl phenotype when grown under blue light, proving to have a role also in the signalling of blue light^[50].

3.4. Hormone Signalling and Stress-Response Related Molecular Functions

One of the plant hormones, whose signalling interacts with *GI*, is gibberellins. In Arabidopsis, *GI* affects the growth of the hypocotyl through the interaction with *SPINDLY* (*SPY*), a gene involved in the regulation of gibberellin signalling. *SPY* is, in fact, a negative regulator of gibberellin signalling and an inhibitor of hypocotyl lengthening^{[25][26]}. *GI* interacts with a protein-protein interaction domain of *SPY*, consisting of 10 tetratricopeptide repeats (TPRs). Three pathways are controlled through *SPY*-*GI* interaction: Flowering, circadian cotyledon movements, and hypocotyl elongation. In the case of the flowering pathway, the reduction of *CO* and *FT* RNA abundance in *gi* plants is partially suppressed in the *spy* mutant.

GI also interacts with gibberellin signalisation through its interaction with REPRESSOR OF *ga1-3* (*RGA*), a DELLA protein. Under LD, warmer temperatures induce accumulation of *GI*, which functions as thermostabiliser of *RGA*, which leads to the attenuation of thermomorphogenesis mediated through PHYTOCHROME INTERACTING FACTOR 4 (*PIF4*). Under SD and lower *GI* levels, *RGA* is degraded through the gibberellic acid-mediated ubiquitination-proteasome pathway^{[54][55]}. An effect of warm temperatures on *GI* accumulation was concluded from observations on *GI*-deficient *gi-2* mutants, which showed hypocotyl overgrowth at 28 °C but not at 23 °C. Overgrowth at warmer temperatures was related to an increased auxin action, as both *YUCCA8* (*YUC8*), an auxin biosynthetic enzyme, and *SMALL AUXIN UPREGULATED RNA 22* (*SAUR22*), an auxin-responsive protein, were upregulated in *gi-2* mutants at warmer temperatures^[54].

The Arabidopsis mutant *abz126* bears a T-DNA insertion into the ninth exon of the *GIGANTEA* (*GI*) gene, resulting in a loss-of-function of *GI*, leading to long petioles, tall plant height, many rosette leaves and late flowering, as well as insensitivity to paclobutrazol and brassinolide, indicating an association of *GI* with brassinosteroid hormone signalling^[56].

Interaction between *GI* and stress response exists concerning tolerance to salt. Under normal conditions, *GI* interacts with the protein kinase SALT OVERLY SENSITIVE 2 (*SOS2*), thus preventing its interaction with the Na^+/H^+ anti-porter *SOS1* to promote Na^{2+} export and salt tolerance. Under saline conditions, *GI* is degraded by the proteasome. Due to *GI* degradation under salt stress, *SOS2* interacts with *SOS3* and this protein kinase complex then activates *SOS1*, salt tolerance and retardation of flowering^{[57][58]}.

Opposite to salt stress, drought stress accelerates flowering time, and *GI* was found to be a key component mediating drought response in *Arabidopsis*. One proposed mode of action predicts that *GI* may regulate chromatin accessibility and/or interfere with repressor activity at the florigen promoters in a plant stress hormone abscisic acid (ABA) dependent manner^[59].

GI also has molecular functions involved in freezing tolerance. *GI* activates the expression of *CO* and *FT*, the key floral regulators, by facilitating the degradation of a family of CDF1, which acts as transcriptional repressors. In *Arabidopsis* and tomato, in a *gi*-mutant background, increased stability and accumulation of CDF proteins and higher transcript levels of stress-responsive genes, including *COR15a*, *RD29A* and *ERD1*, were observed as well as an increased level of protection against cold stress^{[60][61]}. Therefore, proposed that, in addition to flowering, the regulation of CDFs through *GI*, influences responses to freezing temperatures^[61].

4. Biological Functions of Gigantea

The described molecular interactions of *GI* as well as its cellular location, result in a wide range of biological functions analysed in *Arabidopsis*. Part of these functions were found to be conserved in many other plant species, both monocots and dicots. Biological functions include circadian clock regulation, light signalling, flowering time control, which were already described in detail under molecular functions, as well as chlorophyll accumulation, sugar metabolism, stress tolerance, vegetative growth, flower development and floral scent emission described below.

4.1. Chlorophyll Accumulation

The analysis of different *Arabidopsis gi*-mutants hint to a *GI* function in chloroplast biogenesis and chlorophyll accumulation—factors essential for photosynthetic efficiency and therefore crop productivity. The *Arabidopsis gi-2* mutant is characterized by a reduced sensitivity to lincomycin, a chloroplast biogenesis inhibitor, and the mutant maintains high levels of photosynthetic proteins. In contrast, *GI*-overexpressing plants have variegated leaves, reduced photosynthetic protein levels and are sensitive to lincomycin^[62]. A series of *Arabidopsis* mutants in *Ler* background (*gi-3,4,5*) showed a significantly higher chlorophyll content in seedlings^[62]. RNA interference of one *GI* paralogue in *Petunia x hybrida*, *PhGI1*, leads to leaves with a greener appearance in the denser apical region coinciding with a progressive increase in chlorophyll content compared to the wild type^[63] (Figure 3a–c). These results thus point to a general role of *GI* in chlorophyll homeostasis.

4.2. Sugar and Starch Metabolism

Sugars function in plants as a source of energy and as a signal, among others, in the circadian regulation of flowering^[64]. The *Arabidopsis* circadian system was described to be sensitive to sucrose in the dark, indicating feedback between metabolism and circadian clock, and it was predicted that *GI* is required for the full response of the circadian clock to sucrose^[65]. Indeed, *GI* protein is stabilized by sucrose in the night, and this mechanism requires interaction with *ZTL*^[66].

Furthermore, the *Arabidopsis gi-3* mutant is characterized by an enhanced freezing sensitivity, and this was related to a reduction in soluble sugar content in leaves. Transcript levels of the cold-responsive gene *RD29A* and abscisic acid-responsive gene *RAB18* were not affected in this mutant, indicating a direct connection between *GI* gene function and sucrose metabolism^[67]. In contrast to this observation, field-grown rice plants carrying a null mutation in the rice homolog *OsGI* showed a significantly increased sucrose and starch content in the leaves at most time point^[4]. Increased starch content was also observed in *gi-1, 2 and 3* alleles in *Arabidopsis*^[68]. A recent proteomic analysis of interactors of *GI* has identified TREHALOSE-6-PHOSPHATASE SYNTHASE 8 (*TPS8*) as a direct interactor^[69]. This indicates a possible direct link between *GI* and sugar metabolism.

4.3. Stress Tolerance

Plants cope with environmental stresses by activating a series of specific metabolic pathways. Regarding low-temperature stress, they must avoid freezing injury that can occur in both intracellular and extracellular compartments, prevent chilling wounds and cell injuries that could cause tissue death. The response to cold temperatures implies alterations in the expression of genes, followed by increases in the levels of metabolites, including those known to have protective effects

against the damaging effects of cold stress [70]. Plants have the ability to adapt to low temperatures by increasing their freezing tolerance through the gradual exposure to low but non-freezing temperatures, a process identified as cold acclimation [71]. Plant resistance to cold appears to be organ-specific [72]. This process is characterised by complex biochemical and physiological changes, including gene expression [73][74], enzyme activities [75], lipid membrane composition [76] and leaf ultrastructure modification [77]. Chilling injuries cause the collapse of deep cell layers and disruption of antioxidant activity amongst other effects [78]. The biological function of *GI* concerning freezing tolerance results from its interaction with CDF proteins. *GI* facilitates the degradation of CDF proteins. *Arabidopsis gi*-mutants accumulate CDFs accompanied by higher transcript levels of stress-responsive genes and increased cold stress [60][61]. Additionally, enhanced freezing sensitivity in *gi*-mutants is also related to a reduction in soluble sugar content [67]. Another direct biological function of *GI* is related to salt stress. This stress leads to proteasomal degradation of *GI* [57][58]. *GI* function in salt tolerance was found to be conserved in poplar, as overexpressing *PagGIs* in wild type (WT) *Arabidopsis* induces salt-sensitivity.

4.4. Vegetative Growth

Observations on plant development in *Arabidopsis* indicate that *GI* is a negative regulator of vegetative growth as can be inferred from its name as a mutant. Loss of function of *GI* effects hypocotyl growth, but it also results in long petioles, tall plant height and many rosette leaves [68]. The latter is proposed to be related to gibberellin signalling, as SPINDLY (SPY) protein, a negative regulator of gibberellin signalling in *Arabidopsis* and an inhibitor of hypocotyl elongation, interacts with *GI* protein [25]. Similarly, downregulation of the *G11* paralog in *Petunia* by RNA interference results in bigger leaves (Figure 3a–c), an increased basal internode length, and an increased number of axillary meristems. Middle and apical internodes are reduced, indicating the existence of an acropetal gradient with opposite effects during the early stages of development and middle to late stages (Figure 3d,g). Similarly, downregulation of *PagGIs* in poplar by RNA interference leads to vigorous growth, and higher biomass [46].

4.5. Flower Development and Floral Scent Emission

A new role of *GI* in the development of reproductive organs has recently been described in *Petunia x hybrida*. Plants with loss of *G11* function, grown under LD conditions, show a reduction in the total number of floral buds and smaller flowers than wild type. Flowers show a significant reduction in the corolla diameter as well as in the tube length (Figure 3e). There is an additional floral bud at the bifurcation point, where a given shoot that terminates in a flower and a new sympodial shoot separate (Figure 3f). The aborted flowers clearly show early development of carpel and stamen tissues, indicating that the senescence process occurs after the activation of the floral organ identity genes. These results indicate that *PhG11* is a repressor of early flower senescence, a biological function that had not been described before. The significant reduction in the total number of flower buds also indicates an upstream effect related to the flower-meristem-identity genes *PETUNIA FLOWERING GENE (PFG)* and *ALF (ABERRANT LEAF AND FLOWER)* [79][80].

Among the main strategies that plants use to ensure entomophilic reproduction is the emission of floral scents in the form of volatile organic compounds (VOCs). Plant scents are mixtures of volatile lipophilic molecules of benzenoids/phenylpropanoids and fatty acid derivatives with low molecular weight and high vapor pressure at ambient temperature, synthesized in all plant organs, from roots to flowers [81][82]. The plant VOCs are classified into different classes according to their biosynthetic origins, such as: Terpenoids, benzenoids/phenylpropanoids, fatty acid derivatives, and amino acid derivatives as well as few other genus-specific species [83]. In vegetative organs, VOCs are part of the plant's defence system and are mainly synthesised in glandular trichomes [84], single specialized cells [85] or tubes [86] from which they can sprout out in case of breakage. In many Angiosperms, the quantity and composition of the VOCs can fluctuate during the day, mainly in relation to the age of the flower. This rhythmic release of scent is also closely related to the flower hormonal regulation, circadian clock, flower and plant development, nutrient availability, temperature, humidity and general environmental conditions [87][88].

In recent decades, many studies on the production and regulation processes of VOCs in plants have been carried out, but few genes have been characterized involved in the regulation of scent production. The studies conducted by Verdonk et al., [89] in *Petunia x hybrida*, have identified a R2R3 MYB-type gene named *ODORANT1 (ODO1)* which controls the synthesis of the precursor Phe in the shikimate pathway. The *Petunia x hybrida* *CHANEL (PhCHL)* gene, the ortholog of *Arabidopsis thaliana* *ZTL*, has been shown previously to play an important role in regulating both the timing and the quantity of volatile emissions [19]. Methyl benzoate is the dominant floral VOC in *Petunia hybrida*. Recent studies on *PhG11* in *Petunia x hybrida* have revealed thus far unreported functions of this pleiotropic gene concerning the control over VOC emission. *PhG11* loss of function plants exhibited a 20% reduction in the total emission of VOCs during 24 hours. The circadian volatile emission pattern in the silenced lines remained unchanged. However, an important twist in the scent profile was observed, compared to the wild type and non-transgenic siblings with changes in the relative abundance of the

trans-cinnamic acid derivatives benzyl alcohol, ethyl benzoate, benzyl benzoate and isoeugenol, indicating involvement of *GI* in the phenylalanine emission pathway, interfering in the rhythmic regulation of the VOC biosynthesis and their daily emission profile. Interestingly, a null mutation in the rice homolog *OsGI*, affects the production of several metabolites in the phenylpropanoid pathway, which was significantly increased, whereas the pool of Phe, the major chemical precursor in the pathway of phenylpropanoids, was significantly decreased^[4]. These convergent results suggest the role of *GI* in the control of Phe and phenylpropanoids in plants.

Figure 3. Phenotypic characteristics in *Petunia x hybrida* plants with loss of *GI1* function. Apical (a), median (b) and basal (c) leaves of wild type *Petunia* (left) compared to *iRNA: PhGI1* leaves (right). (d) Vegetative growth architecture of *Petunia* wild type (left) compared to the loss of *PhGI1* function (right). *Petunia* inflorescence appearance (e) of wild type (left) and *PhGI1* silenced line (right). Abortive flower (f) and (g) schematic representation of variations in internode length and number of axillary shoot meristems (indicated by the green colour) between wild type plants (left) and *iRNA: PhGI1* plants (right).

References

1. György P. Rédei; Supervital Mutants of Arabidopsis. *Genetics* **1962**, 47, 443-460, .
2. Sarah Fowler; Karen Lee; Hitoshi Onouchi; Alon Samach; Kim Richardson; Bret Morris; George Coupland; Jo Putterill; GIGANTEA: a circadian clock-controlled gene that regulates photoperiodic flowering in Arabidopsis and encodes a protein with several possible membrane-spanning domains. *The EMBO Journal* **1999**, 18, 4679-4688, [10.1093/emboj/18.17.4679](https://doi.org/10.1093/emboj/18.17.4679).
3. D. H. Park; Control of Circadian Rhythms and Photoperiodic Flowering by the Arabidopsis GIGANTEA Gene. *Science* **1999**, 285, 1579-1582, [10.1126/science.285.5433.1579](https://doi.org/10.1126/science.285.5433.1579).
4. Takeshi Izawa; Motohiro Mihara; Yuji Suzuki; Meenu Gupta; Hironori Itoh; Atsushi J. Nagano; Ritsuko Motoyama; Yuji Sawada; Masahiro Yano; Masami Yokota Hirai; et al. Os-GIGANTEA Confers Robust Diurnal Rhythms on the Global Transcriptome of Rice in the Field. *The Plant Cell* **2011**, 23, 1741-1755, [10.1105/tpc.111.083238](https://doi.org/10.1105/tpc.111.083238).
5. Aureliano Bombarely; Michel Moser; Avichai Amrad; Manzhao Bao; Laure Bapaume; Cornelius S. Barry; Mattijs Blik; Maaïke R. Boersma; Lorenzo Borghi; Rémy Bruggmann; et al. Insight into the evolution of the Solanaceae from the parental genomes of *Petunia hybrida*. *Nature Plants* **2016**, 2, 16074, [10.1038/nplants.2016.74](https://doi.org/10.1038/nplants.2016.74).
6. Marta I. Terry; Marta Carrera-Alesina; Julia Weiss; Marcos Egea-Cortines; Transcriptional Structure of *Petunia* Clock in Leaves and Petals. *Genes* **2019**, 10, 860, [10.3390/genes10110860](https://doi.org/10.3390/genes10110860).
7. Mizoguchi, T.; Wright, L.; Fujiwara, S.; Cremer, F.; Lee, K.; Onouchi, H.; Mouradov, A.; Fowler, S.; Kamada, H.; Putterill, J.; et al. Distinct Roles of GIGANTEA in Promoting Flowering and Regulating Circadian Rhythms in Arabidopsis. *Plant Cell* 2005, 17, 2255–2270.
8. Kim, Y.; Lim, J.; Yeom, M.; Kim, H.; Kim, J.; Wang, L.; Kim, W.Y.; Somers, D.E.; Nam, H.G. ELF4 regulates GIGANTEA chromatin access through subnuclear sequestration. *Cell Rep.* 2013, 3, 671–677.
9. Woe-Yeon Kim; Sumire Fujiwara; Sung-Suk Suh; Jeongsik Kim; Yumi Kim; Linqi Han; Karine David; Joanna Putterill; Hong Gil Nam; David E. Somers; et al. ZEITLUPE is a circadian photoreceptor stabilized by GIGANTEA in blue light. *Nature* **2007**, 449, 356-360, [10.1038/nature06132](https://doi.org/10.1038/nature06132).
10. Jeongsik Kim; Ruishuang Geng; Richard A. Gallenstein; David E. Somers; The F-box protein ZEITLUPE controls stability and nucleocytoplasmic partitioning of GIGANTEA.. *Development* **2013**, 140, 4060-4069, [10.1242/dev.096651](https://doi.org/10.1242/dev.096651).
11. Jae-Woong Yu; Vicente Rubio; Na-Yeoun Lee; Sulan Bai; Sun-Young Lee; Sang-Sook Kim; Lijing Liu; Yiyue Zhang; María Luisa Irigoyen; James A. Sullivan; et al. COP1 and ELF3 Control Circadian Function and Photoperiodic Flowering by Regulating GI Stability. *Molecular Cell* **2008**, 32, 617-630, [10.1016/j.molcel.2008.09.026](https://doi.org/10.1016/j.molcel.2008.09.026).
12. Alexandra Pokhilko; Aurora Piñas Fernández; Kieron D Edwards; Megan M Southern; Karen J Halliday; Andrew J Millar; The clock gene circuit in Arabidopsis includes a repressilator with additional feedback loops. *Molecular Systems Biology* **2012**, 8, 574-574, [10.1038/msb.2012.6](https://doi.org/10.1038/msb.2012.6).
13. Sally Adams; Ian Manfield; Peter Stockley; Isabelle A. Carré; Revised Morning Loops of the Arabidopsis Circadian Clock Based on Analyses of Direct Regulatory Interactions. *PLOS ONE* **2015**, 10, e0143943, [10.1371/journal.pone.0143943](https://doi.org/10.1371/journal.pone.0143943).
14. Paloma Más; Woe-Yeon Kim; David E. Somers; Steve A. Kay; Targeted degradation of TOC1 by ZTL modulates circadian function in Arabidopsis thaliana. *Nature* **2003**, 426, 567-570, [10.1038/nature02163](https://doi.org/10.1038/nature02163).

15. Mark R. Doyle; Seth J. Davis; Ruth M. Bastow; Harriet G. McWatters; László Kozma-Bognár; Ferenc Nagy; Andrew J Millar; Richard M. Amasino; The ELF4 gene controls circadian rhythms and flowering time in *Arabidopsis thaliana*. *Nature* **2002**, 419, 74-77, [10.1038/nature00954](https://doi.org/10.1038/nature00954).
16. Claudio Brandoli; César Petri; Marcos Egea-Cortines; Julia Weiss; The clock gene *Gigantea 1* from *Petunia hybrida* coordinates vegetative growth and inflorescence architecture. *Scientific Reports* **2020**, 10, 1-17, [10.1038/s41598-019-57145-9](https://doi.org/10.1038/s41598-019-57145-9).
17. Joon-Yung Cha; Jeongsik Kim; Tae-Sung Kim; Qingning Zeng; Lei Wang; Sang Yeol Lee; Woe-Yeon Kim; David E. Somers; GIGANTEA is a co-chaperone which facilitates maturation of ZEITLUPE in the *Arabidopsis* circadian clock. *Nature Communications* **2017**, 8, 1-12, [10.1038/s41467-016-0014-9](https://doi.org/10.1038/s41467-016-0014-9).
18. David E Somers; Thomas F Schultz; Maureen Milnamow; Steve A Kay; ZEITLUPE Encodes a Novel Clock-Associated PAS Protein from *Arabidopsis*. *Cell* **2000**, 101, 319-329, [10.1016/s0092-8674\(00\)80841-7](https://doi.org/10.1016/s0092-8674(00)80841-7).
19. Marta I. Terry; Fernando Pérez-Sanz; María Victoria Díaz-Galián; Felipe Pérez De Los Cobos; Pedro Navarro; Marcos Egea-Cortines; Julia Weiss; The *Petunia* CHANEL Gene is a ZEITLUPE Ortholog Coordinating Growth and Scent Profiles.. *Cells* **2019**, 8, 343, [10.3390/cells8040343](https://doi.org/10.3390/cells8040343).
20. Suárez-López, P.; Wheatley, K.; Robson, F.; Onouchi, H.; Valverde, F.; Coupland, G. CONSTANS mediates between the circadian clock and the control of flowering in *Arabidopsis*. *Nature* **2001**, 410, 1116–1120.
21. Imaizumi, T.; Kay, S.A. Photoperiodic control of flowering: Not only by coincidence. *Trends Plant Sci.* **2006**, 11, 550–558.
22. Federico Valverde; Aidyn Mouradov; Wim Soppe; Dean Ravenscroft; Alon Samach; George Coupland; Photoreceptor Regulation of CONSTANS Protein in Photoperiodic Flowering. *Science* **2004**, 303, 1003-1006, [10.1126/science.1091761](https://doi.org/10.1126/science.1091761).
23. Mariko Sawa; Dmitri A. Nusinow; Steve A. Kay; Takato Imaizumi; FKF1 and GIGANTEA Complex Formation Is Required for Day-Length Measurement in *Arabidopsis*. *Science* **2007**, 318, 261-265, [10.1126/science.1146994](https://doi.org/10.1126/science.1146994).
24. Takato Imaizumi; Hien G. Tran; Trevor E. Swartz; Winslow R. Briggs; Steve A. Kay; FKF1 is essential for photoperiodic-specific light signalling in *Arabidopsis*. *Nature* **2003**, 426, 302-306, [10.1038/nature02090](https://doi.org/10.1038/nature02090).
25. Tong-Seung Tseng; Patrice A. Salomé; C. Robertson McClung; Neil Olszewski; SPINDLY and GIGANTEA Interact and Act in *Arabidopsis thaliana* Pathways Involved in Light Responses, Flowering, and Rhythms in Cotyledon Movements. *The Plant Cell* **2004**, 16, 1550-1563, [10.1105/tpc.019224](https://doi.org/10.1105/tpc.019224).
26. Stephen M. Swain; Tong-Seung Tseng; Neil E. Olszewski; Altered Expression of SPINDLY Affects Gibberellin Response and Plant Development. *PLANT PHYSIOLOGY* **2001**, 126, 1174-1185, [10.1104/pp.126.3.1174](https://doi.org/10.1104/pp.126.3.1174).
27. Mariko Sawa; Steve A. Kay; GIGANTEA directly activates Flowering Locus T in *Arabidopsis thaliana*. *Proceedings of the National Academy of Sciences* **2011**, 108, 11698-11703, [10.1073/pnas.1106771108](https://doi.org/10.1073/pnas.1106771108).
28. Jae-Hoon Jung; Yeon-Hee Seo; Pil Joon Seo; Jose Luis Reyes; Ju Yun; Nam-Hai Chua; Chung-Mo Park; The GIGANTEA-Regulated MicroRNA172 Mediates Photoperiodic Flowering Independent of CONSTANS in *Arabidopsis*. *The Plant Cell* **2007**, 19, 2736-2748, [10.1105/tpc.107.054528](https://doi.org/10.1105/tpc.107.054528).
29. Young Hun Song; Akane Kubota; Michael S. Kwon; Michael F. Covington; Nayoung Lee; Ella R. Taagen; Dianne Laboy Cintrón; Dae Yeon Hwang; Reiko Akiyama; Sarah K. Hodge; et al. Molecular basis of flowering under natural long-day conditions in *Arabidopsis*. *Nature Plants* **2018**, 4, 824-835, [10.1038/s41477-018-0253-3](https://doi.org/10.1038/s41477-018-0253-3).
30. Zhao, X.Y.; Liu, M.S.; Li, J.R.; Guan, C.M.; Zhang, X.S. The wheat TaGI1, involved in photoperiodic flowering, encodes an *Arabidopsis* GI ortholog. *Plant Mol. Biol.* **2005**, 58, 53–64.
31. Hecht, V.; Knowles, C.L.; Vander Schoor, J.K.; Liew, L.C.; Jones, S.E.; Lambert, M.J.M.; Weller, J.L. Pea LATE BLOOMER1 Is a GIGANTEA ortholog with roles in photoperiodic flowering, deetiolation, and transcriptional regulation of circadian clock gene homologs. *Plant Physiol.* **2007**, 144, 648–661.
32. Wuriyangan, H.; Zhang, B.; Cao, W.-H.; Ma, B.; Lei, G.; Liu, Y.-F.; Wei, W.; Wu, H.-J.; Chen, L.-J.; Chen, H.-W.; et al. The ethylene receptor ETR2 delays floral transition and affects starch accumulation in rice. *Plant Cell* **2009**, 21, 1473–1494.
33. Bendix, C.; Mendoza, J.M.; Stanley, D.N.; Meeley, R.; Harmon, F.G. The circadian clock-associated gene *gigantea1* affects maize developmental transitions: *Gigantea1* regulates maize developmental transitions. *Plant Cell Environ.* **2013**, 36, 1379–1390.
34. Xie, Q.; Lou, P.; Hermand, V.; Aman, R.; Park, H.J.; Yun, D.-J.; Kim, W.Y.; Salmela, M.J.; Ewers, B.E.; Weinig, C.; et al. Allelic polymorphism of GIGANTEA is responsible for naturally occurring variation in circadian period in *Brassica rapa*. *Proc. Natl. Acad. Sci. USA* **2015**, 112, 3829–3834.

35. Karlgren, A.; Gyllenstrand, N.; Källman, T.; Lagercrantz, U. Conserved function of core clock proteins in the Gymnosperm Norway spruce (*Picea abies* L. Karst). *PLoS ONE* 2013, 8, e60110.
36. Tang, W.; Yan, H.; Su, Z.; Park, S.-C.; Liu, Y.; Zhang, Y.; Wang, X.; Kou, M.; Ma, D.; Kwak, S.-S.; et al. Cloning and characterization of a novel GIGANTEA gene in sweet potato. *Plant Physiol. Biochem.* 2017, 116, 27–35.
37. Satoshi Watanabe; Zhengjun Xia; Rumiko Hideshima; Yasutaka Tsubokura; Shusei Sato; Naoki Yamanaka; Ryoji Takahashi; Toyoaki Anai; Satoshi Tabata; Keisuke Kitamura; et al. A Map-Based Cloning Strategy Employing a Residual Heterozygous Line Reveals that the GIGANTEA Gene Is Involved in Soybean Maturity and Flowering. *Genetics* **2011**, 188, 395–407, [10.1534/genetics.110.125062](https://doi.org/10.1534/genetics.110.125062).
38. Yan Wang; Yongzhe Gu; Huihui Gao; Lijuan Qiu; Ruzhen Chang; Shouyi Chen; Chang Ruzhen; Molecular and geographic evolutionary support for the essential role of GIGANTEA in soybean domestication of flowering time. *BMC Evolutionary Biology* **2016**, 16, 1–13, [10.1186/s12862-016-0653-9](https://doi.org/10.1186/s12862-016-0653-9).
39. Tao Wang; Ming-Yang Sun; Xue-Song Wang; Wen-Bin Li; Yong-Guang Li; Over-Expression of GmGla-Regulated Soybean miR172a Confers Early Flowering in Transgenic *Arabidopsis thaliana*. *International Journal of Molecular Sciences* **2016**, 17, 645, [10.3390/ijms17050645](https://doi.org/10.3390/ijms17050645).
40. Shakhira Zakhrebekova; Simon P. Gough; Ilka Braumann; André H. Müller; Joakim Lundqvist; Katharina Ahmann; Christoph Dockter; Izabela Matyszcak; Marzena Kurowska; Arnis Druka; et al. Induced mutations in circadian clock regulator *Mat-a* facilitated short-season adaptation and range extension in cultivated barley. *Proceedings of the National Academy of Sciences* **2012**, 109, 4326–4331, [10.1073/pnas.1113009109](https://doi.org/10.1073/pnas.1113009109).
41. Hayama, R.; Izawa, T.; Shimamoto, K. Isolation of rice genes possibly involved in the photoperiodic control of flowering by a fluorescent differential display method. *Plant Cell Physiol.* 2002, 43, 494–504.
42. Huq, E.; Tepperman, J.M.; Quail, P.H. GIGANTEA is a nuclear protein involved in phytochrome signaling in *Arabidopsis*. *Proc. Natl. Acad. Sci. USA* 2000, 97, 9789–9794.
43. Hayama, R.; Yokoi, S.; Tamaki, S.; Yano, M.; Shimamoto, K. Adaptation of photoperiodic control pathways produces short-day flowering in rice. *Nature* 2003, 422, 719–722.
44. Kojima, S.; Takahashi, Y.; Kobayashi, Y.; Monna, L.; Sasaki, T.; Araki, T.; Yano, M. Hd3a, A rice ortholog of the *Arabidopsis* FT gene, promotes transition to flowering downstream of Hd1 under short-day conditions. *Plant Cell Physiol.* 2002, 43, 1096–1105.
45. Yano, M.; Katayose, Y.; Ashikari, M.; Yamanouchi, U.; Monna, L.; Fuse, T.; Baba, T.; Yamamoto, K.; Umehara, Y.; Nagamura, Y.; et al. Hd1, a major photoperiod sensitivity quantitative trait locus in rice, is closely related to the *Arabidopsis* flowering time gene *CONSTANS*. *Plant Cell* 2000, 12, 2473–2483.
46. Qingbo Ke; Ho Soo Kim; Zhi Wang; Chang Yoon Ji; Jae Cheol Jeong; Haeng-Soon Lee; Young-Im Choi; Bingcheng Xu; Xiping Deng; Dae-Jin Yun; et al. Down-regulation of GIGANTEA-like genes increases plant growth and salt stress tolerance in poplar. *Plant Biotechnology Journal* **2016**, 15, 331–343, [10.1111/pbi.12628](https://doi.org/10.1111/pbi.12628).
47. Sandra Rychel; Michał Książkiewicz; Magdalena Tomaszewska; Wojciech Bielski; Bogdan Wolko; FLOWERING LOCUS T, GIGANTEA, SEPALLATA, and FRIGIDA homologs are candidate genes involved in white lupin (*Lupinus albus* L.) early flowering. *Molecular Breeding* **2019**, 39, 43, [10.1007/s11032-019-0952-0](https://doi.org/10.1007/s11032-019-0952-0).
48. Akane Kubota; Shogo Kita; Kimitsune Ishizaki; Ryuichi Nishihama; Katsuyuki T. Yamato; Takayuki Kohchi; Co-option of a photoperiodic growth-phase transition system during land plant evolution. *Nature Communications* **2014**, 5, 3668, [10.1038/ncomms4668](https://doi.org/10.1038/ncomms4668).
49. Casal, J.J.; Luccioni, L.G.; Oliverio, K.A.; Boccandalro, H.E. Light, phytochrome signalling and photomorphogenesis in *Arabidopsis*. Dedicated to Professor Silvia Bratslavsky, to mark her great contribution to photochemistry and photobiology particularly in the field of photothermal methods. *Photochem. Photobiol. Sci.* 2003, 2, 625.
50. Martin-Tryon, E.L.; Kreps, J.A.; Harmer, S.L. GIGANTEA acts in blue light signaling and has biochemically separable roles in circadian clock and flowering time regulation. *Plant Physiol.* 2007, 143, 473–486.
51. Chentao Lin; Plant blue-light receptors. *Trends in Plant Science* **2000**, 5, 337–342, [10.1016/s1360-1385\(00\)01687-3](https://doi.org/10.1016/s1360-1385(00)01687-3).
52. Priyanka Emishra; Kishore Chandra Sekhar Panigrahi; GIGANTEA – an emerging story. *Frontiers in Plant Science* **2015**, 6, 8, [10.3389/fpls.2015.00008](https://doi.org/10.3389/fpls.2015.00008).
53. Karina Andrea Oliverio; María Crepy; Ellen L. Martin-Tryon; Raechel Milich; Stacey L. Harmer; J. Putterill; Marcelo J. Yanovsky; Jorge J. Casal; GIGANTEA Regulates Phytochrome A-Mediated Photomorphogenesis Independently of Its Role in the Circadian Clock. *Plant Physiology* **2007**, 144, 495–502, [10.1104/pp.107.097048](https://doi.org/10.1104/pp.107.097048).
54. Park, Y.-J.; Kim, J.Y.; Lee, J.-H.; Lee, B.-D.; Paek, N.-C.; Park, C.-M. GIGANTEA shapes the photoperiodic rhythms of thermomorphogenic growth in *Arabidopsis*. *Mol. Plant* 2020, 13, 459–470.

55. Nohales, M.A.; Kay, S.A. GIGANTEA gates gibberellin signaling through stabilization of the DELLA proteins in Arabidopsis. *Proc. Natl. Acad. Sci. USA* 2019, 116, 21893–21899.
56. Hwang, C.; Park, J.; Lee, B.; Cheong, H.; Loss of Function in GIGANTEA Gene is Involved in Brassinosteroid Signaling. *J. Chosun Nat. Sci.* **2011**, 4, 113–120, .
57. Kim, W.-Y.; Ali, Z.; Park, H.J.; Park, S.J.; Cha, J.-Y.; Perez-Hormaeche, J.; Quintero, F.J.; Shin, G.; Kim, M.R.; Qiang, Z.; et al. Release of SOS2 kinase from sequestration with GIGANTEA determines salt tolerance in Arabidopsis. *Nat. Commun.* 2013, 4, 1–13.
58. Kazan, K.; Lyons, R. The link between flowering time and stress tolerance. *J. Exp. Bot.* 2016, 67, 47–60.
59. Matteo Riboni; Alice Robustelli Test; Massimo Galbiati; Chiara Tonelli; Lucio Conti; Environmental stress and flowering time. *Plant Signaling & Behavior* **2014**, 9, e29036, [10.4161/psb.29036](https://doi.org/10.4161/psb.29036).
60. Corrales, A.-R.; Nebauer, S.G.; Carrillo, L.; Fernández-Nohales, P.; Marqués, J.; Renau-Morata, B.; Granell, A.; Pollmann, S.; Vicente-Carbajosa, J.; Molina, R.-V.; et al. Characterization of tomato Cycling Dof Factors reveals conserved and new functions in the control of flowering time and abiotic stress responses. *J. Exp. Bot.* 2014, 65, 995–1012.
61. Fornara, F.; Panigrahi, K.C.S.; Gissot, L.; Sauerbrunn, N.; Rühl, M.; Jarillo, J.A.; Coupland, G. Arabidopsis DOF transcription factors act redundantly to reduce CONSTANS expression and are essential for a photoperiodic flowering response. *Dev. Cell* 2009, 17, 75–86
62. Joon-Yung Cha; Da-Yeon Lee; Imdad Ali; Song Yi Jeong; BoBae Shin; Hyunjung Ji; Jung Sun Kim; Min-Gab Kim; Woe-Yeon Kim; Arabidopsis GIGANTEA negatively regulates chloroplast biogenesis and resistance to herbicide butafenacil. *Plant Cell Reports* **2019**, 38, 793-801, [10.1007/s00299-019-02409-x](https://doi.org/10.1007/s00299-019-02409-x).
63. Jasmina Kurepa; Jan Smalle; Marc Va; N Montagu; Dirk Inze; Oxidative stress tolerance and longevity in Arabidopsis: the late-flowering mutant gigantea is tolerant to paraquat. *The Plant Journal* **1998**, 14, 759-764, [10.1046/j.1365-3113x.1998.00168.x](https://doi.org/10.1046/j.1365-3113x.1998.00168.x).
64. Mohammad Reza Bolouri Moghaddam; Wim Van Den Ende; Sugars, the clock and transition to flowering. *Frontiers in Plant Science* **2013**, 4, null, [10.3389/fpls.2013.00022](https://doi.org/10.3389/fpls.2013.00022).
65. Neil Dalchau; Seong J. Baek; Helen M. Briggs; Fiona C. Robertson; Antony N. Dodd; Michael J. Gardner; Matthew A. Stancombe; Michael J. Haydon; Guy-Bart Stan; Jorge M. Gonçalves; et al. The circadian oscillator gene GIGANTEA mediates a long-term response of the Arabidopsis thaliana circadian clock to sucrose. *Proceedings of the National Academy of Sciences* **2011**, 108, 5104-5109, [10.1073/pnas.1015452108](https://doi.org/10.1073/pnas.1015452108).
66. Michael J. Haydon; Olga Mielczarek; Alexander Frank; Ángela Román; Alex A.R. Webb; Sucrose and Ethylene Signaling Interact to Modulate the Circadian Clock. *Plant Physiology* **2017**, 175, 947-958, [10.1104/pp.17.00592](https://doi.org/10.1104/pp.17.00592).
67. S. Q. Cao; Y. Q. Song; L. Su; Freezing sensitivity in the gigantea mutant of Arabidopsis is associated with sugar deficiency. *Biologia plantarum* **2007**, 51, 359-362, [10.1007/s10535-007-0073-1](https://doi.org/10.1007/s10535-007-0073-1).
68. Klaus Eimert; Shue-Mei Wang; Wei-Ling Lue; Jychian Chen; Monogenic Recessive Mutations Causing Both Late Floral Initiation and Excess Starch Accumulation in Arabidopsis. *The Plant Cell* **1995**, 7, 1703, [10.2307/3870031](https://doi.org/10.2307/3870031).
69. Johanna Krahmer; Greg S. Goraloglia; Akane Kubota; Argyris Zardilis; Richard S. Johnson; Young Hun Song; Michael J. MacCoss; Thierry Ele Bihan; Karen J. Halliday; Takato Imaizumi; et al. Time-resolved interaction proteomics of the GIGANTEA protein under diurnal cycles in Arabidopsis. *FEBS Letters* **2018**, 593, 319-338, [10.1002/1873-3468.13311](https://doi.org/10.1002/1873-3468.13311).
70. Gulzar S. Sanghera; Shabir H. Wani; Wasim Hussain; N. B. Singh; Engineering Cold Stress Tolerance in Crop Plants. *Current Genomics* **2011**, 12, 30-43, [10.2174/138920211794520178](https://doi.org/10.2174/138920211794520178).
71. C. Guy; Cold Acclimation And Freezing Stress Tolerance: Role Of Protein Metabolism. *Annual Review of Plant Biology* **1990**, 41, 187-223, [10.1146/annurev.arplant.41.1.187](https://doi.org/10.1146/annurev.arplant.41.1.187).
72. J. Weiss; Marcos Egea-Cortines; Transcriptomic analysis of cold response in tomato fruits identifies dehydrin as a marker of cold stress. *Journal of Applied Genetics* **2009**, 50, 311-319, [10.1007/bf03195689](https://doi.org/10.1007/bf03195689).
73. Guy, C.L.; Niemi, K.J.; Brambl, R. Altered gene expression during cold acclimation of spinach. *Proc. Natl. Acad. Sci. USA* 1985, 82, 3673–3677.
74. Thomashow, M.F. PLANT COLD ACCLIMATION: Freezing Tolerance Genes and Regulatory Mechanisms. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 1999, 50, 571–599.
75. Matsuo Uemura; Gareth Warren; Peter L. Steponkus; Freezing Sensitivity in the sfr4 Mutant of Arabidopsis Is Due to Low Sugar Content and Is Manifested by Loss of Osmotic Responsiveness. *Plant Physiology* **2003**, 131, 1800-1807, [10.1104/pp.102.013227](https://doi.org/10.1104/pp.102.013227).

76. M. Miquel; D. James; H. Dooner; J. Browse; Arabidopsis requires polyunsaturated lipids for low-temperature survival.. *Proceedings of the National Academy of Sciences* **1993**, 90, 6208-6212, [10.1073/pnas.90.13.6208](https://doi.org/10.1073/pnas.90.13.6208).
77. Z. Ristic; E. N. Ashworth; Changes in leaf ultrastructure and carbohydrates in *Arabidopsis thaliana* L. (Heyn) cv. Columbia during rapid cold acclimation. *Protoplasma* **1993**, 172, 111-123, [10.1007/bf01379368](https://doi.org/10.1007/bf01379368).
78. Perla Gómez; M Ángeles Ferrer; J Pablo Fernández-Trujillo; Antonio Calderón; Francisco Artés; Marcos Egea-Cortines; Julia Weiss; Structural changes, chemical composition and antioxidant activity of cherry tomato fruits (cv. Micro-Tom) stored under optimal and chilling conditions. *Journal of the Science of Food and Agriculture* **2009**, 89, 1543-1551, [10.1002/jsfa.3622](https://doi.org/10.1002/jsfa.3622).
79. Immink, R.G.; Hannapel, D.J.; Ferrario, S.; Busscher, M.; Franken, J.; Lookeren Campagne, M.M.; Angenent, G.C. A petunia MADS box gene involved in the transition from vegetative to reproductive development. *Development* **1999**, 126, 5117–5126.
80. Souer, E.; Rebocho, A.B.; Bliet, M.; Kusters, E.; de Bruin, R.A.M.; Koes, R. Patterning of Inflorescences and Flowers by the F-Box Protein DOUBLE TOP and the LEAFY Homolog ABERRANT LEAF AND FLOWER of Petunia. *Plant Cell* **2008**, 20, 2033–2048
81. Knudsen, J.T.; Tollsten, L.; Bergström, L.G. Floral scents—A checklist of volatile compounds isolated by head-space techniques. *Phytochemistry* **1993**, 33, 253–280.
82. Pichersky, E.; Gershenzon, J. The formation and function of plant volatiles: Perfumes for pollinator attraction and defense. *Curr. Opin. Plant Biol.* **2002**, 5, 237–243.
83. Natalia Dudareva; Antje Klempien; Joëlle K. Muhlemann; Ian Kaplan; Biosynthesis, function and metabolic engineering of plant volatile organic compounds. *New Phytologist* **2013**, 198, 16-32, [10.1111/nph.12145](https://doi.org/10.1111/nph.12145).
84. Alain Tissier; John A Morgan; Natalia Dudareva; Plant Volatiles: Going 'In' but not 'Out' of Trichome Cavities. *Trends in Plant Science* **2017**, 22, 930-938, [10.1016/j.tplants.2017.09.001](https://doi.org/10.1016/j.tplants.2017.09.001).
85. Efraim Lewinsohn; Nativ Dudai; Yaakov Tadmor; Irena Katzir; Uzi Ravid; Eli Putievsky; Daniel M. Joel; Histochemical Localization of Citral Accumulation in Lemongrass Leaves (*Cymbopogon citratus*(DC.) Stapf., Poaceae). *Annals of Botany* **1998**, 81, 35-39, [10.1006/anbo.1997.0525](https://doi.org/10.1006/anbo.1997.0525).
86. Vincent R. Franceschi; Paal Krokene; Erik Christiansen; Trygve Krokling; Anatomical and chemical defenses of conifer bark against bark beetles and other pests. *New Phytologist* **2005**, 167, 353-376, [10.1111/j.1469-8137.2005.01436.x](https://doi.org/10.1111/j.1469-8137.2005.01436.x).
87. Colquhoun, T.A.; Verdonk, J.C.; Schimmel, B.C.J.; Tieman, D.M.; Underwood, B.A.; Clark, D.G. Petunia floral volatile benzenoid/phenylpropanoid genes are regulated in a similar manner. *Phytochemistry* **2010**, 71, 158–167.
88. Cna'Ani, A.; Muhlemann, J.K.; Ravid, J.; Masci, T.; Klempien, A.; Nguyen, T.T.H.; Dudareva, N.; Pichersky, E.; Vainstein, A. Petunia × hybrida floral scent production is negatively affected by high-temperature growth conditions: Ambient temperature and floral scent. *Plant Cell Environ.* **2015**, 38, 1333–1346.
89. Myles P. Fenske; Kristen D. Hewett Hazelton; Andrew K. Hempton; Jae Sung Shim; Breanne M. Yamamoto; Jeffrey A. Riffell; Takato Imaizumi; Circadian clock gene LATE ELONGATED HYPOCOTYL directly regulates the timing of floral scent emission in Petunia. *Proceedings of the National Academy of Sciences* **2015**, 112, 9775-9780, [10.1073/pnas.1422875112](https://doi.org/10.1073/pnas.1422875112).