

Natural Preservatives for Fish

Subjects: **Food Science & Technology**

Contributor: Jun Mei

Fish is extremely perishable as a result of rapid microbial growth naturally present in fish or from contamination. Synthetic preservatives are widely used in fish storage to extend shelf life and maintain quality and safety. Natural preservatives from microorganisms, plants, and animals have been shown potential in replacing the chemical antimicrobials. Bacteriocins and organic acids from bacteria showed good antimicrobial activities against spoilage bacteria. Plant-derived antimicrobials could prolong fish shelf life and decrease lipid oxidation. Animal-derived antimicrobials also have good antimicrobial activities.

natural preservatives

fish

spoilage mechanisms

application

1. Introduction

Fish has high protein content and low saturated fat content, which is considered as highly valuable food ^[1]. In particular, fish is the primary dietary source of omega-3 polyunsaturated fatty acid (PUFA), including docosahexaenoic acid (DHA) and eicosapentaenoic acid (EPA), both of which are well-known for the anti-inflammatory action and protective effects on cardiovascular disease ^{[2][3][4]}. The World Health Organization (WHO) recommends a regular fish consumption of 1–2 servings per week to provide the equivalent of 200–500 mg of omega-3 PUFA ^{[5][6]}. According to statistics, fresh and live fish account for about half of the total seafood consumed by human beings. The word “fresh” refers to fish that have not been frozen, including still alive fish as well as kept in the cold but not frozen section or packed in modified atmosphere ^[7]. Unlike frozen fish, the fresh fish cannot stay as inventory for one month ^[8]. For consumers, freshness is also often associated with safety, reassurance and superior taste. In most cases, the customers believe that the fish sold in market in China are caught recently. However, it could take more than a week to arrive at these stores. Consumers are still unsure whether the “fresh” product is really fresh or has been frozen and then thawed ^{[9][10]}.

Fresh fish can easily deteriorate after being captured due to the endogenous enzyme and rapid microbial growth naturally present in fish or from contamination ^[11]. In the process of fish decay, decomposition of various components and formation of new compounds occur. What is more, changes in composition during fish decay lead to protein degradation and lipid oxidation, as well as changes in fish odor, flavor, and texture. Therefore, it has become inevitable to develop effective treatment methods to extend the shelf life of fish ^[12]. Soft or mushy texture of fish limits the shelf life, thereby impeding its marketing. During postmortem handling and storage, the holding temperature, oxygen, endogenous or microbial proteases, moisture can result in detrimental changes in the color, odor, texture, and flavor of fish ^{[13][14]}. Therefore, fish have traditionally been cooled and stored in flake ice, refrigerated sea water, or ice slurries or they have been preserved by exposure to chemical agents. At the same

time, the fishery industry is always looking for new preservation methods to extend the fish shelf life and provide consumers with the best quality in terms of sensory and nutritional levels [\[15\]](#).

In recent years, researchers have put much effort into searching natural preservatives that could inhibit the growth of bacteria and fungi in food. Meanwhile, a growing number of consumers are aware of the potential negative health effects of chemical preservatives, which has prompted the food industry to find natural products used and developed as alternatives. Natural preservatives are available from a variety of sources including plants, animals, bacteria, algae, and fungi [\[16\]\[17\]](#). Microbial derived preservatives (e.g., bacteriocin), plant derived preservatives (thyme essential oil, tea polyphenols, rosemary extract, etc.), and animal derived preservatives (e.g., chitosan from crab or shrimp shells) have been demonstrated to have antimicrobial or antioxidant properties. In addition, antimicrobial compounds produced by algae and fungal (mushroom) could be served as potential sources of new antimicrobial substances for use as natural preservatives in food.

2. Spoilage Mechanisms

Fish is extremely perishable compared to other muscle foods and will enter into rigor mortis where fish lose their flexibility due to the stiffening of their muscle just a few hours after death [\[18\]](#). Studies show that the spoilage of fish results from three basic factors: enzymatic autolysis, oxidation, and microbial growth [\[19\]](#).

2.1. Autolytic Enzymatic Spoilage

As the degradation process of fish begins with autolytic enzymatic spoilage, chemical and biological changes take place in dead fish shortly after capture due to autolytic enzymatic breakdown of major molecules. The autolytic enzymes had a significant impact on textural deterioration (softening) and therefore on spoilage; however, they did not produce the characteristic spoilage off-flavors and off-odors [\[20\]\[21\]](#). Some studies have demonstrated that the fish quality could deteriorate even though comparatively low levels of autolytic degradation were present and limit the shelf life of fish [\[22\]\[23\]](#).

Some gastrointestinal digestion enzymes as well as endogenous muscular enzymes are found in the viscera and muscle of fish after being caught. These enzymes can be conducive to postmortem degradation in fish muscle and fish products during processing and storage, which can lead to a sensory or product associated change. On the other hand, the autolysis of fish muscle proteins could also result in fish meat spoilage and biogenic amines production. The degree of fish freshness has to be determined in the autolysis stage before the spoilage of fish begins [\[24\]](#). It has been pointed out that low temperature and a_w kept during the storage period can maintain a low activity of the endogenous autolytic enzymes in muscle [\[24\]](#). In addition, the temperature of vacuum-packed cold-smoked salmon storage should be at low temperature to maintain low native enzyme activity in the tissue [\[25\]](#).

2.2. Oxidative Spoilage

Lipid oxidation could result in spoilage and deterioration and contained a three-stage free radical mechanism: initiation, propagation, and termination [\[26\]\[27\]\[28\]](#). Some studies suggested that lipid oxidation in fish may be

initiated and propagated by a number of mechanisms including auto-oxidation, lipoxygenase, microsomal enzymes, photosensitized oxidation, and peroxidase [29]. In addition, iron-bound proteins in fish muscles, such as hemoglobin, ferritin, myoglobin, hemosiderin, and transferrin, as well as other metals, may be released during storage and play an important role in initiating and/or activating lipid oxidation [30].

Lipid oxidation can be divided into enzymatic and non-enzymatic oxidation, both of which can result in serious decreases in fish qualities. It is typically recognized that oxidation involves the reaction of oxygen with the double bonds of fatty acids. Fish is rich in unsaturated fatty acids and prooxidant molecules, which are extremely susceptible to oxidation resulting in rancidity development and quality loss [31]. These compounds result in deteriorations of smell, color, texture, and nutritional values. Lipid oxidation products also have been demonstrated to promote protein denaturation, modification of protein electrophoretic profiles, nutritional losses, endogenous antioxidant systems losses, and developments of fluorescent compounds [32]. Accordingly, lipid oxidation leads to a decrease of fish acceptability by consumers [33].

2.3. Microbial Spoilage

Fish has high contents of free amino acids, a high post mortem pH, high water contents, and many fish species contain trimethylamine oxide (TMAO), which promote bacterial growth in a wide temperature range [34]. The microbial growth is considered to be the major cause of the deterioration of fish quality, causing up to 25–30% loss of such products [17]. It is generally believed that each fish has its unique flora, which is determined by raw materials, processing parameters, subsequent storage conditions, and microbial tolerance to storage conditions [35]. For example, it was reported that spoilage microorganisms for aerobically stored frozen fish including salmon were species within the genera *Shewanella* (S.) and *Pseudomonas* (P.), while the CO₂-resistant *Photobacterium* (Ph.) *phosphoreum* dominated on fish under modified atmosphere packaging [34][35][36].

It is important to distinguish non-spoilage bacteria from spoilage ones as many of the bacteria present do not actually contribute to spoilage during fish storage [37]. A variety of psychrotrophic Gram-negative bacteria comprise the main part of the initial microbiota of fish from temperate sea waters. However, only a small fraction of fish microbiota is responsible for spoilage, known as specific spoilage organisms (SSOs). The SSOs could dominate and produce the metabolites through assimilation of nonprotein-nitrogen of fish muscle with unpleasant and unacceptable off-flavors, which directly affect the organoleptic properties of fish resulting in its rejection by the consumers [38]. The SSOs were different for different fish species and preservation conditions. Research studies demonstrated that *Pseudomonas* spp. was the SSOs for Atlantic salmon (*Salmo salar*) in modified atmosphere packaging and bighead carp (*Aristichthys nobilis*) with 2% salt, whereas *Aeromonas* was the SSOs for unsalted bighead carp [39].

3. Natural Preservatives for Fish

Fresh fish is highly perishable, and even if it can be refrigerated or frozen to extend its shelf life, these processes may not be sufficient to prevent lipid oxidation, rancidity or bacterial growth. In most cases, it is also necessary to

improve the quality of fish [40]. For this reason, it is required that preservatives should be properly added to fish during storage.

With the rapid development of social economy, the application of natural preservatives in food has attracted more and more attention from the public. Generally, the public will choose a food with no preservatives, but if these are not available, the same consumer will choose a food containing natural preservatives over synthetic ones [41]. Natural preservatives guarantee that the food is free of microorganisms and safe to eat. Ideally, natural preservatives should have broad bactericidal and fungicidal activities, be non-toxic, be active at low concentrations, impart no flavor or color to food, have no pharmaceutical applications, label friendly, and finally cost effective [42]. Natural preservatives generally come from three sources: microorganisms, animals, and plants. In addition, various bio-active compounds extracted from algae, mushrooms and so on can also provide a potential source of new natural preservatives in the food industry [43].

3.1. Microbial-Derived Compounds

Specific strains of lactic acid bacteria (LAB) produce some inhibitory substances (such as diacetyl, reutericyclin), antifungal compounds (such as phenyl-lactate, propionate, cyclic dipeptides, hydroxyphenyl-lactate, and 3-hydroxy fatty acid), bacteriocins, and bacteriocin-like inhibitory substances, which have antibacterial activity and can be exploited against spoilage bacteria and food-borne pathogens in fish during storage [44]. Some applications of microbial-derived compounds in fish preservation are depicted in Table 1.

Table 1. Survey of literature dealing with bacteriocin employed biopreservation of fish.

Product	Bacteriocin Employed	Reported Effects	References
Skinless blue shark steak	Pediocin ACCEL	<i>L. monocytogenes</i> ↓ ¹	[45]
Oysters, mussels, clams	BacALP7, bacALP57	<i>L. monocytogenes</i> ↓, <i>L. innocua</i> ↓	[46]
Cold-smoked salmon	Divercin V41	<i>L. monocytogenes</i> ↓	[47][48]
Fresh salmon fillets	Bacteriocin produced by <i>Lb. pentosus</i> 39	<i>A. hydrophila</i> ↓, <i>L. monocytogenes</i> ↓	[49]

Product	Bacteriocin Employed	Reported Effects	References
Reef cod	Enterocin CD1	The total viable count↓	[50]
Reef cod	Bacteriocin PSY2	The total count of spoilage bacteria↓	[51]
Pangasius fish fillets	Bacteriocin 7293	Target microorganisms↓	[52]
Reef cod filets	Bacteriocin GP1	Similar effect with that of sodium benzoate and the nisin B440	[53]
Fish pâté using fresh Nile tilapia	Bacteriocin produced by <i>L. lactis</i> 3MT	<i>Vibrio</i> ↓	[54]
Fresh hake paste	Bacteriocin produced by <i>E. mundtii</i> STw38	Native flora of fish paste↓	[55]

efits of

seafood (fish and shellfish) consumption in relation to health risks associated with exposure to methylmercury. EFSA J. 2016, 12, 3761.

¹Inhibited or decreased.
2. Vlaevert, L.; Borrell, F.; Nadal, M.; Jacobs, S.; Minnens, F.; Verbeke, W.; Marques, A.; Domingo, J.L. Health risk/benefit information for consumers of fish and shellfish: Fish Choice, a new online tool. Food Chem. Toxicol. 2017, 104, 79–84.

3.2. Plant-Derived Compounds

EOs are complex mixtures of volatile organic compounds (VOCs) produced as secondary metabolites in plants and frequently responsible for the characteristic odor of plants [56]. They are characterized by two or three major VOCs at fairly high concentrations (20–70%) compared to other VOCs [57]. Some EOs have antimicrobial and antioxidant properties, and an increasing demand for natural preservatives has led to EOs as potential alternatives for antimicrobials and antioxidants [58]. EOs have been proved to be effective antimicrobials against some foodborne pathogens including *S. Typhimurium*, *E. coli* O157: H7, *Campylobacter*, *L. monocytogenes*, *S. aureus*, and others. Studies show that the efficacy of EOs depends on chemical structure, concentration, matching the antimicrobial activity spectrum with the target microorganism(s), interactions with the food matrix, and application method [59].

5. Tediosi, A.; Fait, G.; Jacobs, S.; Verbeke, W.; Álvarez-Muñoz, D.; Diogene, J.; Reuver, M.; Marques, A.; Caph, E. Insights from an international stakeholder consultation to identify informational needs related to seafood safety. Environ. Res. 2015, 143, 20–28.

6. Wallin, A.; Di Giuseppe, D.; Orsini, N.; Åkesson, A.; Forouhi, N.G.; Wolk, A. Fish consumption and frying of fish in relation to type 2 diabetes incidence: A prospective cohort study of Swedish men. Eur. J. Nutr. 2015, 56, 843–852.

7. Alvarado, W.; Hunt, W.; Storey, J.; Monahan, L.S.; Hill, C.R. Segregation and preference for fresh seafood in an Australian context. *Food Preference* 2014, 38, 146–150. doi:10.1016/j.foodpre.2014.03.002

8. Hicks, D.T. Seafood safety and quality: The consumer's role. *Foods* 2016, 5, 71. doi:10.3390/foods5010071

9. Wakamatsu, H.; Miyata, T. A demand analysis for the Japanese cod markets with unknown structural changes. *Fish. Sci.* 2015, 81, 393–400. doi:10.1007/s12572-015-9288-2

10. Bruhn, C.M. *Consumer Acceptance of High-Pressure Processed Products: American Perspective*; Springer: New York, NY, USA, 2016.

11. Jiang, D.; Liu, Y.; Jiang, H.; Rao, S.; Fang, W.; Wu, M.; Yuan, L.; Fang, W. A novel screen-printed mast cell-based electrochemical sensor for detecting spoilage bacterial quorum signaling molecules (N-acyl-homoserine-lactones) in freshwater fish. *Biosens. Bioelectron.* 2018, 102, 396–402. doi:10.1016/j.bios.2018.05.038

12. Campos, C.A.; Castro, M.P.; Aubourg, S.P.; Velázquez, J.B. *Novel Technologies in Food Science*; Springer: New York, NY, USA, 2012.

13. Sriket, C.; Benjakul, S.; Visessanguan, W.; Kishimura, H. Collagenolytic serine protease in freshwater prawn (*Macrobrachium rosenbergii*): Characteristics and its impact on muscle during iced storage. *Food Chem.* 2010, 124, 29–35. doi:10.1016/j.foodchem.2010.05.048

Figure 1. (a) Bulk EOs and different types of EO delivery systems, including nanoemulsion, liposomes, and biopolymer films. (b) Proposed common mechanisms of action and target sites of essential oils (EOs) or EO delivery systems on bacterial cell [72].

15. Rey, M.S.; Garcíasoto, B.; Fuertesgamundi, J.R.; Aubourg, S.; Barrosvelázquez, J. Effect of a natural organic acid-icing system on the microbiological quality of commercially relevant chilled fish species. *LWT-Food Sci. Technol.* 2012, 46, 217–223. doi:10.1016/j.lwt.2012.05.018

16. Chanbari, M.; Jami, M.; Domig, K.J.; Kneifel, W. Seafood biopreservation by lactic acid bacteria—A review. *LWT-Food Sci. Technol.* 2013, 54, 315–324. doi:10.1016/j.lwt.2012.09.001

17. Hassoun, A.; Gobari, O.E. Essential oils for antimicrobial and antioxidant applications in fish and other seafood products. *Trends Food Sci. Technol.* 2017, 68, 26–36. doi:10.1016/j.tfs.2017.05.002

18. Hong, H.; Regenstein, J.M.; Luo, Y. The importance of ATP-related compounds for the freshness and flavor of post-mortem fish and shellfish muscle: A review. *Crit. Rev. Food Technol.* 2015, 57, 1787–1798. doi:10.1080/10407179.2015.1061111

19. Chai, A.E.; Dave, D.; Budge, S.; Brooks, W.S. Fish spoilage mechanisms and preservation techniques. *Review Am. J. Appl. Sci.* 2010, 7, 859–877. doi:10.5555/ajass.2010070859

20. Lyhs, U.; Lahtinen, J.; Schelvismit, R. Microbiological quality of maatjes herring stored in air and under modified atmosphere at 4 and 10 °C. *Food Microbiol.* 2007, 24, 508–516. doi:10.1016/j.fm.2007.03.002

Additionally, the smoke treatment could increase redness of the fish muscle and stabilized it during frozen storage [84]. It should be noted that wood smoke contains some harmful compounds,

21. Ocaño-Higuera, V.M.; Medina-Ramírez, P.A.S.; Vázquez-Sánchez, E.; Sanizales-Medina, D.F.; Castillo-Ruiz, F.; Ruiz-Quintero, E.; Gracian-Pareda, A.; Pérez-Majano, M.F. Freshness smoke assessment of rayfish stored in trays by biochemical, chemical and physical methods. *Food Chem.* 2011, 125, 49–54.
22. Lakshmanan, R.; Piggott, J.R.; Paterson, A. Potential applications of high pressure for treatment to reduce or eliminate food-borne pathogens on fish products. Antimicrobial efficacy of liquid smoke can be enhanced by vacuum-packaging, essential oils, and NaCl [88][89][90][91][92]. Also, wood smoke can be converted into nanocapsules using chitosan and surface contact area could be increased, which delayed microbial growth in fish fillets at cold storage conditions [93].
23. Sumar, S.; Fraser, O.P. Compositional changes and spoilage in fish (part II) – Microbiological and chemical changes. *Nutr. Food Sci.* 1998, 98, 325–329.
24. Pachecoaguiar, R.; Lugosanchez, M.E.; Roblesburgueno, M.R. Postmortem biochemical and functional characteristic of monterey sardine muscle stored at 0 °C. *J. Food Sci.* 2010, 65, 40–47.
25. Dondero, M.; Cisternas, F.; Cavañal, L.; Simpson, R. Changes in quality of vacuum-packed cold-smoked salmon (*Salmo salar*) as a function of storage temperature. *Food Chem.* 2004, 87, 543–550.
26. Fraser, O.; Sumar, S. Compositional changes and spoilage in fish—an introduction. *Nutr. Food Sci.* 1998, 98, 275–279.
27. Shahidi, F.; Botta, J.R. *Seafoods: Chemistry, Processing Technology and Quality*; Springer: New York, NY, USA, 1994.
28. Frankel, E.N. Chemistry of free radicals and singlet oxidation of lipids. *Prog. Lipid Res.* 1984, 23, 197–221.
29. Hsieh, R.J.; Kinsella, J.E. Oxidation of polyunsaturated fatty acids: Mechanisms, products, and permeability, protein leakage, and binding of bacterial DNA [104]. The antimicrobial activities of mushrooms may be related to a variety of secondary metabolites with biological activity, such as gallic acids, some phenols, volatile compounds, free fatty acids, and their derivatives [95].
30. Alishahi, A.; Aider, M. Applications of chitosan in the seafood industry and aquaculture: A review. *Food Bioprocess Technol.* 2012, 5, 817–820.
31. Losada, V.; Barros-Velázquez, J.; Aubourg, S.P. Rancidity development in frozen pelagic fish: Influence of slurry ice as preliminary chilling treatment. *LWT-Food Sci. Technol.* 2007, 40, 991–999.
32. Lucas, A.; Losada, V.; Hóvári, J.; Lebovics, V.; Jakóczy, L.; Aubourg, S. Effect of pre-soaking whole pelagic fish in a plant extract on sensory and biochemical changes during subsequent frozen storage. *LWT-Food Sci. Technol.* 2007, 40, 930–936.
33. Beck, C. Effects of high pressure processing on lipid oxidation: A review. *Innov. Food Sci. Emerg. Technol.* 2014, 22, 1–10.
34. Chaillou, S.; Chaillot-Talmon, A.; Gaekobeker, H.; Cardinal, M.; Christmann, S.; Denis, C.; Desmonts, M.H.; Doussset, X.; Feurer, C.; Hamon, E. Origin and ecological selection of core and alkyl chain groups generally decreases the activity. The hydrophobic substituents such as prenyl groups, alkyl chain, and hydrogen or oxygen-containing heterocyclic moieties usually enhance the

activity against specific bacterial strains [112]. Antimicrobials associated with meat and seafood spoilage [64]. Use of natural preservatives [113]. people are increasingly interested in finding more natural alternatives.

35. Mørseth, T.; Moen, B.; Heir, E.; Hansen, A.Å.; Langsrud, S. Contamination of salmon fillets and processing plants with spoilage bacteria. *Int. J. Food Microbiol.* 2016, 237, 98–108.

At present, many animal-derived antimicrobial compounds have also been used for fish preservation. Such examples include chitosan from shellfish, lactoperoxidase, and lactoferrin from milk, and lysozymes from hen eggs [113]. However, one of the major problems associated with animal-derived antimicrobials is their allergen risk; the sources of such ingredients are often allergen-containing foods including shellfish, milk, and egg [114].

36. Murhekar, S.; Wright, M.H.; Greene, A.C.; Brownlie, J.C.; Cock, I.E. Inhibition of *Shewanella* spp. growth by *Syzygium australe* and *Syzygium luehmannii* extracts: Natural methods for the prevention of fish spoilage. *J. Food Sci. Technol.* 2017, 54, 3314–3326.
37. Huss, H.H. *Quality and Quality Changes in Fresh Fish*; FAO: Rome, Italy, 1995.
38. Boziaris, I.S.; Parlapani, F.F. *The Microbiological Quality of Food*; Woodhead Publishing: Sawston, UK, 2017.
39. Liu, X.; Zhang, Y.; Li, D.; Luo, Y. Characterization of the microbiota in lightly salted bighead carp (*Aristichthys nobilis*) fillets stored at 4 °C. *Food Microbiol.* 2017, 62, 106–111.
40. Erkan, N.; Doğruyol, H.; Günlü, A.; Genç, İ.Y. Use of natural preservatives in seafood: Plant extracts, edible film and coating. *J. Food Health Sci.* 2015, 1, 33–49.
41. Carocho, M.; Barreiro, M.F.; Morales, P.; Ferreira, I.C.F.R. Adding molecules to food, pros and cons: A review on synthetic and natural food additives. *Compr. Rev. Food Sci. Food Safety* 2014, 13, 377–399.
42. Carocho, M.; Morales, P.; Ferreira, I.C.F.R. Natural food additives: Quo vadis? *Trends Food Sci. Technol.* 2015, 45, 284–295.
43. Gyawali, R.; Ibrahim, S.A. Natural products as antimicrobial agents. *Food Control* 2014, 46, 412–429.
44. Gálvez, A.; Abriouel, H.; Benomar, N.; Lucas, R. Microbial antagonists to food-borne pathogens and biocontrol. *Curr. Opin. Biotechnol.* 2010, 21, 142–148.
45. Yin, L.J.; Chienwei, W.U.; Jiang, S.T. Biopreservative effect of pediocin ACCEL on refrigerated seafood. *Fish. Sci.* 2007, 73, 907–912.
46. Pinto, A.L.; Fernandes, M.; Pinto, C.; Albano, H.; Castilho, F.; Teixeira, P.; Gibbs, P.A. Characterization of anti-*Listeria* bacteriocins isolated from shellfish: Potential antimicrobials to control non-fermented seafood. *Int. J. Food Microbiol.* 2009, 129, 50–58.
47. Brillet-Viel, A.; Pilet, M.F.; Courcoux, P.; Prévost, H.; Leroi, F. Optimization of growth and bacteriocin activity of the food bioprotective *Carnobacterium divergens* V41 in an animal origin protein free medium. *Front. Mar. Sci.* 2016, 3, 128.
48. Brillet, A.; Pilet, M.F.; Prevost, H.; Bouttefroy, A.; Leroi, F. Biodiversity of *Listeria monocytogenes* sensitivity to bacteriocin-producing *Carnobacterium* strains and application in sterile cold-smoked

- salmon. *J. Appl. Microbiol.* 2004, 97, 1029–1037.
49. Anacarso, I.; Messi, P.; Condò, C.; Iseppi, R.; Bondi, M.; Sabia, C.; Niederhäusern, S.D. A bacteriocin-like substance produced from *Lactobacillus pentosus* 39 is a natural antagonist for the control of *Aeromonas hydrophila* and *Listeria monocytogenes* in fresh salmon fillets. *LWT-Food Sci. Technol.* 2014, 55, 604–611.
 50. Sarika, A.R.; Lipton, A.P.; Aishwarya, M.S.; Dhivya, R.S. Efficacy of bacteriocin of *Enterococcus faecalis* CD1 as a biopreservative for high value marine fish reef cod (*Epinephelus diacanthus*) under different storage conditions. *J. Microbiol. Biotechnol. Res.* 2017, 1, 18–24.
 51. Sarika, A.R.; Lipton, A.P.; Aishwarya, M.S.; Dhivya, R.S. Isolation of a bacteriocin-producing *Lactococcus lactis* and application of its bacteriocin to manage spoilage bacteria in high-value marine fish under different storage temperatures. *Appl. Biochem. Biotechnol.* 2012, 167, 1280–1289.
 52. Woraprayote, W.; Pumpuang, L.; Tosukhowong, A.; Zendo, T.; Sonomoto, K.; Benjakul, S.; Visessanguan, W. Antimicrobial biodegradable food packaging impregnated with Bacteriocin 7293 for control of pathogenic bacteria in pangasius fish fillets. *LWT-Food Sci. Technol.* 2018, 89, 427–433.
 53. Sarika, A.R.; Lipton, A.P.; Aishwarya, M.S. Biopreservative efficacy of bacteriocin GP1 of *Lactobacillus rhamnosus* GP1 on stored fish filets. *Front. Nutr.* 2019, 6, 29.
 54. Kaktcham, P.M.; Tchamani Piame, L.; Sandjong Sileu, G.M.; Foko Kouam, E.M.; Temgoua, J.-B.; Zambou Ngoufack, F.; de Lourdes Pérez-Chabela, M. Bacteriocinogenic *Lactococcus lactis* subsp. *lactis* 3MT isolated from freshwater Nile Tilapia: Isolation, safety traits, bacteriocin characterisation, and application for biopreservation in fish pâté. *Arch. Microbiol.* 2019.
 55. Delcarlo, S.B.; Parada, R.; Schelegueda, L.I.; Vallejo, M.; Marguet, E.R.; Campos, C.A. From the isolation of bacteriocinogenic LAB strains to the application for fish paste biopreservation. *LWT-Food Sci. Technol.* 2019, 110, 239–246.
 56. Fleita, D.; El-Sayed, M.; Rifaat, D. Evaluation of the antioxidant activity of enzymatically-hydrolyzed sulfated polysaccharides extracted from red algae; *Pterocladia capillacea*. *LWT-Food Sci. Technol.* 2015, 63, 1236–1244.
 57. Bakkali, F.; Averbeck, S.; Averbeck, D.; Idaomar, M. Biological effects of essential oils—A review. *Food Chem. Toxicol.* 2008, 46, 446–475.
 58. Calo, J.R.; Crandall, P.G.; O'Bryan, C.A.; Ricke, S.C. Essential oils as antimicrobials in food systems—A review. *Food Control* 2015, 54, 111–119.
 59. Tiwari, B.K.; Valdramidis, V.P.; O'Donnell, C.P.; Muthukumarappan, K.; Bourke, P.; Cullen, P.J. Application of natural antimicrobials for food preservation. *J. Agric. Food Chem.* 2009, 57, 5987–6000.

60. Wan, J.; Zhong, S.; Schwarz, P.; Chen, B.; Rao, J. Enhancement of antifungal and mycotoxin inhibitory activities of food-grade thyme oil nanoemulsions with natural emulsifiers. *Food Control* 2019, 106, 106709.
61. Kang, J.; Jin, W.; Wang, J.; Sun, Y.; Wu, X.; Liu, L. Antibacterial and anti-biofilm activities of peppermint essential oil against *Staphylococcus aureus*. *LWT-Food Sci. Technol.* 2019, 101, 639–645.
62. Zhang, L.L.; Zhang, L.F.; Hu, Q.P.; Hao, D.L.; Xu, J.G. Chemical composition, antibacterial activity of *Cyperus rotundus* rhizomes essential oil against *Staphylococcus aureus* via membrane disruption and apoptosis pathway. *Food Control* 2017, 80, 290–296.
63. Ziaee, E.; Razmjooei, M.; Shad, E.; Eskandari, M.H. Antibacterial mechanisms of *Zataria multiflora* Boiss. essential oil against *Lactobacillus curvatus*. *LWT-Food Sci. Technol.* 2018, 87, 406–412.
64. Hącwydro, K.; Flasiński, M.; Romańczuk, K. Essential oils as food eco-preservatives: Model system studies on the effect of temperature on limonene antibacterial activity. *Food Chem.* 2017, 235, 127–135.
65. Wang, Y.; Zhang, Y.; Shi, Y.Q.; Pan, X.H.; Lu, Y.H.; Cao, P. Antibacterial effects of cinnamon (*Cinnamomum zeylanicum*) bark essential oil on *Porphyromonas gingivalis*. *Microb. Pathog.* 2018, 116, 26–32.
66. Guo, L.; Sun, Q.; Gong, S.; Bi, X.; Jiang, W.; Xue, W.; Fei, P. Antimicrobial activity and action approach of the olive oil polyphenol extract against *Listeria monocytogenes*. *Front. Microbiol.* 2019, 10, 1586.
67. Cui, S.; Ma, X.; Wang, X.; Zhang, T.-A.; Hu, J.; Tsang, Y.F.; Gao, M.-T. Phenolic acids derived from rice straw generate peroxides which reduce the viability of *Staphylococcus aureus* cells in biofilm. *Ind. Crop. Prod.* 2019, 140, 111561.
68. Meira, N.; Holley, R.A.; Bordin, K.; Macedo, R.; Luciano, F.B. Combination of essential oil compounds and phenolic acids against *Escherichia coli* O157: H7 in vitro and in dry-fermented sausage production. *Int. J. Food Microbiol.* 2017, 260, 59–64.
69. Friedman, M.; Levin, C.E.; Henika, P.R. Addition of phytochemical-rich plant extracts mitigate the antimicrobial activity of essential oil/wine mixtures against *Escherichia coli* O157:H7 but not against *Salmonella enterica*. *Food Control* 2017, 73, 562–565.
70. Hyltdgaard, M.; Mygind, T.; Meyer, R.L. Essential oils in food preservation: Mode of action, synergies, and interactions with food matrix components. *Front. Microbiol.* 2012, 3, 12.
71. Wen, P.; Zhu, D.H.; Wu, H.; Zong, M.H.; Jing, Y.R.; Han, S.Y. Encapsulation of cinnamon essential oil in electrospun nanofibrous film for active food packaging. *Food Control* 2016, 59, 366–376.

72. Rao, J.; Chen, B.; McClements, D.J. Improving the efficacy of essential oils as antimicrobials in foods: Mechanisms of action. *Annu. Rev. Food Sci. Technol.* 2019, 10, 365–387.
73. Akter, S.; Netzel, M.E.; Tinggi, U.; Osborne, S.A.; Fletcher, M.T.; Sultanbawa, Y. Antioxidant rich extracts of *Terminalia ferdinandiana* inhibit the growth of foodborne bacteria. *Foods* 2019, 8, 281.
74. Bouarab Chibane, L.; Degraeve, P.; Ferhout, H.; Bouajila, J.; Oulahal, N. Plant antimicrobial polyphenols as potential natural food preservatives. *J. Sci. Food Agric.* 2019, 99, 1457–1474.
75. Snchez, E.; García, S.; Heredia, N. Extracts of edible and medicinal plants damage membranes of *vibrio cholerae*. *Appl. Environ. Microb.* 2010, 76, 6888–6894.
76. Chew, Y.L.; Chan, E.W.L.; Tan, P.L.; Lim, Y.Y.; Stanslas, J.; Goh, J.K. Assessment of phytochemical content, polyphenolic composition, antioxidant and antibacterial activities of Leguminosae medicinal plants in Peninsular Malaysia. *BMC Complem. Altern. Med.* 2011, 11, 12.
77. Ashrafi, A.; Jokar, M.; Mohammadi Nafchi, A. Preparation and characterization of biocomposite film based on chitosan and kombucha tea as active food packaging. *Int. J. Biol. Macromol.* 2018, 108, 444–454.
78. Kharchoufi, S.; Licciardello, F.; Siracusa, L.; Muratore, G.; Hamdi, M.; Restuccia, C. Antimicrobial and antioxidant features of ‘Gabsi’ pomegranate peel extracts. *Ind. Crop. Prod.* 2018, 111, 345–352.
79. Ghayempour, S.; Montazer, M.; Mahmoudi, R.M. Tragacanth gum as a natural polymeric wall for producing antimicrobial nanocapsules loaded with plant extract. *Int. J. Biol. Macromol.* 2015, 81, 514–520.
80. Cando, D.; Morcuende, D.; Utrera, M.; Estévez, M. Phenolic-rich extracts from Willowherb (*Epilobium hirsutum* L.) inhibit lipid oxidation but accelerate protein carbonylation and discoloration of beef patties. *Eur. Food Res. Technol.* 2014, 238, 741–751.
81. Kumar, S.; Pandey, A.K. Chemistry and biological activities of flavonoids: An overview. *Sci. World J.* 2013, 2013, 162750.
82. Negi, P.S. Plant extracts for the control of bacterial growth: Efficacy, stability and safety issues for food application. *Int. J. Food Microbiol.* 2012, 156, 7–17.
83. Lingbeck, J.M.; Cordero, P.; O’Bryan, C.A.; Johnson, M.G.; Ricke, S.C.; Crandall, P.G. Functionality of liquid smoke as an all-natural antimicrobial in food preservation. *Meat Sci.* 2014, 97, 197–206.
84. Kristinsson, H.G.; Danyali, N.; Ua-Angkoon, S. Effect of filtered wood smoke treatment on chemical and microbial changes in mahi mahi fillets. *J. Food Sci.* 2007, 72, C16–C24.
85. Hokkanen, M.; Luhtasela, U.; Kostamo, P.; Ritvanen, T.; Peltonen, K.; Jestoi, M. Critical effects of smoking parameters on the levels of polycyclic aromatic hydrocarbons in traditionally smoked fish

- and meat products in Finland. *J. Chem.* 2018, 2018, 2160958.
86. Bomfeh, K.; Jacxsens, L.; Amoa-Awua, W.K.; Tandoh, I.; Afoakwa, E.O.; Gamarro, E.G.; Ouadi, Y.D.; De Meulenaer, B. Reducing polycyclic aromatic hydrocarbon contamination in smoked fish in the Global South: A case study of an improved kiln in Ghana. *J. Sci. Food Agric.* 2019, 99, 5417–5423.
 87. Holley, R.A.; Patel, D. Improvement in shelf-life and safety of perishable foods by plant essential oils and smoke antimicrobials. *Food Microbiol.* 2005, 22, 273–292.
 88. Chatzikyriakidou, K.; Katsanidis, E. Effect of liquid smoke dipping and packaging method on the keeping quality of raw and cooked chub mackerel (*Scomber japonicus*) fillets. *J. Aquat. Food Prod. Technol.* 2012, 21, 445–454.
 89. Suñen, E.; Aristimuño, C.; Fernandez-Galian, B. Activity of smoke wood condensates against *Aeromonas hydrophila* and *Listeria monocytogenes* in vacuum-packaged, cold-smoked rainbow trout stored at 4 °C. *Food Res. Int.* 2003, 36, 111–116.
 90. Goulas, A.E.; Kontominas, M.G. Effect of salting and smoking-method on the keeping quality of chub mackerel (*Scomber japonicus*): Biochemical and sensory attributes. *Food Chem.* 2005, 93, 511–520.
 91. Kristinsson, H.G.; Crynen, S.; Yagiz, Y. Effect of a filtered wood smoke treatment compared to various gas treatments on aerobic bacteria in yellowfin tuna steaks. *LWT-Food Sci. Technol.* 2008, 41, 746–750.
 92. Rabiey, S.; Hosseini, H.; Rezaei, M. The hurdle effect of *Bunium persicum* essential oil, smoke and NaCl for controlling the *Listeria monocytogenes* growth in fish model systems. *J. Food Safety* 2013, 33, 137–144.
 93. Ceylan, Z.; Sengor, G.F.U.; Yilmaz, M.T. Nanoencapsulation of liquid smoke/thymol combination in chitosan nanofibers to delay microbiological spoilage of sea bass (*Dicentrarchus labrax*) fillets. *J. Food Eng.* 2018, 229, 43–49.
 94. Saloko, S.; Darmadji, P.; Setiaji, B.; Pranoto, Y. Antioxidative and antimicrobial activities of liquid smoke nanocapsules using chitosan and maltodextrin and its application on tuna fish preservation. *Food Biosci.* 2014, 7, 71–79.
 95. Sousa, W.M.; Silva, R.O.; Bezerra, F.F.; Bingana, R.D.; Barros, F.C.N.; Costa, L.E.C.; Sombra, V.G.; Soares, P.M.G.; Feitosa, J.P.A.; Paula, R.C.M.D. Sulfated polysaccharide fraction from marine algae *Solieria filiformis*: Structural characterization, gastroprotective and antioxidant effects. *Carbohydr. Polym.* 2016, 152, 140–148.
 96. Pane, G.; Cacciola, G.; Giacco, E.; Mariottini, G.L.; Coppo, E. Assessment of the antimicrobial activity of algae extracts on bacteria responsible of external otitis. *Mar. Drugs* 2015, 13, 6440–6452.

97. Wang, J.H.; Xu, J.L.; Zhang, J.C.; Liu, Y.; Sun, H.J.; Zha, X. Physicochemical properties and antioxidant activities of polysaccharide from floral mushroom cultivated in Huangshan Mountain. *Carbohydr. Polym.* 2015, 131, 240–247.
98. Smolskaitė, L.; Venskutonis, P.R.; Talou, T. Comprehensive evaluation of antioxidant and antimicrobial properties of different mushroom species. *LWT-Food Sci. Technol.* 2015, 60, 462–471.
99. Heleno, S.A.; Barros, L.; Martins, A.; Morales, P.; Fernández-Ruiz, V.; Glamoclija, J.; Sokovic, M.; Ferreira, I.C.F.R. Nutritional value, bioactive compounds, antimicrobial activity and bioaccessibility studies with wild edible mushrooms. *LWT-Food Sci. Technol.* 2015, 63, 799–806.
100. Pinapérez, M.C.; Rivas, A.; Martínez, A.; Rodrigo, D. Antimicrobial potential of macro and microalgae against pathogenic and spoilage microorganisms in food. *Food Chem.* 2017, 235, 34–44.
101. Shen, H.S.; Shao, S.; Chen, J.C.; Zhou, T. Antimicrobials from mushrooms for assuring food safety. *Compr. Rev. Food Sci. Food Safety* 2017, 16, 316–329.
102. Watson, S.B.; Cruz-Rivera, E. Algal chemical ecology: An introduction to the special issue. *Phycologia* 2003, 42, 319–323.
103. Wang, Y.; Xu, Z.; Bach, S.J.; Mcallister, T.A. Sensitivity of *Escherichia coli* to seaweed (*Ascophyllum nodosum*) phlorotannins and terrestrial tannins. *Asian Austral. J. Anim.* 2009, 22, 238–245.
104. Amorim, R.D.N.D.S.; Rodrigues, J.A.G.; Holanda, M.L.; Quinderé, A.L.G.; Paula, R.C.M.D.; Melo, V.M.M.; Benevides, N.M.B. Antimicrobial effect of a crude sulfated polysaccharide from the red seaweed *Gracilaria ornata*. *Braz. Arch. Biol. Technol.* 2012, 55, 171–181.
105. Bala, N.; Aitken, E.A.; Cusack, A.; Steadman, K.J. Antimicrobial potential of Australian macrofungi extracts against foodborne and other pathogens. *Phytother. Res.* 2012, 26, 465–469.
106. Borah, B.; Phukon, P.; Hazarika, M.P.; Ahmed, R.; Sarmah, D.K.; Wann, S.B.; Das, A.; Bhau, B.S. *Calamus leptospadix* Griff. a high saponin yielding plant with antimicrobial property. *Ind. Crop. Prod.* 2016, 82, 127–132.
107. Alcázar, M.; Kind, T.; Gschaedler, A.; Silveria, M.; Arrizon, J.; Fiehn, O.; Vallejo, A.; Higuera, I.; Lugo, E. Effect of steroidal saponins from *Agave* on the polysaccharide cell wall composition of *Saccharomyces cerevisiae* and *Kluyveromyces marxianus*. *LWT-Food Sci. Technol.* 2017, 77, 430–439.
108. Ribeiro, B.D.; Alviano, D.S.; Barreto, D.W.; Coelho, M.A.Z. Functional properties of saponins from sisal (*Agave sisalana*) and juá (*Ziziphus joazeiro*): Critical micellar concentration, antioxidant and antimicrobial activities. *Colloids Surf. A Physicochem. Eng. Asp.* 2013, 436, 736–743.

109. Ahmad, A.; Kaleem, M.; Ahmed, Z.; Shafiq, H. Therapeutic potential of flavonoids and their mechanism of action against microbial and viral infections—A review. *Food Res. Int.* 2015, 77, 221–235.
110. Tripoli, E.; Guardia, M.L.; Giammanco, S.; Majo, D.D.; Giammanco, M. Citrus flavonoids: Molecular structure, biological activity and nutritional properties: A review. *Food Chem.* 2007, 104, 466–479.
111. Seleem, D.; Pardi, V.; Murata, R.M. Review of flavonoids: A diverse group of natural compounds with anti-*Candida albicans* activity in vitro. *Arch. Oral Biol.* 2017, 76, 76–83.
112. Xie, Y.; Yang, W.; Tang, F.; Chen, X.; Ren, L. Antibacterial activities of flavonoids: Structure-activity relationship and mechanism. *Curr. Med. Chem.* 2014, 22, 132–149.
113. Zheng, Z. Ingredient technology for food preservation. *Ind. Biot.* 2014, 10, 28–33.
114. Ford, L.S.; Taylor, S.L.; Pacenza, R.; Niemann, L.M.; Lambrecht, D.M.; Sicherer, S.H. Food allergen advisory labeling and product contamination with egg, milk, and peanut. *J. Allergy Clin. Immun.* 2010, 126, 384–385.

Retrieved from <https://encyclopedia.pub/entry/history/show/26659>