

Pulse Wave Velocity

Subjects: [Nursing](#) | [Cardiac & Cardiovascular Systems](#)

Contributor: Irene Sequí-Domínguez , Iván Cavero-Redondo , Celia Álvarez-Bueno , Diana P Pozuelo-Carrascosa , Sergio Nuñez de Arenas-Arroyo , Vicente Martínez-Vizcaíno

Pulse wave velocity (PWV) is a simple, reproducible and non-invasive technique to assess arterial stiffness. It estimates the velocity of arterial wave propagation to travel a known distance between two anatomic sites within the arterial system ^[1]. PWV has been established as an independent predictor of cardiovascular mortality and is consequently used for diagnosis and prognosis in patients at risk of atherosclerotic cardiovascular disease ^[2]; however, PWV's significance relies on the fact that arterial stiffness, specially of muscular arteries can be modified by interventions that improve endothelial function such as exercise training, which makes PWV an important tool for cardiovascular risk management ^[3].

pulse wave velocity

cardiovascular risk

arterial stiffness

vascular function

1. Definition

Pulse wave velocity (PWV) is a simple, reproducible and non-invasive technique to assess arterial stiffness. It estimates the velocity of arterial wave propagation to travel a known distance between two anatomic sites within the arterial system ^[1]. PWV has been established as an independent predictor of cardiovascular mortality and is consequently used for diagnosis and prognosis in patients at risk of atherosclerotic cardiovascular disease ^[2]; however, PWV's significance relies on the fact that arterial stiffness, specially of muscular arteries can be modified by interventions that improve endothelial function such as exercise training, which makes PWV an important tool for cardiovascular risk management ^[3].

2. Introduction

Pulse wave velocity (PWV) is considered the gold standard method for assessing aortic stiffness ^{[4][5]}. Arterial stiffness measures, and carotid femoral PWV (cfPWV) in particular, are being included in the routine clinical assessment of patients and within the framework of large-scale clinical studies ^[4] as new instrumental solutions that allow the PWV assessment, such as photoplethysmography or magnetic resonance emerge ^[6] (Table 1). Nevertheless, an introduction into clinical practice has not been implemented further due to the fact that there is a lack of established reference values based on a large population and due to the absence of a standardized methodology for PWV assessment ^[7].

3. Methods used to Determine PWV

Table 1. Methods used to determine PWV. aPWV: aortic pulse wave velocity; baPWV: brachial-ankle pulse wave velocity; cfPWV: carotid-femoral pulse wave velocity; DVP: digital volume pulse; ECG: electrocardiogram; PWV: pulse wave velocity.

	Method	Description	Measure
Non-invasive methods	Applanation tonometry	Apply a pressure sensor through the skin and applanate a superficial artery by applying a downward pressure sufficient to flatten the artery.	baPWV, cfPWV
	Computerized oscillometry	Simultaneous acquisition and analysis of the pulsation of the artery, which is caused by the heart, as the pressure oscillation in the cuff.	Heart-brachial PWV, heart-ankle PWV, brachial-ankle PWV, cfPWV
	Mechanotransducer	Two dedicated piezoelectric pressure mechanotransducers directly applied to the skin in a simultaneous measurement of pressure pulses	carotid–femoral, carotid–brachial or femoral–dorsalis pedis PWV
	Ultrasound	Doppler pulses are recorded sequentially in 2 different arterial sites and compared using the R-wave of the ECG	baPWV, cfPWV
	Photoplethysmography	DVP measured by the photoplethysmography transducer	DVP associated with aPWV
	Magnetic Resonance Imaging	Assessment of the blood flow velocity with an enough temporal and spatial resolution to study the propagation of the aortic systolic flow wave	Local PWV

Invasive methods	Aortic angiography	Intra-aortic catheter measurements	Local PWV
------------------	--------------------	------------------------------------	-----------

Previous meta-analyses have attempted to calculate quantitative estimates of the predictive value of PWV for different outcomes. However, to the best of our knowledge, no previous meta-analysis has estimated the predictive performance (diagnostic odds ratio (dOR), sensitivity, specificity, positive likelihood ratio (PLR), and negative likelihood ratio (NLR)) of PWV considering the thresholds for a higher risk of cardiovascular or all-cause mortality estimated using hierarchical summary receiver operating characteristic (HSROC) models. Moreover, reference values for PWV have been established through cross-sectional studies [7] or expert consensus [8], in which subjects by age and blood pressure categories with no additional identified cardiovascular risk factors were considered.

References

1. James Oliver; David J. Webb; Noninvasive Assessment of Arterial Stiffness and Risk of Atherosclerotic Events. *Arteriosclerosis, Thrombosis, and Vascular Biology* **2003**, 23, 554-566, 10.1161/01.atv.0000060460.52916.d6.
2. Tomoki Shokawa; Michinori Imazu; Hideya Yamamoto; Mamoru Toyofuku; Naohito Tasaki; Tomokazu Okimoto; Kiminori Yamane; Nobuoki Kohno; Pulse Wave Velocity Predicts Cardiovascular Mortality. *Circulation Journal* **2005**, 69, 259-264, 10.1253/circj.69.259.
3. Michael F. O'Rourke; Junichiro Hashimoto; Arterial Stiffness. *Journal of Cardiopulmonary Rehabilitation and Prevention* **2008**, 28, 225-237, 10.1097/01.hcr.0000327179.21498.38.
4. G. Mancia; G. De Backer; Anna Dominiczak; Renata Cífková; R. Fagard; Giuseppe Germano; Guido Grassi; Anthony Heagerty; S. E. Kjeldsen; Stéphane Laurent; et al. Krzysztof Narkiewicz Luis Miguel Ruilope Andrzej Rynkiewicz R. E. Schmieder H. A. J. Struijker-Boudier A. Zanchetti Alec Vahanian John Camm Raffaele De Caterina Veronica Dean Kenneth Dickstein G. Filippatos Christian Funck-Brentano Irene Hellems Steen Dalby Kristensen Keith McGregor Udo Sechtem Sigmund Silber Michal Tendera Petr Widimsky J. L. Zamorano S. Erdine W. Kiowski E. Agabiti-Rosei E. Ambrosioni Lars H. Lindholm A. J. Manolis Peter M. Nilsson J. Redon M. Viigimaa Stamatis Adamopoulos Vicente Bertomeu Denis Clement Csaba Farsang Dan Gaiță Gregory Lip Jean-Michel Mallion E. O'Brien Piotr Ponikowski Frank Ruschitzka Juan Tamargo Pieter Van Zwieten Bernard Waeber Bryan Williams ESC Committee for Practice Guidelines (CPG): Esh Scientific Council: Document Reviewers: Athanasios J. Manolis 2007 Guidelines for the management of arterial hypertension: The Task Force for the Management of Arterial Hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). *European Heart Journal* **2006**, 28, 1462-1536, 10.1093/eurheartj/ehm236.

5. Raymond Townsend; Ian B. Wilkinson; Ernesto L. Schiffrin; Alberto P. Avolio; Julio A. Chirinos; John R. Cockcroft; Kevin S. Heffernan; Edward G. Lakatta; Carmel M. McEniery; Gary F. Mitchell; et al.Samer S. NajjarWilmer W. NicholsElaine M. UrbinaThomas WeberAmerican Heart Association Council on Hypertension Recommendations for Improving and Standardizing Vascular Research on Arterial Stiffness: A Scientific Statement From the American Heart Association.. *Hypertension* **2015**, 66, 698-722, 10.1161/HYP.0000000000000033.
6. Tania Pereira; Carlos Correia; João Cardoso; Novel Methods for Pulse Wave Velocity Measurement. *Journal of Medical and Biological Engineering* **2015**, 35, 555-565, 10.1007/s40846-015-0086-8.
7. Reference Values for Arterial Stiffness' Collaboration; The Reference Values for Arterial Stiffness' Collaboration; Determinants of pulse wave velocity in healthy people and in the presence of cardiovascular risk factors: 'establishing normal and reference values'.. *European Heart Journal* **2010**, 31, 2338-50, 10.1093/eurheartj/ehq165.
8. Luc M. Van Bortel; Stéphane Laurent; Pierre Boutouyrie; Phil Chowienczyk; J.K. Cruickshank; Tine De Backer; Jan Filipovsky; Sofie Huybrechts; Francesco U.S. Mattace-Raso; Athanase D Protogerou; et al.Giuseppe SchillaciPatrick SegersSebastian VermeerschThomas Weber Expert consensus document on the measurement of aortic stiffness in daily practice using carotid-femoral pulse wave velocity. *Journal of Hypertension* **2012**, 30, 445-448, 10.1097/hjh.0b013e32834fa8b0.

Retrieved from <https://encyclopedia.pub/entry/history/show/2769>