

# Biomedical Effects of Graviola

Subjects: [Food Science & Technology](#) | [Integrative & Complementary Medicine](#) | [Nutrition & Dietetics](#)

Contributor: Yamixa Delgado

A member of the Annonaceae (Custard-apple family), *Annona muricata*, commonly known as soursop, graviola, paw-paw, or “guanabana”, is a tree native to Central America and West Indies that is abundant at altitudes lower than 900 m above sea level.

Phytonutrients

Phytochemicals

Graviola

Lipinski's rule of 5

Veber's rules

Ghose filter

Anticancer

Antioxidant

Antimicrobial

Anti-inflammatory

Immunomodulatory

## 1. Botanical Description

It is cultivated in tropical and subtropical climates in countries such as Angola, Brazil, Colombia, Costa Rica, Puerto Rico, India, and Venezuela <sup>[1]</sup>. The graviola tree is mainly appreciated for its edible fruit. Still, its parts (leaves, fruit, bark, root, etc.) have been commonly used in traditional pharmacopeia in the form of macerations, decoction, or as a topical medication <sup>[2][3]</sup>. While the graviola tree can grow in a large variety of soils, it prefers deep soils with good oxygenation <sup>[1]</sup>. Botanically speaking, its leaves are large and obovate to elliptically shaped, are green on top, and paler under the top with short petioles and a pungent smell. The tree produces yellow-greenish flowers and lags about two years in producing heart-shaped fruits. It usually bears fruits yearly from that point on (12–24 per year) and can produce up to 50 fruits from its fifth year <sup>[4]</sup>.

## 2. Phytochemicals

More than two hundred (>200) bioactive compounds have been isolated from the leaves, seeds, root, bark, fruit, and fruit peel of the graviola tree <sup>[3]</sup>. Most frequently identified are alkaloids, phenolics, and terpenoids <sup>[5][6]</sup>. Acetogenins are considered the main bioactive compound in the *Annonaceae* family, with over 120 acetogenins identified from the root, leaves, stems, fruit pulp, and the seed of the family members <sup>[7][8]</sup>. Acetogenins are a particular class of secondary metabolites that could be considered part of the phenolics integrating polyketides and polyethers found exclusively in the plants of the *Annonaceae* family <sup>[9]</sup>. The structure of acetogenins is composed of a long carbon chain (35–38 carbons) as a fatty acid derivative. Graviola leaves contain key medically relevant polyphenolics compounds, including quercetin, rutin, and gallic acid <sup>[10][11][12]</sup>. The leaves of graviola also contain close to eighty (80) essential oils, including bioactive sesquiterpenes, and compounds such as potassium; calcium; zinc, phosphorus; magnesium; carbohydrates; vitamin A, B, and C; phytosterol; and calcium oxalate <sup>[13][14]</sup>.

## 3. Biomedical Effects

### 3.1. Anticancer

Graviola anticancer activity has been extensively studied, and the cytotoxicity of graviola has been reported for several cancer types e.g., breast, colorectal, skin, head and neck, lung, liver, pancreatic, prostate cancer, and leukemia [\[1\]\[15\]\[16\]\[17\]](#). Most of the antiproliferative properties of the extracts are suggested to be mediated by the graviola acetogenins. The acetogenins exert an inhibitory activity on the NADPH mitochondrial complex 1, a component of the energy transport chain, which is crucial to the synthesis of high quantities of ATP in cancer cells [\[16\]\[18\]\[19\]](#). Acetogenins have also been shown to target several critical cancer metabolic pathways by inhibiting the  $\text{Na}^+/\text{K}^+$  ATPase pump and the hypoxic and glycolytic pathways, inducing apoptosis and cell cycle arrest [\[19\]\[20\]\[21\]](#).

### 3.2. Antioxidant

Studies conducted in vitro and in vivo suggest that graviola contains antioxidant compounds that act as free-radical scavengers and increase the activity of the antioxidant enzymes superoxide dismutase and catalase and downregulate the function of mitochondrial NADPH oxidase complex I [\[22\]\[23\]\[24\]](#). The leaf and the fruit pulp of graviola are the parts of the tree with the highest antioxidant properties [\[5\]](#). The antioxidant activity of graviola is believed to be mediated by the following phenolic phytochemicals: quercetin, gallic acid, and graviola leaf polysaccharides [\[25\]\[26\]](#).

### 3.3. Antimicrobial

#### 3.3.1. Antiviral

It has been suggested that the phytochemicals polyphenolics in graviola exert some antiviral activity against RNA and DNA viruses [\[12\]\[27\]](#). A study by Wahab et al. showed that pretreating monkey kidney epithelial cells with a graviola leaf extract 24 h prior to infecting them with the dengue virus serotype 2 inhibited the virus replication. The treatment also increased the survival of the dengue-infected cells [\[28\]](#). A recent clinical study conducted by Le Donne et al. investigated the antiviral properties of graviola on human papillomavirus (HPV)-infected patients who were supplemented with ellagic acid and graviola extract twice a day for six months. Results showed a 74% HPV clearance in treated patients compared to the 25% clearance for the placebo group [\[29\]](#). Furthermore, recent in silico studies suggest that rutin, a phytonutrient abundant in graviola, could act as strong ligands and inhibit the function of proteins of the SARS-CoV and SARS-CoV-2 virus, thus suggesting potential therapeutic benefits against the COVID-19 infection [\[30\]\[31\]](#).

#### 3.3.2. Antibacterial

Graviola leaf extracts have been shown to exert in vitro antibacterial activity against oral pathogenic strains such as *S. mutants*, *S. mitis*, *P. gingivalis*, *P. intermedia*, *P. intermedia*, and *C. albicans* [\[32\]\[33\]](#). An in vivo study conducted in albino rats demonstrated the efficiency of graviola unripe fruit extracts to inhibit the growth of *S.*

*typhi* [34]. Furthermore, aqueous leaf extract and fruit-skin ethanolic graviola extracts showed a strong antibacterial effect against *K. pneumoniae*, *S. aureus*, and *P. aeruginosa* bacteria, i.e., the pathogens responsible for respiratory infections in the human immunodeficiency virus (HIV/AIDS) patients [35].

### 3.3.3. Antifungal

Researchers did not find studies testing the antifungal activity of any graviola extract. However, researchers found studies evaluating this property in some of graviola's phytochemicals. In 2017, a research group found that gallic acid has in vitro antifungal activity against dermatophyte strains (between 43.75 and 83.33 µg/mL), and *Candida* strains (*C. albicans* MIC = 12.5 µg/mL, and *Trichophyton rubrum* MIC = 43.75 µg/mL) by inhibiting the ergosterol synthesis. They also confirmed this activity after in vivo studies administering 80 mg/kg d of gallic acid [36]. In another study, researchers found that quercetin induces apoptosis in *Candida albicans* through mitochondrial dysfunction by increasing intracellular magnesium [37].

### 3.4. Anti-Inflammatory

The anti-inflammatory properties of graviola have been extensively studied in vitro and in vivo [38]. Cercato et al. reported that a topical application of a graviola leaf extract (0.3, 1, or 3 mg/ear) significantly reduced ear edema and myeloperoxidase activity in Swiss mice with 12-O-tetradecanoylphorbol-13-acetate (TPA)-induced ear inflammation. The authors were also able to show that the anti-inflammatory effect of the extract was associated with a reduction in the total amount of hydroperoxides and with modulation of catalase antioxidant activity [39]. While studying the anti-inflammatory response in lipopolysaccharide (LPS)-stimulated murine macrophage cell line RAW264.7 treated with graviola ethanolic leaf extracts, Laksmitawati et al. reported a downregulation in pro-inflammatory protein markers, such as tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ), interleukin-1 $\beta$ , interleukin-6, in the treated macrophages cells compared to untreated controls [40]. Furthermore, graviola aqueous extract suppresses nitric oxide production [41]. Similarly, an in vivo study conducted in rodents by Ishola et al. showed that the administration of a lyophilized graviola fruit extract inhibits the activity of the pro-inflammatory biomarkers cyclooxygenase (COX)-1 and COX-2 in a dose-dependent manner [42].

### 3.5. Immunomodulatory

As previously described, the specific bioactive constituents responsible for the major antioxidant, anti-inflammatory, and antimicrobial properties of graviola include different classes of annonaceous acetogenins (metabolites and products of the polyketide pathway), alkaloids, flavonoids, and sterols. Several studies have reported that graviola possesses immunomodulatory properties. For example, a study conducted in rodents by Umayra et al. shows that administration of an ethanolic graviola leaf extract boosts the immunological response through the activation of phagocytic cells [43]. Furthermore, an immune-enhancing activity of graviola leaf extracts has been observed in RAW 264.7 macrophage cells in vitro, a phenomenon which is believed to be mediated by the activation of the mitogen-activated protein kinase (MAPK) pathways [44].

### 3.6. Predicted gastrointestinal absorption (GIA)

The physicochemical properties for the main six bioactive phytochemicals in graviola (benzylisoquinoline, annonacin/acetogenin, cinnamic acid, coumaric acid, caffeic acid and rutin) were calculated based on the combination of Lipinski's, Ghose's, and Veber's rules (L-Ro5, GF, VR). The range of pharmacokinetics data for the molecules are summarized as follow: molecular weight (160-500 Da); hydrogen bond donors  $\leq 5$ ; hydrogen bond acceptors  $\leq 10$ ; molar refractivity (40-130); lipophilicity (-0.4–5.6); rotatable bonds  $\leq 10$ ; polar surface area  $< 140$ ; the total number of atoms (20-70), lipophilicity considering ionizable groups at pH 7.4 [LogD] (1-5) [45], [46], [47], [48]. 50 % of the graviola's bioactive phytochemicals (benzylisoquinoline, coumaric acid, caffeic acid) comply with all of the "drug-likeness" rules. The remaining 50% (annonacin/acetogenin, cinnamic acid, and rutin) violate most of the parameters of the L-Ro5, GF and VR. Accordingly, rutin is predicted to have the lowest GIA, followed by annonacin/acetogenin and then, cinnamic acid.

## 4. Contraindications

Studies from patients of the French West Indies, whose diet is rich in graviola, showed the development of a type of Parkinsonism resistant to the common anti-parkinsonism drug to treat tremors, levodopa [15]. This observation must lead scientists to further study if the acetogenins and alkaloids present in the graviola fruit could be toxic to brain cells, specifically dopaminergic neurons, which are the main cells affected in Parkinson's disease. In general, patients with neurological ailments should avoid the consumption of graviola supplements. Given that graviola is already widely used in traditional medicine, it could potentially be used against many health conditions if properly tested in further clinical studies. In conclusion, the benefits and side-effects of graviola should be carefully evaluated on a case-by-case basis.

This entry is adapted from [10.3390/app11188477](https://doi.org/10.3390/app11188477)

## References

1. Rethinam, P.; Sundararaj, P. *Annona muricata* L., soursop (graviola)-nature's gift to mankind with amazing medicinal benefits. *Int. J. Innov. Hortic.* 2016, 5, 73–80.
2. Chatrou, L.W.; Pirie, M.; Erkens, R.; Couvreur, T.; Neubig, K.M.; Abbott, J.R.; Mols, J.B.; Maas, J.W.; Saunders, R.; Chase, M.W. A new subfamilial and tribal classification of the pantropical flowering plant family Annonaceae informed by molecular phylogenetics. *Bot. J. Linn. Soc.* 2012, 169, 5–40.
3. Coria-Téllez, A.V.; Montalvo-González, E.; Yahia, E.M.; Obledo-Vázquez, E.N. *Annona muricata*: A comprehensive review on its traditional medicinal uses, phytochemicals, pharmacological activities, mechanisms of action and toxicity. *Arab. J. Chem.* 2018, 11, 662–691.
4. *Annona muricata* Sour Sop PFAF Plant Database. 2021. Available online: <https://pfaf.org/user/Plant.aspx?LatinName=Annona+muricata> (accessed on 7 July 2021).

5. Agu, K.C.; Okolie, P.N. Proximate composition, phytochemical analysis, and in vitro antioxidant potentials of extracts of *Annona muricata*(Soursop). *Food Sci. Nutr.* 2017, 5, 1029–1036.
6. Viraraghavan, S.S.N. Phytochemical screening of hydroalcohol fruit extract of *annona muricata*. *Int. J. Adv. Res. Dev.* 2018, 3, 681–685.
7. Anaya Esparza, L.M.; Montalvo-González, E. Bioactive compounds of soursop (*annona muricata* L.) fruit. In *Bioactive Com-Pounds in Underutilized Fruits and Nuts*; Murthy, H.N., Bapat, V.A., Eds.; Springer International Publishing: Cham, Switzerland, 2020; pp. 175–189.
8. Rady, I.; Bloch, M.; Chamcheu, R.-C.N.; Mbeumi, S.B.; Anwar, R.; Mohamed, H.; Babatunde, A.S.; Kuate, J.-R.; Noubissi, F.K.; El Sayed, K.A.; et al. Anticancer Properties of Graviola (*Annona muricata*): A Comprehensive Mechanistic Review. *Oxidative Med. Cell. Longev.* 2018, 2018, 1–39.
9. Simpson, D.; Amos, S. Other plant metabolites. In *Pharmacognosy*; Badal, S., Delgoda, R., Eds.; Academic Press: Boston, MA, USA, 2017; Chapter 12; pp. 267–280.
10. Jan, A.T.; Kamli, M.R.; Murtaza, I.; Singh, J.B.; Ali, A.; Haq, Q. Dietary Flavonoid Quercetin and Associated Health Benefits—An Overview. *Food Rev. Int.* 2010, 26, 302–317.
11. Kahkeshani, N.; Farzaei, F.; Fotouhi, M.; Alavi, S.S.; Bahramsoltani, R.; Naseri, R.; Momtaz, S.; Abbasabadi, Z.; Rahimi, R.; Farzaei, M.H.; et al. Pharmacological effects of gallic acid in health and disease: A mechanistic review. *Iran. J. Basic Med. Sci.* 2019, 22, 225–237.
12. Seo, D.J.; Choi, C. Viral disease and use of polyphenolic compounds. In *Polyphenols: Prevention and Treatment of Human Disease*, 2nd ed.; Watson, R.R., Preedy, V.R., Zibadi, S., Eds.; Academic Press: Cambridge, MA, USA, 2018; Chapter 25; pp. 301–312.
13. Enweani, I.; Obroku, J.; Enahoro, T.; Omoifo, C. The biochemical analysis of soursop (*annona muricata* L.) and sweetsop (*a.Squamosa* L.) and their potential use as oral rehydration therapy. *J. Food Agric. Environ.* 2004, 22, 39–43.
14. Gyesei, J.N.; Opoku, R.; Borquaye, L.S. Chemical Composition, Total Phenolic Content, and Antioxidant Activities of the Essential Oils of the Leaves and Fruit Pulp of *Annona muricata* L. (Soursop) from Ghana. *Biochem. Res. Int.* 2019, 2019, 1–9.
15. Lannuzel, A.; Michel, P.P.; Caparros-Lefebvre, D.; Abaul, J.; Hocquemiller, R.; Ruberg, M. Toxicity of Annonaceae for dopaminergic neurons: Potential role in atypical parkinsonism in Guadeloupe. *Mov. Disord.* 2002, 17, 84–90.
16. Deep, G.; Kumar, R.; Jain, A.K.; Dhar, D.; Panigrahi, G.K.; Hussain, A.; Agarwal, C.; El-Elmat, T.; Sica, V.P.; Oberlies, N.H.; et al. Graviola inhibits hypoxia-induced NADPH oxidase activity in prostate cancer cells reducing their proliferation and clonogenicity. *Sci. Rep.* 2016, 6, 23135.
17. Qazi, A.K.; Siddiqui, J.; Jahan, R.; Chaudhary, S.; Walker, L.; Sayed, Z.; Jones, D.T.; Batra, S.K.; Macha, M. Emerging therapeutic potential of graviola and its constituents in cancers.

- Carcinogenesis 2018, 39, 522–533.
18. McLaughlin, J.L. Paw Paw and Cancer: Annonaceous Acetogenins from Discovery to Commercial Products. *J. Nat. Prod.* 2008, 71, 1311–1321.
  19. Torres, M.P.; Rachagani, S.; Purohit, V.; Pandey, P.; Joshi, S.; Moore, E.D.; Johansson, S.L.; Singh, P.K.; Ganti, A.K.; Batra, S.K. Graviola: A novel promising natural-derived drug that inhibits tumorigenicity and metastasis of pancreatic cancer cells in vitro and in vivo through altering cell metabolism. *Cancer Lett.* 2012, 323, 29–40.
  20. Chung, I.; Yap, C.V.; Subramaniam, K.S.; Khor, S.W. Annonacin exerts antitumor activity through induction of apoptosis and extracellular signal-regulated kinase inhibition. *Pharmacogn. Res.* 2017, 9, 378–383.
  21. Yiallouris, A.; Patrikios, I.; Johnson, E.O.; Sereti, E.; Dimas, K.; de Ford, C.; Fedosova, N.U.; Graier, W.F.; Sokratous, K.; Kyriakou, K.; et al. Annonacin promotes selective cancer cell death via nka-dependent and serca-dependent pathways. *Cell Death Dis.* 2018, 9, 764.
  22. Byun, E.-B.; Song, H.-Y.; Kim, W.S. Polysaccharides from *Annona muricata* leaves protect normal human epidermal keratinocytes and mice skin from radiation-induced injuries. *Radiat. Phys. Chem.* 2020, 170, 108672.
  23. George, V.C.; Kumar, D.; Rajkumar, V.; Suresh, P.; Kumar, R.A. Quantitative assessment of the relative antineoplastic potential of the n-butanolic leaf extract of *annona muricata* linn. In normal and immortalized human cell lines. *Asian Pac. J. Cancer Prev.* 2012, 13, 699–704.
  24. Zamudio-Cuevas, Y.; Díaz-Sobac, R.; Vázquez-Luna, A.; Landa-Solís, C.; Cruz-Ramos, M.; Santamaría-Olmedo, M.; Martínez-Flores, K.; Fuentes-Gómez, A.J.; López-Reyes, A. The antioxidant activity of soursop decreases the expression of a member of the NADPH oxidase family. *Food Funct.* 2014, 5, 303–309.
  25. Baskar, R.; Rajeswari, V.; Sathishkumar, T. In vitro antioxidant studies in leaves of *Annona* species. *Indian J. Exp. Boil.* 2007, 45, 480–485.
  26. Biba, V.; Amily, A.; Sangeetha, S.; Remani, P. Anticancer, antioxidant and antimicrobial activity of annonaceae family. *World J. Pharm. Pharm. Sci.* 2014, 3, 1595–1604.
  27. Padma, P.; Pramod, N.P.; Thyagarajan, S.P.; Khosa, R.L. Effect of the extract of *annona muricata* and *petunia nyctagini-flora* on herpes simplex virus. *J. Ethnopharmacol.* 1998, 61, 81–83.
  28. Wahab, N.Z.A.; Ibrahim, N.; Kamarudin, M.K.A.; Lananan, F.; Juahir, H.; Ghazali, A.; Ireana Yusra, A.F. Cytotoxicity and antiviral activity of *annona muricata* aqueous leaves extract against dengue virus type. *J. Fundam. Appl. Sci.* 2018, 10, 580–589.
  29. Le Donne, M.; Lentini, M.; Alibrandi, A.; Salimbeni, V.; Giuffre', G.; Mazzeo, F.; Triolo, O.; D'Anna, R. Antiviral activity of Ellagic acid and *Annona Muricata* in cervical HPV related pre-neoplastic

- lesions: A randomized trial. *J. Funct. Foods* 2017, 35, 549–554.
30. Elmi, A.; Sayem, S.A.-J.; Ahmed, M.; Abdoul-Latif, F. NATURAL COMPOUNDS FROM DJIBOUTIAN MEDICINAL PLANTS AS INHIBITORS OF COVID-19 BY IN SILICO INVESTIGATIONS. *Int. J. Curr. Pharm. Res.* 2020, 52–57.
  31. Jianfeng, Y.; Shengxi, S.; Bin, L.; Zhihao, W.; Yi-Zhou, J.; Yunqing, L.; Feng, C.; Bing, L. Emergency antiviral drug discovery during a pandemic-a case study on the application of natural compounds to treat covid-19. *ChemiRxiv* 2020.
  32. Pai, B.M.; Rajesh, G.; Shenoy, R.; Rao, A. Anti-microbial efficacy of soursop leaf extract (*annona muricata*) on oral patho-gens: An in-vitro study. *JCDR* 2016, 10, ZC01–ZC04.
  33. Rodriguez-Perez, J.L.; A Millones-Gomez, P. Antibacterial Effect of *Annona muricata* L. Leaves on *Streptococcus mutans* ATCC 25175 Strains. *J. Clin. Diagn. Res.* 2019, 13.
  34. Dada, E.O.; Faleye, O.S. In-vitro and in-vivo Activities of the Ethanol Pulp Extract of *Annona muricata* (Linn) Fruit in Albino Rats Infected with *Salmonella typhi*. *J. Adv. Med. Pharm. Sci.* 2016, 5, 1–12.
  35. Iyanda-Joel, W.O.; Omonigbehin, E.A.; Iweala, E.E.J.; Chinedu, S.N. Antibacterial studies on fruit-skin and leaf extracts of *Annona muricata* in Ota, Nigeria. *IOP Conf. Ser. Earth Environ. Sci.* 2019, 331, 012029.
  36. Li, Z.-J.; Liu, M.; Dawuti, G.; Dou, Q.; Ma, Y.; Liu, H.-G.; Aibai, S. Antifungal Activity of Gallic Acid In Vitro and In Vivo. *Phytother. Res.* 2017, 31, 1039–1045.
  37. Kwun, M.S.; Lee, D.G. Quercetin-induced yeast apoptosis through mitochondrial dysfunction under the accumulation of magnesium in *Candida albicans*. *Fungal Biol.* 2020, 124, 83–90.
  38. Wahab, S.M.A.; Jantan, I.; Haque, A.; Arshad, L. Exploring the Leaves of *Annona muricata* L. as a Source of Potential Anti-inflammatory and Anticancer Agents. *Front. Pharmacol.* 2018, 9, 661.
  39. Cercato, L.M.; Araújo, J.M.D.; Oliveira, A.S.; Melo, A.J.O.; Lima, B.S.; dos Santos, E.W.P.; Dos S Neto, A.G.; de Albu-querque-Júnior, R.L.C.; Duarte, M.C.; Araujo, A.A.S. Reduced cutaneous inflammation associated with antioxidant action after topical application of the aqueous extract of *annona muri-cata* leaves. *Inflammopharmacology* 2021, 29, 307–315.
  40. Laksmiawati, D.R.; Prasanti, A.P.; Larasinta, N.; Syauta, G.A.; Hilda, R.; Ramadaniati, H.U.; Widyastuti, A.; Karami, N.; Afni, M.; Rihibiha, D.D.; et al. Anti-Inflammatory Potential of Gandarusa (*Gendarussa vulgaris* Nees) and Soursoup (*Annona muricata* L) Extracts in LPS Stimulated-Macrophage Cell (RAW264.7). *J. Nat. Remedies* 2016, 16, 73–81.
  41. Choi, M.; Kang, Y.-R.; Zu, H.D.; Lim, I.-S.; Jung, S.K.; Chang, Y.H. Effects of Time on Phenolics and in vitro Bioactivity in Autoclave Extraction of Graviola (*Annona muricata*) Leaf. *Biotechnol. Bioprocess. Eng.* 2020, 25, 9–15.

42. Ishola, I.O.; Awodele, O.; Olusayero, A.M.; Ochieng, C.O. Mechanisms of Analgesic and Anti-Inflammatory Properties of *Annona muricata* Linn. (Annonaceae) Fruit Extract in Rodents. *J. Med. Food* 2014, 17, 1375–1382.
43. Umayra, A.N.; Qasim, M.J.; Al-Yasari, A.K.S.; Dakheel, M.H. Immunological Role of *Annona muricata* (Dietary Supplement of graviola) Against Cypermethrin's Toxic Effects in Female Rats. *Ann. Trop. Med. Public Health* 2020, 23, 103–109.
44. Kim, G.-T.; Tran, N.K.S.; Choi, E.-H.; Song, Y.-J.; Song, J.-H.; Shim, S.-M.; Park, T.-S. Immunomodulatory efficacy of standardized *annona muricata* (graviola) leaf extract via activation of mitogen-activated protein kinase pathways in raw 264.7 macrophages. *Evid. Based Complementary Altern. Med.* 2016, 2016, 2905127.

---

Retrieved from <https://encyclopedia.pub/entry/history/show/37141>