

Irrigation-Water and Cotton

Subjects: **Agricultural Engineering**

Contributor: Komlan Koudahe , Koffi Djaman

A decrease in water resources, as well as changing environmental conditions, calls for efficient irrigation-water management in cotton-production systems. Cotton (*Gossypium* sp.) is an important cash crop in many countries, and it is used more than any other fiber in the world.

cotton

water management

yield

water productivity

1. Introduction

The global production of cotton fiber is estimated to be 24.65 million tons ^[1]. In 2018, the production was estimated to be 6.71 million tons in the Americas; 0.38 million tons in Europe; 1.56 million tons in Africa; 0.95 million tons in Oceania; and 15.06 million tons in Asia ^[1]. In 2020, across the United States, the production of cotton was estimated to be 3.26 million tons, and cotton was grown on about 3.52 million hectares ^[2]. The United States is the world's leading cotton exporter, supplying about 35% of global cotton exports in recent years ^[3]. Cotton is a valuable, natural-textile fiber and the purest source of cellulose ^[4]. Aside from the fibers, cotton also is produced as a source of seeds that provide edible oil, seed by-products, and other products to industries. Its residues provide organic matter to the soil. Zhang and Dong ^[5] estimated that cotton fiber contributes to half of the world's clothes.

Therefore, improved irrigation-water management practices that optimize the lint and seed yield of cotton and promote water-use efficiency, while maintaining maximum quality, are critical for the future sustainability of cotton production.

2. Cotton Water-Use Efficiency

The concept of water-use efficiency (WUE) was introduced more than 100 years ago by Briggs and Shantz ^[6] indicating a relationship between plant productivity and water use. Cotton lint yield is found to rise with increasing crop water use ^[7]. Hatfield and Dold ^[8] defined WUE as the quantity of assimilated carbon in terms of biomass or grain per unit of water used by the crop. In plant breeding, it has been proposed to use WUE to select water-use-efficient genotypes under changing environments, heat and water stress, and the interactions between them. Research results have revealed variations between genotypes for WUE in upland cotton and pima cotton ^{[9][10][11]}. Snowden et al. ^[12] carried out a study on WUE and irrigation response of cotton cultivars under subsurface drip irrigation in West Texas, USA. They reported that WUE differed among the six cultivars and the deficit strategies used. In 2010, the cotton variety FM9160 had the greatest WUE of 0.20 kg m^{-3} under severe deficit irrigation; DP1044 had the greatest WUE of 0.32 kg m^{-3} under mild deficit irrigation; and DP0912 had the greatest WUE of

0.33 kg m⁻³ under full irrigation (Table 1). Among the irrigation regimes, full irrigation provided the highest WUE while severe deficit gave the lowest WUE in 2010. Moreover, a study conducted in Australia found that the water-use efficiency increased by 40% over a ten-year period along with developments in plant breeding, the utilization of genetically modified varieties, and improved water management practices, and they resulted in yield increases [13]. These results are in line with the findings of Hatfield and Dold [8] who reported that WUE is dependent upon genotype and management practices. Evett et al. [14] reported that evapotranspiration water-use efficiency (ETWUE) ranged from 0.15 kg/m³ to 0.33 kg/m³. The improvement in ETWUE is most probably attributable to increased yield as well as to reduced soil evaporation and transpiration. It was deduced that management practices that lessen soil water evaporation and move the water for crop water use (more transpiration) reduce crop exposure to water stress and maintain water-use efficiency at the maximum level possible. For instance, Hatfield and Dold [8] reported that the adoption of drip irrigation reduced by 23% wheat water use, but, at the same time, it improved yield by 37%. In cotton, this practice reduced water use by 37% and diminished yield by 21%. Therefore, the use by farmers of micro-irrigation systems, such as drip-irrigation, lessens not only the soil water evaporation from between plant rows early in the season but also prevents almost all the evaporation component from the canopy. These management practices have a positive effect on WUE in areas where crops are micro-irrigated and show that WUE can be improved by water management practices.

Table 1. Irrigation regimes, yield, and water-use efficiency (WUE) (SE—severe deficit, MD—mild deficit, F.Irr—full irrigation, DI—drip irrigation, SDI—subsurface drip irrigation, CEF—closed-end furrow).

Location	Year	Watering Regime	Water-Use Efficiency (kg m ⁻³)				Cotton Variety References
			Lint	Seed	WUE	IWUE	
West Texas, USA	2010	SE	712		0.20		FM9160
		MD	1436		0.32		DP1044
		F.Irr	1743		0.33		DP0912
	2011	SE	596		0.15		DP0935
		MD	1268		0.23		DP1044
		F.Irr	1537		0.22		DP0935
Texas, California and Uzbekistan		DI			0.15– 0.33		Evett et al. [14]
Turkey	2016– 2017	SDI	4082	0.83	0.84		Cetin and Kara, [15]
Arizona, USA	1988– 1999	N/A ¹	1280– 1420		0.127– 0.138	upland cotton	Grismar [16]

Location	Year	Watering Regime	Watering Yield (kg ha ⁻¹)		Water-Use Efficiency (kg m ⁻³)		Cotton Variety References
			Lint	Seed	WUE	IWUE	
California, USA	1988–1999	N/A	910–120		0.09–0.109		pima cotton
			1110–1440		0.134–0.210		upland cotton
			1170–1340		0.151–0.177		pima cotton
Bushland, Texas, USA	2003	MESA 100%	1229		0.164	0.492	Paymaster2280 BG RR
		MESA 75%	1001		0.142	0.491	
		MESA 50%	536		0.089	0.288	
		MESA 25%	213		0.045	0.024	
		LESA 100%	1208		0.160	0.482	
		LESA 75%	984		0.143	0.480	
		LESA 50%	575		0.098	0.321	
		LESA 25%	288		0.058	0.130	
		LEPA 100%	1153		0.158	0.456	
		LEPA 75%	1149		0.164	0.581	
		LEPA 50%	685		0.109	0.415	
		LEPA 25%	362		0.072	0.234	
		SDI 100%	1150		0.159	0.454	

Location	Year	Watering Regime	Water-Use Efficiency (kg m ⁻³)			Cotton Variety References	
			Lint	Seed	WUE		
Turkey	2013	SDI 75%	1082		0.152	0.540	
		SDI 50%	844		0.135	0.549	
		SDI 25%	491		0.092	0.416	
		CEF 100%		5640	0.64	0.81	
		CEF 70%		4460	0.63	0.91	
	2014	CEF 50%		3720	0.64	1.06	
		CEF 30%		3210	0.71	1.52	
		CEF(0%)		1820	0.67	-	Nazilli-84
	2014	CEF 100%		5340	0.62	0.74	Dagdelen et al. [18]
		CEF 70%		3990	0.62	0.79	
		CEF 50%		3590	0.73	0.99	
Bornova- Izmir, Turkey	1992– 1994	CEF 30%		2800	0.74	1.29	
				1740	0.72	-	
		CEF 0%					N84
		Furrow			0.38– 0.46	0.48– 0.65	Anac et al. [19]
Australia	1991– 1994	Drip		2.23			Hodgson et al. [20]
		Furrow		1.89			
		Drip		0.487			
Anatolia, Turkey	1991– 1994	Furrow		1	0.387		Sayar-314
		Sprinkler			0.236		Cetin and Bidgel [21]

Similarly, Even et al. [14] revealed in several experimental studies at different locations of Texas and California in USA that water productivity (lint/evapotranspiration) and lint yield were improved by adopting drip-irrigation systems instead of furrow irrigation. In the same line, Fan et al. [24] found from a metadata analysis the highest cotton evapotranspiration water-use efficiency of 0.88 kg/m³, and this can be achieved by lessening by 5.5% the crop

Location	Year	Watering Regime	Watering Yield (kg ha^{-1})		Water-Use Efficiency (kg m^{-3})		Cotton Variety	References
			Lint	Seed	WUE	IWUE		
Harran plain, Turkey		LEPA			0.55– 0.67	0.58– 0.77	Yazar et al. [22]	optimum CWP) for on actual, it varied
		Drip			0.50– 0.74	0.60– 0.81		
Cukurova, Turkey		Furrow			1.9– 5.9	1.5– 5.1	Kanber et al. [23]	$\text{kg ha}^{-1} \text{ mm}^{-1}$ and [18] reported

WUE values varied from 1.59 to 2.30 kg m^{-3} for corn and from 0.61 to 0.72 kg m^{-3} for cotton in two years. WUE values of 0.38–0.46 kg m^{-3} were obtained by Anac et al. [19] in the coastal part of the Aegean region, Turkey.

It is important to highlight that WUE varies also according to the irrigation technology used. Some irrigation devices are found to limit water to the root zone, while others provide water to all the soil surface. Hodgson et al. [20] compared furrow and drip-irrigation methods for cotton and found that the WUEs were 2.23 and 1.89 kg m^{-3} for drip and furrow irrigation methods, respectively. Under drip, furrow, and sprinkler irrigation, Cetin and Bidgel [21] found water-use efficiencies of 4.87, 3.87, and 2.36 kg/ha-mm , respectively, proving that drip irrigation provides a greater yield per unit drop. Yazar et al. [22] reported that WUE values of cotton irrigated by LEPA and the drip method were, respectively, 0.55–0.67 kg m^{-3} and 0.50–0.74 kg m^{-3} in the Harran Plain in Turkey. Moreover, Kanber et al. [23] determined WUE values of 1.9–5.9 $\text{kg ha}^{-1} \text{ mm}^{-1}$ under furrow irrigation in the Cukurova Plain in southern Turkey and found irrigation water-use efficiency (IWUE) values for furrow irrigated cotton ranged from 1.5 to 5.1 kg m^{-3} . According to Anac et al. [19], IWUE values were 0.48–0.65 kg m^{-3} . In addition, IWUE values for LEPA and drip-irrigated cotton were 0.58–0.77 kg m^{-3} and 0.60–0.81 kg m^{-3} , respectively, in the Harran Plain of Turkey [22]. Ertek and Kanber [25] determined IWUE values for drip-irrigated cotton of 0.75–0.94 kg m^{-3} in the Cukurova Plain in Turkey. In Queensland, Australia, furrow irrigation has been optimized and tested in the field for cotton. Results showed an increase in WUE and a decline in labor requirement [26]. The water-use efficiency fluctuates between farming fields and across regions due to many factors. Therefore, site-specific measurements are crucial for decision making and improvements in WUE.

3. Cotton Yield and Yield Components under Different Irrigation Techniques

Cotton can be cultivated under rainfed conditions only in a limited number of regions, and usually an optimum yield cannot be achieved without irrigation [21]. Therefore, irrigation is necessary for cotton production. For instance, in the Mississippi Delta region, USA, Pinnamaneni et al. [27] reported that irrigation is a crucial factor in achieving both high fiber yield and seed quality, while Sui et al. [28] found out that irrigation augmented cotton yield and improved fiber length. Different irrigation technologies are widely used to produce cotton, with most common being:

- low-energy precision application (LEPA),

- low-elevation spray application (LESA),
- mid-elevation spray application (MESA),
- mobile drip irrigation (MDI),
- surface irrigation (SI),
- subsurface drip irrigation (SDI), and
- furrow irrigation (FI).

Various results have been obtained under different irrigation practices depending on local climates, soil conditions, genotypes, and management practices. In northern Texas and southwestern Kansas, USA, Colaizzi et al. [17] carried out a study in 2003 on cotton production with surface drip irrigation (SDI), LEPA, and spray irrigation, and found that the highest lint yield and water-use efficiency were achieved with SDI at low irrigation rates. Similar results were found by Colaizzi et al. [29] and Segarra et al. [30], who reported that SDI performed better than any other spray irrigation system (MESA, LESA, and LEPA). Moreover, the same study in 2004 revealed that lint yield and gross returns were improved with SDI at any irrigation rate. Bordovsky [31] found that under irrigation treatments with less than 50% of full irrigation, LEPA induced a 16% yield increase over sprinkler irrigation, but SDI resulted in a 14% higher yield over LEPA. At irrigation levels greater than 50% of full irrigation, yield was slightly smaller in sprinkler compared to LEPA, and SDI was found to provide a 7% greater yield than LEPA. However, Bordovsky et al. [32] carried out a study where soil matric potential was used to schedule irrigation and found that LEPA and drip irrigation provided the same yields for cotton, corn, and soybeans.

In Turkey, Cetin and Bidgel [21] carried out a study with three different irrigation methods on seed cotton yield and yield components and reported that maximum seed yield was 4380, 3630, and 3380 kg/ha under drip, furrow, and sprinkler irrigation, respectively. Drip irrigation generated 21% more yield than furrow irrigation and 30% more yield than sprinkler irrigation. In southeastern Turkey, Cetin et al. [33] did a similar study and compared different irrigation methods for effective water use on cotton. The highest seed cotton yield was found in drip-irrigated plots, and it was 4650 kg ha^{-1} . It was followed by furrow irrigation, which had a yield of 3120 kg ha^{-1} . In terms of lint yield, lint quality, and water-use efficiency, SDI has been found to slightly surpass LEPA and spray irrigation [30][34]. In India, Choudhary et al. [35] found that drip irrigation increased plant height, number of bolls per plant, boll weight, and number of monopods and sympods per plant. Further, water-use efficiency was greatest under drip irrigation as compared to other irrigation systems in all four cotton cultivars that Choudhary et al. [35] studied. According to Sezan et al. [36], for cotton production drip irrigation was more advantageous compared to conventional practices of irrigation. In China, Wang et al. [37] compared traditional flood irrigation and mulched drip irrigation and found that mulched drip irrigation promoted the root growth of cotton and improved the production of fine roots after the full-boll stage. The boll number per plant and yield were increased with mulched drip irrigation.

Drip irrigation has been found to be the most effective water-saving system. It can conserve soil, aggregate structure, successfully prevent deep water loss and surface water loss, and therefore, decrease exposure of the soil to degradation and salinization [38][39][40][41]. Fereres et al. [42] reported that an early and increased cotton yield could be achieved by drip irrigation. Mateos et al. [43] stated that drip irrigation was more beneficial than furrow irrigation. In the same line, Ibragimov et al. [44] in Uzbekistan reported that, with drip irrigation used for cotton production, 18–42% of the irrigation water was saved in contrast to furrow irrigation. According to Ward and Pulido-Velazquez [45], compared to flood irrigation, drip irrigation increased cotton yields by about 25% and helped to save water by 40–50%. In the Harran Plain in Turkey, Cetin and Bilgel [21] found that drip irrigation improved seed cotton yield by 21 and 30% over furrow and sprinkler irrigation, respectively. Similarly, in the Texas High Plain, Colaizzi et al. [46] showed that SDI had the best cotton productivity and gross returns, followed by LEPA and spray irrigation. However, Cetin and Kara [15] reported that the use of SDI is limited, because it has adverse effects on cotton seed germination, if during sowing there is no moisture in the soil. For this reason, an alternative irrigation technology, such as sprinkler irrigation, is advised for better cotton germination.

Field-based studies are critical to identify a technology that can provide an optimum yield and quality of cotton, and, at the same time, maximize water-use efficiency.

References

1. FAOSTAT. 2021. Available online: <http://www.fao.org/faostat/en/#data/QCL>. (accessed on 25 July 2021).
2. USDA. Crop Production 2020; National Agricultural Statistics Service; USDA: Washington, DC, USA, 2021; 123p. Available online: <https://downloads.usda.library.cornell.edu/usda-esmis/files/k3569432s/w3764081j/5712n018r/cropan21.pdf> (accessed on 6 May 2021).
3. OECD/FAO. “OECD-FAO Agricultural Outlook”, OECD Agriculture Statistics (Database). 2019. Available online: <http://www.agri-outlook.org/commodities/Cotton.pdf> (accessed on 8 May 2021).
4. Hsieh, Y.L. Chemical structure and properties of cotton. In Cotton: Science and Technology; Chapter 1; Woodhead: Sawston, UK, 2007; 34p.
5. Zhang, Y.; Dong, H. Yield and fiber quality of cotton. *Encycl. Renew. Sustain. Mater.* 2020, 2, 356–364.
6. Briggs, L.J.; Shantz, H.L. The Water Requirement of Plants II—A Review of the Literature; U.S. Department of Agriculture: Washington, DC, USA, 1913; pp. 1–96.
7. Baker, J.T.; Dennis, C.; Gitz, D.C.; John, E.; Stout, J.E.; Lascano, R.J. Cotton water use efficiency under two different deficit irrigation scheduling methods. *Agronomy* 2015, 5, 363–373.
8. Hatfield, J.L.; Dold, C. Water-Use Efficiency: Advances and Challenges in a Changing Climate. *Front. Plant Sci.* 2019, 10, 103.

9. Quisenberry, J.E.; McMichael, B.L. Genetic variation among cotton germplasm for water-use efficiency. *Environ. Exp. Bot.* 1991, 31, 453–460.
10. Fish, D.A.; Earl, H.J. Water-use efficiency is negatively correlated with leaf epidermal conductance in cotton (*Gossypium* spp.). *Crop Sci.* 2009, 49, 1409–1415.
11. Saranga, Y.; Jiang, C.X.; Wright, R.J.; Yakir, D.; Paterson, A.H. Genetic dissection of cotton physiological responses to arid conditions and their inter-relationships with productivity. *Plant Cell Environ.* 2004, 27, 263–277.
12. Snowden, C.; Ritchie, G.; Thompson, T. Water Use Efficiency and Irrigation Response of Cotton Cultivars on Subsurface Drip in West Texas. *J. Cotton Sci.* 2013, 17, 1–9.
13. Roth, G.; Harris, G.; Gillies, M.; Montgomery, J.; Wigginton, D. Water-use efficiency and productivity trends in Australian irrigated cotton: A review. *Crop Past. Sci.* 2013, 64, 1033–1048.
14. Evett, S.R.; Baumhardt, R.L.; Howell, T.A.; Ibragimov, N.M.; Hunsaker, D.J. Cotton. *Crop Yield Response to Water; FAO irrigation and drainage paper. No. 66*; FAO: Rome, Italy, 2012; pp. 152–161.
15. Cetin, O.; Kara, A. Assessment of water productivity using different drip irrigation systems for Cotton. *Agric. Water Manag.* 2019, 223, 105693.
16. Grismer, M.E. Regional cotton lint yield, ET_C, and water value in Arizona and California. *Agric. Water Manag.* 2002, 54, 227–242.
17. Colaizzi, P.D.; Evett, S.R.; Howell, T.A. Cotton production with SDI, LEPA, and spray irrigation in a thermally-limited climate. In *Proceedings of the Irrigation Association Conference*, Phoenix, Arizona, 6–8 November 2005.
18. Dagdelen, N.; Yılmaz, E.; Sezgin, F.; Gurbuz, T. Water-yield relation and water use efficiency of cotton (*Gossypium hirsutum* L.) and second crop corn (*Zea mays* L.) in western Turkey. *Agric. Water Manag.* 2006, 82, 63–85.
19. Anac, S.; Ul, M.A.; Tuzel, I.H.; Anac, D.; Okur, B.; Hakerler, H. Optimum irrigation scheduling for cotton under deficit irrigation conditions. In *Crop Yield Response to Deficit Irrigations*; Kırda, C., Moutonnet, P., Hera, C., Nielsen, D.R., Eds.; Kluwer Academic Publishers: Dordrecht, The Netherlands, 1999.
20. Hodgson, A.S.; Constable, G.A.; Duddy, G.R.; Danieles, I.G. A comparison of drip and furrow irrigated cotton on a cracking clay soil. *Soils Fertil.* 1992, 55, 2.
21. Cetin, O.; Bilgel, L. Effects of different irrigation methods on shedding and yield of cotton. *Agric. Water Manag.* 2002, 54, 1–15.
22. Yazar, A.; Sezen, S.M.; Sesveren, S. LEPA and trickle irrigation of cotton in the Southeast Anatolia Project (GAP) area in Turkey. *Agric. Water Manag.* 2002, 54, 189–203.

23. Kanber, R.; Onder, S.; Unlu, M.; Koksal, H.; Ozekici, B.; Sezen, S.M.; Yazar, A.; Koc, K. Optimization of surface irrigation methods for cotton and comparison with siprnikler irrigation. In Research Report No: 18, GAP Research Projects. Faculty of Agriculture Publication No: 155; Cukurova University: Adana, Turkey, 1996; p. 148.

24. Fan, Y.; Wang, C.; Nan, Z. Determining water use efficiency for wheat and cotton: A meta-regression analysis. In Proceedings of the 2016 Agricultural & Applied Economics Association Annual Meeting, Boston, MA, USA, 31 July–2 August 2016; p. 40.

25. Ertek, A.; Kanber, R. Water-use efficiency (WUE) and change in the yield response factor (ky) of cotton irrigated by a drip irrigation system. *Turkish J. Agric. Forest.* 2001, 25, 111–118.

26. Koech, R.; Smith, R.; Gillies, M. A real-time optimisation system for automation of furrow irrigation. *Irrig. Sci.* 2014, 32, 319–327.

27. Pinnamaneni, S.R.; Anapalli, S.S.; Sui, R.; Bellaloui, N.; Reddy, K.N. Effects of irrigation and planting geometry on cotton (*Gossypium hirsutum* L.) fiber quality and seed composition. *J. Cotton Res.* 2021, 4, 2.

28. Sui, R.; Byler, R.K.; Delhom, C.D. Effect of nitrogen application rates on yield and quality in irrigated and rainfed cotton. *J. Cotton Sci.* 2017, 21, 113–121.

29. Colaizzi, P.D.; Evett, S.R.; Howell, T.A. Comparison of SDI, LEPA, and spray irrigation performance for cotton in the North Texas High Plains. *Trans. ASAE* 2004, 47, 1477–1492.

30. Segarra, E.; Almas, L.; Bordovsky, J.P. Adoption of advanced irrigation technology: LEPA vs. drip in the Texas High Plains. *Proc. Beltwide Cotton Conf.* 1999, 1, 324–328.

31. Bordovsky, J.P. Low-Energy Precision Application (LEPA) Irrigation: A Forty-Year Review. *Trans. ASABE* 2019, 62, 1343–1353.

32. Bordovsky, J.P.; Lyle, W.M.; Bender, D.A.; Lipe, W.N.; Vrubel, L.W.; Lorenz, D.C. LEPA vs. Drip Irrigation Methods; Annual Progress Report; Texas Agricultural Experiment Station: Halfway, TX, USA, 1984; pp. 7–12.

33. Cetin, O.; Ozyurt, E.; Sener, S. The effects of different irrigation methods on the yield and water use efficiency of cotton in Harran Plain. In Proceedings of the 17th European Regional Conference on Efficient and Ecologically Sound Use of Irrigation Water with Special Reference to European Countries, Varna, Bulgaria, 16–22 May 1994.

34. Bordovsky, J.P.; Porter, D. Cotton response to pre-plant irrigation level and irrigation capacity using spray, LEPA, and subsurface drip irrigation. In Proceedings of the 2003 ASAE International Meeting, Las Vegas, NV, USA, 27–30 July 2003.

35. Choudhary, K.K.; Dahiya, R.; Phogat, V.K. Effect of drip and furrow irrigation methods on yield and water use efficiency in cotton. *Res. Crops* 2016, 17, 823–828.

36. Sezan, S.M.; Yazar, A.; Akyildiz, A.; Dasgan, H.Y.; Gencel, B. Yield and quality response of drip irrigated green beans under full and deficit irrigation. *Sci. Hort.* 2008, 117, 95–102.
37. Wang, J.; Du, G.; Tian, J.; Zhang, Y.; Jiang, C.; Zhang, W. Effect of irrigation methods on root growth, root-shoot ratio and yield components of cotton by regulating the growth redundancy of root and shoot. *Agric. Water Manag.* 2020, 234, 106120.
38. Wang, R.S.; Kang, Y.H.; Wan, S.Q.; Hu, W.; Liu, S.P.; Liu, S.H. Salt distribution and the growth of cotton under different drip irrigation regimes in a saline area. *Agric. Water Manag.* 2011, 100, 58–69.
39. Ayars, J.E.; Phene, C.J.; Hutmacher, R.B.; Davis, K.R.; Schoneman, R.A.; Vail, S.S.; Mead, R.M. Subsurface drip irrigation of row crops: A review of 15 years of research at the water management research laboratory. *Agric. Water Manag.* 1999, 42, 1–27.
40. Batchelor, C.; Lovell, C.; Murata, M. Simple microirrigation techniques for improving irrigation efficiency on vegetable gardens. *Agric. Water Manag.* 1996, 32, 37–48.
41. Karlberg, L.; Frits, W.T.P.V. Exploring potentials and constraints of low-cost drip irrigation with saline water in sub-Saharan Africa. *Phys. Chem. Earth.* 2004, 29, 1035–1042.
42. Fereres, E.; Cuevas, R.; Orgaz, F. Drip Irrigation of Cotton in Southern Spain; American Society of Agricultural Engineers: St. Joseph, MICH, USA, 1985; pp. 185–192.
43. Mateos, L.; Berengena, J.; Orgaz, F.; Diz, J.; Fereres, E.A. comparison between drip and furrow irrigation in cotton at two levels of water supply. *Agric. Water Manag.* 1991, 19, 313–324.
44. Ibragimov, N.; Evet, S.R.; Esanbekov, Y.; Kamilov, B.; Mirzaev, L.; Lamers, J.P.A. Water use efficiency of irrigated cotton in Uzbekistan under drip and furrow irrigation. *Agric. Water Manag.* 2007, 90, 112–120.
45. Ward, F.A.; Pulido-Velazquez, M. Water conservation in irrigation can increase water use. *Proc. Natl. Acad. Sci. USA* 2008, 105, 18215–18220.
46. Colaizzi, P.D.; Gowda, P.H.; Marek, T.H.; Porter, D.O. Irrigation in the Texas high plains: A brief history and potential reductions in demand. *Irrig. Drain.* 2008, 58, 257–274.

Retrieved from <https://encyclopedia.pub/entry/history/show/34392>