

# 2-Oxocarboxylic Acids

Subjects: **Chemistry**, **Physical**

Contributor: Marcelo Guzman

Atmospheric organic aerosols play a major role in climate, demanding a better understanding of their formation mechanisms by contributing multiphase chemical reactions with the participation of water. The sunlight driven aqueous photochemistry of small 2-oxocarboxylic acids is a potential major source of organic aerosol, which prompted the investigations into the mechanisms of glyoxylic acid and pyruvic acid photochemistry reviewed here. While 2-oxocarboxylic acids can be contained or directly created in the particles, the majorities of these abundant and available molecules are in the gas phase and must first undergo the surface uptake process to react in, and on the surface, of aqueous particles.

pyruvic acid

glyoxylic acid

cross-photoreaction

## 1. Introduction

The Earth's atmosphere is not just made up of gaseous chemicals but also contains very small particles known as aerosols. Atmospheric aerosols, suspensions of fine solid, semisolid, or liquid particles in air (e.g., smog, haze, smoke, fog, and mist) can have very large and important effects on the environment. A well-known example of these environmental effects is smog formation in large cities, where the particle concentration has become too high. Breathing in the smog caused by these particles can aggravate respiratory problems like asthma, and even be correlated to causing cancer in the long term. The particles also have major effects on the Earth's climate by directly preventing sunlight from reaching the ground or by forming clouds that block incoming sunlight.

Much like how meteorologists can predict tomorrow's weather, climate scientists can predict what effects the particles will have on Earth's future climate and on the health of humans. This work can contribute to improving these predictions from new knowledge and understanding of chemical reactions that occur within the particles and on their surface. Specifically, the work covered in this review studies what happens to two chemicals commonly found in water based atmospheric particles, specifically glyoxylic acid and pyruvic acid, when they are exposed to sunlight <sup>[1]</sup>. The discoveries presented below show that both chemicals undergo complex reaction steps in water under a variety of conditions to form products that can contribute to an increase in the number of atmospheric particles on Earth. The quantum efficiency measured for each reaction is available in the literature discussed below to be used by climate scientists in models with improved predictive accuracy. The improved understanding provided by the work reviewed below should help to protect the health of humans by preventing harmful pollution events. The work should also contribute toward guiding decisions about environmental policy.

### 1.1. Defining the Importance of Studying Small 2-Oxocarboxylic Acids

The aqueous photochemistry of small 2-oxocarboxylic acids is a topic that has garnered much attention in the literature recently [2][3][4][5][6][7][8] because of the complex mechanisms of reaction involved for these ubiquitous chemical species. The chemical behavior of these small molecules is of interest to a wide variety of disciplines because they are common waypoints in the chain of oxidation of various organic molecules, for example the atmospheric processing of aromatic emissions [9][10][11][12][13], of both biogenic and anthropogenic origin. 2-Oxocarboxylic acids are also key species in the chain of reactions during glycolysis [14] and the reductive tricarboxylic acid cycle [15]. The prevalence of small 2-oxocarboxylic acids across many different chemical environments, from atmospheric water to the inside of cells, demands a thorough understanding of their transformation pathways. This review explores the behavior and photochemical transformations of small 2-oxocarboxylic acids in the context of atmospheric aqueous chemistry.


## 2. Atmospheric Photochemistry and Small 2-Oxocarboxylic Acids

Sunlight is overwhelmingly the largest source of energy input on the Earth [16]. As such, it is the driving force for most of the chemical transformations that occur in the atmospheric gas and aqueous phases [17]. The majority of organic aerosol can be found in the troposphere where sunlight of wavelengths greater than 290 nm penetrates [18]. The troposphere is the lowest layer of the atmosphere that extends from the Earth's surface to about 10–16 km depending on the current temperature, underlying surface configuration, latitude, and time of day [19]. The primary reason for the 290 nm cut off in the troposphere is that ozone ( $O_3$ ) in the layer directly above the troposphere, called the stratosphere, filters out much of the light between 220 and 330 nm [19]. The total solar irradiance that reaches the Earth's atmosphere is on the order of  $1360 \text{ W m}^{-2}$  [16], a huge amount of energy that allows many varied light-dependent reactions to occur.

In atmospheric chemistry, organic molecules typically undergo two types of photochemical reactions: direct and indirect photolysis. Direct photolysis may occur if the species in question can absorb photons of the wavelengths available in sunlight. Typically, the importance of direct photolysis is evaluated by comparing quantum yields ( $\Phi$ ), which is defined as a number of events per photon absorbed [20]. For this work,  $\Phi$  would be the number of molecules of reactant destroyed or of product generated per photon absorbed.  $\Phi$  for many processes in the gas phase and in organic solvents have been reported in the literature, but aqueous photochemical processes have received little attention until recently, leading to a lack of  $\Phi$  values for atmospheric models to evaluate the relative importance of each photochemical process [21]. If the molecule cannot absorb sunlight, indirect photooxidation can occur through reaction with photochemically produced hydroxyl radicals ( $HO^\bullet$ ), which is the most important atmospheric oxidant in both the gas phase and the aqueous phase. For most organics, reaction with  $HO^\bullet$  controls their primary fate due to absorption bands that are below 290 nm, or to the small molar absorption coefficients that make the typically fast  $HO^\bullet$  reaction outcompete direct photolysis [21].

In this regard, small 2-oxocarboxylic acids are a special case. Because of the conjugation of the two adjacent carbonyls, the  $n \rightarrow \pi^*$  transition of 2-oxocarboxylic acids' absorption profiles are shifted to low enough energy that they can absorb sunlight directly, leading to complex radical chemistry. **Figure 1** shows the UV-visible absorption

spectra of aqueous GA (dashed green trace) and PA (solid blue trace) overlapped with the direct solar irradiance spectrum (red trace) recorded at noon on a sunny cloudless day at 230 m above ground level on 19 August 2018 in Lexington, KY, USA (elevation of 298 m above the sea level). The vertical dashed line in **Figure 1** indicates the minimum wavelength of solar radiation that can penetrate to ground level, 290 nm. The overlap between the spectra of GA and PA with the solar spectrum indicates that both 2-oxocarboxylic acids should be readily photoactivated to possibly form higher complexity, more oxidized products, contributing to SOA production in the atmosphere. Furthermore, two studies based on a combination of rate constants and absorption profiles have predicted that direct aqueous photochemistry should be a significant fate for PA, and may even be the dominant fate for PA rather than gas phase photolysis or gas/aqueous reaction with  $\text{HO}^\bullet$  [2][21]. The removal by  $\text{HO}^\bullet$  of GA and PA is relatively fast in the aqueous phase [22][23] with average lifetimes of five days for both. Instead, the photochemical lifetime of GA and PA in atmospheric water are 11 days and 22 min, respectively [3][4]. Thus, GA can also be expected to undergo direct photolysis, albeit the slower rate. However, with the higher atmospheric production of GA than PA, its direct photolysis, while not the primary sink, has a significant impact on atmospheric composition, making both GA and PA contributors to SOA production.



**Figure 1.** The molar absorption coefficients ( $\epsilon$ , left axis) in the UV-visible absorption spectra of (dashed green) glyoxylic acid and (solid blue) pyruvic acid in water, and (red and right axis) direct solar irradiance spectrum from a sunny and cloudless day (19 August 2018) in Lexington, KY (USA) at 230 m above ground level. The vertical dashed line indicates the ozone cut-off at a wavelength  $\lambda_{\text{cut-off}} \geq 290$  nm.

Thus, given their potential importance to SOA production, this review encompasses work to identify the products of photoreactions and propose mechanisms for the aqueous photolysis of GA and PA under atmospherically relevant

conditions. The quantum yields recently available in the literature are reported to incorporate these processes into atmospheric models of SOA production.

## 2.1. Future Directions for Aqueous Phase Photoreactions Studies in Atmosphere

We provided a brief description of some representative photochemical studies with small 2-oxocarboxylic acids above, where water played a key role in the observed chemistry. Indeed, aqueous tropospheric chemistry can be influenced by photochemical reactions, which provide reactive species, an emerging issue that deserves more attention. Despite the substantial progress that has been made in this field, the current comprehension of water solvation effects in the photochemistry is still in progress. Future laboratory experiments, computational quantum mechanical modeling, and molecular dynamic simulations could advance our understanding of effects such as solvation, stabilization, and solute position and orientation [24][25] in the photoprocesses in bulk and interface chemistry. Because the photochemical products of these reactions can contribute to the reservoir of light absorbing species in atmospheric waters, they can also play an active role in the aging of aqueous SOA, i.e., by triggering photosensitized reactions. Thus, quantitative understanding of the radicals and reactive oxygen species (ROS) produced from the photochemistry of 2-oxocarboxylic acids and of their implications in multiphase ageing of particulates is vital, as should also be the comparison with other competitive loss pathways. Studying how the generated excited triplet states directly and indirectly alter the organic surface-active components typically coating the air–water interface of atmospheric particles is of major importance. Addressing such problems should inform us of new processes that can modify the morphology and properties of atmospheric particles and their influence on climate forcing processes [26].

## References

1. Guzman, Marcelo I.; Eugene, Alexis J. Aqueous Photochemistry of 2-Oxocarboxylic Acids: Evidence, Mechanisms, and Atmospheric Impact . *Molecules* **2021**, 26, 5278, <https://doi.org/10.3390/molecules26175278>.
2. Epstein, S.A.; Tapavicza, E.; Furche, F.; Nizkorodov, S.A. Direct photolysis of carbonyl compounds dissolved in cloud and fog droplets. *Atmos. Chem. Phys.* 2013, 13, 9461–9477.
3. Eugene, A.J.; Guzman, M.I. Reactivity of ketyl and acetyl radicals from direct solar actinic photolysis of aqueous pyruvic acid. *J. Phys. Chem. A* 2017, 121, 2924–2935. <https://doi.org/10.1021/acs.jpca.6b11916>
4. Eugene, A.J.; Xia, S.-S.; Guzman, M.I. Aqueous photochemistry of glyoxylic acid. *J. Phys. Chem. A* 2016, 120, 3817–3826. <https://doi.org/10.1021/acs.jpca.6b00225>
5. Griffith, E.C.; Carpenter, B.K.; Shoemaker, R.K.; Vaida, V. Photochemistry of aqueous pyruvic acid. *Proc. Nat. Acad. Sci. USA* 2013, 110, 11714–11719.

6. Guzman, M.I.; Colussi, A.J.; Hoffmann, M.R. Photoinduced oligomerization of aqueous pyruvic acid. *J. Phys. Chem. A* 2006, 110, 3619–3626. <https://doi.org/10.1021/jp056097z>
7. Rapf, R.J.; Perkins, R.J.; Dooley, M.R.; Kroll, J.A.; Carpenter, B.K.; Vaida, V. Environmental processing of lipids driven by aqueous photochemistry of  $\alpha$ -keto acids. *ACS Cent. Sci.* 2018, 4, 624–630.
8. Reed Harris, A.E.; Ervens, B.; Shoemaker, R.K.; Kroll, J.A.; Rapf, R.J.; Griffith, E.C.; Monod, A.; Vaida, V. Photochemical kinetics of pyruvic acid in aqueous solution. *J. Phys. Chem. A* 2014, 118, 8505–8516.
9. Pillar, E.A.; Camm, R.C.; Guzman, M.I. Catechol oxidation by ozone and hydroxyl radicals at the air-water interface. *Environ. Sci. Technol.* 2014, 48, 14352–14360. <https://doi.org/10.1021/es504094x>
10. Pillar, E.A.; Zhou, R.; Guzman, M.I. Heterogeneous oxidation of catechol. *J. Phys. Chem. A* 2015, 119, 10349–10359. <https://doi.org/10.1021/acs.jpca.5b07914>
11. Andreae, M.O.; Browell, E.V.; Garstang, M.; Gregory, G.; Harriss, R.; Hill, G.F.; Jacob, D.J.; Pereira, M.; Sachse, G.; Setzer, A. Biomass-burning emissions and associated haze layers over Amazonia. *J. Geophys. Res. Atmos.* 1988, 93, 1509–1527.
12. Martín-Reviejo, M.; Wirtz, K. Is benzene a precursor for secondary organic aerosol? *Environ. Sci. Technol.* 2005, 39, 1045–1054.
13. Rana, M.S.; Guzman, M.I. Oxidation of phenolic aldehydes by ozone and hydroxyl radicals at the air–water interface. *J. Phys. Chem. A* 2020, 124, 8822–8833. <https://doi.org/10.1021/acs.jpca.0c05944>
14. Garrett, R.H.; Grisham, C.M. *Biochemistry*, 2nd ed.; Saunders College Publishing: Fort Worth, TX, USA; Orlando, FL, USA, 1999.
15. Guzman, M.I.; Martin, S.T. Photo-production of lactate from glyoxylate: How minerals can facilitate energy storage in a prebiotic world. *Chem. Comm.* 2010, 46, 2265–2267. <https://doi.org/10.1021/jp3011316>
16. Kopp, G.; Lean, J.L. A new, lower value of total solar irradiance: Evidence and climate significance. *Geophys. Res. Lett.* 2011, 38, L01706.
17. George, C.; Ammann, M.; D’Anna, B.; Donaldson, D.J.; Nizkorodov, S.A. Heterogeneous photochemistry in the atmosphere. *Chem. Rev.* 2015, 115, 4218–4258.
18. Schwarzenbach, R.P.; Gschwend, P.M.; Imboden, D.M. *Environmental Organic Chemistry*, 2nd ed.; John Wiley & Sons: Hoboken, NJ, USA, 2003.
19. Manahan, S.E. *Environmental Chemistry*, 9th ed.; CRC press: Boca Raton, FL, USA, 2010; p. 783.

20. Hoque, M.; Guzman, M. Photocatalytic activity: Experimental features to report in heterogeneous photocatalysis. *Materials* 2018, 11, 1990. <https://doi.org/10.3390/molecules26175278>
21. Epstein, S.A.; Nizkorodov, S.A. A comparison of the chemical sinks of atmospheric organics in the gas and aqueous phase. *Atmos. Chem. Phys.* 2012, 12, 8205–8222.
22. Ervens, B.; Gligorovski, S.; Herrmann, H. Temperature-dependent rate constants for hydroxyl radical reactions with organic compounds in aqueous solutions. *Phys. Chem. Chem. Phys.* 2003, 5, 1811–1824.
23. Schöne, L.; Herrmann, H. Kinetic measurements of the reactivity of hydrogen peroxide and ozone towards small atmospherically relevant aldehydes, ketones and organic acids in aqueous solutions. *Atmos. Chem. Phys.* 2014, 14, 4503–4514.
24. Eugene, A.J.; Pillar, E.A.; Colussi, A.J.; Guzman, M.I. Enhanced acidity of acetic and pyruvic acids on the surface of water. *Langmuir* 2018, 34, 9307–9313. <https://doi.org/10.3390/ma15207378>
25. Guzman, M.I.; Athalye, R.R.; Rodriguez, J.M. Concentration effects and ion properties controlling the fractionation of halides during aerosol formation. *J. Phys. Chem. A* 2012, 116, 5428–5435. <https://doi.org/10.1021/jp3011316>
26. Pillar-Little, E.A.; Guzman, M.I. An overview of dynamic heterogeneous oxidations in the troposphere. *Environments* 2018, 5, 104. <https://doi.org/10.3390/environments5090104>

---

Retrieved from <https://encyclopedia.pub/entry/history/show/104695>