

L19-TNF

Subjects: [Biochemistry & Molecular Biology](#) | [Oncology](#)

Contributor: Sheila Dakhel

Tumor necrosis factor (TNF) is used as a pro-inflammatory payload to trigger haemorrhagic necrosis and boost anti-cancer immunity at the tumor site. There is a depotentiated version of TNF (carrying the single point mutation I97A), which displayed reduced binding affinity to its cognate receptor tumor necrosis factor receptor 1 (TNFR-1) and lower biocidal activity.

tumor targeting
fibronectin

immunocytokines

activity-on-demand

targeted-TNF

EDB-

1. Introduction

Cytokines are small proteins which modulate the activity of the immune system in health and in disease. Certain proinflammatory cytokines are used for cancer immunotherapy ^{[1][2]}, while other cytokines with anti-inflammatory properties may be considered as tools for the treatment of chronic inflammatory and autoimmune conditions ^[3]. Targeting an active cytokine to a tumor-associated antigen by means of a fusion with a suitable monoclonal antibody may represent a strategy for the enhancement of the cytokines' therapeutic index ^{[4][5]}.

Tumor necrosis factor (TNF) is a homotrimeric cytokine which shows direct cytotoxic activity on certain types of cancer cells ^[6], stimulates anti-tumor immunity ^{[7][8]}, and may also be active on the tumor endothelium, leading to intravascular blood coagulation ^[9] or to haemorrhagic necrosis ^[10]. Recombinant human TNF (Beromun[®]) is an approved drug in Europe for the treatment of metastatic melanoma and soft tissue sarcoma in the setting of isolated limb perfusion. The systemic use of non-targeted TNF is limited by its narrow therapeutic index and high toxicity at very low doses ^[11]. Therefore, TNF has been used as an active payload for the generation of tumor-specific immunocytokines ^[12].

L19-TNF (Fibromun, Philogen S.p.A., Siena, Italy) is a fully-human immunomodulatory product consisting of a non-covalent homotrimeric immunocytokine featuring human TNF fused to the L19 antibody (specific to EDB of fibronectin, a marker of tumor neovasculature) ^[5]. L19-TNF has been systemically administered to patients with solid tumors ^[13], as well as in isolated limb perfusion procedures to patients with in-transit melanoma metastases ^[14].

Now, there is the generation of a depotentiated version of Fibromun product, capable of inducing potent in vivo anti-tumor activity with reduced systemic toxicity. The new molecule, named L19-TNFI97A, was selected among a series of candidates displaying single-point mutations of amino acid residues involved in surface interactions between TNF monomers. The substitution of isoleucine for alanine in position 95 of the human TNF, hence named L19-TNFI97A, showed decreased activity (~100-fold) when compared to wild type L19-TNF (L19-TNFWT), while its biological activity was restored upon binding to the target antigen EDB both in vitro and in vivo. Moreover, the binding affinity of L19-TNFI97A towards its cognate receptor, TNFR1, was decreased compared to the wild type molecule. Preclinical characterization of L19-TNFI97A in tumor-bearing

2. Production and Characterization of L19-TNF Mutant Fusion Proteins

L19-TNF is a recombinant fusion protein composed by the fully human L19 antibody in scFv format

conjugate to the human TNF by a (S₄G)₃ linker (**Figure 1A**) forming a natural homotrimer (**Figure 1B**). To identify products with promising features for subsequent in vivo investigations, 11 different mutant versions of L19-TNF were expressed in mammalian cells and purified by Protein A chromatography, as the VH domain of the L19 antibody is compatible with affinity capture modality [15]. Six of the eleven products exhibited favorable size exclusion and sodium dodecyl-sulfate polyacrylamide gel electrophoresis (SDS-PAGE) profiles (**Figure 1C** and [Supplementary Information Figure S1](#)). Deconvolution of the liquid chromatography-mass spectrometry (LC-MS) profile allowed identification of the expected experimental masses for all the targeted TNF-constructs which were compatible with the predicted values in the absence of any protein glycosylation. Sequences of the fusion proteins are listed in the [Supplementary Information](#).




Figure 1. Cloning, expression and quality control analyses of L19-TNF mutants. **(A)** Schematic representation of L19-TNF^{WT} vector used as a template for the generation of mutants. Leader sequence: MGWSLILLFLVAVATGVHS **(B)** General scheme of L19-TNF mutants. **(C)** Quality control of L19-TNF^{WT} and of L19-TNF^{Y59F}, L19-TNF^{I97A}, L19-TNF^{Q102N}, L19-TNF^{Y119S}, L19-TNF^{S147A} mutants by SDS-PAGE, Size Exclusion Chromatography (pic eluting at 12 mL corresponding to the homotrimer form) and LC/MS analyses. NR = non reducing conditions, R = reducing conditions.

3. Conditional In Vitro Cytotoxicity Effect of L19-TNF Mutants upon EDB Concentration

TNF is known to induce direct killing on several tumor cell lines [10]. To evaluate the biocidal activity of L19-TNF mutants upon antigen concentration, a panel of human tumor cell lines were analyzed for the EDB expression by immunofluorescence (**Figure 2A**). The antigen was not detectable in the human melanoma cell line A-375 and, therefore, the L19-TNF mutants were tested for their cytotoxic activity on these cells in the presence or absence of EDB coating (**Figure 2B**). Cell killing of L19-TNF^{WT} and the majority of the mutants tested was not affected by the presence of EDB. However, in the absence of EDB, the mutein carrying the single substitution of the isoleucine at position 97 by alanine (hence named L19-TNF^{I97A}) showed 10 times less potency

compared to the wild type version. This effect was re-established upon EDB presence (**Figure 2C**). EC₅₀ values for all the tested proteins are reported in **Table 1** and [Supplementary Information Table S4](#).




Figure 2. Assessment of EDB expression by different human tumor cell lines and in vitro biocidal activity of L19-TNF mutants. **(A)** Immunofluorescence analysis of EDB expression on different human tumor cells. All tumor cells showed high EDB expression, with the exception of the A-375 cell line. Green = EDB, Blue = Nucleus. Scale bar = 100 Åµm. **(B)** Immunofluorescence EDB staining results on A-375 cultured in absence or presence of recombinant EDB coating. **(C)** In vitro cytotoxicity of L19-TNF wild type and mutants assayed on A-375 cells in the presence (dashed line) or absence (continuous line) of EDB coating. L19-TNF^{I97A} showed conditional cytotoxicity in the presence of EDB. Data points represent cell viability (mean Å± SD, n = 3).

Table 1. Calculated EC₅₀ values for L19-TNF wild type and mutants.

	No EDB Coating (M)	100 nM EDB Coating (M)
L19-TNF ^{WT}	1.076 Å 10 ^{Å10}	1.033 Å 10 ^{Å10}
L19-TNF ^{Y59F}	4.172 Å 10 ^{Å10}	9.501 Å 10 ^{Å10}
L19-TNF ^{I97A}	2.273 Å 10 ^{Å9}	2.254 Å 10 ^{Å10}
L19-TNF ^{Q102N}	5.012 Å 10 ^{Å11}	5.986 Å 10 ^{Å11}
L19-TNF ^{Y119S}	3.245 Å 10 ^{Å11}	6.417 Å 10 ^{Å11}
L19-TNF ^{S147A}	1.261 Å 10 ^{Å11}	1.58 Å 10 ^{Å11}

4. The Mutant I97A Has Reduced Binding Affinity towards TNFR1

Affinity of L19-TNF^{WT} and of L19-TNF^{I97A} was analyzed by Surface Plasmon Resonance. The

proteins were reversibly coated on independent EDB-precoated chips. TNFR1 was injected at different concentrations and SPR response was measured (**Figure 3A**). L19-TNF^{WT} showed a fast k_{on} and very slow k_{off} towards TNFR1 (experimental calculated $KD = 10$ nM) and the binding kinetics of the mutant L19-TNF^{Q102N} (selected as a mutant with similar in vitro activity to the wild type protein) were comparable (**Figure 3B,D**). By contrast, the L19-TNF^{I97A} mutant was still a stable homotrimer in gel filtration analysis but exhibited a faster dissociation from the receptor, indicative of a depotentiation of TNFR1 binding (**Figure 3C**). Fitting and calculated binding constants are depicted in the [Supplementary Information Tables S5 and S6](#).




Figure 3. Real time binding analysis of TNFR1 on immobilized L19-TNF mutants. **(A)** Schematic representation of affinity measurement methodology. EDB was covalently immobilized on the chip, while different L19-TNF mutants were coated on the solid surface thanks to non-covalent L19-EDB interaction. TNFR1 was sequentially injected at different concentrations (starting concentration of 1000 nM, followed by 1:2 dilution series). SPR sensorgrams of **(B)** L19-TNF^{WT}, **(C)** L19-TNF^{Q102N} and **(D)** L19-TNF^{I97A} are presented.

5. The I97A Mutant Accumulates at the Tumor Site and Shows High Levels in Blood

In order to test immunoreactivity of the L19 antibody function and EDB in L19-TNF^{I97A}, immunofluorescence staining was performed on non-treated WEHI-164 tumor cryosections. L19-TNF^{WT} and L19-TNF^{I97A} were both able to bind to EDB present in tumor specimens (**Figure 4A**). The tumor-targeting performance of L19-TNF^{I97A} was evaluated in immunocompetent BALB/c mice bearing subcutaneously-grafted WEHI-164 tumors. The protein was radiolabeled with ¹²⁵I and injected intravenously at the recommended dose of 250 μ g/Kg. The mutant preferentially localized at the site of the disease, with a high tumor uptake value and excellent tumor-to-normal organs ratio (average tumor-to-organs ratio 5.5:1). Surprisingly, high levels of the mutant were found in blood 24 h post-administration (average tumor-to-blood ratio of 2-to-1) (**Figure 4B**).




Figure 5. Tumor-targeting performance of L19-TNF^{I97A} and in vivo therapy study in tumor-bearing mice. **(A)** Immunofluorescence staining of WEHI-164 tumors with L19-TNF^{I97A} (L19-TNF^{WT} was used as the positive control of the experiment). Green = L19-TNF^{WT} or L19-TNF^{I97A}, Red = CD31. Scale bar = 100 μm . **(B)** In vivo biodistribution with radioiodinated L19-TNF^{I97A} in BALB/c mice bearing WEHI-164 subcutaneous tumors, 24 h after intravenous administration. The product shows selective accumulation in tumor lesions (n = 3). Assessment of therapeutic efficacy and toxicity in BALB/c mice bearing WEHI-164 subcutaneous tumors. Mice were injected with a high dose of L19-TNF products (375 $\mu\text{g}/\text{kg}$, black arrows indicate IV administration). Tumor volume **(C)** and changes in animal body weight **(D)** induced by the administered therapeutic proteins were monitored throughout the study (mean \pm SEM; n = 3). The statistical analysis for the tumor volume was performed using a two-way ANOVA analyses with Bonferroni post-test **** p < 0.0001).

References

1. Lee, S.; Margolin, K. Cytokines in cancer immunotherapy. *Cancers* 2011, 3, 3856–3893.
2. Dranoff, G. Cytokines in cancer pathogenesis and cancer therapy. *Nat. Rev. Cancer* 2004, 4, 11–22.
3. Moudgil, K.D.; Choubey, D. Cytokines in autoimmunity: Role in induction, regulation, and treatment. *J. Interferon Cytokine Res.* 2011, 31, 695–703.
4. Carnemolla, B.; Borsi, L.; Balza, E.; Castellani, P.; Meazza, R.; Berndt, A.; Ferrini, S.; Kosmehl, H.; Neri, D.; Zardi, L. Enhancement of the antitumor properties of interleukin-2 by its targeted delivery to the tumor blood vessel extracellular matrix. *Blood* 2002, 99, 1659–1665.
5. Borsi, L.; Balza, E.; Carnemolla, B.; Sassi, F.; Castellani, P.; Berndt, A.; Kosmehl, H.; Birnbaumer, A.; Siri, A.; Orecchia, P.; et al. Selective targeted delivery of TNF α to tumor blood vessels. *Blood* 2003, 102, 4384–4392.
6. Fiers, W. Tumor necrosis factor Characterization at the molecular, cellular and in vivo level. *FEBS Lett.* 1991, 285, 199–212.
7. Calcinotto, A.; Grioni, M.; Jachetti, E.; Curnis, F.; Mondino, A.; Parmiani, G.; Corti, A.; Bellone, M. Targeting TNF α to Neoangiogenic Vessels Enhances Lymphocyte Infiltration in Tumors and Increases the Therapeutic Potential of Immunotherapy. *J. Immunol.* 2012, 188, 2687–2694.

8. Elia, A.R.; Grioni, M.; Basso, V.; Curnis, F.; Freschi, M.; Corti, A.; Mondino, A.; Bellone, M. Targeting tumor vasculature with TNF leads effector t cells to the tumor and enhances therapeutic efficacy of immune checkpoint blockers in combination with adoptive cell therapy. *Clin. Cancer Res.* 2018, 24, 2171â2181.
9. Lejeune, F.J.; LiÃ©nard, D.; Matter, M.; RÃ¼egg, C. Efficiency of recombinant human TNF in human cancer therapy. *Cancer Immun.* 2006, 6, 6.
10. Carswell, E.A.; Old, L.J.; Kassel, R.L.; Green, S.; Fiore, N.; Williamson, B. An endotoxin-induced serum factor that causes necrosis of tumors (activated macrophage). *Proc. Natl. Acad. Sci. USA* 1975, 72, 3666â3670.
11. Creaven, P.J.; Brenner, D.E.; Cowens, J.W.; Huben, R.P.; Wolf, R.M.; Takita, H.; Arbuck, S.G.; Razack, M.S.; Proefrock, A.D. A phase I clinical trial of recombinant human tumor necrosis factor given daily for five days. *Cancer Chemother. Pharmacol.* 1989, 23, 186â191.
12. Spitaleri, G.; Berardi, R.; Pierantoni, C.; De Pas, T.; Noberasco, C.; Libbra, C.; GonzÃ¡lez-Iglesias, R.; Giovannoni, L.; Tasciotti, A.; Neri, D.; et al. Phase I/II study of the tumour-targeting human monoclonal antibody-cytokine fusion protein L19-TNF in patients with advanced solid tumours. *J. Cancer Res. Clin. Oncol.* 2013, 139, 447â455.
13. Papadia, F.; Basso, V.; Patuzzo, R.; Maurichi, A.; Di Florio, A.; Zardi, L.; Ventura, E.; GonzÃ¡lez-Iglesias, R.; Lovato, V.; Giovannoni, L.; et al. Isolated limb perfusion with the tumor-targeting human monoclonal antibody-cytokine fusion protein L19-TNF plus melphalan and mild hyperthermia in patients with locally advanced extremity melanoma. *J. Surg. Oncol.* 2013, 107, 173â179.
14. Neri, D. AntibodyâCytokine Fusions: Versatile Products for the Modulation of Anticancer Immunity. *Cancer Immunol. Res.* 2019, 7, 348â354.
15. Halin, C.; Rondini, S.; Nilsson, F.; Berndt, A.; Kosmehl, H.; Zardi, L.; Neri, D. Enhancement of the antitumor activity of interleukin-12 by targeted delivery to neovasculature. *Nat. Biotechnol.* 2002, 20, 264â269.

Retrieved from <https://encyclopedia.pub/entry/15292>