

Surgical Techniques for Rectal Prolapse

Subjects: Gastroenterology & Hepatology

Contributor: Bengi Balci

Complete rectal prolapse or rectal procidentia is a debilitating disease that presents with fecal incontinence, constipation, and rectal discharge. Definitive surgical techniques described for this disease include perineal procedures such as mucosectomy and rectosigmoidectomy, and abdominal procedures such as rectopexy with or without mesh and concomitant resection. The debate over these techniques regarding the lowest recurrence and morbidity rates, and the best functional outcomes for constipation or incontinence, has been going on for decades. The heterogeneity of available studies does not allow us to draw firm conclusions.

Keywords: rectal prolapse ; perineal approach ; abdominal approach ; laparoscopic approach

1. Introduction

Complete rectal prolapse or rectal procidentia is defined as a full-thickness protrusion of the rectum through the anal canal, and should be distinguished from the mucosal prolapse ^[1]. Although the definite etiology of this disease is unclear, the most common related pathologies are the redundant sigmoid colon, a deep cul-de-sac, and the diastasis of levator ani muscles ^{[2][3]}.

The incidence is higher in females, with a peak in the seventh decade ^[4]. Patients usually complain of fecal incontinence, which is thought to be a result of a chronic stretch of the anal sphincter and continuous stimulation of the rectoanal inhibitory reflex by the prolapsed tissue ^{[5][6]}. Other symptoms include constipation, pain, bloody or mucous rectal discharge ^[7].

The diagnosis can be made by demonstrating the protrusion of the rectal circumferential radial folds with straining maneuvers on clinical examination ^[7]. However, additional diagnostic tests such as dynamic pelvic magnetic resonance imaging (MRI), anal manometry, and endoanal ultrasonography (EUS) are usually required to detect co-existing anatomical abnormalities and decide on the optimal surgical treatment method ^{[8][9][10][11][12]}.

Described surgical techniques are varied, and whether the approach is abdominal or perineal, the treatment aims to correct anatomical and functional abnormalities by the fixation of the rectum to the sacrum and/or the resection of the redundant bowel ^[13]. In this article, these different techniques are reviewed for their

2. Surgical Techniques

As a general consideration, perineal procedures have been associated with reduced complication rates and chosen in frail patients with co-morbidities. However, with recent randomized controlled trials and reviews, perineal and abdominal procedures have been reported to have similar outcomes in expert hands regarding perioperative complications, recurrence rate, and quality of life ^[14].

A recent Cochrane Database review demonstrated no significant difference in recurrence rates ^[15], and Mustain et al. reported no difference in terms of postoperative complications between these two approaches ^[16]. Moreover, Emile et al. revealed that similar recurrence and complication rates were seen, but increased length of hospital stay in perineal procedures and longer operative time in abdominal procedures were noted ^[17].

2.1. Abdominal Approach

Described techniques, whether rectopexy using a mesh or suture and/or resection, have been proven to be effective and safe via laparoscopy ^[13]. The laparoscopic approach for rectal prolapse has been associated with shorter hospital stay and less postoperative pain. Moreover, there was no statistical difference in recurrence rates, postoperative constipation, or incontinence scores ^{[18][19][20]}.

2.1.1. Ripstein Procedure (Anterior Mesh Rectopexy)

This procedure was first described by Ripstein in 1952 [21]. Reported results ranged from 0 to 13% for recurrence and 0 to 2.8% for mortality, with high complication rates of 33% [22]. Its well-recognized complication was the obstruction and narrowing of the rectum; therefore, patients may complain from newly developed or worsened constipation in the postoperative setting [23].

2.1.2. Wells Procedure (Posterior Mesh Rectopexy)

With the use of absorbable materials instead of non-absorbable materials, as initially described by Wells, recurrence and complications such as pelvic sepsis have been lower with posterior mesh rectopexy [24].

Laparoscopic posterior mesh rectopexy (LPMR) was investigated by several prospective studies [25][26][27]. Dyrberg et al. reported a recurrence rate of 11%, as well as improvements in continence and constipation scores (74% and 65%, respectively) [26]. In a comparative study with a mean follow-up of 46 months, recurrence rates were found to be 3%, and this was similar to the laparoscopic ventral mesh rectopexy (LVMR) group, whereas patients with LPMR had less improvement in constipation scores [28].

2.1.3. Suture Rectopexy

Suture rectopexy was first described by Cutait in 1959, and it aimed to form scar tissue between the rectum and sacrum [29][30]. Presumably, due to extensive rectal mobilization, postoperative constipation rates are increased [15][31].

2.1.4. Frykman-Goldberg Procedure (Resection Rectopexy)

In addition to rectopexy, sigmoid resection has been considered effective in terms of recurrence and functional outcomes, particularly in patients with complaints of constipation [14][15][32]. For patients with the main symptom of fecal incontinence, this procedure seems to be unnecessary [33].

Laparoscopic resection rectopexy (LRR) is the most commonly chosen technique for rectal prolapse in the United States [34]. Ashari et al. demonstrated similar improvement rates of continence and constipation (62% and 69%, respectively) and a recurrence rate of 3% with a median follow-up of 62 months [35]. Moreover, a retrospective comparative study between LRR and LVMR reported no significant differences in recurrence and functional outcomes, but higher complication rates in the LRR group [36].

2.1.5. Ventral Mesh Rectopexy

Ventral mesh rectopexy was first described by D'Hoore in 2004 [37]. The main advantage of this technique is the limited dissection anteriorly to the rectum, which enables reinforcement of the rectovaginal septum, preservation of the autonomic nerves, and concomitant colpopexy. High improvement rates in constipation and decreased de novo constipation compared to posterior rectopexy are attributed to the avoidance of complete rectal mobilization [38][39]. Based on these favorable outcomes, ventral mesh rectopexy has become a standard treatment for complete rectal prolapse in European countries [40][41].

2.2. Perineal Approach

2.2.1. Delorme Mucosectomy

Delorme procedure was first described in 1900 [42], and it is usually preferred in patients with short segments of rectal prolapse [43]. Initial studies reported that this procedure was associated with high recurrence rates and worse functional outcomes [44]. Marchal et al. demonstrated an improvement rate of 42% and 54% of continence and constipation, respectively, and 20% morbidity in patients with a mean follow-up of 73 months [45]. In a prospective study including 113 patients with a mean follow-up of 36 months, an improvement of continence was found to be much higher (89%), with an increased morbidity rate of 30% [46].

On the other hand, Fleming et al. reported similar results between Altemeir and Delorme procedures in terms of functional outcomes, postoperative complications, and mortality [47]. Moreover, in a randomized prospective study comparing patients undergoing Delorme procedure vs. Delorme with levatoroplasty, it was revealed that the combined procedure was associated with decreased recurrence rates and better functional outcomes [48].

2.2.2. Altemeir Procedure (Perineal Rectosigmoidectomy)

Altemeier procedure has been commonly performed in patients with long segments of rectal prolapse (>5 cm) [14]. It has also been described as a preferred method for strangulated rectal prolapse combined with diverting ileostomy [49][50]. There are mostly retrospective studies reporting contradicted results regarding functional outcomes, morbidity, and mortality rates of this procedure. In a large retrospective study including 518 patients, morbidity and recurrence rates were 9% and 23%, respectively [51]. On the other hand, Trompetto et al. reported a significantly higher morbidity rate of 38% and recurrence rate of 35% in patients with a median follow-up of 49 months [52].

2.2.3. Perineal Stapled Prolapse Resection (PSPR)

PSPR is faster and easier than conventional perineal procedures and it is suitable for high-risk patients under spinal anesthesia. However, high rates of early recurrence have been reported by several studies [53][54]. Tschuor et al. reported a recurrence rate of 44% in a median follow-up of 40 months [55]. Even though the recurrence rates were comparable to Altemeier's and Delorme's procedures in a recent meta-analysis, the authors emphasized the necessity of further randomized prospective studies comparing PSPR with other techniques in complete rectal prolapse [56].

References

1. Lowry, A.C.; Simmang, C.L.; Boulos, P.; Farmer, K.C.; Finan, P.J.; Hyman, N.; Killingback, M.; Lubowski, D.Z.; Moore, R.; Penfold, C.; et al. Consensus statement of definitions for anorectal physiology and rectal cancer. *ANZ J. Surg.* 2001, 71, 603–605.
2. Moschcowitz, A.V. The pathogenesis, anatomy, and cure of prolapse of the rectum. *Dis. Colon Rectum* 1983, 26, 553–565.
3. Devadhar, D.S.C. A new concept of mechanism and treatment of rectal procidentia. *Dis. Colon Rectum* 1965, 8, 75–77.
4. Wilson, J.; Engledow, A.; Crosbie, J.; Arulampalam, T.; Motson, R. Laparoscopic nonresectional suture rectopexy in the management of full-thickness rectal prolapse: Substantive retrospective series. *Surg. Endosc.* 2011, 25, 1062–1064.
5. Hawkins, A.; Olariu, A.G.; Savitt, L.R.; Gingipally, S.; Wakamatsu, M.M.; Pulliam, S.; Weinstein, M.M.; Bordeianou, L. Impact of rising grades of internal rectal intussusception on fecal continence and symptoms of constipation. *Dis. Colon Rectum* 2016, 59, 54–61.
6. Kim, D.-S.; Tsang, C.B.S.; Wong, D.W.; Lowry, A.C.; Goldberg, S.M.; Madoff, R.D. Complete rectal prolapse: Evolution of management and results. *Dis. Colon Rectum* 1999, 42, 460–466.
7. Bordeianou, L.; Hicks, C.W.; Kaiser, A.M.; Alavi, K.; Sudan, R.; Wise, P.E. Rectal Prolapse: An overview of clinical features, diagnosis, and patient-specific management strategies. *J. Gastrointest. Surg.* 2014, 18, 1059–1069.
8. Pannu, H.K.; Kaufman, H.S.; Cundiff, G.W.; Genadry, R.; Bluemke, D.A.; Fishman, E.K. Dynamic MR imaging of pelvic organ prolapse: Spectrum of abnormalities. *Radiographics* 2000, 20, 1567–1582.
9. Comiter, C.V.; Vasavada, S.P.; Barbaric, Z.L.; Gousse, A.; Raz, S. Grading pelvic prolapse and pelvic floor relaxation using dynamic magnetic resonance imaging. *Urology* 1999, 54, 454–457.
10. Pucciani, F. Anorectal manometry. In *Colon, Rectum and Anus: Anatomic, Physiologic and Diagnostic Bases for Disease Management*; Ratto, C., Parello, A., Donisi, L., Litta, F., Eds.; Springer: Cham, Switzerland, 2017; pp. 277–283.
11. Glasgow, S.; Birnbaum, E.H.; Kodner, I.J.; Fleshman, J.W.; Dietz, D.W. Preoperative anal manometry predicts continence after perineal proctectomy for rectal prolapse. *Dis. Colon Rectum* 2006, 49, 1052–1058.
12. Dvorkin, L.S.; Chan, C.L.H.; Knowles, C.H.; Williams, N.S.; Lunniss, P.J.; Scott, M.S. Anal sphincter morphology in patients with full-thickness rectal prolapse. *Dis. Colon Rectum* 2004, 47, 198–203.
13. Tsunoda, A. Surgical treatment of rectal prolapse in the laparoscopic era; A review of the literature. *J. Anus Rectum Colon* 2020, 4, 89–99.
14. Gallo, G.; Martellucci, J.; Pellino, G.; Ghiselli, R.; Infantino, A.; Pucciani, F.; Trompetto, M. Consensus statement of the Italian Society of Colorectal Surgery (SICCR): Management and treatment of complete rectal prolapse. *Tech. Coloproctology* 2018, 22, 919–931.
15. Tou, S.; Brown, S.R.; Nelson, R.L. Surgery for complete (full-thickness) rectal prolapse in adults. *Cochrane Database Syst. Rev.* 2015, 2015, CD001758.
16. Mustain, W.C.; Davenport, D.L.; Parcells, J.P.; Vargas, H.D.; Hourigan, J.S. Abdominal versus perineal approach for treatment of rectal prolapse: Comparable safety in a propensity-matched cohort. *Am. Surg.* 2013, 79, 686–692.

17. Emile, S.H.; Elbanna, H.; Youssef, M.; Thabet, W.; Omar, W.; Elshobaky, A.; El-Hamed, T.M.A.; Farid, M. Laparoscopic ventral mesh rectopexy vs. Delorme's operation in management of complete rectal prolapse: A prospective randomized study. *Color. Dis.* 2017, 19, 50–57.
18. Baker, R.; Senagore, A.J.; Luchtefeld, M.A. Laparoscopic-assisted vs. open resection. Rectopexy offers excellent results. *Dis. Colon Rectum* 1995, 38, 199–201.
19. Solomon, M.J.; Young, C.J.; Eysers, A.A.; Roberts, R.A. Randomized clinical trial of laparoscopic versus open abdominal rectopexy for rectal prolapse. *BJS* 2002, 89, 35–39.
20. Demirbas, S.; Akin, M.L.; Kalemoglu, M.; Ogün, I.; Celenk, T. Comparison of laparoscopic and open surgery for total rectal prolapse. *Surg. Today* 2005, 35, 446–452.
21. Ripstein, C.B. Treatment of massive rectal prolapse. *Am. J. Surg.* 1952, 83, 68–71.
22. Hrabe, J.; Gurland, B. Optimizing treatment for rectal prolapse. *Clin. Colon Rectal Surg.* 2016, 29, 271–276.
23. Tjandra, J.J.; Fazio, V.W.; Church, J.M.; Milsom, J.W.; Oakley, J.R.; Lavery, I.C. Ripstein procedure is an effective treatment for rectal prolapse without constipation. *Dis. Colon Rectum* 1993, 36, 501–507.
24. Sayfan, J.; Pinho, M.; Alexander-Williams, J.; Keighley, M.R.B. Sutured posterior abdominal rectopexy with sigmoidectomy compared with Marlex® rectopexy for rectal prolapse. *Br. J. Surg.* 1990, 77, 143–145.
25. Dulucq, J.-L.; Wintringer, P.; Mahajna, A. Clinical and functional outcome of laparoscopic posterior rectopexy (Wells) for full-thickness rectal prolapse. A prospective study. *Surg. Endosc.* 2007, 21, 2226–2230.
26. Thejeswi, P. Evaluation of clinical outcomes after abdominal rectopexy and Delorme's procedure for rectal prolapse: A prospective study. *J. Clin. Diagn. Res.* 2014, 8, NC04–NC07.
27. Dyrberg, D.; Nordentoft, T.; Rosenstock, S. Laparoscopic posterior mesh rectopexy for rectal prolapse is a safe procedure in older patients: A prospective follow-up study. *Scand. J. Surg.* 2015, 104, 227–232.
28. Madbouly, K.M.; Youssef, M. Laparoscopic ventral rectopexy versus laparoscopic wells rectopexy for complete rectal prolapse: Long-term results. *J. Laparoendosc. Adv. Surg. Tech.* 2018, 28, 1–6.
29. Cutait, D. Sacro-promontory fixation of the rectum for complete rectal prolapse. *Proc. R. Soc. Med.* 1959, 52, 105.
30. Gourgiotis, S.; Baratsis, S. Rectal prolapse. *Int. J. Color. Dis.* 2006, 22, 231–243.
31. Aitola, P.T.; Hiltunen, K.-M.; Matikainen, M.J. Functional results of operative treatment of rectal prolapse over an 11-year period: Emphasis on transabdominal approach. *Dis. Colon Rectum* 1999, 42, 655–660.
32. Sohn, V.; Steele, S.R.; Mellgren, A. Rectopexy with mesh according to the Ripstein technique. In *Rectal Prolapse: Diagnosis and Clinical Management*; Altomare, D.F., Pucciani, F., Eds.; Springer: Milan, Italy, 2008; pp. 121–129.
33. Hsu, A.; Brand, M.I.; Saclarides, T.J. Laparoscopic rectopexy without resection: A worthwhile treatment for rectal prolapse in patients without prior constipation. *Am. Surg.* 2007, 73, 858–861.
34. Jonkers, H.A.F.; Draaisma, W.A.; Wexner, S.D.; Broeders, I.A.M.J.; Bemelman, W.A.; Lindsey, I.; Consten, E.C.J. Evaluation and surgical treatment of rectal prolapse: An international survey. *Color. Dis.* 2012, 15, 115–119.
35. Ashari, L.H.S.; Lumley, J.W.; Stevenson, A.R.L.; Stitz, R.W. Laparoscopically-assisted resection rectopexy for rectal prolapse: Ten years' experience. *Dis. Colon Rectum* 2005, 48, 982–987.
36. Jonkers, H.A.F.; Maya, A.; Draaisma, W.A.; Bemelman, W.A.; Broeders, I.A.; Consten, E.C.J.; Wexner, S.D. Laparoscopic resection rectopexy versus laparoscopic ventral rectopexy for complete rectal prolapse. *Tech. Coloproctology* 2014, 18, 641–646.
37. D'Hoore, A.; Cadoni, R.; Penninckx, F. Long-term outcome of laparoscopic ventral rectopexy for total rectal prolapse. *BJS* 2004, 91, 1500–1505.
38. Samaranayake, C.; Luo, C.; Plank, A.W.; Merrie, A.E.H.; Plank, L.; Bissett, I.P. Systematic review on ventral rectopexy for rectal prolapse and intussusception. *Color. Dis.* 2009, 12, 504–512.
39. Boons, P.; Collinson, R.; Cunningham, C.; Lindsey, I. Laparoscopic ventral rectopexy for external rectal prolapse improves constipation and avoids de novo constipation. *Color. Dis.* 2009, 12, 526–532.
40. Gurland, B. Ventral Mesh Rectopexy: Is this the new standard for surgical treatment of pelvic organ prolapse? *Dis. Colon Rectum* 2014, 57, 1446–1447.
41. Panis, Y. Laparoscopic ventral rectopexy: Resection or no resection? That is the question. *Tech. Coloproctology* 2014, 18, 611–612.
42. Delorme, R. Sur le traitement des prolapsus du rectum totaux pour l'excision de la musculature rectale ou rectocolique. *Bull. Mem. Soc. Paris* 1900, 26, 499–518.

43. Varma, M.; Rafferty, J.; Buie, W.D. Practice parameters for the management of rectal prolapse. *Dis. Colon Rectum* 2011, 54, 1339–1346.
44. Elagili, F.; Gurland, B.; Liu, X.; Church, J.; Ozuner, G. Comparing perineal repairs for rectal prolapse: Delorme versus Altemeier. *Tech. Coloproctology* 2015, 19, 521–525.
45. Marchal, F.; Bresler, L.; Ayav, A.; Zarnegar, R.; Brunaud, L.; Duchamp, C.; Boissel, P. Long-term results of Delorme's procedure and orr-loygue rectopexy to treat complete rectal prolapse. *Dis. Colon Rectum* 2005, 48, 1785–1790.
46. Watts, A.M.; Thompson, M.R. Evaluation of Delorme's procedure as a treatment for full-thickness rectal prolapse. *Br. J. Surg.* 2000, 87, 218–222.
47. Fleming, F.J.; Kim, M.J.; Gunzler, D.; Messing, S.; Monson, J.R.T.; Speranza, J.R. It's the procedure not the patient: The operative approach is independently associated with an increased risk of complications after rectal prolapse repair. *Color. Dis.* 2012, 14, 362–368.
48. Youssef, M.; Thabet, W.; El Nakeeb, A.; Magdy, A.; Alla, E.A.; El Nabeey, M.A.; Fouda, E.Y.; Omar, W.; Farid, M. Comparative study between Delorme operation with or without postanal repair and levateroplasty in treatment of complete rectal prolapse. *Int. J. Surg.* 2013, 11, 52–58.
49. Cernuda, R.B.; Ángel, J.P.; Fernández, N.T.; Sánchez-Farpón, J.H.; Pérez, J.A. Álvarez Perineal rectosigmoidectomy (altemeier procedure) as treatment of strangulated rectal prolapse. *J. Gastrointest. Surg.* 2016, 20, 2102–2103.
50. Voulimeneas, I. Perineal rectosigmoidectomy for gangrenous rectal prolapse. *World J. Gastroenterol.* 2010, 16, 2689–2691.
51. Tiengtianthum, R.; Jensen, C.C.; Goldberg, S.M.; Mellgren, A. Clinical outcomes of perineal proctectomy among patients of advanced age. *Dis. Colon Rectum* 2014, 57, 1298–1303.
52. Trompetto, M.; Tutino, R.; Luc, A.R.; Novelli, E.; Gallo, G.; Clerico, G. Altemeier's procedure for complete rectal prolapse; outcome and function in 43 consecutive female patients. *BMC Surg.* 2019, 19, 1.
53. Mistrangelo, M.; Tonello, P.; Contul, R.B.; Arnone, G.; Passera, R.; Grasso, L.; Rapetti, L.; Borroni, R.; Pozzo, M.; Roveroni, M.; et al. Perineal stapled prolapse resection for full-thickness external rectal prolapse: A multicentre prospective study. *Color. Dis.* 2016, 18, 1094–1100.
54. Sehmer, D.; Marti, L.; Wolff, K.; Hetzer, F.H. Midterm results after perineal stapled prolapse resection for external rectal prolapse. *Dis. Colon Rectum* 2013, 56, 91–96.
55. Tschuor, C.; Limani, P.; Nocito, A.; Dindo, D.; Clavien, P.-A.; Hahnloser, D. Perineal stapled prolapse resection for external rectal prolapse: Is it worthwhile in the long-term? *Tech. Coloproctology* 2013, 17, 537–540.
56. Fan, K.; Cao, A.M.; Barto, W.; De Lacavalerie, P. Perineal stapled prolapse resection for external rectal prolapse: A systematic review and meta-analysis. *Color. Dis.* 2020, 22, 1850–1861.