

Phosphorus-Substituted Pyridines

Subjects: **Chemistry, Organic**

Contributor: Yulia Volkova , Igor Zavarzin

Phosphorus(V)-substituted pyridine was first synthesized by Plazek's research group in 1936 by a reaction of 2-dimethylaminopyridine with phosphorus trichloride under oxidative conditions. Later, related compounds were obtained by reactions of metalated pyridines with phosphorus-halogen compounds, of pyridinediazonium tetrafluoroborate with phosphorus trichloride, of *N*-alkoxypyridines with sodium diethyl phosphite and phosphines, and of *N*-pyridylpyridines with phosphonic acid, by the Michaelis–Arbuzov reaction, Pd(II)-catalyzed phosphorylation of halopyridines, cyclization of phosphorus-containing 3-azatrienes, and the Diels–Alder reaction involving 3-phosphoryl-1-azadienes.

heterocycles

pyridine

Michael reaction


phosphorylation

carbenoid-mediated reactions

formal cycloaddition


1. Cyclizations Based on the Michael Reaction

In recent years, synthesis of pyridines via the Michael reaction using phosphoryl-substituted Michael acceptors and donors has gained significant attention. Allais et al. [1][2][3] described three-component condensation of 1,3-dicarbonyl compounds **3** with P(O)Et₂-bearing vinyl ketones **2** and ammonium acetate (**1**) in the presence of oxygen, leading to pyridine-2-phosphonates **4** (Scheme 1). The authors suggested that the reaction occurs through successive addition of compound **3** to Michael acceptors **2**, **providing** ketone **5** followed by enamination of the latter with ammonia (**1'**) to form intermediate **6**, which undergoes intramolecular cyclization into dihydropyridine **7**. Complete oxidation of intermediate **7** to pyridine **4** was achieved using oxygen in the presence of activated carbon. The reaction was general with respect to β-oxo esters and β-oxoamides, providing products in 49–80% yields.



Scheme 1. Synthesis of pyridine-2-phosphonates.


Hanashalshahaby and Unaleroglue [4] showed that pyridine-3-phosphonates **10** can be obtained by three-component oxidative coupling of diethyl (2-oxobutyl)phosphonate (**8**) with Mannich bases **9** and ammonium acetate (**1**) in the presence of catalyst K-10 (Scheme 2). The product yields are reasonable both for aryl- and alkyl-substituted Mannich bases. The authors hypothesized that β -keto phosphonate **8** reacts with ammonia **1'** generated in situ from ammonium acetate (**1**) to form enamine **11**, which is accompanied by thermal decomposition of Mannich base **9**, giving α,β -unsaturated carbonyl compound **9'**. These two intermediates are subjected to the Michael addition to form ketoamine **12**, which undergoes intramolecular cyclization to dihydropyridine **13**, followed by oxidation of the latter with atmospheric oxygen to provide final product **10**.

**Scheme 2.** Reaction of diethyl (2-oxobutyl)phosphonate with Mannich bases.

Further, Abdou et al. [5] found that diethyl (2-amino-2-thioxoethyl)phosphonate acts as an efficient Michael donor in the addition reaction with β -(dimethylamino)vinyl ketone, thus leading to 2-thioxopyridin-3-ylphosphonate. Liao et al. [6] described one example of the aza-Michael reaction between diethyl (3-phenyl-3-oxopropyn-1-yl)phosphonate with methyl 3-aminocrotonate, providing phosphonate-ester-containing pyridine moiety under mild conditions.


2. Carbenoid-Mediated Reactions

Recently, several studies by Park and co-workers demonstrated the prospects of using metal carbenoids in synthesis of phosphoryl-substituted pyridines. They described [7] Rh(II)-catalyzed intramolecular cyclization of δ -diazo oximes to pyridines and showed that this is a facile method for synthesis of pyridine-2-phosphonate **15** (Scheme 3). $\text{Rh}_2(\text{CF}_3\text{CONH})_4$ was used as the catalyst of choice. The authors proposed a mechanism that involves reaction of the diazo group of oxime **14** with Rh(II), providing rhodium carbenoid **16**, followed by insertion of the latter into the N–O bond to form dihydropyridinone **17**, which undergoes aromatization to final pyridine **15** via elimination of methanol.



Scheme 3. Rh(II)-catalyzed cyclization of δ -diazo oxime ether.


In another work by Park and co-workers [8], synthesis of pyridine-2-phosphonate **20** was accomplished using phosphorylated vinyl carbenoid **21** generated in situ from diazophosphonate **19** and dirhodium(II) catalyst $\text{Rh}_2(\text{esp})_2$ (Scheme 4). Addition of compound **21** to 2*H*-azirine **18** affords intermediate **22**. The latter undergoes three-membered ring opening accompanied by elimination of the Rh(II) catalyst to provide 3-azahexatriene **23**, followed by cyclization to yield pyridine **20**.



Scheme 4. Reaction of phosphorylated carbenoid with 2*H*-azirine.

3. Formal [2+2+2]-Cycloaddition

An approach towards phosphoryl-substituted pyridines, which is probably one of the most versatile, is based on the formal [2+2+2]-cycloaddition. Tanaka's research group [9][10] described synthesis of annulated pyridine-2-phosphonates **26** based on rhodium(I)/biaryl-bisphosphine-complex-catalyzed cycloaddition of 1,6- and 1,7-diynes **24** with diethyl phosphorocyanidate (**25**) (Scheme 5). The reaction has a broad scope with respect to diynes since quaternary carbon-, methylene-, nitrogen-, and oxygen-linked internal 1,6-diynes and terminal biaryl-linked 1,7-diynes can be involved in the heterocyclization. Steric and electronic variations in diynes had minimal impact on the efficacy of the reaction, but, in some cases, using unsymmetrically substituted diynes, the reactions afforded mixtures of regioisomers. The authors proposed rhodium cyclopentadiene **27** or rhodium azacyclopentadiene **28** as two possible key intermediates in the reaction (Scheme 5). Ring expansion of both cyclopentadienes can provide seven-membered intermediate **29**, and reductive elimination of Rh(I)⁺ from the latter accomplishes formation of the final product **26**. Based on the outcome of the enantioselective version of the reaction, the authors were inclined to believe that the main reaction pathway involves intermediate **28**.




Scheme 5. Formal [2+2+2]-cycloaddition of 1,6- and 1,7-diynes with diethyl phosphorocyanidate.

4. Phosphorylation of Pyridines

Phosphorylation is definitely the most general route to phosphorus-substituted pyridines. Due to a wide range of available phosphorylating agents and the possibility of performing the reaction in a nucleophilic, electrophilic, or radical manner, phosphorylation has attracted great attention. Such methods as the Arbuzov reaction, Hirao coupling, palladium-catalyzed cross-coupling of diethyl phosphonate with halogen-substituted heterocycles, and C(sp²)H-phosphorylation of heterocycles with diethyl phosphonates promoted by one-electron oxidants were extensively developed in past decades. Their specific applications in recent years are discussed in more detail later in the text.

4.1. Radical Phosphorylation of Pyridines


P-centered radicals can easily be generated through hydrogen atom transfer or single-electron transfer using peroxides, metal salts, and photocatalysts. Thus, radical phosphorylation has become an efficient strategy for synthesis of structurally diverse phosphorus(V)-substituted pyridines. Due to environmental friendliness and potential industrial application, photocatalytic radical phosphorylation of the pyridine ring received extra attention. In 2018, Yuan et al. [11] reported synthesis of 2- and 3-phosphine oxide-substituted pyridines **31** and **33** (Scheme 6, lines a,b) from 2- and 3-halopyridines **30** and **32** by photocatalytic functionalization with secondary phosphine oxides in the presence of ^tBuOK. The reaction occurred under mild conditions using irradiation with a blue-light-emitting diode. The scope of this transformation is somewhat limited to pyridines with electron-donating substituents and their analogs with an extended π -system. The plausible reaction mechanism involves formation of a complex of halopyridine with potassium *tert*-butoxide **34**, absorption of a light quantum, transition to an excited state **35**, and electron transfer from *tert*-butoxide to a pyridine ring to form the *tert*-butoxy radical and halopyridine radical anion **36**. Elimination of halide from radical anion **36** affords aryl radical **37**. Simultaneously, the *tert*-butoxy radical causes the proton abstraction from secondary phosphine to form a phosphorus-containing radical. Recombination of the latter radical with an aryl radical affords the final product. Recently, the approach proposed by Yuan et al. was expanded [12][13], including electron-primed photoredox [14] and visible light-induced nickel-catalyzed photoredox [15] conditions for radical generation.



Scheme 6. Photocatalytic phosphorylation of halopyridines: (a) Synthesis of 2-phosphine oxide-substituted pyridines; (b) Synthesis of 3-phosphine oxide-substituted pyridines.


In 2019, the photocatalytic radical reaction of pyridylazo sulfones **38** with triphenyl phosphite giving pyridine-3-phosphonates **39** was described by Qiu et al. [16] (Scheme 7). The reaction occurred in the presence of water. A variety of substituted pyridines can be efficiently employed in this reaction. The proposed mechanism for this transformation involves excitation of arylazo sulfone **38** under visible light to provide radical **40**, its decomposition

into sulfonyl and aryl radicals, along with extrusion of a nitrogen molecule. Then, aryl radical **41** reacts with triphenyl phosphate to form phosphorus-centered radical **42**, followed by its oxidation with the sulfonyl radical and elimination of phenol to provide the final product **39**.




Scheme 7. Synthesis of pyridine-3- and pyridine-4-phosponates from pyridylazo sulfones.

Independently, in 2019, Kim et al. [17] described site-selective synthesis of pyridines **44** bearing phosphine oxide moieties at C-4 using radical coupling of *N*-ethoxypyridinium salts **43** with secondary phosphine oxides under photocatalytic conditions (Scheme 8). Further, 3-Diphenylphosphoryl-6-methoxy-1-methyl-2(1*H*)-quinolinone under blue-light-emitting diode illumination was used as a photocatalyst and potassium persulfate as an oxidant. Examination of the reaction scope revealed that a variety of electron-withdrawing and electron-donating substituted pyridines, as well various aryl-substituted phosphine oxides, were tolerated. The plausible mechanism of this reaction involves single-electron transfer from the photocatalyst to *N*-ethoxypyridinium **43**, giving radical **45**, which undergoes decomposition accompanied by elimination of pyridine. The remaining ethoxy radical abstracts a proton from phosphine, followed by addition of the resulting phosphinyl radical **46** to *N*-ethoxypyridinium **43**. The subsequent deprotonation of intermediate **47** and elimination of the new ethoxy radical from intermediate **48** afford target product **44**. The origin of reaction chemoselectivity was revealed by DFT calculations, showing that phosphinoyl radicals are too large for providing an electrostatic attraction between the its oxo functionality and the pyridine nitrogen crucial for *ortho* functionalization.



Scheme 8. Photocatalytic reaction of *N*-ethoxypyridinium salts with phosphine oxides.

Apart from the approaches based on photocatalytic radical phosphorylation, several examples using metal salts for P-radical generation were described. Huang and co-workers [18] developed a CH-phosphorylation method for synthesis of pyridinyl-2-phosphonates **50** based on Ag(I)-catalyzed reaction of pyridines **49** with dialkyl phosphonates using potassium persulfate as an oxidant (Scheme 9). An interesting feature of this reaction is that it involves subsequent treatment of the reaction mixture with sodium thiosulfate, which makes it possible to significantly increase the product yield due to a reduction of pyridine *N*-oxide formed as a by-product. In the authors' opinion, this transformation proceeds through a radical pathway and begins with oxidation of Ag(I) to Ag(II) with persulfate, followed by oxidation of dialkyl phosphite with Ag(II) to form radical cation **5**. Addition of the latter at 2 position of the pyridine ring provides intermediate **52**. Subsequent abstraction of two protons accompanied by oxidation of radical **52** affords the final product. Recently, Kittikool et al. [19] expanded the scope of this phosphorylation approach to 2-pyridones using $Mn(OAc)_2$ as the catalyst.




Scheme 9. Oxidative coupling of pyridines with dialkyl phosphites in the presence of Ag(I).

Noteworthy also is a one-pot three-step protocol for synthesis of diphenyl(pyridin-2-yl)phosphine oxide based on the KOH-promoted oxidative radical phosphorylation of 2-bromopyridine with diphenylphosphine developed by Chen et al. [20].


4.2. Nucleophilic Phosphorylation of Pyridines

Studies by Trofimov's research group [21][22][23] have contributed to recent progress in nucleophilic phosphorylation of pyridines at the 4 position. They accomplished synthesis of 4-phosphine oxide-substituted pyridines **55** by coupling of pyridines **53** with secondary phosphine oxides using diphenyl ethynyl ketone (**54**) as the oxidant (Scheme 10). A variety of substituted aryl and alkyl phosphine oxides can efficiently be employed in this reaction. The proposed mechanism for this transformation involves aza-Michael reaction of pyridine **53** with acetylene **54**, providing intermediate **56**, and deprotonation of phosphine, resulting in formation of the anion. Subsequent addition of a phosphine anion at the 4 position of the activated pyridine ring affords dihydropyridine **57**. The latter undergoes isomerization to intermediate **58**, followed by elimination of alkene **59** to form the final product. In this approach, 3-phenyl-2-propynenitrile can also be applied as an oxidant [22]. The reaction can be stopped at the step of 1,4-dihydropyridines **57** using terminal acylacetylenes [23]. It is interesting that 1,4-dihydropyridines are shown to be produced through 2-4 migration of the POR₂ group in 1,2-dihydro adducts during vinylation/phosphorylation of pyridines [23]. DFT calculations supported the hypotheses that 1,4-dihydropyridines are the thermodynamic products while their 1,2-regioisomers are the kinetic ones [23].




Scheme 10. Coupling of pyridines with secondary phosphine oxides in the presence of acetylenes.

Direct nucleophilic phosphorylation at the 2 position of the pyridine ring was described in 2012 by Oka et al. [24] A series of 2-phosphinate-substituted pyridines **61** were prepared from *N*-methoxypyridinium tosylates **60** by reaction with secondary phosphinates (Scheme 11). Presumably, the reaction proceeds via the $S_N\text{Ar}$ mechanism. Further, 1,8-Diazabicyclo[5.4.0]undec-7-ene (DBU) in acetonitrile at low temperature was found to be the optimal base for chemoselective transformation. The procedure was general for thiophosphinate, providing 2-pyridyl thiophosphinate. The 2-pyridyl phosphinate derivatives produced by this method are P-chiral, and the reaction can be accomplished with high diastereoselectivity using chiral phosphines.




Scheme 11. Reaction between *N*-methoxypyridinium tosylates and secondary phosphines.

In 2014, Wang et al. [25] showed that synthesis of pyridine-2-phosphonates can be accomplished directly from pyridine *N*-oxides and dimethyl phosphate over long-term heating in toluene. Independently, Lee et al. [26] developed an approach to synthesize diethyl pyridine-2-phosphonates **63** by reaction of pyridine *N*-oxides **62** with triethyl phosphite in the presence of ethyl chloroformate (Scheme 12). This reaction presumably starts with activation of pyridine *N*-oxide **62** by chloroformate, followed by addition of triethyl phosphite to the resulting pyridinium salt **64** to form intermediate **65**. The nucleophilic attack of the chloride anion at the ethyl moiety followed by elimination of ethyl carbonate from 1,2-dihydropyridine **66** affords the final product. This approach is general for preparation of quinolin-2-ylphosphonates. An apparent limitation of this approach is formation of phosphorylated isomeric mixtures in the case of 3-bromopyridine *N*-oxide.



Scheme 12. Reaction of pyridine *N*-oxides with triethyl phosphite.

Recently, Tsantrizos and co-workers [27] proposed a route to P-chiral 2-phosphine oxide-substituted pyridines (Scheme 13). Using the reaction of (*R*)-*N*-(1-(5-chloro-2-hydroxyphenyl)ethyl)-4-methylbenzenesulfonamide (**67**) with Grignard to generate chiral phosphorylating agent **68**, they accomplished enantioselective synthesis of products **70** from 2-fluoropyridines **69**.




Scheme 13. Asymmetric synthesis of tertiary pyridine-containing phosphine oxides.

4.3. Transition-Metal-Catalyzed Phosphorylation of Pyridines

Transition metals are widely used to increase efficiency of conventional non-catalyzed phosphorylation of heterocycles [28]. In the past decade, significant progress in the field of phosphorylation of pyridines was made due to application of palladium and nickel catalysis, which enabled expansion of the range of substrates active in phosphorylation reactions to involve halogen-, hydroxyl-, boronic acid-, triflate-, nonaflate-, ester-substituted, trimethylammonium pyridines, etc. It was also shown that transition-metal-catalyzed phosphorylation with phosphine oxides, phosphates, and phosphites could be efficiently used for construction of functionalized pyridines, including chiral structures. Meanwhile, transition-metal-catalyzed C–H phosphorylation of pyridines still remains an unsolved challenge.


Halopyridines (Hal = Cl, Br, and I) are efficient precursors for synthesis of 2,3,4-POR₂-substituted pyridines under palladium catalysis conditions. In 2017, Han et al. [29] reported stereoselective palladium-catalyzed cross-coupling

of 2-bromopyridine with chiral *tert*-butyl-containing phosphine oxides (Scheme 14). Use of $\text{Pd}_2(\text{dba})_3$ with the dppp ligand enabled synthesis of tertiary pyridine-containing phosphine oxides **71** with excellent selectivity. In 2016, Dziuganowska et al. [30] used the same catalytic system to prepare a series of phosphonopyridinecarboxylic acid esters from appropriate bromides.



Scheme 14. Synthesis of chiral tertiary pyridine-containing phosphine oxides.

Mykhailiuk and co-workers [31] expanded the scope of this reaction using the $\text{Pd}_2(\text{dba})_3$ /Xantphos catalytic system in the reaction of bromo(iodo)pyridines with dimethylphosphine oxide (Scheme 15). The scope of this reaction was found to be quite general and enables facile preparation of 2-, 3-, and 4-dimethylphosphine-oxide-substituted pyridines **72** in up to good yields. A variety of electron-withdrawing and electron-donating aryl-substituted pyridines successfully participated in the reaction.



29 examples

2-phosphorylated products:

$\text{R} = \text{H}, 3\text{-NH}_2, 4\text{-NH}_2, 5\text{-NH}_2, 6\text{-NH}_2, 5\text{-CN}, 4\text{-CO}_2\text{Me}, 5\text{-CO}_2\text{Me}, 4\text{-CH(OMe)}_2, 5\text{-CH(OMe)}_2$

3-phosphorylated products:

$\text{R} = \text{H}, 2\text{-F}, 5\text{-F}, 6\text{-F}, 5\text{-Cl}, 6\text{-Cl}, 2\text{-NH}_2, 4\text{-NH}_2, 5\text{-NH}_2, 6\text{-NH}_2, 2\text{-CN}, 5\text{-CN}, 6\text{-CN}, 5\text{-CO}_2\text{Me}$


4-phosphorylated products:

$\text{R} = \text{H}, 2\text{-F}, 2\text{-Cl}, 2\text{-NH}_2, 2\text{-CN}, 2\text{-CO}_2\text{Me}$

Scheme 15. Cross-coupling of halopyridines with dimethylphosphine oxide.


Borisova and co-workers [32] accomplished synthesis of bis(phosphoryl)pyridines and 2,2'-bipyridines using palladium acetate/dppf-catalyzed cross-coupling of chloropyridines with secondary phosphine oxides. Ligand-free microwave-assisted $\text{Pd}(\text{OAc})_2$ -catalyzed phosphorylation of bromopyridines with diphenylphosphine oxide or diethyl phosphite was described by Henyecz et al. [33]. Catalyst $\text{Pd}(\text{OAc})_2$ in the presence of sodium iodide as the promoter was also shown to be an effective catalyst for ligand-free coupling of diphenylphosphine oxide with pyridinium nonaflate [34]. An example of $\text{Pd}(\text{PPh}_3)_4$ -catalyzed coupling of 2-bromo-6-pyrrolylpyridine with diethyl phosphite was recently reported by Ti et al. [35].

The palladium-catalyzed Hirao coupling of bromopyridines with triethyl phosphite enables preparation of pyridinephosphonates. Adam et al. [36] used the palladium-catalyzed cross-coupling of 3-bromopyridines **73** with triethyl phosphite to synthesize 2-aminopyridine-3-phosphonates **74** (Scheme 16). It is worth noting that the reaction was performed in the absence of the solvent during short-term heating to 160–180 °C.




Scheme 16. Pd(II)-catalyzed cross-coupling of 3-bromopyridines with triethyl phosphite.

Apart from halopyridines, pyridine boronic acids and hydroxypyridines were involved in phosphorylation under palladium catalysis. Zou, Wu, and co-workers [37] constructed a C–P bond by means of cross-coupling of pyridine boronic acids **75** with dialkyl phosphonates under PdCl_2 catalysis conditions (Scheme 17). The reaction requires the presence of Ag_2O as the oxidant. The method can be used to prepare structurally diverse pyridine-3- and pyridine-4-phosphonates **76**.




Scheme 17. Cross-coupling of pyridine carboxylic acids with dialkyl phosphites.

Recently, Ding and co-workers [38] described the general palladium-catalyzed one-pot procedure for synthesis of phosphonates, phosphinates, and phosphine oxides from phenols mediated by sulfuryl fluoride. The reaction was efficient for functionalization of 2- and 3-hydroxypyridines **77** with diethyl phosphonate (Scheme 18). According to the proposed mechanism, the fluorosulfates generated in situ are key intermediates in synthesis of pyridines **78**.




Scheme 18. Replacement of the OH group of hydroxypyridines mediated by sulfonyl fluoride.

Before these works, in 2012, direct replacement of an OH group of hydroxypyridines by a diphenylphosphoryl moiety was achieved by Zhao et al. [39] under Ni(II) catalysis. A C–O bond was activated using bromotripyrrolidinophosphonium hexafluorophosphate. The activated complex of hydroxypyridine **77** with this salt reacts with diphenylphosphine oxide in the presence of dichloro[1,3-bis(diphenylphosphino)propane]nickel salts as the catalyst to produce pyridines **79** (Scheme 19).

**Scheme 19.** Replacement of the OH group of hydroxypyridines by the diphenylphosphoryl moiety.


Later, the substrate scope of Ni-catalyzed phosphorylation of pyridines with phosphine oxides was expanded to pyridine carboxylic acid phenyl esters [40] and trimethylammonium triflate of pyridine [41]. Cross-coupling of pyridine tosylates [42] and aryltrimethylammonium tetrafluoroborate [43] with phosphites was also implemented.

A relatively general method was proposed by Yamaguchi's research group [40]. They performed cross-coupling of pyridine carboxylic acid phenyl esters **80** with secondary phosphine oxides accompanied by decarbonylation (Scheme 20). The reaction is catalyzed by nickel acetate and requires high temperatures (150–170 °C). This approach is applicable to synthesis of 2- and 3-phosphine-oxide-substituted pyridines **81**, the position of the POR₂ substituent in the product being defined by the position of the carbonyl group in the starting compound. The authors suggested that this transformation initially proceeds through oxidative insertion of Ni(0) into the C–O bond, providing intermediate **82**, and exchange of the phenoxide ligand by phosphine provides **83**. Subsequent decarbonylation of intermediate **83** and reductive elimination from intermediate **84** afford the final product.



Scheme 20. Cross-coupling of aryl carboxylic acid esters with phosphine oxides.

Of interest as well is Ni(II)-catalyzed reaction of tosylates **85** with H-phosphonate diesters proposed by Chun-Jing Li ^[42] as an approach to 2- and 3-phosphorylpyridines **86**; $\text{NiCl}_2(\text{cod})_2$ was an optimal catalyst for this transformation (Scheme 21).



Scheme 21. Ni(II)-catalyzed replacement of the tosyl group in pyridines.

References

- Allais, C.; Liéby-Muller, F.; Rodriguez, J.; Constantieux, T. Metal-free michael-addition-initiated three-component reaction for the regioselective synthesis of highly functionalized pyridines: Scope, mechanistic investigations and applications. *Eur. J. Org. Chem.* 2013, 2013, 4131–4145.
- Allais, C.; Constantieux, T.; Rodriguez, J. Use of beta,gamma-unsaturated alpha-ketocarbonyls for a totally regioselective oxidative multicomponent synthesis of polyfunctionalized pyridines. *Chemistry* 2009, 15, 12945–12948.
- Allais, C.; Liéby-Muller, F.; Constantieux, T.; Rodriguez, J. Dual heterogeneous catalysis for a regioselective three-component synthesis of bi- and tri(hetero)arylpyridines. *Adv. Synth. Catal.* 2012, 354, 2537–2544.
- Hanashalshahaby, E.H.; Unaleroglu, C. Mannich bases as enone precursors for water-mediated efficient synthesis of 2,3,6-trisubstituted pyridines and 5,6,7,8-tetrahydroquinolines. *ACS Comb. Sci.* 2015, 17, 374–380.

5. Abdou, W.M.; Shaddy, A.A.; Kamel, A.A. Structure-based design and synthesis of acyclic and substituted heterocyclic phosphonates linearly linked to thiazolobenzimidazoles as potent hydrophilic antineoplastic agents. *Chem. Pap.* 2017, 71, 1961–1973.
6. Liao, L.; Zhang, H.; Zhao, X. Selenium- π -acid catalyzed oxidative functionalization of alkynes: Facile access to ynones and multisubstituted oxazoles. *ACS Catal.* 2018, 8, 6745–6750.
7. Qi, X.; Dai, L.; Park, C.M. Carbenoid-mediated N-O bond insertion and its application in the synthesis of pyridines. *Chem. Commun.* 2012, 48, 11244–12246.
8. Loy, N.S.; Singh, A.; Xu, X.; Park, C.M. Synthesis of pyridines by carbenoid-mediated ring opening of 2H-azirines. *Angew. Chem. Int. Ed.* 2013, 52, 2212–2216.
9. Kashima, K.; Ishii, M.; Tanaka, K. Synthesis of pyridylphosphonates by rhodium-catalyzed cycloaddition of 1,6- and 1,7-diynes with diethyl phosphorocyanidate. *Eur. J. Org. Chem.* 2015, 2015, 1092–1099.
10. Kashima, K.; Teraoka, K.; Uekusa, H.; Shibata, Y.; Tanaka, K. Rhodium-catalyzed atroposelective cycloaddition of ortho-substituted phenyl diynes with nitriles: Effect of ortho substituents on regio- and enantioselectivity. *Org. Lett.* 2016, 18, 2170–2173.
11. Yuan, J.; To, W.P.; Zhang, Z.Y.; Yue, C.D.; Meng, S.; Chen, J.; Liu, Y.; Yu, G.A.; Che, C.M. Visible-light-promoted transition-metal-free phosphinylation of heteroaryl halides in the presence of potassium tert-butoxide. *Org. Lett.* 2018, 20, 7816–7820.
12. Zeng, H.; Dou, Q.; Li, C.-J. Photoinduced transition-metal-free cross-coupling of aryl halides with H-phosphonates. *Org. Lett.* 2019, 21, 1301–1305.
13. Liu, Y.; Guo, W. Visible-light driven C-P bond formation with recyclable carbon nitride photocatalyst. *ChemCatChem* 2022, 14, e202200449.
14. Cowper, N.G.; Chernowsky, C.P.; Williams, O.P.; Wickens, Z.K. Potent reductants via electron-primed photoredox catalysis: Unlocking aryl chlorides for radical coupling. *J. Am. Chem. Soc.* 2020, 142, 2093–2099.
15. Zhu, D.-L.; Jiang, S.; Wu, Q.; Wang, H.; Chai, L.-L.; Li, H.-Y.; Li, H.-X. Visible-light-induced nickel-catalyzed P(O)–C(sp²) coupling using thioxanthen-9-one as a photoredox catalysis. *Org. Lett.* 2020, 23, 160–165.
16. Qiu, D.; Lian, C.; Mao, J.; Ding, Y.; Liu, Z.; Wei, L.; Fagnoni, M.; Protti, S. Visible Light-Driven, Photocatalyst-free Arbuzov-like reaction via arylazo sulfones. *Adv. Synth. Catal.* 2019, 361, 5239–5244.
17. Kim, I.; Kang, G.; Lee, K.; Park, B.; Kang, D.; Jung, H.; He, Y.T.; Baik, M.H.; Hong, S. Site-selective functionalization of pyridinium derivatives via visible-light-driven photocatalysis with quinolinone. *J. Am. Chem. Soc.* 2019, 141, 9239–9248.

18. Xiang, C.B.; Bian, Y.J.; Mao, X.R.; Huang, Z.Z. Coupling reactions of heteroarenes with phosphites under silver catalysis. *J. Org. Chem.* 2012, 77, 7706–7710.
19. Kittikool, T.; Phakdeeyothin, K.; Chantarojsiri, T.; Yotphan, S. Manganese-promoted regioselective direct C3-phosphinoylation of 2-pyridones. *Eur. J. Org. Chem.* 2021, 2021, 3071–3078.
20. Zhou, Y.; Liu, C.; Shen, Z.; Dai, B.; Chen, J. Efficient potassium hydroxide promoted P-arylation of aryl halides with diphenylphosphine. *J. Organomet. Chem.* 2021, 949, 121932.
21. Trofimov, B.A.; Volkov, P.A.; Khrapova, K.O.; Telezhkin, A.A.; Ivanova, N.I.; Albanov, A.I.; Gusarova, N.K.; Chupakhin, O.N. Metal-free site selective cross-coupling of pyridines with secondary phosphine chalcogenides using acylacetylenes as oxidants. *Chem. Commun.* 2018, 54, 3371–3374.
22. Volkov, P.A.; Gusarova, N.K.; Khrapova, K.O.; Telezhkin, A.A.; Albanov, A.I.; Vasilevskiy, S.F.; Trofimov, B.A. A mechanistic insight into the chemoselectivity of the reaction between 3-phenyl-2-propynenitrile, secondary phosphine oxides and pyridinoids. *Mendeleev Commun.* 2021, 31, 670–672.
23. Volkov, P.A.; Telezhkin, A.A.; Khrapova, K.O.; Ivanova, N.I.; Albanov, A.I.; Gusarova, N.K.; Trofimov, B.A. Metal-free S HN cross-coupling of pyridines with phosphine chalcogenides: Polarization/deprotonation/oxidation effects of electron-deficient acetylenes. *New J. Chem.* 2021, 45, 6206–6219.
24. Oka, N.; Ito, K.; Tomita, F.; Ando, K. Synthesis of 2-pyridylphosphinate and thiophosphinate derivatives by nucleophilic aromatic substitution of N-methoxypyridinium tosylates. *Chem. Lett.* 2012, 41, 1630–1632.
25. Wang, H.; Cui, X.; Pei, Y.; Zhang, Q.; Bai, J.; Wei, D.; Wu, Y. Direct regioselective phosphonation of heteroaryl N-oxides with H-phosphonates under metal and external oxidant free conditions. *Chem. Commun.* 2014, 50, 14409–14411.
26. Lee, S.-J.; Kim, H.-S.; Yang, H.-W.; Yoo, B.-W.; Yoon, C.M. Synthesis of diethyl pyridin-2-ylphosphonates and quinolin-2-ylphosphonates by deoxygenative phosphorylation of the corresponding N-oxides. *Bull. Korean Chem. Soc.* 2014, 35, 2155–2158.
27. Li, S.G.; Yuan, M.; Topic, F.; Han, Z.S.; Senanayake, C.H.; Tsantrizos, Y.S. Asymmetric library synthesis of P-chiral t-butyl-substituted secondary and tertiary phosphine oxides. *J. Org. Chem.* 2019, 84, 7291–7302.
28. Budnikova, Y.H.; Sinyashin, O.G.D. Phosphorylation of C–H bonds of aromatic compounds using metals and metal complexes. *Russ. Chem. Rev.* 2015, 84, 917–951.
29. Han, Z.S.; Wu, H.; Xu, Y.; Zhang, Y.; Qu, B.; Li, Z.; Caldwell, D.R.; Fandrick, K.R.; Zhang, L.; Roschangar, F.; et al. General and stereoselective method for the synthesis of sterically

- congested and structurally diverse P-stereogenic secondary phosphine oxides. *Org. Lett.* 2017, 19, 1796–1799.
30. Dziuganowska, Z.A.; Slepokura, K.; Volle, J.-N.; Virieux, D.; Pirat, J.-L.; Kafarski, P. Structural analogues of Selfotel. *J. Org. Chem.* 2016, 81, 4947–4954.
31. Stambirskiy, M.V.; Kostiuik, T.; Sirobaba, S.I.; Rudnichenko, A.; Titikaiev, D.L.; Dmytriv, Y.V.; Kuznietsova, H.; Pishel, I.; Borysko, P.; Mykhailiuk, P.K. Phosphine oxides (– POME₂) for medicinal chemistry: Synthesis, properties, and applications. *J. Org. Chem.* 2021, 86, 12783–12801.
32. Zakirova, G.G.; Mladentsev, D.Y.; Borisova, N.E. Synthesis of chelating tertiary phosphine oxides via palladium-catalysed C–P bond formation. *Tetrahedron Lett.* 2017, 58, 3415–3417.
33. Henyecz, R.; Oroszy, R.; Keglevich, G. Microwave-assisted Hirao reaction of heteroaryl bromides and >P(O)H reagents using Pd(OAc)₂ as the catalyst precursor in the absence of added P-ligands. *Curr. Org. Chem.* 2019, 23, 1151–1157.
34. McErlain, H.; Riley, L.M.; Sutherland, A. Palladium-catalyzed C–P bond-forming reactions of aryl nonaflates accelerated by iodide. *J. Org. Chem.* 2021, 86, 17036–17049.
35. Ma, F.; Zhang, T.-T.; Zhang, Z.-H.; Tong, H.-X.; Yi, X.-Y. Photorelease of nitric oxide in water-soluble diruthenium nitrosyl complexes with phosphonate substituted pyridylpyrrole. *Inorg. Chim. Acta* 2022, 534, 120826.
36. Adam, M.S.S.; Kindermann, M.K.; Köckerling, M.; Heinicke, J.W. Phosphonylation of 2-amino- and 2-amido-3-bromopyridines and 2-amino-3-chloroquinoxalines with triethyl phosphite. *Eur. J. Org. Chem.* 2009, 2009, 4655–4665.
37. Geng, Z.; Zhang, Y.; Zheng, L.; Li, J.; Zou, D.; Wu, Y.; Wu, Y. Pd-catalyzed C–P coupling of heteroaryl boronic acid with H-phosphonate diester. *Tetrahedron Lett.* 2016, 57, 3063–3066.
38. Zhang, G.; Wang, J.; Guan, C.; Zhao, Y.; Ding, C. Nickel- and palladium-catalyzed cross-coupling of aryl fluorosulfonates and phosphites: Synthesis of aryl phosphonates. *Eur. J. Org. Chem.* 2021, 2021, 810–813.
39. Zhao, Y.L.; Wu, G.J.; Han, F.S. Ni-catalyzed construction of C–P bonds from electron-deficient phenols via the in situ aryl C–O activation by PyBroP. *Chem. Commun.* 2012, 48, 5868–5870.
40. Isshiki, R.; Muto, K.; Yamaguchi, J. Decarbonylative C–P bond formation using aromatic esters and organophosphorus compounds. *Org. Lett.* 2018, 20, 1150–1153.
41. Yang, B.; Wang, Z.-X. Ni-Catalyzed C–P coupling of aryl, benzyl, or allyl ammonium salts with P(O)H compounds. *J. Org. Chem.* 2019, 84, 1500–1509.
42. Li, C.-J. Nickel-catalyzed phosphorylation of tosylates. *Russ. J. Gen. Chem.* 2020, 90, 725–730.

43. Li, C.J. Aryltrimethylammonium tetrafluoroborates in nickel-catalyzed C–P bond-forming reactions. *Russ. J. Org. Chem.* 2021, 57, 954–960.
-

Retrieved from <https://encyclopedia.pub/entry/history/show/95496>