

# Non-Thermal Plasma Treatment on Seed

Subjects: Agriculture, Dairy & Animal Science

Contributor: Jaroslav Julak

The use of non-thermal plasmas in agriculture or plant biology has also been widely reported in the last few years. The topics, related to the decontamination of seeds, modification of surface properties, metabolomic pathways, and enzymatic activity, enhancing seed germination and the initial growth, are summarized e.g., in Plants 2021, 10, 1616; <https://doi.org/10.3390/plants10081616> (<https://doi.org/10.3390/plants10081616>). Disease control and mycotoxin degradation were also reported.

Keywords: legumes ; low temperature plasma ; plasma treatment ; seed ; seedling

---

## 1. Introduction

Plasma, called also the fourth state of matter, is a partially or fully ionized gas. A distinction is made between thermal (high temperature, equilibrium) and non-thermal (cold, low temperature, non-equilibrium) plasma. The thermal plasma reaches the temperatures of thousands of Kelvins and occurs in the Sun, lightning, electric sparks, tokamaks, etc. and it is therefore not applicable in biological applications. On the other hand, the non-thermal plasma (NTP), also called low-temperature or cold plasma, occurs at nearly ambient temperature and the high kinetic energy is stored in electrons only. Its biological and also medical applications are very wide and include, among others, disinfection processes, acceleration of blood coagulation and improved wound and infection healing, dental applications or cancer therapy. These are summarized in numerous reviews, such as [1][2][3][4][5][6], or in the comprehensive books of Shintani and Sakudo [7] and Metelmann et al. [8].

The use of non-thermal plasmas in agriculture or plant biology has also been widely reported in the last few years. The topics, related to the decontamination of seeds, modification of surface properties, metabolomic pathways, and enzymatic activity, enhancing seed germination and the initial growth, are summarized e.g., in [9][10][11][12][13][14][15][16]. Plant disease control [17] or mycotoxin degradation [18][19] were also reported. The nature of chemical reactions in NTP is rather complex, see e.g., [20][21][22][23].

NTP may be easily generated in various electric discharges, among which the most commonly used ones are corona discharges, plasma jets (called also plasma needle, plasma torch or plasma pen), dielectric barrier discharges, gliding arcs and microwave discharges. For a general description of plasma sources, see e.g., [24][25][26][27]. In addition, the described effects are not constrained to the direct NTP treatment, but on a lower scale are also mediated by the effects of plasma-activated water (PAW) or air, i.e., the water exposed to NTP prior to the application to desired objects. The described effects can persist for many months after exposure, mainly due to the presence of stable reactive oxygen and nitrogen particles as described e.g., in [28][29][30]. The use of NTP has already been addressed by the authors in relation to wheat (*Triticum aestivum*) [30] and seeds that can be used as raw seed [31].

## 2. Surface Seed and Sprout Decontamination

Both microbial and toxin decontamination of seed surface and plant sprout were included. Runtzel et al. [32] reported the effective fungal inactivation of *Aspergillus parasiticus* and *Penicillium* sp. on the surface of common bean after 10–30 min exposure of dielectric barrier discharge (DBD). Selcuk et al. [33] reported the inactivation of pathogenic fungi — *Aspergillus* sp. and *Penicillium* sp.— by NTP in a SF6 atmosphere on artificially contaminated seeds of common bean, chickpea, lentil and soybean without affecting the cooking time and other food qualities. A significant reduction of 3-log was achieved within 15 min.

A significant reduction of the seed-borne microbial contamination on pea was observed by Khatami and Ahmadiania [34], where the amount of microorganisms decreased by ca 3-log from initial 5.5 to 2.5 log CFU/mL/cm<sup>2</sup> (the units are not explained in the original paper) after 60 s of exposure. Peanut decontamination was reported by Basaran et al. [35], where 1-log and 5-log reductions of *Aspergillus parasiticus* after 5 min treatment in air or a SF 6 atmosphere were observed,

respectively. The decontamination of several leguminous species (blue lupine, goat-rue, honey clover, soybean, and pea) has been studied [36][37], where the plasma treatment contributed to better fungicidal effect against of *Fusarium* sp., *Alternaria* sp., and *Stemphiliium* sp. on seeds. On the other hand, a possible reduction of toxin production on peanut kernels was reported [38], where a significant reduction of aflatoxin levels without any negative sensory effect was reported.

The decontamination of soybean seeds contaminated with bacteria using PAW was reported by Lee et al. [39]. PAW reduced the overall 4.3-log CFU/mL amount of aerobic microbes and 7.0-log CFU/mL of artificially inoculated *Salmonella* Typhimurium within 5 min and 2 min, respectively. Two following works reported the decontamination of sprouts, both by PAW only. Schnabel et al. [40] contaminated mung beans sprouts with the bacteria *Escherichia coli*, *Pseudomonas fluorescens*, *Pseudomonas marginalis*, *Pectobacterium carotovorum*, and *Listeria innocua*. The experimental results showed a reduction from 2.5-log to 3.5-log of bacteria and better growth of the mung bean sprouts, while untreated samples became strongly glassy and cell liquor was released, no influence of treated samples was observed. Similar results with mung bean were reported also by Xiang et al. [41], where reductions of 2.3 to 2.8-log were observed in aerobic bacteria, yeasts and moulds.

### **3. Effects on Seed Surface Properties**

The applications of NTP or PAW also affect the properties of samples. These changes appeared to be beneficial and are listed below, and may be divided into seed surface properties and seed internal content properties. The primary effect on the surface is a decrease of the surface energy leading to better wettability or higher hydrophilicity, as measured by the contact angle of water droplets. This change in the wetting properties of seeds is at least partially due to oxidation of their surface by NTP.

Bormashenko et al. [42][43] reported a contact angle decrease in common beans and their markedly accelerated water absorption after tens of seconds of cold radiofrequency plasma treatment. The treatment leads to hydrophilization of the cotyledon and tissues constituting the seed coat when they are exposed to plasma separately. On the contrary, when the entire seed is exposed to plasma treatment, only the external surface of the common bean is hydrophilized by the cold plasma. All these statements are in agreement with the results of [44][45], where the authors observed pea seeds. SEM and Fourier transform infrared (FTIR) surface analyses showed small changes in the surface layer caused by the oxidation of lipids and polysaccharides (the consequences are mentioned in the original work). Moreover, the result of performed genotoxicological tests also confirmed that the level of DNA damage is minimal. A significant increase in water imbibition was also reported for soybean seeds [46][47].

Surface modification was also reported after treatment by PAW. Fan et al. [48] reported a water absorption rate increase from 65% for control samples to 75% for treated mung beans. Sajib et al. [49] reported a similar lipid or wax coat alteration of black gram due to the interactions of NO<sup>2-</sup> and H<sub>2</sub>O<sub>2</sub> with wax. Zhou et al. [50] confirmed by SEM that the seed coat of mung bean is chapped and that it improves the water and nutrients absorption, which is a condition that could enhance the germination rate of mung bean and promote the growth of hypocotyls and radicles.

### **4. Field Production and Quality Crop Yield**

The following three works enhance the studies up to the level of field experiments. In Tarrad et al. [51], seeds of Egyptian clover (*T. alexandrium* cv. Gemmiza 1 and cv. Fahl) were treated by pulsed atmospheric-pressure plasma jet, that increased the final yield. The total dry matter yield increased by about 15 % and 9 % over the non-treated control for Gemmiza 1 and Fahl cultivars, respectively.

In [52], the authors reported for blue lupine NTP treated seed that due to the decrease of seed infection and stimulation of field germination, to early seedling growth and to plant resistance to pathogens, the yield increased by 27 %. In [53], the authors exposed soybean seeds to NTP in various atmospheres of air, O<sub>2</sub> or N<sub>2</sub>. Under greenhouse conditions, dry weight of roots, plant height, stem diameter and yield of plants grown from either healthy or infected seeds were improved. The plant height, stem diameter and root dry weight of plants from plasma-treated seeds showed increases of 3 %, 8 % and 12 %, respectively; the NTP treatment had positive effects on all the monitored parameters, as compared with either infected plants or fungicide control.

## 5. Conclusions

Non-thermal plasma (NTP) has become a widely used technique in various fields of biology, medicine, food processing and others.

The exposure to NTP or to plasma-activated water (PAW) can significantly affect the different properties of legume seeds. Namely, germination starts from water uptake, and the capability of water absorption could be significantly influenced by the action of plasma. Important surface properties and some physiological parameters of seeds could be also modified. Oxidation processes of plasmatic reactive species may increase water adsorption capability by increasing wettability of seed coats and could also be associated to gas exchanges and to electrolyte leakage by the seed. It is likely that NTP can effectively change dormancy of hard seeds by affecting seed permeability and triggering subsequent processes. NTP can positively influence the germination and growth of the seed, and subsequently also the properties of the seedlings. NTP treatment could reduce the hardness associated with mechanical dormancy of many *Fabaceae* species (alfalfa, blue lupine, grass pea, honey clover, *Mimosa* sp., *Trifolium* sp., etc.). NTP can be advantageously used in decontamination of plant seed surfaces or legume products. Legumes tolerate this physico-chemical treatment well, and the mild stress it causes appears to have a positive effect on them. Changes in physiological factors can then have a positive effect on the number of crops in the field and their yield.

Acknowledgement: This study was supported by Charles University research program Progress Q25.

---

## References

1. Tendero, C.; Tixier, C.; Tristant, P.; Desmaison, J.; Leprince, P. Atmospheric Pressure Plasmas: A Review. *Spectrochim. Acta B*. 2006, 61, 2–30.
2. Bourke, P.; Ziuzina, D.; Boehm, D.; Cullen, P.J.; Keener, K. The Potential of Cold Plasma for Safe and Sustainable Food Production. *Trends Biotechnol.* 2018, 36, 615–626.
3. Julák, J.; Scholtz, V. The Potential for use of Non-thermal Plasma in Microbiology and Medicine. *Epidemiol. Microbiol. Immunol.* 2020, 69, 28–36.
4. Scholtz, V.; Pazlarova, J.; Souskova, H.; Khun, J.; Julák, J. Non-thermal Plasma—A Tool for Decontamination and Disinfection. *Biotechnol. Adv.* 2015, 33, 1108–1119.
5. Zhu, Y.L.; Li, C.Z.; Cui, H.Y.; Lin, L. Feasibility of Cold Plasma for the Control of Biofilms in Food Industry. *Trends Food Sci. Technol.* 2020, 99, 142–151.
6. von Woedtke, T.; Schmidt, A.; Bekeschus, S.; Wende, K.; Weltmann, K.D. Plasma Medicine: A Field of Applied Redox Biology. *In Vivo* 2019, 33, 1011–1026.
7. Shintani, H.; Sakudo, A. Gas Plasma Sterilization in Microbiology: Theory, Applications, Pitfalls and New Perspectives; Caister Academic Press: Poole, UK, 2016.
8. Metelmann, H.R.; von Woedtke, T.; Weltmann, K.D. Comprehensive Clinical Plasma Medicine: Cold Physical Plasma for Medical Application; Springer: Cham, Switzerland, 2018; ISBN 331967627X.
9. Holubová, L.; Kyzek, S.; Ďurovcová, I.; Fabová, J.; Horváthová, E.; Ševčíčková, A.; Gálová, E. Non-Thermal Plasma—A New Green Priming Agent for Plants? *Int. J. Mol. Sci.* 2020, 21, 9466.
10. Scholtz, V.; Šerá, B.; Khun, J.; Šerý, M.; Julák, J. Effects of Nonthermal Plasma on Wheat Grains and Products. *J. Food Qual.* 2019, 1–10.
11. Magallanes Lopez, A.M.; Simsek, S. Pathogens Control on Wheat and Wheat Flour: A Review. *Cereal Chem.* 2021, 98, 17–30.
12. Siddique, S.S.; Hardy, G.S.J.; Bayliss, K.L. Cold Plasma: A Potential New Method to Manage Postharvest Diseases Caused by Fungal Plant Pathogens. *Plant Pathol.* 2018, 67, 1011–1021.
13. Han, Y.; Cheng, J.H.; Sun, D.W. Activities and Conformation Changes of Food Enzymes Induced by Cold Plasma: A Review. *Crit. Rev. Food Sci. Nutr.* 2019, 59, 794–811.
14. Ekezie, E.F.G.; Chizoba, F.G.; Sun, D.W.; Cheng, J.H. A Review on Recent Advances in Cold Plasma Technology for the Food Industry: Current Applications and Future Trends. *Trends Food Sci. Technol.* 2017, 69, 46–58.
15. Pankaj, S.K.; Wan, Z.; Keener, K.M. Effects of Cold Plasma on Food Quality: A Review. *Foods* 2018, 7, 4.

16. Attri, P.; Ishikawa, K.; Okumura, T.; Koga, K.; Masaharu, S.M. Plasma Agriculture from Laboratory to Farm: A Review. *Processes* 2020, 8, 1002.
17. Adhikari, B.; Pangomm, K.; Veerana, M.; Mitra, S.; Park, G. Plant Disease Control by Non-Thermal Atmospheric-Pressure Plasma. *Front. Plant Sci.* 2020, 11, 77.
18. Ten Bosch, L.; Pfohl, K.; Avramidis, G.; Wieneke, S.; Viöl, W.; Karlovsky, P. Plasma-Based Degradation of Mycotoxins Produced by *Fusarium*, *Aspergillus* and *Alternaria* Species. *Toxins* 2017, 9, 97.
19. Čolović, R.; Puvača, N.; Cheli, F.; Avantaggiato, G.; Greco, D.; Duragić, O.; Kos, J.; Pinotti, L. Decontamination of Mycotoxin-Contaminated Feedstuffs and Compound Feed. *Toxins* 2019, 11, 617.
20. Graves, D.B. The Emerging Role of Reactive Oxygen and Nitrogen Species in Redox Biology and Some Implications for Plasma Applications to Medicine and Biology. *J. Phys. D Appl. Phys.* 2012, 45, 263001.
21. Kelly, S.; Turner, M. Atomic oxygen patterning from a biomedical needle-plasma source. *J. Appl. Phys.* 2013, 114, 123301.
22. Sysolyatina, E.; Mukhachev, A.; Yurova, M.; Grushin, M.; Karalnik, V.; Petryakov, A.; Trushkin, N.; Ermolaeva, S.; Akish, Y. Role of the Charged Particles in Bacteria Inactivation by Plasma of a Positive and Negative Corona in Ambient Air. *Plasma Process. Polym.* 2014, 11, 315–334.
23. Liao, X.Y.; Cullen, P.J.; Muhammad, A.I.; Jiang, Z.M.; Ye, X.Q.; Liu, D.H.; Ding, T. Cold Plasma-Based Hurdle Interventions: New Strategies for Improving Food Safety. *Food Eng. Rev.* 2020, 12, 321.
24. Yousfi, M.; Merbahi, N.; Sarrette, J.P.; Eichwald, O.; Ricard, A.; Gardou, J.P.; Ducasse, O.; Benhenni, M. Non Thermal Plasma Sources of Production of Active Species for Biomedical Uses: Analyses, Optimization and Prospect. In *Biomedical Engineering Frontiers and Challenges*; Reza, F.R., Ed.; Intech Europe: Rijeka, Croatia, 2011; pp. 99–124.
25. Ehlbeck, J.; Schnabel, U.; Polak, M.; Winter, J.; von Woedtke, T.; Brandenburg, R.; von dem Hagen, T.; Weltmann, K.D. Low Temperature Atmospheric Pressure Plasma Sources for Microbial Decontamination. *J. Phys. D Appl. Phys.* 2011, 44, 013002.
26. Simoncicova, J.; Krystofova, S.; Medvecká, V.; Durisova, K.; Kalinakova, B. Technical Applications of Plasma Treatments: Current State and Perspectives. *Appl. Microbiol. Biotechnol.* 2019, 103, 5117–5129.
27. Khun, J.; Scholtz, V.; Hozák, P.; Fitl, P.; Julák, J. Various DC-driven Point-to-plane Discharges as Non-Thermal Plasma Sources and their Bactericidal Effects. *Plasma Sources Sci. Technol.* 2018, 27, 065002.
28. Julák, J.; Hujacova, A.; Scholtz, V.; Khun, J.; Holada, K. Contribution to the Chemistry of Plasma-Activated Water. *Plasma Phys. Rep.* 2018, 44, 125–136.
29. Al-Sharify, Z.T.; Al-Sharify, T.A.; Al-Azawi, A.M. Investigative Study on the Interaction and Applications of Plasma Activated Water (PAW). *IOP Conf. Ser. Mater. Sci. Eng.* 2020, 870, 012042.
30. Zhou, R.; Zhou, R.; Wang, P.; Xian, Y.; Prochnow, A.M.; Lu, X.; Cullen, J.P.; Ostrikov, K.; Bazaka, K. Plasma-Activated Water: Generation, Origin of Reactive Species and Biological Applications. *J. Phys. D Appl. Phys.* 2020, 53, 303001.
31. Sera, B.; Sery, M. Non-Thermal Plasma Treatment as a New Biotechnology in Relation to Seeds, Dryfruits, and Grains. *Plasma Sci. Technol.* 2018, 20, 044012.
32. Runtzel, C.L.; da Silva, J.R.; da Silva, B.A.; Moecke, E.S.; Scussel, V.M. Effect of Cold Plasma on Black Beans (*Phaseolus vulgaris*, L.), Fungi Inactivation and Micro-Structures stability. *Emir. J. Food Agric.* 2019, 31, 864–873.
33. Selcuk, M.; Oksuz, L.; Basaran, P. Decontamination of Grains and Legumes Infected with *Aspergillus* spp. and *Penicillium* spp. by Cold Plasma Treatment. *Bioresour. Technol.* 2008, 99, 5104–5109.
34. Khatami, S.; Ahmadiania, A. Increased Germination and Growth Rates of Pea and Zucchini Seed by FSG Plasma. *J. Theor. Appl. Phys.* 2018, 12, 33–38.
35. Basaran, P.; Basaran-Akgul, N.; Oksuz, L. Elimination of *Aspergillus Parasiticus* from nut Surface with Low Pressure Cold Plasma (LPCP) Treatment. *Food Microbiol.* 2008, 25, 626–632.
36. Filatova, I.; Azharonok, V.; Kadyrov, M.; Beljavsky, V.; Gvozdev, A.; Shik, A.; Antonuk, A. The Effect of Plasma Treatment of Seeds of Some Grain and Legumes on their Sowing Quality and Productivity. *Rom. J. Phys.* 2011, 56, 139–143.
37. Filatova, I.; Azharonok, V.; Shik, A.; Antoniuk, A.; Terletskaia, N. Fungicidal Effects of Plasma and Radio-Wave Pre-treatments on Seeds of Grain Crops and Legumes. In *Nanomaterials for Security*; Springer Science and Business Media LLC: Cham, Switzerland, 2011; pp. 469–479.
38. Iqdam, B.M.; Abuagela, M.O.; Boz, Z.; Marshall, S.M.; Goodrich-Schneider, R.; Sims, C.A.; Marshall, M.R.; MacIntosh, A.J.; Welt, B.A. Effects of Atmospheric Pressure Plasma Jet Treatment on Aflatoxin Level, Physicochemical Quality, and Sensory Attributes of Peanuts. *J. Food Process. Preserv.* 2020, 44, e14305.

39. Lee, E.J.; Khan, M.S.I.; Shim, J.; Kim, Y.J. Roles of Oxides of Nitrogen on Quality Enhancement of Soybean Sprout During Hydroponic Production Using Plasma Discharged Water Recycling Technology. *Sci. Rep.* 2018, 8, 16872.
40. Schnabel, U.; Sydow, D.; Schlüter, O.; Andrasch, M.; Ehlbeck, J. Decontamination of Fresh-cut Iceberg Lettuce and Fresh Mung Bean Sprouts by Non-Thermal Atmospheric Pressure Plasma Processed Water (PPW). *Mod. Agric. Sci. Technol.* 2015, 1, 23–39.
41. Xiang, Q.; Liu, X.; Liu, S.; Ma, Y.; Xu, C.; Bai, Y. Effect of Plasma-Activated Water on Microbial Quality and Physicochemical Characteristics of Mung Bean Sprouts. *Innov. Food Sci. Emerg. Technol.* 2019, 52, 49–56.
42. Bormashenko, E.; Grynyov, R.; Bormashenko, Y.; Drori, E. Cold Radiofrequency Plasma Treatment Modifies Wettability and Germination Speed of Plant Seeds. *Sci. Rep.* 2012, 2, 741.
43. Bormashenko, E.; Shapira, Y.; Grynyov, R.; Whyman, G.; Bormashenko, Y.; Drori, E. Interaction of Cold Radiofrequency Plasma with Seeds of Beans (*Phaseolus vulgaris*). *J. Exp. Bot.* 2015, 66, 4013–4021.
44. Stolárik, T.; Henselová, M.; Martinka, M.; Novák, O.; Zahoranová, A.; Černák, M. Effect of Low-Temperature Plasma on the Structure of Seeds, Growth and Metabolism of Endogenous Phytohormones in Pea (*Pisum sativum* L.). *Plasma Chem. Plasma Process.* 2015, 35, 659–676.
45. Svubova, R.; Kyzek, S.; Medvecká, V.; Slovakova, L.; Galova, E.; Zahoranova, A. Novel Insight at the Effect of Cold Atmospheric Pressure Plasma on the Activity of Enzymes Essential for the Germination of Pea (*Pisum sativum* L. cv. Prophet) Seeds. *Plasma Chem. Plasma Process.* 2020, 40, 1221–1240.
46. Li, L.; Jiang, J.; Li, J.; Shen, M.; He, X.; Shao, H.; Dong, Y. Effects of Cold Plasma Treatment on Seed Germination and Seedling Growth of Soybean. *Sci. Rep.* 2014, 4, 5859.
47. Perez-Piza, M.C.; Cejas, E.; Zilli, C.; Prevosto, L.; Mancinelli, B.; Santa-Cruz, D.; Yannarelli, G.; Balestrasse, K. Enhancement of Soybean Nodulation by Seed Treatment with Non-Thermal Plasmas. *Sci. Rep.* 2020, 10, 4917.
48. Fan, L.M.; Liu, X.F.; Ma, Y.F.; Xiang, Q.S. Effects of Plasma-Activated Water Treatment on Seed Germination and Growth of Mung Bean Sprouts. *J. Taibah Univ. Sci.* 2020, 14, 823–830.
49. Sajib, S.A.; Billah, M.; Mahmud, S.; Miah, M.; Hossain, F.; Omar, F.B.; Roy, N.C.; Hoque, K.M.F.; Talukder, M.R.; Kabir, A.H.; et al. Plasma Activated Water: The Next Generation Eco-Friendly Stimulant for Enhancing Plant Seed Germination, Vigor and Increased Enzyme Activity, a Study on Black Gram (*Vigna mungo* L.). *Plasma Chem. Plasma Process.* 2020, 40, 119–143.
50. Zhou, R.; Li, J.; Zhou, R.; Zhang, X.; Yang, S. Atmospheric-Pressure Plasma Treated Water for Seed Germination and Seedling Growth of Mung Bean and its Sterilization Effect on Mung Bean Sprouts. *Innov. Food Sci. Emerg. Technol.* 2019, 53, 36–44.
51. Tarrad, M.M.; Ahmed, G.; Zayed, E.M. Response of Egyptian Clover Ecotypes to the Non-Thermal Plasma Radiation. *Range Manag. Agrofor.* 2011, 32, 9–14.
52. Filatova, I.; Lyushkevich, V.; Goncharik, S.; Zhukovsky, A.; Krupenko, N.; Kalatskaja, J. The Effect of Low-Pressure Plasma Treatment of Seeds on the Plant Resistance to Pathogens and Crop Yields. *J. Phys. D Appl. Phys.* 2020, 53, 244001.
53. Perez-Piza, M.C.; Prevosto, L.; Grijalba, P.E.; Zilli, C.G.; Cejas, E.; Mancinelli, B.; Balestrasse, K.B. Improvement of Growth and Yield of Soybean Plants Through the Application of Non-Thermal Plasmas to Seeds with Different Health Status. *Heliyon* 2019, 5, e01495.