Physical Activity Patterns

Subjects: Sport Sciences

Contributor: Thayse Natacha Gomes , Peter T. Katzmarzyk , Sara Pereira , Mabliny Thuany , Martyn Standage , José Maia

Although "physical activity pattern" (PAP) is a widely used expression, its precise meaning remains vague. Indeed, PAP is sometimes used to describe physical activity (PA) levels/intensities, however the term is also applied to express how PA averages differ among group, between different days (i.e., week and weekend days), across seasons (e.g., summer, winter), and between school and vacation time. A lack of a formal definition for PAP has not prevented researchers from using a vast array of instruments to measure PA (accelerometers, pedometers, questionnaires, direct observations, and heart rate monitors) in an attempt to capture PAP. Discrepancies also exist in how PAP has been expressed with researchers using daily mean PA, heart rate frequency, number of daily steps, time engaged in certain types of activity, and/or engagement in sports activities. Researchers have also used numerous statistical procedures in their attempts to capture PAP. Notwithstanding this diversity, we suggest that PAP should be used to jointly capture similarities/dissimilarities as well as stabilities/changes in children's PA at an intra-personal level. PAP should be used to best describe individual streams of behaviours, and not exclusively PA levels/intensities. A formal and clear definition of PAP would support important, systematic, and coherent lines of empirical enquiry.

physical activity pattern

children health

1. Introduction

It is well-acknowledged that organized and systematic physical activity (PA) engagement provides numerous health benefits to the paediatric population^{[1][2]}. Notwithstanding such evidence, epidemiological data show an apparent non-random decline in youth PA levels. Further, a large amount of children/adolescents do not comply with the international recommendations^[3]. Since children do not engage in PA in a similar manner^{[4][5][6][7][8][9]}, a large number of studies not only dedicate their efforts to trying to understand PA patterns, but also on trying to explain differences in PA conditioned on their manifold predictors, and linked them to health benefits^{[10][11]}.

2. Precise Definition of PAP

To date, there is no consensus regarding a precise definition of PAP. Yet, this has not prevented researchers from using different instruments, procedures, and protocols in an attempt to capture PAP^{[7][8][11][12]}, nor from relying on a variety of statistical procedures to analyse their data. It is evident from the extant literature that researchers have yet to conceptualise, capture, assess, and operationalise PAP in a manner that will allow systematic lines of research and practice to develop^{[8][11][12][13][14][15]}.

Therefore, we conducted a systematic review on studies involving children aged 6-11 years. Our focus was on: (1) definitions of PAP; (2) instruments and techniques used to describe PAP; (3) statistical procedures used to analyse PAP; and (4) implications of PAP to children's health.

From reviewing the available papers (n=76), it is evident that authors do not use a common definition of PAP in their studies, with no paper providing a formal definition of PAP. Yet, this did not prevent authors from attempting to research PAP by considering children's involvement in different PA intensities (sedentary, light, moderate, vigorous, very vigorous), the type of activity children performed, or even their sports participation, for example. Further, most papers considered the compliance with PA guidelines as an indicator of PAP. We also identified studies that explored PA averages, and their differences into distinct daily segments across week and weekend days, or even across different seasons suggesting that this way PAP could be identified and comprehensively described.

A wide range of instruments were also used to measure PAP, including accelerometers^{[8][9][12][16][17][18][19][20][21][22]} [23][24][25][26][27][28][29][30][31][32][33][34][35][36][37][38][39][40][41][42][43][44], pedometers^{[45][46][47][48][49][50][51]}, heart rate monitors^{[52][53][54][55][56][57][58][59][60]}, questionnaires/diaries^{[10][11][61][62][63][64][65][66][67][68][69][70][71][72][73]}, observational methods^{[74][75]}, and GPS^[6]. Yet, most of them did not explicitly describe a formal statistical method/model to determine children's PAP. This absence and/or difficulty did not prevent researchers from linking their putative PAP with health indicators. It is important to note that papers (n=24) were seemingly more interested in studying the relationship between PA levels and health (or health indicators), rather than PAP *per se*. This is the main reason behind the use of PA markers such as moderate to vigorous PA, different PA intensities children are engaged with, and the compliance with PA guidelines with health indicators. Notwithstanding this commendable effort, it is important to highlight that we are still void of robust data, as well as putative explanatory mechanisms on how PAP influence children's physical and psychological health and well-being.

A pattern can be generically defined as "the regular way in which something happens, develops, or is done"^[76], and as such children's PAP should be examined at an intra-personal level beyond a thoroughly description of their PA levels. We contend that it is important to present a formal and clear definition of PAP to guide future research in children and adolescents. Further, we suggest that PAP may also search for streams of activity/sedentary behaviours to identify differences, or not, across a variety of conditions that rule children and adolescents' lives, meaning that PAP should be used to best describe individual streams of personal behaviours, not only PA intensities. In addition, it is of relevance that researchers working with PA to probably focus on providing more extensive answers to issues relating to the relationship between different PAP manifestations and health (risk and benefits) in children. For example, is it better to (1) be engaged in 60 min of continuous MVPA or in various time fractions alongside the day or (2) to have a more erratic PAP profile (i.e., with random bursts of MVPA)? These and other questions could be answered if a consensual definition of PAP would exist, which in all likelihood would guide research in a more enriching and impactful way, aiming to provide children opportunities to be safely active, during different moments of their days.

References

- 1. Poitras, V.J., et al., Systematic review of the relationships between objectively measured physical activity and health indicators in school-aged children and youth. Applied Physiology, Nutrition, and Metabolism 2016. 41(6 Suppl 3): p. S197-239.
- 2. World Health Organization, Global recommendation on physical activity for health. 2010, Geneva: WHO Press.
- 3. Hallal, P.C., et al., Global physical activity levels: surveillance progress, pitfalls, and prospects. Lancet, 2012. 380(9838): p. 247-57.
- 4. Bauman, A.E., et al., Correlates of physical activity: why are some people physically active and others not? Lancet, 2012. 380(9838): p. 258-71.
- Van Hecke, L., et al., Variation in population levels of physical activity in European children and adolescents according to cross-European studies: a systematic literature review within DEDIPAC. International Journal of Behavioral Nutrition & Physical Activity, 2016. 13: p. 70.
- Bürgi, R., et al., Spatial physical activity patterns among primary school children living in neighbourhoods of varying socioeconomic status: a cross-sectional study using accelerometry and Global Positioning System. BMC Public Health, 2016. 16(1): p. 1-12.
- 7. Ruiz, R.M., et al., A novel approach to characterize physical activity patterns in preschool-aged children. Obesity, 2013. 21(11): p. 2197-203.
- 8. Remmers, T., et al., Daily Weather and Children's Physical Activity Patterns. Medicine & Science in Sports & Exercise, 2017. 49(5): p. 922-929.
- 9. Ridgers, N.D., J. Salmon, and A. Timperio, Too hot to move? Objectively assessed seasonal changes in Australian children's physical activity. International Journal of Behavioral Nutrition & Physical Activity, 2015. 12(1): p. 1-8.
- Vergara-Castaneda, A., et al., Overweight, obesity, high blood pressure and lifestyle factors among Mexican children and their parents. Environmental Health and Preventive Medicine, 2010. 15(6): p. 358-66.
- Muntaner-Mas, A., et al., Obesity and physical activity patterns among Balearic Islands children and adolescents: A cross-sectional study. Journal of Human Sport and Exercise, 2017. 12(2): p. 333-348.
- 12. Laguna, M., et al., Obesity and physical activity patterns in children and adolescents. Journal of Paediatrics and Child Health, 2013. 49(11): p. 942-9.
- 13. Rowlands, A.V. and D.R. Hughes, Variability of physical activity patterns by type of day and season in 8-10-year-old boys. Research Quarterly for Exercise and Sport, 2006. 77(3): p. 391-5.

- 14. Jago, R., et al., Physical activity and sedentary behaviour typologies of 10-11 year olds. International Journal of Behavioral Nutrition & Physical Activity, 2010. 7: p. 59-68.
- 15. Davison, K.K., T.M. Cutting, and L.L. Birch, Parents' activity-related parenting practices predict girls' physical activity. Medicine & Science in Sports & Exercise, 2003. 35(9): p. 1589-95.
- 16. Miller, W.C., J.G. Redmond, and A.T. Vaux-Bjerke, Activity patterns and perceptions about active transport to school. American Journal of Health Behavior, 2013. 37(2): p. 190-8.
- 17. Arto, G., Children's segment specific light physical activity across two years of school-based program. Journal of Physical Education and Sport, 2015. 15(1): p. 88-95.
- 18. Aznar, S., et al., Patterns of physical activity in Spanish children: a descriptive pilot study. Child Care, Health and Development, 2011. 37(3): p. 322-8.
- Baquet, G., et al., Improving physical activity assessment in prepubertal children with highfrequency accelerometry monitoring: A methodological issue. Preventive Medicine, 2007. 44(2): p. 143-147.
- Bielemann, R.M., et al., Objectively Measured Physical Activity in Children From a Southern Brazilian City: A Population-Based Study. Journal of Physical Activity & Health, 2013. 10(8): p. 1145-1152.
- Blaes, A., et al., Physical activity patterns in French youth--from childhood to adolescence-monitored with high-frequency accelerometry. American Journal of Human Biology, 2011. 23(3): p. 353-8.
- 22. Butte, N.F., et al., Longitudinal effects of parental, child and neighborhood factors on moderatevigorous physical activity and sedentary time in Latino children. International Journal of Behavioral Nutrition & Physical Activity, 2014. 11: p. 1-21.
- 23. Fairclough, S.J., et al., School day segmented physical activity patterns of high and low active children. BMC Public Health, 2012. 12(1): p. 406-417.
- 24. Guerra, S., et al., Assessment of Children's and Adolescents' Physical Activity Levels. European Physical Education Review, 2003. 9(1): p. 75-85.
- 25. Hoos, M.B., et al., Physical activity pattern of children assessed by triaxial accelerometry. European Journal of Clinical Nutrition, 2004. 58(10): p. 1425-8.
- 26. Jago, R., et al., Better with a Buddy: Influence of Best Friends on Children's Physical Activity. Medicine & Science in Sports & Exercise, 2011. 43(2): p. 259-265.
- 27. Jáuregui, A., et al., The physical activity level of Mexican children decreases upon entry to elementary school. Salud Pública de México, 2011. 53(3): p. 228-236.

- 28. Loprinzi, P.D., et al., Markers of adiposity among children and adolescents: Implications of the isotemporal substitution paradigm with sedentary behavior and physical activity patterns. Journal of Diabetes and Metabolic Disorders, 2015. 14(1): p. 46.
- 29. McGall, S.E., M.R. McGuigan, and C. Nottle, Contribution of free play towards physical activity guidelines for New Zealand primary school children aged 7-9 years. British Journal of Sports Medicine, 2011. 45(2): p. 120-4.
- 30. Mooses, K., et al., Different Methods Yielded Two-Fold Difference in Compliance with Physical Activity Guidelines on School Days. PLoS One, 2016. 11(3): p. e0152323.
- Muthuri, S.K., et al., Comparative study of physical activity patterns among school children in Kenya and Canada: Results from the ISCOLE Project. African Journal for Physical, Health Education, Recreation & Dance, 2014. 20(2.2): p. 765-779.
- 32. Nyberg, G.A., et al., Physical activity patterns measured by accelerometry in 6- to 10-yr-old children. Medicine & Science in Sports & Exercise, 2009. 41(10): p. 1842-8.
- 33. Page, A., et al., Physical activity patterns in nonobese and obese children assessed using minuteby-minute accelerometry. International Journal of Obesity, 2005. 29(9): p. 1070-1076.
- 34. Pau, M., et al., Influence of School Schedules on Physical Activity Patterns in Primary School Children: A Case Study in Italy. Journal of Physical Activity & Health, 2017. 14(7): p. 501-505.
- 35. Ridgers, N.D., et al., Examining influences on boy's and girls' physical activity patterns: the A-CLASS project. Pediatric Exercise Science, 2010. 22(4): p. 638-50.
- Stone, M.R., et al., Physical activity patterns of children in Toronto: the relative role of neighbourhood type and socio-economic status. Canadian Journal of Public Health, 2012. 103(9 Suppl 3): p. eS9-14.
- Walter, C.M., In-school physical activity patterns of primary school learners from disadvantaged schools in South Africa. African Journal for Physical, Health Education, Recreation & Dance, 2011: p. 779-788.
- 38. Williams, K.D., et al., Quantitative physical activity assessment of children and adolescents in a rural population from Eastern Nepal. American Journal of Human Biology, 2016. 28(1): p. 129-37.
- 39. Aadland, E., et al., Associations of volumes and patterns of physical activity with metabolic health in children: A multivariate pattern analysis approach. Preventive Medicine, 2018. 115: p. 12-18.
- 40. Fetter, D.S., et al., Effect of the Shaping Healthy Choices Program, a Multicomponent, School-Based Nutrition Intervention, on Physical Activity Intensity. Journal of the American College of Nutrition, 2018. 37(6): p. 472-478.
- 41. Saint-Maurice, P.F., et al., Youth Physical Activity Patterns During School and Out-of-School Time. Children, 2018. 5(9).

- 42. Briceño, G., et al., Physical activity patterns associated with a pedagogical intervention in six to eight year old children in an urban school. Cogent Education, 2019. 6(1): p. 1-10.
- 43. Huang, W.Y. and S.H.S. Wong, Prospective associations between weekend catch-up sleep, physical activity, and childhood obesity. Childhood Obesity, 2019. 15(1): p. 40-47.
- 44. McLellan, G., et al., Segmented sedentary time and physical activity patterns throughout the week from wrist-worn ActiGraph GT3X+ accelerometers among children 7–12 years old. Journal of Sport and Health Science, 2019.
- 45. Brusseau, T.A., et al., Pedometer-determined segmented physical activity patterns of fourth- and fifth-grade children. Journal of Physical Activity & Health, 2011. 8(2): p. 279-86.
- 46. Brusseau, T.A., The Intricacies of Children's Physical Activity. Journal of Human Kinetics, 2015. 47: p. 269-75.
- 47. Brusseau, T.A., et al., Daily physical activity patterns of children living in an American Indian community. Journal of Physical Activity & Health, 2013. 10(1): p. 48-53.
- 48. Brusseau, T.A., et al., Children's step counts on weekend, physical education, and non-physical education days. Journal of Human Kinetics, 2011. 27(1): p. 123-134.
- 49. Czajka, K., et al., Assessment of physical activity by pedometer in polish preschool children. Human Movement, 2015. 16(1): p. 15-19.
- Dauenhauer, B.D. and X.D. Keating, The influence of physical education on physical activity levels of urban elementary students. Research Quarterly for Exercise and Sport, 2011. 82(3): p. 512-20.
- 51. Johnson, T.G., et al., School day physical activity patterns of Pima Indian children in two communities. Research Quarterly for Exercise and Sport, 2007. 78(4): p. 364-8.
- 52. Armstrong, N. and S. Bray, Primary school children's physical activity patterns during autumn and summer. Bulletin of Physical Education, 1990. 26(3): p. 23-26.
- 53. Armstrong, N. and S. Bray, Physical activity patterns defined by continuous heart rate monitoring. Archives of Disease in Childhood, 1991. 66(2): p. 245-7.
- 54. Armstrong, N., et al., Physical activity patterns and aerobic fitness among prepubescents. / Types d'activite physique et aptitude physique aerobie chez des enfants prepuberes. European Physical Education Review, 1996. 2(1): p. 19-29.
- 55. Benefice, E., Physical activity and anthropometric and functional characteristics of mildly malnourished Senegalese children. Annals of Tropical Paediatrics, 1992. 12(1): p. 55-66.
- 56. Benefice, E. and G. Ndiaye, Relationships between anthropometry, cardiorespiratory fitness indices and physical activity levels in different age and sex groups in rural Senegal (West Africa).

Annals of Human Biology, 2005. 32(3): p. 366-82.

- 57. Loftin, M., et al., Comparison and relationship of VO2peak and physical activity patterns in elementary and high school females. Pediatric Exercise Science, 1998. 10(2): p. 153-163.
- 58. Massin, M.M., et al., Patterns of physical activity defined by continuous heart rate monitoring among children from Liege. Acta Clinica Belgica, 2004. 59(6): p. 340-5.
- Zarrouk, F., et al., Physical activity patterns and estimated daily energy expenditures in normal and overweight tunisian schoolchildren. Journal of Sports Science & Medicine, 2009. 8(1): p. 83-8.
- 60. Gilliam, T.B., et al., Exercise programs for children: a way to prevent heart disease? Physician & Sportsmedicine, 1982. 10(9): p. 96-101.
- 61. Brewer, M. and R.T. Kimbro, Neighborhood context and immigrant children's physical activity. Social Science & Medicine, 2014. 116: p. 1-9.
- Ha, A., et al., Eating and physical activity practices in risk of overweight and overweight children: Compliance with US Department of Agriculture food guide pyramid and with National Association for Sport and Physical Activity guidelines for children. Nutrition Research, 2005. 25(10): p. 905-915.
- 63. Amini, M., et al., Childhood obesity and physical activity patterns in an urban primary school in thailand. Rawal Medical Journal, 2009. 34(2): p. 203-206.
- 64. Antonogeorgos, G., et al., Physical activity patterns and obesity status among 10- to 12-year-old adolescents living in Athens, Greece. Journal of Physical Activity & Health, 2010. 7(5): p. 633-40.
- 65. Bathrellou, E., et al., Physical activity patterns and sedentary behaviors of children from urban and rural areas of Cyprus. Central European Journal of Public Health, 2007. 15(2): p. 66-70.
- Colin-Ramirez, E., et al., Outcomes of a school-based intervention (RESCATE) to improve physical activity patterns in Mexican children aged 8-10 years. Health Education Research, 2010. 25(6): p. 1042-9.
- 67. Swaminathan, S., et al., Longitudinal trends in physical activity patterns in selected urban south Indian school children. The Indian Journal of Medical Research, 2011. 134(8): p. 174-80.
- 68. Schmidt, G.J., J.J. Walkuski, and D.J. Stensel, The Singapore Youth Coronary Risk and Physical Activity Study. Medicine & Science in Sports & Exercise, 1998. 30(1): p. 105-13.
- 69. Dickenson, B., The physical activity patterns of young people the implications for P.E. Bulletin of Physical Education, 1986. 22(1): p. 36-38.
- 70. Riddoch, C., et al., Long term health implications of fitness and physical activity patterns. Archives of Disease in Childhood, 1991. 66(12): p. 1426-33.

- 71. Esht, V., et al., A preliminary report on physical activity patterns among children aged 8-14 years to predict risk of cardiovascular diseases in Malwa region of Punjab. Indian Heart Journal, 2018.
- 72. Bosch, L.S.M.M., et al., Associations of extracurricular physical activity patterns and body composition components in a multi-ethnic population of UK children (the Size and Lung Function in Children study): a multilevel modelling analysis. BMC Public Health, 2019. 19(1): p. N.PAG-N.PAG.
- 73. Nwogu, U.B., et al., Bone mineral density in an urban and a rural children population—A comparative, population-based study in Enugu State, Nigeria. BONE, 2019. 127: p. 44-48.
- 74. Berman, N., et al., Spectral and bout detection analysis of physical activity patterns in healthy, prepubertal boys and girls. American Journal of Human Biology, 1998. 10(3): p. 289-297.
- 75. Corbin, C.B. and P. Pletcher, Diet and physical activity patterns of obese and nonobese elementary school children. Research Quarterly of the American Association for Health, Physical Education and Recreation, 1968. 39(4): p. 922-928.
- 76. Longman Exams Dictionary. 2006, Pearson Education Limited: Essex.

Retrieved from https://encyclopedia.pub/entry/history/show/8990