# <sup>13</sup>C Solid-State NMR Application to Waste Composting

Subjects: Green & Sustainable Science & Technology Contributor: Silvia Pizzanelli , Lucia Calucci , Claudia Forte , Silvia Borsacchi

<sup>13</sup>C solid-state Nuclear Magnetic Resonance (SSNMR) has often been applied to follow the transformation of organic matter during waste composting to produce soil amendments, as well as to assess the quality of the products and the effectiveness of the treatment.

humic substances humic acids 13C CP-MAS

#### 1. Introduction

Recycling has become a key issue in waste management. In fact, the globally produced waste presently amounts to 2 billion tons/year and, given the continuous population growth and increase in industrial development, it is expected to reach 3.40 billion tons/year by 2050 <sup>[1][2]</sup>. Considering that the organic fraction represents approximately 45% of the total waste and that it is rich in nutrients and biochemicals, its recycling is certainly of great relevance from a circular economy perspective.

In the past, organic waste was mainly disposed of in landfills with the now well-known environmental impacts <sup>[3]</sup>; alternatively, it was reused as food for animals or directly applied on agricultural lands as fertilizer. More recently, it has been recognized that organic waste can be used as feedstock for platform chemicals <sup>[4][5][6][7]</sup>. Among the techniques suggested for the conversion of organic waste into valuable products, biological treatments for the production of energy (e.g., biogas from anaerobic digestion) and organic soil improvers or fertilizers (compost from aerobic composting and digestate from anaerobic digestion) are those most commonly used <sup>[5][8]</sup>. In the case of application to soil, important features of the product are its stability and maturity, the former indicating low biological activity and high resistance to decomposition, the latter referring to the stabilization of the physicochemical characteristics of the material, the disappearance of phytotoxicity, and the ability to favor seed germination and plant growth <sup>[9][10]</sup>.

Given its ease of implementation and low operational costs, the most commonly used organic waste treatment is composting <sup>[11]</sup>. Composting is the decomposition process of organic matter (OM) performed in aerobic conditions by microorganisms <sup>[12]</sup>, which allows the transformation of organic wastes of different kinds, both vegetal and animal, into a stable organic substance rich in "humus". Humus is constituted by the high molecular weight association of amphiphilic compounds, rich in carboxylic groups, slow to decompose under natural conditions, and is responsible for many physicochemical properties that ensure soil fertility. The composting process is naturally driven by microorganisms, which use OM and oxygen for their metabolism, producing heat,  $CO_2$ ,  $H_2O$ , minerals

and stable OM <sup>[13]</sup>. This process starts with a bio-oxidative phase characterized by a rising temperature, which reaches 45–70 °C. During this phase, which lasts for days, thermophilic microorganisms decompose OM very fast, using the most easily available substances. The high temperature ensures sanitation, i.e., the inactivation of most pathogens and invasive plant seeds. Afterwards, the temperature progressively decreases to mesophilic conditions, and the fresh compost enters a curing phase, which can last for weeks, in which decomposition continues, involving the more recalcitrant compounds (such as lignin) and bringing to the formation of humic compounds (fulvic and humic acids, and humin) with high maturity and stability. Compost has great value as a soil amendment and fertilizer, with applications in agriculture, horticulture, and gardening, and hence represents a perfect example of a circular economy; in fact, the last decades have seen an increase in compost production. Today, in Europe 42 million tons of waste per annum, that is 59% of the separately collected biowaste, both municipal and commercial/industrial, is composted

Understanding the transformations of OM that take place during the composting process is fundamental for the agricultural application of the stabilized waste and for the optimization of process parameters, also considering that national and international regulations have established stringent requirements for the use of this product as fertilizer or soil amendment in terms of organic carbon content, humic substance content, etc. For these reasons, since the 1990s, many studies have been published in the scientific literature testing and comparing different chemical and instrumental methods for assessing compost stability and/or maturity, as, for example, measurement of pH, electrical conductivity, CO<sub>2</sub> evolution rate, seed germination rate, dissolved organic carbon, volatile organic carbon, C/N ratio, and humic acid content (see for instance [15][16][17][18][19][20][21]). To this end, besides conventional analytical methods, advanced instrumental techniques, such as thermal analyses, Fourier transform infrared spectroscopy (FTIR), UV–Vis and fluorescence spectroscopies, nuclear magnetic resonance (NMR) spectroscopy, gas chromatography–mass spectrometry, and scanning electron microscopy, can be used to assess the stabilization of organic matter and the quality of the final product. In fact, these techniques provide detailed information on compositional, functional, and behavioral properties of organic materials, even complex ones such as compost. Nevertheless, it is generally agreed that a combination of analyses is advised for a thorough characterization.

Among the techniques that can give access to compositional information of OM, NMR, and solid-state NMR (SSNMR) in particular, stands out as a powerful method to access comprehensive qualitative and quantitative structural information not available from other techniques. Compared to liquid-state NMR, which is useful for studying soluble components, especially those with low molecular weight (see Refs. <sup>[22][23]</sup> for a comprehensive review), SSNMR has many advantages when applied to OM: (i) it allows bulk samples to be analyzed without pretreatments or the use of solvents, which may introduce artifacts; (ii) it can be used to investigate insoluble OM samples or fractions; (iii) it allows a much higher sample concentration, enhancing signals and saving instrument time; (iv) it is conservative, i.e., it does not consume sample; (vi) it allows the investigation of anisotropic interactions, averaged out in solution NMR by the fast isotropic tumbling of molecules; (v) it allows domains and heterogeneities to be identified within OM structures. The vast majority of SSNMR studies of OM reported in the literature involve the observation of <sup>13</sup>C nuclei, with only a few exploiting <sup>15</sup>N and <sup>31</sup>P. Indeed, <sup>13</sup>C SSNMR yields useful information on OM composition in terms of organic functional groups <sup>[18][24]</sup>.

## 2. <sup>13</sup>C Solid-State NMR Spectroscopy for the Investigation of Organic Matter in Compost

<sup>13</sup>C has a natural abundance of 1.11% <sup>[25],[26]</sup>, which results in a good balance of sensitivity and spectral "simplicity" since complex interactions among <sup>13</sup>C nuclei can be neglected. The most important nuclear property used for the identification of functional groups is chemical shielding. On the basis of this property, nuclei in different chemical environments give rise to signals at different resonance frequencies, expressed as chemical shifts. Experimental techniques providing high-resolution <sup>13</sup>C SSNMR spectra are at present routinely available on SSNMR instruments, the most important being Magic Angle Spinning (MAS) and high-power decoupling from abundant <sup>1</sup>H nuclei. In the most basic experiment, usually called direct polarization MAS or direct excitation MAS (DE-MAS), these techniques are combined with a radiofrequency excitation pulse on <sup>13</sup>C. The <sup>13</sup>C DE-MAS experiment gives the most accurate quantification of the different species in an OM solid sample, provided that it is performed with a sufficiently long recycle delay between consecutive scans (five times the spin-lattice relaxation time,  $T_1$ , of the <sup>13</sup>C nuclei). Since  $T_1$  values for <sup>13</sup>C nuclei in the solid state can be considerably long (even hundreds of seconds), extremely long measurement times are required for acquiring quantitative <sup>13</sup>C DE-MAS spectra with an acceptable signal-to-noise ratio. For this reason <sup>1</sup>H-<sup>13</sup>C cross polarization MAS (CP-MAS) experiments are most commonly used in the investigation of OM samples. In CP-MAS experiments, the <sup>13</sup>C signal is created by magnetization transfer from the abundant <sup>1</sup>H nuclei during a time interval called contact time, with an enhancement by up to a factor of four. Moreover, in CP experiments the recycle delay is determined by the longitudinal relaxation of the <sup>1</sup>H nuclei, which is usually much faster than that of <sup>13</sup>C. This allows recycle delays of a few seconds to be used, with a substantial reduction in the experimental time. It is worth noting that, while CP-MAS is an excellent technique for investigating the gualitative composition of a sample, the relative peak intensities may not be fully preserved. However, it has been reported that in the case of compost <sup>13</sup>C CP-MAS NMR spectra recorded with a contact time of about 1 ms and a recycle delay of a few seconds show a similar intensity loss (<10%), with respect to the quantitative spectrum, for signals of all functional groups <sup>[27]</sup>, allowing these conditions to be used for relative quantitation. This also generally applies in the comparison of a series of spectra of samples with similar composition. Therefore, <sup>13</sup>C CP-MAS NMR experiments with a contact time of 1 ms and a recycle delay of a few seconds are usually employed in works on OM in composts to investigate and compare OM composition in the feedstock (biomasses or wastes) and throughout (or only at the end of) the decomposition process.

The spectra are typically analyzed by distinguishing OM carbons from different functional groups on the basis of the chemical shift. Given the complex and heterogeneous chemical nature of OM, spectra span a chemical shift range of over 200 ppm, with a strong overlap of signals. Therefore, spectral regions, rather than individual signals, are usually identified and assigned to specific organic species: (i) the 0–45 ppm spectral region is ascribed to alkyl carbons, with short chain aliphatic carbons from volatile fatty acids and steroid-like molecules resonating between 0 and 28 ppm <sup>[28]</sup> and long chain aliphatic carbons from plant aliphatic biopolymers (suberin, cutin, waxes, fatty acids) <sup>[29]</sup> and proteins <sup>[30]</sup> between 28 and 45 ppm; (ii) O-alkyl and N-alkyl carbon signals, ascribable to lignin, carbohydrates (i.e., cellulose and hemicellulose), and proteins <sup>[31]</sup>, are observed between 45 and 110 ppm; (iii) the 110–160 spectral region is assigned to aromatic carbons present, for instance, in lignin and polyphenols <sup>[32]</sup>; (iv)

signals of carboxyl carbons in esters, acids, and amides (160–190 ppm) and carbonyl carbons in aldehydes and ketones (190–220 ppm) are observed between 160 and 220 ppm. It must be noted that slightly different spectral region ranges are indicated in the literature by different authors. Some resolved signals are also assigned to functional groups belonging to specific biomacromolecules present in biomass samples <sup>[33],[34]</sup>.

**Table 1.** Spectral regions used in the analysis of <sup>13</sup>C MAS NMR spectra of OM and assignment to carbons indifferent functional groups.

Chemical shift range	Assignment	Conventional region name
0-28 ppm	$\mbox{CH}_3$ and $\mbox{CH}_2$ in short chain and simple aliphatics	Alkyl C
28-45 ppm	$\mbox{CH}_2$ and $\mbox{CH}$ in long aliphatic chains	
45-60 ppm	O-CH <sub>3</sub> ; CH-N; aliphatic quaternary C	
60-95 ppm <sup>a</sup>	C2-C6 in cellulose and hemicellulose; alcohols; ethers	O-Alkyl C
95-110 ppm <sup>b</sup>	C1 of cellulose and hemicellulose; anomeric carbon of polysaccharides	
110-145 ppm <sup>c</sup>	Unsubstituted or alkyl-substituted aromatic C	Aromatic C
145-160 ppm <sup>d</sup>	O,N-substituted aromatics	
160-190 ppm	Carboxylic acids; esters; amides	Carboxyl/carbonyl C
190-220 ppm	Ketones; aldehydes	

### al 3. 13 Cd Solida State NMR Applications to composting as ary C region;

<sup>d</sup>also referred to as O-aryl C region. The literature reports many examples of the application of <sup>13</sup>C MAS NMR to the investigation of compost, covering a very broad range of diverse feedstocks, including, often mixed, animal (cattle, swine, chicken, buffalo) manure <sup>[40]</sup> At the test of the test of the test of the sector of the test of t

Different indices are also used to express OM composition, the most common ones being the Alkyl C/O-Alkyl C (A/OA) ratio, the aromaticity index a enerally defined as:



Fitting to obtain eremaser while spectra of composts (dashed line) and corresponding feedstocks (solid line).

Feedstock compositions are: (a) exhausted grape marc and cattle manure 76:24 (*w*:*w*); (b) grape marc and cattle Several studies on composts, however, instead of investigating the bulk material, focused on extracts Different manure 72:28 (*w*:*w*); (c) exhausted grape marc and poultry manure 67:33 (*w*:*w*). Reproduced from Ref. bulk types of extracts were investigated, that is, humic substances (HSs), dissolved organic matter (DOM), or compost

table 2) Relative substances areastable optime under with gors and with carboxyl and hydroxyl groups bonded to the rings and the alkyl chains. On the basis of their solubility, two different extractable HS fractions can be obtained,

Ref.	Feedstock <sup>b</sup>	Sample	Alkvi C	O-Alk	vl C	Aroma	atic C	Carboxyl/ Carbonvl	of HSs is
u		Ū	<b>,</b> -				O_And	С	1.0. HAs
				CH₃O/CHN	alkyl C	Aryl C	C C		are richer
[ <u>41</u> ]	Olive mill waste/orchard pruning residues	C 200 d	17.39	11.63	45.86	12.31	5.75	7.06	s tend to
	Olive mill waste/animal manure/wool residues	C 200 d	26.82	15.32	38.59	6.93	4.0	8.34	l is a very
	ChM/saw dust (3:1 w:w)	C 56 d	28.38	32.2	26	16.	51		place in
	CM/saw dust (3:1 w:w)	C 56 d	22.13	32.2	20	10.	49		er divided
[ <u>52</u> ]	SM/saw dust (3:1 <i>w</i> : <i>w</i> )	C 56 d	31.62	41.3	36	17.	71		t in water
	Soybean meal/saw dust (3:1 <i>w:w</i> )	C 56 d	37.94	43.	16	18.	40		
	Lemon peel/saw dust (3:1 <i>w:w</i> )	C 56 d	36.23	41.	10	17.	43		
		C1	33.9	13.0	31.3	12.6	2.1	7.2	
		C2	38.2	12.3	26.8	11.5	2.9	8.2	
		C3	23.4	11.7	44.0	12.1	3.7	5.1	
		C4	21.0	11.8	41.1	14.3	3.8	8.1	
		C5	19.0	11.4	42.4	15.5	3.8	7.9	
[ <u>58</u> ]	Wood chips/vegetable R/aromatic plant R	C6	24.6	10.9	38.7	15.1	3.3	7.4	
		C7	35.7	11.8	31.0	11.8	2.5	7.2	
		C8	37.8	12.0	30.1	11.4	2.2	6.5	
		C9	43.6	10.5	28.0	9.5	2.2	6.3	
		C10	34.7	11.5	30.7	10.4	3.1	9.5	
		C11	30.2	11.3	33.6	12.0	5.3	7.7	
[ <u>43</u> ]	SM/ rice straw (4:1 w:w)	FS	24.6	57.	9	7.	7	9.8	
		C 40 d	23.2	55.	2	9.	8	11.8	
	SM/ rice straw (8:1 w:w)	FS	28.8	50.	5	7.	9	12.8	

Ref.	Feedstock <sup>b</sup>	Sample c	Alkyl C	CO-Alk	yl C	Aroma	atic C	Carboxyl/ Carbonyl C
				CH <sub>3</sub> O/CHN	O-/Di-O- alkyl C	Aryl C	O-Aryl C	
		C 40 d	25.6	48.9	9	10	.6	15.0
	SM/rice straw/biochar (8:1:1 <i>w:w</i> )	FS	24.7	40.9	9	24	.7	9.8
		C 40 d	19.2	28.8	8	42	.5	9.5
[ <u>10</u> ]	COF/fresh grass/mature compost (18:80:2 <i>w</i> : <i>w</i> )	C 6 d	33.5	15.2	23.3	14.8	3.2	10
	Solid pig slurry/cotton gin waste (4:3 <i>v</i> : <i>v</i> )	FS	13.5	5.9	74.9	3.7	0.9	1.1
		BT	17.5	8.2	62.7	5.3	1.8	4.5
		AT	26.0	12.3	51.2	4.2	2.5	3.9
		EB	30.1	10.5	48.1	4.4	1.9	5.0
		AM	35.5	13.1	43.8	3.6	0.7	3.4
	Solid pig slurry/cotton gin waste (3:4 <i>v</i> : <i>v</i> )	FS	19.4	9.3	71.6	0.3	0.0	0.0
		BT	13.6	7.1	71.6	5.5	1.0	1.2
<u>[45</u> ]		AT	18.7	4.7	61.9	7.4	3.2	4.1
		EB	26.4	11.4	53.8	5.7	4.5	6.8
		AM	34.1	12.5	44.3	3.8	1.3	4.1
	Solid pig slurry/cotton gin waste (3:7 <i>v:v</i> )	FS	22.9	7.4	63.6	2.3	1.1	2.7
		BT	24.9	9.3	59.1	0.9	2.0	3.9
		AT	21.0	9.3	65.5	1.1	0.9	2.2
		EB	18.4	8.1	69.7	2.1	1.4	0.3
		AM	21.8	9.0	58.8	2.6	2.6	5.2
[ <u>51</u> ]	BM/CM/maize straw/PT (70:30 <i>w</i> : <i>w</i> )	FS	15.4	10.5	58.4	8.6	3.0	4.1

Ref.	Feedstock <sup>b</sup>	Sample c	Alkyl C	O-Alky	/I C	Aroma	atic C	Carboxyl/ Carbonyl C
				CH₃O/CHN	O-/Di-O- alkyl C	Aryl C	O-Aryl C	
		C 108 d	18.5	11.4	48.0	12.0	3.4	6.7
	+bioplastic (1 wt%)	C 108 d	17.6	11.1	47.9	12.3	4.0	7.1
	+bioplastic (2 wt%)	C 108 d	18.9	11.0	49.2	11.6	3.6	5.7
	SM/pumice	FS	34.6	48.5	5	7.	3	9.7
		C 10 d	21.5	60.4	Ļ	10	.5	7.7
		C 20 d	23.6	57.7	7	10	.9	7.7
		C60 d	26.2	48.3	3	12	.9	12.7
	CM/pumice	FS	14.6	64.8	}	12	.4	8.3
[ <u>49]</u>		C 10 d	9.0	70.4	Ļ	12	.9	7.7
		C 20 d	10.	69.2	2	12	.7	7.7
		C60 d	11.7	66.6	6	13	.3	8.5
	ChM/pumice	FS	29.5	52.3	3	8.	2	10.1
		C 10 d	21.5	66.0	)	6.	4	6.1
		C 20 d	21.7	61.6	6	9.	3	7.4
		C60 d	26.8	52.0	)	10	.5	10.7
[ <u>50</u> ]	Exhausted grape marc/CM (76:24 w:w)	FS	30.7	17.0	39.5	3.3	3.5	6.1
		C 28 d	36.4	15.9	31.2	5.7	3.7	7.3
		C105 d	37.9	16.1	28.9	6.3	3.9	6.9
		C 168 d	37.3	14.3	22.4	7.1	4.8	14.2
	Grape marc/CM (72:28 w:w)	FS	33.6	15.3	37.1	2.9	3.7	7.4

Ref.	Feedstock <sup>b</sup>	Sample c	Alkyl C	CO-Alk	kyl C	Aroma	tic C	Carboxyl/ Carbonyl C
				CH <sub>3</sub> O/CHN	O-/Di-O- alkyl C	Aryl C	O-Aryl C	
		C 28 d	34.3	13.7	37.3	3.6	3.6	7.5
		C105 d	33.3	14.8	38.6	3.8	3.0	6.6
		C 168 d	34.9	15.2	34.4	3.3	4.0	8.2
	Exhausted grape marc/PM (67:33 <i>w</i> : <i>w</i> )	FS	30.8	14.2	41.0	3.5	3.5	7.0
		C 28 d	34.4	14.6	35.4	3.2	4.0	8.5
		C105 d	33.9	15.5	35.7	3.7	3.9	7.5
		C 168 d	34.3	15.5	31.5	3.4	5.0	10.2
	СМ	СС	18.2	7.6	48.8	8.9	5.0	11.6
	Broiler litter	СС	17.9	9.1	48	8.0	5.2	11.7
[ <u>42</u> ]	Green waste	СС	22.4	10.7	40.4	8.2	6.3	12.0
	Nitro-humus	CC	19.6	8.8	43	9.6	6.7	12.4
	MSW	CC	23.0	7.6	48.7	7.3	3.6	9.8
		FS	2.15	25.8	67.4	2.9	8	0.99
		C 13 d	1.98	25.6	68.9	3.0	7	0.51
[56]	Rice husk/rice	C 34 d	0.66	25.6	70.8	2.9	8	-
[ <u>00</u> ]	bran/BEM/molasses	C 53 d	1.10	22.6	73.1	3.2	9	-
		C 61 d	1.03	24.0	71.9	3.0	9	-
		C 116 d	0.88	25.7	69.9	2.6	3	0.88
[ <u>55</u> ]	DOW/plant trimming/vegetal R	C 60 d		37.6	50.8	7.2	2	4.3
	(00.40.10 00.00)	C 90 d		30.6	56.1	7.3	3	6.1

Ref.	Feedstock <sup>b</sup>	Sample c	Alkyl C	C O-Alkyl C	Aromatic C	Carboxyl/ Carbonyl C
				CH <sub>3</sub> O/CHN O-/I	Di-O- Aryl C O-Ary C C	1
		C 150 d		45.3 3	7.6 9.6	7.4
		FS	14.9	54.9	20.3	10.0
		C 60 d	17.4	57.4	20.9	10.6
		C 120 d	17.4	48.6	18.6	9.9
		C 240 d	15.2	34.1	16.1	9.2
		C 365 d	13.9	25.7	12.4	6.5
		C 548 d	12.8	22.8	11.3	6.6
[ <u>46</u> ]	CM/rice straw	FS	13.5	56.7	20.1	9.7
		C 148 d	15.2	39.6	18.7	10.0
		FS	15.2	56.4	18.8	9.6
		C 60 d	13.4	47.5	16.5	8.4
		C 120 d	12.6	42.0	17.5	8.5
		C 240 d	12.1	37.2	15.9	8.1
		C 365 d	12.6	35.6	15.3	7.7
		FS	27.2	55.6	9.2	8.0
		C 7 d	18.6	66.1	9.8	5.5
[ <u>44</u> ]	SM/wheat straw (95:5 <i>w</i> : <i>w</i> )	C 14 d	15.2	65.3	12.2	7.3
		C 21 d	16.8	65.2	11.5	6.5
		C 28 d	14.7	63.3	13.6	8.3

Ref.	Feedstock <sup>b</sup>	Sample	Alkyl C	0-/	Alkyl C	Arom	atic C	Carboxyl/ Carbonyl C	
			CI	H₃O/CH	IN O-/Di-O- alkvl C	Aryl C	O-Aryl C	-	of OM
	MSW (composted in spring)	FS	15. <mark>82</mark> ]		59.5	14	.9	7.9	d Baldoc
[ <u>63</u> ]		C 28 d	17.7		59.2	13	6.6	7.4	lucing th
	13	C 42 d	17.5		55.5	16	.2	7.2	V/C ratio
[61]		C 49 d	17.3		55.5	17	.7	5.4	es, amin
	MSW (composted in summer)	FS	16.2		60.4	12	.1	7.8	erimenta
		C 28 d	16.8		60.2	13	.9	6.9	raction on ns, lignin
		C 42 d	[ <u>44]</u> 8.0		56.1	15	.3	6.8	ingly, th
		C 49 d	18.2		56.9	15	.8	6.0	
		C1	28.4	13	45.6	7.4	4.6	14.0	either fo
		C2	25.4		48.6	9.8	4.8	11.4	of matur
		C3	30.3		32.5	11.1	6.3	19.8	
		C4	32.4		38.2	7.9	4.9	<sup>13</sup> 16.6	IAS NMI
[ <u>60</u> ]	Kitchen waste/garden waste	C5	19.2		53.0	12.8	8.2	6.8	spectra
		C6	25.5		40.8	12.6	6.6	14.5	variety o
	[ <u>64</u> ]	C7	27.4		42.8	11.3	6.3	12.2	se in th
		C8	26.7		42.6	9.0	7.9	13.8	ase as a
		C9	14.1		43.6	8.1	12.0	22.2	sition. It i
	[65]	FS	26.9		13 47.9	11.6	4.3	8.3	propose
	. <u> </u>	C 34 d	25.5		52.1	10.5	3.6	8.2	methoxy
[ <u>62</u> ]	MSW	C 76 d	24.7		46.5	13.6	4.9	10.3	to whic
		C 90 d	23.6		42.4	19.0	6.6	11.4	researc r <i>LiaR</i> ) c
		C132 d	23.6		40.4	16.9	7.6	11.4	s variabl
[ <u>59</u> ]	Grape skin	FS	8.7		53.4	17	.5	18.7	ulation o

itrant compounds, such as suberin and cutin molecules, lipids, and waxes. The intensity of the carboxyl region often increases due to the oxidation processes occurring during composting and this is generally related to the increase in the compost cation exchange capacity, of which carboxyl groups are mainly responsible.

Ref		Sample		13		Carboxyl/	es and to	
a	Feedstock <sup>b</sup>	C	Alkyl C	O-Alkyl C	Aromatic C	Carbonyl C	ratio as a	
				CH <sub>3</sub> O/CHN O-/Di-O- alkyl C	Aryl C O-Aryl C C		common	
		C 160					Instance	
	<u>[45][66]</u>	d	10.9	51.8	18.9	17.0	index of	
	Grape seeds	FS	25.0	41.3	13.1	18.7		
	13	C 160					between	
		d	21.7	38.4	[ <u>60]</u> 15.7	22.3	backyard	
	Grape skin and seeds	FS	16.2	49.7	13.2	18.8	ste at the	
		C 160					······	
		d	18.6	43.7	15.9	20.3	) vigorous	
biolog								

Anda et al. [56], in contrast to most of the literature, observed that composting of rice husk resulted in a high O-alkyl References are reported in reverse chronological order. BEM = beneficial effective microorganisms; BM = C content, followed by N-alkyl/methoxyl C, and di-O-alkyl C, attributed to cellulose, with trace amounts of phenolic, buffalo manure; ChM = chicken manure; CM = cattle manure; COF = coffee grounds; DOW = domestic organic carboxyl, and alkyl C, indicating that rice husk is quite resistant to microbial degradation. Such a compost applied waste; MSW = municipal solid waste; PM = poultry manure; P1 = poplar trimmings; R = residues; SM = Swine to an Oxisol significantly increased soil pH and the Ca, Mg, K, Na, and Si content of in situ soil solution, but manure. AM = after maturation; AT = advanced thermophilic phase; B1 = beginning of thermophilic phase; C = decreased the amount of toxic ions (AI, Mn, and Fe). Moreover, this compost was found to increase cocoa growth compost; CC = commercial compost; d = days; EB = end of biooxidative phase; FS = feedstock. up to 37%. It was hypothesized that the high content of organic C of the composted rice husk and the chemical resistance to microbial decomposition could provide special benefit in sustaining an organic C reserve if used as a soil ameliorant, especially in tropical soils where organic matter decomposes rapidly.

Torres-Climent et al. <sup>[50]</sup> examined the composting process of winery-distillery waste, i.e., grape marc (the primary waste of wine production obtained after the pressing process) and exhausted grape marc (the solid waste of alcohol distillation), showing the importance of combining thermal analysis, and FTIR and <sup>13</sup>C SSNMR spectroscopies with classical analytical techniques, the latter being insufficient to study the humification processes of these kinds of feedstocks. Areas of the <sup>13</sup>C CP-MAS NMR spectra, evaluated at the beginning and after two months of composting, showed a relative decrease in the carbohydrates and an increase in the alkyl groups. Interestingly, the obtained compost showed good properties for agronomic uses.

Pane et al. <sup>[58]</sup> used <sup>13</sup>C CP-MAS NMR spectra to characterize the OM composition of eleven different composts obtained from waste derived from aromatic plant production and compared the NMR results with the suppressive properties against *Rhizoctonia solani* and *Sclerotinia sclerotiorum* on garden cress, a vegetable crop appreciated for its nutraceutical and taste-giving components, but threatened by soil-borne pathologies due to its very short life-cycle. The study demonstrated good potential for several of the investigated composts as sustainable remedies, substituting for synthetic chemicals. The analysis of <sup>13</sup>C CP-MAS NMR spectra showed correlations between the best performing composts and the higher humification level. In particular, the removal of essential oils resulted in an improved humification, a preservation of apolar compounds, and a decrease in *LigR*. This beneficial effect was

attributed to the known antioxidant and antimicrobial properties of essential oils, which could have hampered the microbial activity in the processed biomass.

lacomino et al. <sup>[41]</sup> investigated the effect on soil fertility and crop yields in a volcanic Mediterranean soil of two different composts, one made from olive mill waste and orchard pruning residues, and one from olive mill waste, animal manure and wool residues, as well as of a biochar and of mixtures of the biochar and the composts. <sup>13</sup>C CP-MAS NMR spectra were recorded for the two composts and the biochar, and the areas of the characteristic spectral regions showed that biochar had the lowest content of O-alkyl, methoxyl and alkyl C and the highest content of aromatic species. Overall, the authors concluded that the effects of these organic amendments on crop yield are largely variable and influenced by the interaction with soil chemistry.

<sup>13</sup>C SSNMR has been applied in a few cases to study compost from MSW <sup>[27][61][62]</sup>, which generally has a large organic fraction (40%), but, due to its origin, can also show peculiar spectral features. In particular, Pichler et al. <sup>[27]</sup> applied a complete set of quantitative and selective <sup>13</sup>C SSNMR experiments to MSW after various composting times and, besides observing the expected decomposition of carbohydrates, the presence of plastics (especially polystyrene, but also polyethylene and polypropylene) was recognized from characteristic signals in both the aliphatic and aromatic regions. Both aromatic and aliphatic signals increased with composting time because of an accumulation of these non-biodegradable materials.

The effect of composting on plastics was also investigated by Spaccini et al. <sup>[51]</sup> by means of different <sup>13</sup>C SSNMR measurements. In this case the decomposition of a bio-degradable starch-based thermoplastic mulching film for horticultural crops was followed in a real on-farm composting process. By recording variable contact time <sup>13</sup>C CP-MAS NMR spectra of the film before and after composting, the authors detected changes in the <sup>1</sup>H longitudinal relaxation time in the rotating frame,  $T_{1p}$ , values that were ascribed to an alteration of the inter-molecular linkages among the organic components of the plastic, due to the progressive decomposition of its constituents.

The effect of specific additives on different composts was also investigated. Skene et al. <sup>[67]</sup> studied the decomposition of straw in the presence and absence of inorganic matrices, such as sand, kaolin, and loamy sand, and found that straw incubated in inorganic matrices decomposes faster than straw alone and that sand seems to preserve alkyl groups. Considering that biochar in ruminant diets is being assessed as a method for simultaneously improving animal production and reducing enteric CH<sub>4</sub> emissions, Romero et al. <sup>[40]</sup> examined the effects of feeding a pine-based biochar to beef cattle on manure composting. <sup>13</sup>C CP-MAS NMR showed that biochar increases compost aromaticity, without altering the bulk C speciation of manure. Liu et al. <sup>[43]</sup> investigated the effect of adding biochar to a swine manure compost, and found a promoting influence of biochar on the aromatization process.

<sup>13</sup>C SSNMR has also been applied to freeze-dried extracts of compost to aid the understanding of the chemical structure of this complex material, as well as the changes occurring in OM during composting. Results on the contribution of the different functional groups to the <sup>13</sup>C CP-MAS NMR spectra of different types of extracts obtained from various composts and feedstocks are reported in **Table 3**.

**Table 3.** Relative areas (% of total area) of the different regions of <sup>13</sup>C CP-MAS NMR spectra for extracts of composts and corresponding feedstocks reported in the literature.

Ref. a	Feedstock <sup>b</sup>	Sample <sup>c</sup>	Alkyl C	O-Alky	/I C	Aroma	atic C	Carboxyl/ Carbonyl C	2
			c	H₃O/CHN	O-/Di- O-Alkyl C	Aryl C	O- ( Aryl C	CarboxylCarb C C	onyl ;
	SM/poplar sawdust (5:3 <i>w</i> : <i>w</i> )	HA 60 d	30.3	33.9	)	16.8	7.1	11.9	
	+sepiolite 3 wt%	HA 60 d	26.7	29.9	)	18.3	8.4	16.8	
[ <u>72</u> ]	+sepiolite 6 wt%	HA 60 d	34.7	22.6	6	17.4	7.9	17.4	
	+sepiolite 9 wt%	HA 60 d	24.3	29.4	Ļ	18.3	10.4	17.6	
	+sepiolite 12 wt%	HA 60 d	24.0	29.9	)	20.8	8.7	16.5	
	CYN/corn straw and WC (70:30 <i>w:w</i> )	HS 100 d	20.93	15.98	29.78	16.26	5.51	11.54	
[ <u>69</u> ]	COF/corn straw and WC (70:30 <i>w:w</i> )	HS 100 d	30.63	13.07	24.69	15.33	3.42	12.85	
	PEP/corn straw and WC (70:30 <i>w:w</i> )	HS 100 d	20.06	14.22	35.75	14.40	4.49	11.08	
[ <u>87</u> ]	CYN/WC (70:30 w:w)	HA 100 d	22.2	13.8	27.5	20.0	5.7	10.9	
	SM/rice straw 4:1	HA FS	46.8	19.6	6	18	.1	15.5	
		HA 40d	28.4	21.7	7	31	.3	18.7	
	SM/rice straw 8:1	HA FS	45.8	17.2	2	14	.7	22.3	
[ <u>43</u> ]		HA 40d	28.9	26.8	3	31	.6	12.7	
	SM/rice straw/biochar 8:1:1	HA FS	43.2	19.9	)	18	.4	18.5	
		HA 40d	27.9	23.4	ļ	30	.5	18.3	
[ <u>71</u> ]	CYN/corn straw (70:30 <i>w:w</i> )	HS 100 d	16.3	13.8	24.7	28.9	5.6	10.6	
[ <u>88</u> ]	Coffee husks/lettuce residues at (60:40 <i>w:w</i> )	CT 100 d	26.9	11.9	26.4	16.6	4.3	14	

Ref.	Feedstock <sup>b</sup>	Sample <sup>c</sup>	Alkyl C	O-Alky	/I C	Aroma	atic C	Carboxyl/ Carbonyl C
				CH <sub>3</sub> O/CHN	O-/Di- O-Alkyl C	Aryl C	O- Aryl C	CarboxylCarbonyl C C C
	CYN with maize straw/WC (70:30 <i>w:w</i> )	CT 100 d	27.6	12.1	27.3	15.0	4.2	13.9
	PEP with maize straw/WC (70:30 <i>w:w</i> )	CT 100 d	17.4	14.3	31.1	19.2	5.8	12.3
	Agricultural crop plants/NH4NO3 (66:34 <i>w</i> :w)	HA 90 d	41.63	24.89	9	19.3	31	14.16
[ <u>68</u> ]	Date palm fronds/NH4NO3 (66:34 <i>w</i> :w)	HA 90 d	36.39	29.05	ō	23.	85	10.70
	Animal waste/NH4NO3 (66:34 w:w)	HA 90 d	31.39	29.0	)	25.4	45	14.33
	Tomato R/escarole R/WC/CS (17.5:20.5:60:2)	CT 105 d	23.50	15.60	28.65	16.80	5.25	10.20
[ <u>89]</u>	Tomato R/escarole R/WC/CS (37:11:50:2)	CT 105 d	22.55	14.95	26.65	18.25	5.90	11.75
	Tomato R/escarole R/WC/CS (50:0:48:2)	CT 105 d	23.40	14.70	26.80	17.10	6.90	11.15
	Commercial compost from biowaste	CT 105 d	26.95	12.85	32.70	10.35	2.90	14.25
[ <u>83</u> ]	DOW/COF/pine needles and WT (1:1:1)	DOM FS	26.0	6.3	42.1	9.5	4.9	11.1
		DOM 90 d	30.8	7.4	27.1	13.4	7.3	14.0
	DOW/GT/FR (2:1:1)	DOM FS	31.8	6.2	38.5	7.9	4.3	11.4
		DOM 90 d	34.6	7.3	25.9	13.1	4.9	14.2
	GT/COF/spent yeast	DOM	30.8	5.3	41.2	6.2	2.8	13.6

Ref.	Feedstock <sup>b</sup>	Sample <sup>c</sup>	Alkyl C	O-Alky	I C	Aroma	atic C	Carboxyl/ Carbonyl C
			(	CH₃O/CHN(	O-/Di- D-Alkyl C	Aryl C	O- Aryl C	CarboxylCarbonyl C C
	(1:1:1)	FS						
		DOM 90 d	35.2	8.4	22.7	11.8	5.8	16.2
	GT/COF/FR/sewage sludge (4:2:2.5:0.25)	DOM FS	30.8	6.2	38.2	8.6	4.9	11.3
		DO 90d	33.9	7.8	25.0	13.2	5.4	14.7
		HA 60 d	28.0	11.3	32.4	16.	8	11.5
[ <u>78</u> ]	DOW/GT/vegetal R from tobacco	HA 90 d	34.9	10.8	28.6	15.	7	10.0
	(50:30:20)	HA 150 d	34.5	9.3	23.1	19.	8	13.4
		CT 120 d	31.0	9.0	23.1	13.	3	23.6
[ <u>85</u> ]	OFMSW/GT/foliage R from tobacco (55:30:15)	HoDOM 120 d	34.6	12.6	19.5	19.	3	14.0
		HiDOM 120 d	30.3	9.8	36.8	9.2	1	14.0
[ <u>70</u> ]	OvM/straw	HS FS	18.6	49.7		8.2	2	23.5
		HS 120 d	17.7	25.0		22.	3	35.0
	Mixture of animal manures	HS FS	33.7	13.8		11.	8	40.7
		HS 120 d	30.4	24.0		9.6	6	36.0
	Solid olive mill wastes	HS FS	23.2	56.4		11.	2	9.2
		HS 120 d	22.2	29.3		19.	7	28.8
	Solid wastes of wineries	HS FS	3.5	46.9		1.5	ō	48.1
		HS 120 d	21.6	13.1		26.	8	38.5

Ref.	Feedstock <sup>b</sup>	Sample <sup>c</sup> A	alkyl C	O-Alkyl C	Aroma	tic C	Carbo Carbo	oxyl/ nyl C	
				O-/Di- CH <sub>3</sub> O/CHNO-Alkyl C	Aryl C	O-( Aryl C	Carboxyl C	Carbonyl C	
	Domestic wastes	HS FS	23.7	28.2	20.	3	27	.8	
		HS 120 d	25.4	34.9	16.	4	23	.3	
		HA FS	43.4	25.9	10.	3	16.4	4.0	
[ <u>79</u> ]	MSW/vegetal wastes	HA 28 d	44.7	22.1	11.	0	17.9	4.3	
	(1.1 0.0)	HA 100 d	42.9	20.3	11.	4	17.9	7.5	
[ <u>61</u> ]	MSW	HA FS	32	45	13	3	10	C	
	1013 00	HA 49 d	48	26	16	ò	10	C	
[ <u>82</u> ]	MSW	HA 6 d	38	31	13	5	11	2	e
	<u>36]</u>	<sup>3]</sup> HA 19 d <sup>13</sup>	45	23	15	5	11	1	ſ(
		HA 33 d	44	23	16	6	11	1	n D
		HA 62 d	42	24	15	6	12	2	t
		HA 105 d	39	25	16	6	12	2	e
		HA 187 d	38	26	17	6	12	3	
	13	Core- HA 6 d	34	24	21	8	12	2	gi gi
		Core- HA 19 d	33	23	23	9	11	2	e
		Core- HA 33 d	34	23	23	9	11 13	1	IP
<u>9]</u>		Core- HA 62 d	38	23 [ <mark>71</mark> ]	19	7	11	3	D
		Core- HA 105 d	43	21	19	7	10	0	21
								J	14

anure and sawdust with the addition of different amounts of sepiolite, a clay mineral, which is indeed expected to affect the composting process due to its high absorption capacity and cation exchange ability, possibly favoring the humification process. The <sup>13</sup>C CP-MAS NMR spectra of HAs extracted from compost obtained from samples with increasing amounts of sepiolite confirmed the increasing extent of humification, as

Ref. a	Feedstock <sup>b</sup>	Sample <sup>c</sup> Alkyl C		O-Alkyl C	Aromatic C		Carboxyl/ Carbonyl C		alkyl and
			C	O-/Di- CH <sub>3</sub> O/CHNO-Alky C	l Aryl C	O- ( Aryl C	Carboxyl C	Carbony C	a that the amended
		Core- HA 187 d	35	22	20	8	13	3	····
[ <u>81</u> ]	OFMSW	HA FS	37.20	34.34 <sub>[73]</sub>	16.	28	12.23		e positive on <sup>[74][75]</sup> .
		HA C	30.07	34.71	22.	67	12.	58	s, as well

as on compost, as previously mentioned. Three mixtures of manure, straw as bulking agent, and biochar in

different proportions, i.e., 4:1:0, 8:1:0 and 8:1:1 by wet weight, were investigated. In **Table 3** the data relative to the "References are reported in reverse chronological order." COF = coffee grounds; CS = compost starters; CYN = different NMR spectral regions for HAs obtained for the three different samples before and after composting are artichoke residues; DOW = domestic organic waste; FR = fresh green vegetable residues; GT = garden trimmings; reported. The authors observed a positive effect of biochar on HA formation on the basis of the higher overall MSW = municipal solid waste; OFMSW = organic fraction of MSW; OvM = ovine manure; PEP = pepper waste; R = spectral intensity. In all cases, the alkyl C contribution sensibly decreased after 40 days of composting, while the residues; SM= swine manure; WC = wood chips; WT = wood trimmings. CT = compost tea; d = days; DOM = aromatic and O-alkyl C contents increased. This was taken as an indication of the decomposition of dissolved organic matter; FS = feedstock; HA = humic acid; HiDOM = hydrophilic fraction of DOM; HoDOM = macromolecules, such as lignin, hemicellulose and cellulose, contributing to the formation of HAs <sup>TG</sup>. The amount hydrophobic fraction of DOM; HS = humic substance. of carboxyl C in the HA sample with the biochar added was higher with respect to the other samples before composting and also remained high after composting, with a value similar to that of the sample with a higher amount of bulking agent. A high proportion of both aromatic and carboxyl groups in compost HAs, as observed in the presence of biochar and with a higher amount of bulking agent, has been suggested to have positive agronomic effects <sup>[77]</sup>.

<sup>13</sup>C SSNMR was more extensively applied to monitor the changes occurring in HAs during composting, although, again, the changes observed were strongly dependent on the feedstock. In fact, although it is generally stated that, during composting, carbohydrates are decomposed and new aromatic groups are formed, no fixed trends were reported in the case of compost HAs.

Data in agreement with the expected trend were reported by Spaccini and Piccolo <sup>[78]</sup> on a mixture of domestic waste, plant trimmings and tobacco residues. After 2 months of composting, the HA spectrum was dominated by alkyl and O-alkyl signals, the former probably deriving from lipids, waxes, plant polyesters, and peptidic compounds, the latter from sugars, but also from ether and epoxy groups of plant polyesters. The aromatic signals were mainly associated with H- or C-substituted carbons, with a low content of O-aryl carbons. Dipolar dephasing NMR experiments showed a low contribution from lignin. On the other hand, there was a large contribution from carboxyl groups in aliphatic acids and/or amino acid amide groups. After 5 months of composting, the extracted HAs showed a significantly lower carbohydrate content and a higher proportion of alkyl C due to the incorporation of stable lipids. Observable, but minor increases were found for the aromatic and carboxyl C signal intensities. The results suggested a preferential preservation in the HA fraction of hydrophobic alkyl components, as well as the inclusion of labile peptidic compounds.

Castaldi et al. <sup>[79]</sup> studied the transformation of HAs during the composting of MSW with the addition of agricultural wastes. The <sup>13</sup>C CP-MAS NMR spectra of HAs, extracted at the beginning and after 100 days of composting, showed that the composting process caused a reduction in carbohydrates and aliphatic structures, and an increase in the aromatic, carboxyl and carbonyl carbons, which suggested a partial neo-formation of humified material.

Fuentes et al. <sup>[70]</sup> followed the composting process by analyzing HSs in five composted materials of diverse origin, that is, solid wastes of wineries, solid olive mill wastes, domestic wastes, ovine manure with the addition of straw, and a mixture of animal manures. Except for domestic wastes and the mixture of animal manures, humification was somewhat as expected, that is, there was an increase in the number of aromatic carbons, with a large contribution of phenolic groups, and a reduction in O-alkyl groups, while the changes in carboxyl/carbonyl C contents were at variance. On the contrary, in the case of domestic wastes and mixed manures, the distribution of functional groups did not undergo significant changes, but there was an unexpected increase in O-alkyl C and a slight decrease in aromatic and carbonyl C. The authors ascribed this unusual behavior to the presence of non-aromatic recalcitrant material and of tannins, these exerting a limiting effect on the decomposition of aliphatic biomolecules, as observed by Almendros et al. <sup>[80]</sup>.

Contrary to the expected trend, the study by Liu et al. <sup>[43]</sup> on the composting of swine manure with straw and biochar, discussed above, showed in all HA samples a significant decrease in the contribution of alkyl C, and an increase in the contribution of all the other carbon types (except for the carboxyl signal in the pig manure/straw with an 8:1 ratio), the major increase involving the aromatic region. As already stated, this trend was explained by assuming the formation of new aromatic structures contributing to HAs resulting from the decomposition of other macromolecular components.

Gonzáles-Vila et al. <sup>[61]</sup> studied HAs extracted from MSW and from its compost obtained after 7 weeks. The <sup>13</sup>C CP-MAS NMR spectra of the HAs isolated before and after the composting process showed a significant reduction in carbohydrate content with a large increase in alkyl carbon upon composting, which indicated a concentration of recalcitrant polymethylenic structures (i.e., waxes and polyesters), while no major increase in aromatic structures or carboxyl groups was observed. The authors pointed out that, given the type of feedstock, most of the signals in the aromatic region were probably due to unsaturated fatty acids, physically occluded and/or esterified with other aliphatic macromolecules. At variance with those results are those obtained by García et al. <sup>[81]</sup>, who investigated HA extracts from the organic fraction of MSW and from its compost. In this case, the only changes observed were a reduction in alkyl C and an increase in aromatic C content upon composting, while the intensity in the carbohydrate region and in the carboxyl/carbonyl C region did not change.

With the aim of better understanding the transformations of HSs during the composting of MSW, Chefetz et al. <sup>[82]</sup> extracted two types of HAs, i.e., HAs and core-HAs, the latter obtained by organic solvent and sulfuric acid extractions prior to the alkaline extraction. Core-HAs represent a recalcitrant fraction, whereas HAs also contain labile components, some of which coat core-HA structures. <sup>13</sup>C CP-MAS NMR spectra indicated that, while core-HAs, being more stable, practically did not change during the process, HAs showed a decrease in carbohydrates and an increase in aromatic structures. Essentially, the "coating" materials (polysaccharides, peptides, and lipids)

bonded to the core-HA structure degraded during composting, leaving HA structures of higher aromaticity. In fact, as composting proceeded, HAs tended to resemble core-HAs. Besides relative intensities, the authors also determined absolute intensities, taking into account the carbon content reduction in the process. Considering the absolute data, the authors noted that, besides degradation of coating compounds, new HAs were formed. Thus, they concluded that during composting carbohydrates were lost by degradation while the residual HAs were enriched with aromatic structures.

Changes of OM in wastes were also followed by observing the changes in DOM. Caricasole et al. <sup>[83]</sup> compared the spectra of DOM extracted from different mixtures of domestic and garden wastes and their composts. In all cases, composting gave rise to a decrease in O-alkyl C, counterbalanced by an increase in all other functional groups. The increase in alkyl C was attributed to the breakdown of polysaccharides yielding simple alkyl chains and/or to the accumulation of highly aliphatic biopolymers occurring in cuticles of higher plants and/or some cell wall products highly resistant to decomposition <sup>[18]</sup>. The relative enrichment in aromatic C was supposed to be related to the degradation of non-aromatic cell wall compounds <sup>[84]</sup>, while carboxyl and carbonyl groups probably arose from oxidative transformations of lignin<sup>[31]</sup>. The <sup>13</sup>C CP-MAS NMR spectra obtained on DOM extracted from fresh substrates and from the final composts indicated that, as observed for HSs, DOM composition depends on the initial feedstock. The highest increase in alkyl degradable pine needles. The highest carbohydrates reduction associated with the highest increase in the *A/OA* ratio and to the lowest carboxyl C increase was observed for the sample originating from the mixture containing spent yeast from beer production, which is a highly degradable material.

Comparison of DOM spectra among different samples and composts can indeed give important information on the composting process. However, the determination of DOM composition is difficult since it is a mixture of lowmolecular weight compounds (free amino acids and sugars) and chemically heterogeneous macromolecules (enzymes, amino-sugar complexes, polyphenols, and HSs). With the aim of better understanding DOM composition, <sup>13</sup>C SSNMR was used on separated hydrophilic (HiDOM) and hydrophobic (HoDOM) fractions [85] and on fractions further separated on the basis of solubility in acid or basic solutions [38]. The two studies highlighted the composition of the two main fractions confirming the hydrophilic nature of HiDOM, richer in oxygenated functional groups, and the hydrophobic nature of HoDOM, richer in alkyl and aromatic carbons. Chefetz et al. [38] followed the fate of three fractions of HoDOM (Ho-acid, Ho-base and Ho-neutral) and three fractions of HiDOM (Hi-acid, Hi-base and Hi-neutral) through the composting process. The Ho-acid fraction, characterized by a fulvic acid composition, was stable throughout the process and became the dominant DOM component in the final compost. The amount of the Ho-neutral fraction, constituted of highly apolar compounds, was considered to represent an intermediate stage in the transformation of DOM compounds to HSs. On the other hand, the Hi-neutral fraction, mainly containing poly- and oligosaccharides, significantly decreased during composting, and was of relevance in OM degradation. On the basis of their findings, the authors suggested possible reasons for the beneficial effects of compost on plant growth: (i) higher amounts of Ho-neutral and Hoacid in the compost imply an increase in the concentration of functional groups which are able to chelate metals,

thus favoring their uptake by plants; (ii) the reduced concentration of Hi-neutral results in the removal of compounds toxic to plants.

Several studies focused on the relationship between structure and bioactivity of compost HSs. Verrillo et al. <sup>[71]</sup> studied the effects of HSs extracted from a compost obtained from artichoke biomass. The use of <sup>13</sup>C CP-MAS NMR, combined with pyrolysis gas chromatography mass spectrometry, indicated that the biostimulation activity was related to the presence of hydrophobic domains containing numerous bioactive molecules, such as ligninderived molecules and saccharidic and aminoacidic components. Of relevance for understanding the structure– activity relationship were the *ARM*, *A/OA*, *HB/HI*, and *LigR* indices. The relatively high *HB/HI* index (=0.90) observed for HSs from artichoke residues and the relatively high aromatic content revealed a significant incorporation of hydrophobic molecules in the humic supramolecular structure, mainly represented by aromatic compounds. The *A/OA* ratio of 0.65 indicated the predominance of bioavailable polar O-alkyl groups with respect to apolar alkyl components, and the high *LigR* value (=2.5) indicated that most of the intensity in the 45–60 ppm region was due to lignin units and, hence, confirmed their incorporation in HAs.

Similar results were found for HSs extracted from composted artichoke and pepper wastes, and from coffee grounds <sup>[69]</sup>. These extracts were also investigated for their antibacterial activity and antioxidant properties, the former being related to the aromatic and phenolic content and the latter only to the phenolic content <sup>[86]</sup>. The antiinflammatory properties of HAs were also shown to be related to the content of aromatic, phenolic and lignin components in a study by Verrillo et al. <sup>[87]</sup>.

The molecular features responsible for bioactivity were also investigated on compost tea. Verrillo et al.<sup>[88]</sup> studied CTs from different composts of recycled agricultural biomasses (artichoke and pepper with maize straw, coffee husks with lettuce residues). Germination tests performed using basil seeds showed that CT from coffee husks and pepper caused an increase in both root and epicotyl length, while artichoke CT showed positive effects only on root development. As in the case of HSs, this biostimulation effect was ascribed by the authors to a higher amount of phenolic and aromatic units, as observed by <sup>13</sup>C CP-MAS NMR.

Pane et al. <sup>[89]</sup> studied three CTs, produced from agricultural wastes (tomato and escarole residues) with the addition of different proportions of woodchips, and a CT from a commercial biowaste compost, investigating their effects in a processing tomato cropping system. The <sup>13</sup>C CP-MAS NMR spectra of CTs obtained from recycled agricultural biomasses were mainly characterized by prominent signals attributable to the incorporation of lignin molecules and polyphenol derivatives, while that of commercial compost showed a composition dominated by carbohydrates. Notwithstanding the chemical differences, the general bioactivity observed for all CT treatments, independently from the sources, was attributed to a correct balance of molecular components giving rise to an equilibrium between the hydrophobic and the hydrophilic domains <sup>[90]</sup>.

#### 4. Conclusions and Perspectives

<sup>13</sup>C SSNMR, widely used in the field of materials science, has revealed as a powerful tool also for the investigation of organic natural matter of relevance in agriculture and environmental science. In fact, <sup>13</sup>C SSNMR gives access to information on the composition of complex organic materials that cannot be obtained otherwise and which can be of great relevance for the optimization of waste recycling processes and the exploitation of waste materials that presently are directly disposed of in landfills.

<sup>13</sup>C SSNMR has been applied to investigate the composting of materials of varied nature, from relatively homogeneous garden wastes to widely heterogenous municipal solid wastes. Most studies have shown that the compositional features of compost depend on the starting feedstock and this is important for the application of these materials, which are often intended to be used as fertilizers or amendments in agricultural soils. Nevertheless, <sup>13</sup>C SSNMR has highlighted the degradation of labile OM components (mainly carbohydrates) and the preservation of the recalcitrant ones (lipids, waxes, lignin). The relative amounts of the different components obtained from NMR data have been correlated with stability and maturity indices determined by analytical methods. In some cases, the results indicate that waste pretreatments may be required, as for example in the case of municipal solid waste where non-biodegradable plastic materials remain in the compost.

Although not conclusive alone, <sup>13</sup>C SSNMR has contributed to the understanding of structure-bioactivity relationship of composts, highlighting the role of hydrophobic and polar chemical functionalities in determining the biostimulation properties of compost of interest in agriculture and the antibacterial, antiflammatory and antioxidant properties of extracts of relevance for their application in pharmacology.

Most of the studies cited relied on the spectral analysis based on the division of the <sup>13</sup>C MAS NMR spectra in regions related to specific functional groups. This is certainly useful for understanding organic matter evolution during composting. However, the results of the different studies indicate that not always the standard decomposition pathway is followed and no univocal indices of humification and maturity, as those typically used in soil science, can be defined. This is probably due to the variable and complex composition of waste materials, especially in the case of municipal solid waste. Nonetheless, comparison of <sup>13</sup>C MAS NMR spectra before and after the composting process yields detailed information on the transformations that have occurred, which is important for process optimization. To this end, molecular models can be of help, especially for the characterization and quality assessment of the final products, and would certainly deserve more consideration and possibly further refinement. Moreover, given the complex nature of OM in waste transformation, a combined <sup>13</sup>C SSNMR investigation of the whole sample and different extracts is advisable for a better comprehension of the degradation processes and the chemical structure of the final products.

As evidenced by the growing literature in the field and by the results therein reported, <sup>13</sup>C SSNMR is an important instrument for the investigation of organic waste transformation within a circular economy model. It can be easily foreseen that research in this field will experience a significant growth given the increased public awareness of the importance of waste recycling, and that <sup>13</sup>C SSNMR will become even more important in the future in this field thanks to the rapid hardware and software advancements that NMR technology is experiencing, which will extend the information that can be obtained on complex materials as those considered.

#### References

- Bandini, F.; Taskin, E.; Bellotti, G.; Vaccari, F.; Misci, C.; Guerrieri, M.C.; Cocconcelli, P.S.; Puglisi, E. The treatment of the organic fraction of municipal solid waste (OFMSW) as a possible source of micro- and nano-plastics and bioplastics in agroecosystems: A review. Chem. Biol. Technol. Agric. 2022, 9, 4.
- 2. Available online: https://datatopics.worldbank.org/what-awaste/trends\_in\_solid\_waste\_management.html (accessed on 20 January 2023).
- 3. Tejaswini, M.S.S.R.; Pathak, P.; Gupta, D.K. Sustainable approach for valorization of solid wastes as a secondary resource through urban mining. J. Environ. Manag. 2022, 319, 115727.
- Coma, M.; Martinez-Hernandez, E.; Abeln, F.; Raikova, S.; Donnelly, J.; Arnot, T.C.; Allen, M.J.; Hong, D.D.; Chuck, C.J. Organic waste as a sustainable feedstock for platform chemicals. Faraday Discuss. 2017, 202, 175–195.
- Lohri, C.R.; Diener, S.; Zabaleta, I.; Mertenat, A.; Zurbrügg, C. Treatment technologies for urban solid biowaste to create value products: A review with focus on low- and middle-income settings. Rev. Environ. Sci. Biotechnol. 2017, 16, 81–130.
- Shah, V.; Singh, A.; Mohanty, S.S.; Srivastava, V.K.; Varjani, S. Organic solid waste: Biorefinery approach as a sustainable strategy in circular bioeconomy. Bioresour. Technol. 2022, 349, 126835.
- De Corato, U.; De Bari, I.; Viola, E.; Pugliese, M. Assessing the main opportunities of integrated biorefining from agro-bioenergy co/by-products and agroindustrial residues into high-value added products associated to some emerging markets: A review. Renew. Sustain. Energy Rev. 2018, 88, 326–346.
- 8. Kumar Khanal, S.; Lü, F.; Wong, J.W.C.; Wu, D.; Oechsner, H. Anaerobic digestion beyond biogas. Bioresour. Technol. 2021, 337, 125378.
- Albrecht, R.; Ziarelli, F.; Alarcón-Gutiérrez, E.; Le Petit, J.; Terrom, G.; Perissol, C. 13C solid-state NMR assessment of decomposition pattern during co-composting of sewage sludge and green wastes. Eur. J. Soil Sci. 2008, 59, 445–452.
- Papale, M.; Romano, I.; Finore, I.; Lo Giudice, A.; Piccolo, A.; Cangemi, S.; Di Meo, V.; Nicolaus, B.; Poli, A. Prokaryotic diversity of the composting thermophilic phase: The case of ground coffee compost. Microorganisms 2021, 9, 218.
- 11. Jara-Samaniego, J.; Perez-Murcia, M.D.; Bustamante, M.A.; Perez-Espinosa, A.; Paredes, C.; Lopez, M.; López-Lluch, D.B.; Gavilanes-Teráne, I.; Moral, R. Composting as sustainable strategy

for municipal solid waste management in the Chimborazo Region, Ecuador: Suitability of the obtained composts for seedling production. J. Clean. Prod. 2017, 141, 1349–1358.

- Cesaro, A.; Conte, A.; Belgiorno, V.; Siciliano, A.; Guida, M. The evolution of compost stability and maturity during the full-scale treatment of the organic fraction of municipal solid waste. J. Environ. Manag. 2019, 232, 264–270.
- Palaniveloo, K.; Amran, M.A.; Norhashim, N.A.; Mohamad-Fauzi, N.; Peng-Hui, F.; Hui-Wen, L.; Kai-Lin, Y.; Jiale, L.; Chian-Yee, M.G.; Jing-Yi, L.; et al. Food waste composting and microbial community structure profiling. Processes 2020, 8, 723.
- 14. Available online: https://www.compostnetwork.info/wordpress/wp-content/uploads/ECN-rapport-2022.pdf (accessed on 20 January 2023).
- 15. Wu, L.; Ma, L.Q.; Martinez, G.A. Comparison of methods for evaluating stability and maturity of biosolids compost. J. Environ. Qual. 2000, 29, 424–429.
- 16. Ouatmane, A.; Provenzano, M.R.; Hafidi, M.; Senesi, N. Compost maturity assessment using calorimetry, spectroscopy and chemical analysis. Compost Sci. Util. 2000, 8, 124–134.
- 17. Butler, T.A.; Sikora, L.J.; Steinhilber, P.M.; Douglass, L.W. Compost age and sample storage effects on maturity indicators of biosolids compost. J. Environ. Qual. 2001, 30, 2141–2148.
- 18. Chen, Y. Nuclear magnetic resonance, infra-red and pyrolysis: Application of spectroscopic methodologies to maturity determination of composts. Compost Sci. Util. 2003, 11, 152–168.
- 19. Francou, C.; Poitrenaud, M.; Houot, S. Stabilization of organic matter during composting: Influence of process and feedstocks. Compost Sci. Util. 2005, 13, 72–83.
- 20. Wichuk, K.M.; McCartney, D. Compost stability and maturity evaluation—A literature review. Can. J. Civ. Eng. 2010, 37, 1505–1523.
- Oviedo-Ocaña, E.R.; Torres-Lozada, P.; Marmolejo-Rebellon, L.F.; Hoyos, L.V.; Gonzales, S.; Barrena, R.; Komilis, D.; Sanchez, A. Stability and maturity of biowaste composts derived by small municipalities: Correlation among physical, chemical and biological indices. Waste Manag. 2015, 44, 63–71.
- Simpson, A.J.; Simpson, M.J. Nuclear magnetic resonance analysis of natural organic matter. In Biophysico-Chemical Processes Involving Natural Nonliving Organic Matter in Environmental Systems; Senesi, N., Xing, B., Huang, P.M., Eds.; John Wiley & Sons: Hoboken, NJ, USA, 2009; pp. 589–651.
- Simpson, A.J.; McNally, D.J.; Simpson, M.J. NMR spectroscopy in environmental research: From molecular interactions to global processes. Prog. Nucl. Magn. Reson. Spectrosc. 2011, 58, 97– 175.

- Baldock, J.A.; Oades, J.M.; Nelson, P.N.; Skene, T.M.; Golchin, A.; Clarke, P. Assessing the extent of decomposition of natural organic materials using solid-state 13C NMR spectroscopy. Aust. J. Soil Res. 1997, 35, 1061–1083.
- Meija, J.; Coplen, T. B.; Berglund, M.; Brand, W. A.; De Bièvre, P.; Gröning, M.; Holden, N. E.; Irrgeher, J.; Loss, R. D.; Walczyk, T.; Prohaska, T. Isotopic compositions of the elements 2013 (IUPAC technical report). Pure Appl. Chem. 2016, 88, 293-306.
- Holden, N. E.; Coplen, T. B.; Böhlke, J. K.; tarbox, L. V.; Benefield, J.; de Laeter, J. R.; Mahaffy, P. G.; O'Connorr, G.; Roth, E.; Tepper, D. H.; Walczyk, T.; Wieser, M. E.; Yoneda, S. IUPAC periodic table of the elements and isotopes (IPTEI) for the educa-tion community (IUPAC technical report). Pure Appl. Chem. 2018, 90, 1833-2092.
- 27. Pichler, M.; Knicker, H.; Kögel-Knabner, I. Changes in the chemical structure of municipal solid waste during composting as studied by solid-state dipolar dephasing and PSRE 13C NMR and solid-state 15N NMR spectroscopy. Environ. Sci. Technol. 2000, 34, 4034–4038.
- 28. Réveillé, V.; Mansuy, L.; Jardé, E.; Garnier-Sillam, E. Characterization of sewage–sludge derived organic matter: lipids and humic acids. Org. Geochem. 2003, 34, 615–627.
- 29. Pereira, M. A.; Pires, O. C.; Mota, M.; Alves, M. M. Anaerobic biodegradation of oleic and palmitic acids: evidence of mass transfer limitations caused by long chain fatty acid accumulation onto the anaerobic sludge. Biotechnol. Bioeng. 2005, 92, 15-23.
- Dignac, M. F.; Derenne, S.; Ginestet, P.; Bruchet, A.; Kniker, H.; Largeau, C. Determination of structure and origin of refractory organic matter in biodepurated wastewater via spectroscopic methods. Comparison of conventional and ozonation treat-ment. Environ. Sci. Technol. 2000, 34, 3389–3394.
- 31. Kögel-Knabner, I. The macromolecular organic composition of plant and microbial residues as inputs to soil organic matter. Soil Biol. Biochem. 2002, 34, 139-162.
- 32. Ussiri, A. A. N.; Johnson, C. E. Characterization of organic matter in a northern hardwood forest soil by 13C NMR spectros-copy and chemicals methods. Geoderma 2003, 11, 123–149.
- 33. Atalla, R. H.; VanderHart, D. L. The role of solid state 13C NMR spectroscopy in studies of the nature of native cellulose. Solid State Nucl. Magn. Reson. 1999, 15, 1–19.
- 34. Hatcher, P. G. Chemical structural studies of natural lignin by dipolar dephasing solid-state 13C nuclear magnetic resonance. Org. Geochem. 1987, 11, 31–39.
- 35. Manu, M. K.; Kumar, R.; Garg, A. Performance assessment of improved composting system for food waste with varying aer-ation and use of microbial inoculum. Bioresource Technol. 2017, 234, 167-177.

- Bustamante, M. A.; Paredes, C.; Marhuenda-Egea, F. C.; Perez-Espinosa, A.; Bernal, M. P.; Moral, R. Co-composting of distill-ery wastes with animal manures: carbon and nitrogen transformations in the evaluation of compost stability. Chemosphere 2008, 72, 551–557.
- Zhou, Y.; Selvam, A.; Wong, J. W. C. Evaluation of humic substances during co-composting of food waste, sawdust and Chi-nese medicinal herbal residues, Bioresource Technol. 2014, 168, 229-234.
- 38. Chefetz, B.; Hadar, Y.; Chen, Y. Dissolved organic carbon fractions formed during composting of municipal solid waste: properties and significance. Acta Hydrochim. Hydrobiol. 1998, 26, 172-179.
- 39. Eudoxie, G.; Martin, M. Compost tea quality and fertility, In Organic fertilizers: history, production and applications, Lar-ramendy, M., Soloneski, S., Eds. IntechOpen: London, UK, 2019.
- 40. Romero, C.M.; Redman, A.-A.P.H.; Owens, J.; Terry, S.A.; Ribeiro, G.O.; Gorzelak, M.A.; Oldenburg, T.B.P.; Hazendonk, P.; Larney, F.J.; Hao, X.; et al. Effects of feeding a pine-based biochar to beef cattle on subsequent manure nutrients, organic matter composition and greenhouse gas emissions. Sci. Total Environ 2022, 812, 152267.
- 41. Iacomino, G.; Sarker, T.C.; Ippolito, F.; Bonanomi, G.; Vinale, F.; Staropoli, A.; Idbella, M. Biochar and compost application either alone or in combination affects vegetable yield in a volcanic Mediterranean soil. Agronomy 2022, 12, 1996.
- 42. Eldridge, S.M.; Chen, C.R.; Xu, Z.H.; Nelson, P.N.; Boyd, S.E.; Meszaros, I.; Chan, K.Y. Molecular composition of recycled organic wastes, as determined by solid-state 13C NMR and elemental analyses. Waste Manag. 2013, 33, 2157–2169.
- 43. Liu, H.; Guo, H.; Guo, X.; Wu, S. Probing changes in humus chemical characteristics in response to biochar addition and varying bulking agents during composting: A holistic multi-evidence-based approach. J. Environ. Manag. 2021, 300, 113736.
- 44. Veeken, A.H.M.; Adani, F.; Nierop, K.G.J.; de Jager, P.A.; Hamelers, H.V.M. Degradation of biomacromolecules during high-rate composting of wheat straw–amended feces. J. Environ. Qual. 2001, 30, 1675–1684.
- Martín-Mata, J.; Lahoz-Ramos, C.; Bustamante, M.A.; Marhuenda-Egea, F.C.; Moral, R.; Santos, A.; Sáez, J.A.; Bernal, M.P. Thermal and spectroscopic analysis of organic matter degradation and humification during composting of pig slurry in different scenarios. Environ. Sci. Pollut. Res. 2016, 23, 17357–17369.
- 46. Tang, J.-C.; Maie, N.; Tada, Y.; Katayama, A. Characterization of the maturing process of cattle manure compost. Process Biochem. 2006, 41, 380–389.
- 47. Inbar, Y.; Chen, Y.; Hadar, Y. Solid-state carbon-13 nuclear magnetic resonance and infrared spectroscopy of composted organic matter. Soil Sci. Soc. Am. J. 1989, 53, 1695–1701.

- 48. Chen, Y.; Inbar, Y.; Hadar, Y.; Malcolm, R.L. Chemical properties and solid-state CPMAS 13C-NMR of composted organic matter. Sci. Total Environ. 1989, 81–82, 201–208.
- 49. Wang, K.; He, C.; You, S.; Liu, W.; Wang, W.; Zhang, R.; Qi, H.; Ren, N. Transformation of organic matters in animal wastes during composting. J. Hazard. Mater. 2015, 300, 745–753.
- Torres-Climent, A.; Gomis, P.; Martín-Mata, J.; Bustamante, M.A.; Marhuenda-Egea, F.C.; Pérez-Murcia, M.D.; Pérez-Espinosa, A.; Paredes, C.; Moral, R. Chemical, thermal and spectroscopic methods to assess biodegradation of winery-distillery wastes during composting. PLoS ONE 2015, 10, e0138925.
- 51. Spaccini, R.; Todisco, D.; Drosos, M.; Nebbioso, A.; Piccolo, A. Decomposition of bio-degradable plastic polymer in a real on-farm composting process. Chem. Biol. Technol. Agric. 2016, 3, 4.
- 52. Lin, Y.-H.; Lin, Y.-Z.; Lin, Y.-H. Preliminary design for establishing compost maturity by using the spectral characteristics of five organic fertilizers. Sci. Rep. 2022, 12, 15721.
- 53. Spaccini, R.; Piccolo, A. Spectroscopic characterization of compost at different maturity stages. Clean 2008, 36, 152–157.
- 54. Knicker, H.; Lüdemann, H.-D. N-15 and C-13 CPMAS and solution NMR studies of N-15 enriched plant material during 600 days of microbial degradation. Org. Geochem. 1995, 23, 329–341.
- Spaccini, R.; Piccolo, A. Molecular characterization of compost at increasing stages of maturity. 2. Thermochemolysis-GC-MS and 13C-CPMAS-NMR spectroscopy. J. Agric. Food Chem. 2007, 55, 2303–2311.
- 56. Anda, M.; Syed Omar, S.R.; Shamshuddin, J.; Fauziah, C.I. Changes in properties of composting rice husk and their effects on soil and cocoa growth. Commun. Soil Sci. Plant Anal. 2008, 39, 2221–2249.
- 57. Caricasole, P.; Provenzano, M.R.; Hatcher, P.G.; Senesi, N. Evolution of organic matter during composting of different organic wastes assessed by CPMAS 13C NMR spectroscopy. Waste Manag. 2011, 31, 411–415.
- 58. Pane, C.; Spaccini, R.; Caputo, M.; De Falco, E.; Zaccardelli, M. Multi-parameter characterization of disease-suppressive bio-composts from aromatic plant residues evaluated for garden cress (Lepidium sativum L.) cultivation. Horticulturae 2022, 8, 632.
- Inbar, Y.; Chen, Y.; Hadar, Y. Carbon-13 CPMAS NMR and FTIR spectroscopic analysis of organic matter transformations during composting of solid wastes from wineries. Soil Sci. 1991, 152, 272– 282.
- 60. Preston, C.M.; Cade-Menun, B.J.; Sayer, B.G. Characterization of Canadian backyard composts: Chemical and spectroscopic analyses. Compost Sci. Util. 1998, 6, 53–66.

- 61. González-Vila, F.J.; Almendros, G.; Madrid, F. Molecular alterations of organic fractions from urban waste in the course of composting and their further transformation in amended soil. Sci. Total Environ. 1999, 236, 215–229.
- 62. Chefetz, B.; Hatcher, P.G.; Hadar, Y.; Chen, Y. Chemical and biological characterization of organic matter during composting of municipal solid waste. J. Environ. Qual. 1996, 25, 776–785.
- 63. Nelson, P.N.; Baldock, J.A. Estimating the molecular composition of a diverse range of natural organic materials from solid-state 13C NMR and elemental analyses. Biogeochemistry 2005, 72, 1–34.
- 64. Zmora-Nahum, S.; Hadar, Y.; Chen, Y. Physico-chemical properties of commercial composts varying in their source materials and country of origin. Soil Biol. Biochem. 2007, 39, 1263–1276.
- 65. Haw, J.F.; Maciel, G.E.; Schroeder, H.A. Carbon-13 nuclear magnetic resonance spectrometric study of wood and wood pulping with cross polarization and magic-angle spinning. Anal. Chem. 1984, 56, 1323–1329.
- 66. Huang, J.; Yu, Z.; Gao, H.; Yan, X.; Chang, J.; Wang, C.; Hu, J.; Zhang, L. Chemical structures and characteristics of animal manures and composts during composting and assessment of maturity indices. PLoS ONE 2017, 12, e0178110.
- 67. Skene, T.M.; Skjemstad, J.O.; Oades, J.M.; Clarke, P.J. The influence of inorganic matrices on the decomposition of straw. Aust. J. Soil Res. 1996, 34, 413–426.
- 68. Al-Faiyz, Y.S.S. CPMAS 13C NMR characterization of humic acids from composted agricultural Saudi waste. Arab. J. Chem. 2017, 10, 5839–5853.
- 69. Verrillo, M.; Salzano, M.; Savy, D.; di Meo, V.; Valentini, M.; Cozzolino, V.; Piccolo, A. Antibacterial and antioxidant properties of humic substances from composted agricultural biomasses. Chem. Biol. Technol. Agric. 2022, 9, 28.
- 70. Fuentes, M.; Baigorri, R.; González-Gaitano, G.; García-Mina, J.M. The complementary use of 1H NMR, 13C NMR, FTIR and size exclusion chromatography to investigate the principal structural changes associated with composting of organic materials with diverse origin. Org. Geochem. 2007, 38, 2012–2023.
- Verrillo, M.; Cozzolino, V.; Spaccini, R.; Piccolo, A. Humic substances from green compost increase bioactivity and antibacterial properties of essential oils in basil leaves. Chem. Biol. Technol. Agric. 2021, 8, 28.
- Zheng, W.; Yang, Z.; Huang, L.; Chen, Y. Roles of organic matter transformation in the bioavailability of Cu and Zn during sepiolite-amended pig manure composting. J. Environ. Manag. 2022, 314, 115046.

- 73. Guo, X.; Liu, H.; Zhang, J. The role of biochar in organic waste composting and soil improvement: A review. Waste Manag. 2020, 102, 884–899.
- 74. Ngo, P.; Rumpel, C.; Ngo, Q.; Alexis, M.; Vargas, G.V.; Mora Gil, M.D.L.L.; Dang, D.; Jouquet, P. Biological and chemical reactivity and phosphorus forms of buffalo manure compost, vermicompost and their mixture with biochar. Bioresour. Technol. 2013, 148, 401–407.
- 75. Chen, Y.; Huang, X.; Han, Z.; Huang, X.; Hu, B.; Shi, D.; Wu, W. Effects of bamboo charcoal and bamboo vinegar on nitrogen conservation and heavy metals immobility during pig manure composting. Chemosphere 2010, 78, 1177–1181.
- 76. Jurado, M.M.; Suárez-Estrella, F.; López, M.J.; Vargas-García, M.C.; López-González, J.A.; Moreno, J. Enhanced turnover of organic matter fractions by microbial stimulation during lignocellulosic waste composting. Bioresour. Technol. 2015, 186, 15–24.
- 77. Guo, X.; Liu, H.; Wu, S. Humic substances developed during organic waste composting: Formation mechanisms, structural properties, and agronomic functions. Sci. Total Environ. 2019, 662, 501–510.
- 78. Spaccini, R.; Piccolo, A. Molecular characteristics of humic acids extracted from compost at increasing maturity stages. Soil Biol. Biochem. 2009, 41, 1164–1172.
- 79. Castaldi, P.; Alberti, G.; Merella, R.; Melis, P. Study of the organic matter evolution during municipal solid waste composting aimed at identifying suitable parameters for the evaluation of compost maturity. Waste Manag. 2005, 25, 209–213.
- Almendros, G.; Dorado, J.; González-Vila, F.J.; Blanco, M.J.; Lankes, U. 13C NMR assessment of decomposition patterns during composting of forest and shrub biomass. Soil Biol. Biochem. 2000, 32, 793–804.
- 81. García, C.; Hernández, T.; Costa, F. Comparison of humic acids derived from city refuse with more developed humic acids. J. Soil Sci. Plant Nutr. 1992, 38, 339–346.
- 82. Chefetz, B.; Adani, F.; Genevini, P.; Tambone, F.; Hadar, Y.; Chen, Y. Humic-acid transformation during composting of municipal solid waste. J. Environ. Qual. 1998, 27, 794–800.
- Caricasole, P.; Provenzano, M.R.; Hatcher, P.G.; Senesi, N. Chemical characteristics of dissolved organic matter during composting of different organic wastes assessed by 13C CPMAS NMR spectroscopy. Bioresour. Technol. 2010, 101, 8232–8236.
- Vinceslas-Akpa, M.; Loquet, M. Organic matter transformations in lignocellulosic waste products composted or vermicomposted (Eisenia Fetida Andrei): Chemical analysis and 13C CPMAS NMR spectroscopy. Soil Biol. Biochem. 1997, 29, 751–758.
- 85. Spaccini, R.; Baiano, S.; Gigliotti, G.; Piccolo, A. Molecular characterization of a compost and its water-soluble fractions. J. Agric. Food Chem. 2008, 56, 1017–1024.

- 86. Açıkgöz, M.A. Evaluation of phytochemical compositions and biological properties of Achillea gypsicola at different phenological stages. Chem. Biodivers. 2019, 16, e1900373.
- Verrillo, M.; Parisi, M.; Savy, D.; Caiazzo, G.; di Caprio, R.; Luciano, M.A.; Cacciapuoti, S.; Fabbrocini, G.; Piccolo, A. Antiflammatory activity and potential dermatological applications of characterized humic acids from a lignite and a green compost. Sci. Rep. 2022, 12, 2152.
- 88. Verrillo, M.; Salzano, M.; Cozzolino, V.; Spaccini, R.; Piccolo, A. Bioactivity and antimicrobial properties of chemically characterized compost teas from different green composts. Waste Manage. 2021, 120, 98–107.
- Pane, C.; Palese, A.M.; Spaccini, R.; Piccolo, A.; Celano, G.; Zaccardelli, M. Enhancing sustainability of a processing tomato cultivation system by using bioactive compost teas. Sci. Hortic. 2016, 202, 117–124.
- Canellas, L.P.; Piccolo, A.; Dobbss, L.B.; Spaccini, R.; Olivares, F.L.; Zandonadi, D.B.; Façanha, A.R. Chemical composition and bioactivity properties of size-fractions separated from a vermicompost humic acid. Chemosphere 2010, 78, 457–466.

Retrieved from https://www.encyclopedia.pub/entry/history/show/96095