

# Application of Barley Tweak Spike Mutants

Subjects: Biology

Contributor: Raimondas Šiukšta

Barley developmental mutants tweaky spike (*tw*) with disturbed auxin pathways possess a unique feature of an increased level of mouldy germinating grains (MGG), which serves as a convenient model to investigate the effects of plant immunity-related substances. The effects of the auxin 2,4-dichlorophenoxyacetic acid (2,4-D), auxin inhibitors, salicylic acid (SA), and trans-cinnamic acid (TCA) were studied using the *tw*-WT system in surface-sterilized and unsterilized germinating grains under high rates of natural infection. Significant differences among the allelic *tw* mutants were revealed at the natural MGG level and in response to 2,4-D, SA, and TCA. The most effective means against MGG were sterilization and TCA. 2,4-D inhibited root growth in *tw* and *tw2* mutants, occurring only in unsterilized and not sterilized germinating grains, while the opposite was observed for TCA and SA. The *tw* mutations influenced variations in the seed-borne fungal spectra, decreasing the frequency of *Bipolaris sorokiniana* and increasing *Fusarium* spp. Hypochlorite-based surface sterilization methods should be used with caution in studies where the action of exogenous 2,4-D will be analysed in germinating grains. Auxin pathway disturbances specific for pleiotropic *tw* mutants are generally restricted to organogenesis but not to germination events.

Keywords: barley ; MGG assay ; salicylic acid ; trans-cinnamic acid ; tweaky mutants ; 2 ; 4-D


## 1. Introduction

Grain contamination with fungi and their produced mycotoxins is not only a problem for organic producers but also for conventional agriculture [1]. For barley, special attention is required for malt and beer contamination with mycotoxins and their proper control and avoidance [2].

## 2. Analysis on Results


### 2.1. The Action of 2,4-D, SA and TCA on MGG and Root Growth of Allelic Tweak Spike Mutants

Naturally, grain sterilization before germination is an effective treatment against MGG. In most cases studied, the MGG level was significantly lower in surface-sterilized grains than in unsterilized grains under the same experimental conditions, and independent of the plant genotype and the studied 2,4-D concentrations (**Figure 1a** and **Figure 2a**, [Tables S1a](#) and [S2a](#)).



**Figure 1.** Effects of 2,4-D over a 10–50 mg L<sup>-1</sup> range on barley *tweaky*-type mutants. (a) The frequency of mouldy grains (%) and (b) the root lengths (cm) of unsterilized and sterilized germinating grains measured after 5 days. For mouldy

grains,  $n = 10$  Petri dishes (100 grains); for root length  $n = 30$ ; for  $tw_2$ ,  $n = 15$  Petri dishes (150 grains) and  $n = 60$ .  $WT_{2016}$  and  $tw_2_{2016}$ , grain reproduction in 2016; other material, grain reproduction in 2013; the MGG assay was performed in 2017. The asterisks represent significant differences (\*  $p < 0.05$ ; \*\*  $p < 0.01$ ; \*\*\*  $p < 0.001$ ) between the control and the 2,4-D treatment. The numbers denote significant differences (<sup>1</sup>  $p < 0.05$ ; <sup>2</sup>  $p < 0.01$ ) between the controls of the  $tw$ -type mutant and the  $WT$ .



**Figure 2.** Effects of 2,4-D over a range of 50–800 mg L<sup>-1</sup> and the auxin inhibitors HFCA and PCIB on (a) the frequency of mouldy grains (%) and (b,c) the root length (cm) of unsterilized germinating grains of  $tw$  mutants and the  $WT$ , as measured after 5 days. For mouldy grains,  $n = 10$  Petri dishes (100 grains); for root length,  $n = 30$ . Grain was produced in 2013, and the MGG assay was performed in 2017. The asterisks represent significant differences (\*  $p < 0.05$ ; \*\*  $p < 0.01$ ; \*\*\*  $p < 0.001$ ) between the control and the 2,4-D treatment. The numbers denote significant differences (<sup>1</sup>  $p < 0.05$ ; <sup>2</sup>  $p < 0.01$ ) between the controls of the  $tw$ -type mutant and the  $WT$ .


In the range of 10–50 mg L<sup>-1</sup>, an appreciable inhibitory effect of 2,4-D on the root growth of allelic  $tw$ -type mutants depended on (1) the plant genotype and (2) the sterilization status of grains (**Figure 1b** and **Table S1b**). In most cases, 2,4-D significantly inhibited root growth only in unsterilized germinating grains of all tested genotypes, including the nonallelic  $twN18$  and  $twmk$  mutants. The  $twmk$  mutant was the most sensitive to 2,4-D among all tested  $tw$ -type mutants. In contrast, 2,4-D-induced root growth inhibition in sterilized germinating grains mostly in a nonsignificant manner (**Figure 1b** and **Table S1b**).

The tested lower 2,4-D concentrations (10–50 mg L<sup>-1</sup>) revealed an interesting dependence of 2,4-D-induced root growth inhibition on grain sterilization status; consequently, the effect of elevated 2,4-D concentrations was investigated in further experiments. In the range of 50–800 mg L<sup>-1</sup> 2,4-D, a significant effect of 2,4-D on MGG was observed only in unsterilized grains of  $WT$  ( $p = 0.0026$ ), in which 2,4-D increased the level of MGG (**Figure 2a** and **Table S2a**), while in the range of 10–50 mg L<sup>-1</sup> 2,4-D, a significant increase in MGG was observed only in sterilized grains of the  $tw_2$  genotype ( $p = 0.024$ ; **Figure 1a** and **Table S1a**).

In contrast to the effect of the lower concentrations, the higher concentrations (50–800 mg L<sup>-1</sup>) of 2,4-D induced a significant decrease in root length independent of the plant genotype and the grain sterilization conditions, and the inhibitory effect of 2,4-D on root growth was dose-dependent (**Figure 2b,c** and **Table S2b**). The root growth of the  $tw$  and  $tw_2$  allelic mutants, but not the  $tw_1$  allelic mutant, was weaker than that of  $WT$  germinating grains. An unsterilized grain background better revealed the inhibitory effect of 2,4-D on the germination rate, which was uniform independent of the plant genotype. In turn, grain sterilization revealed better differences among allelic mutants in the response to 2,4-D according to the germination rate (**Table S2c**). In general, 2,4-D in the range of 50–800 mg L<sup>-1</sup> inhibited root length independent of the plant genotype (**Figure 2c**), while in the range of 10–50 mg L<sup>-1</sup> 2,4-D, differences between  $WT$  and  $tw$ -type mutants and among allelic  $tw$  mutants themselves were observed (**Figure 1**).

Despite the proposed opposite effects to the action of 2,4-D, the auxin inhibitors HFCA and PCIB did not show a significant effect on MGG, except PCIB in sterilized grains of the  $tw_2$  mutant, in which the MGG level decreased (**Figure 2a** and **Table S2a**). However, both auxin inhibitors suppressed root growth similarly to 2,4-D (**Figure 2c** and **Table S2b**).

Theoretically, the effects of TCA and SA are supposed to be opposite to those of 2,4-D, and TCA decreased the frequency of MGG in unsterilized germinating grains of the *WT* and all allelic *tw*-type mutants (**Figure 3** and [Table S3](#)).




**Figure 3.** Effects of salicylic (SA) and *trans*-cinnamic (TCA) acids on (a) the frequency of mouldy grains (%) and (b) the root lengths (cm) of unsterilized and sterilized germinating grains of barley *tw* mutants and the *WT*, as measured after 5 days. For mouldy grains,  $n = 10$  (100 grains); for root length,  $n = 30$ . Grain was produced in 2016, and the MGG test was performed in 2020. The asterisks represent significant differences ( $* p < 0.05$ ;  $** p < 0.01$ ) between the control and the SA or TCA treatment. The numbers denote significant differences ( $^2 p < 0.01$ ;  $^3 p < 0.001$ ) between the controls of the *tw*-type mutant and the *WT*.

However, in sterilized germinating grains, TCA decreased MGG at a significant level only in the allelic *tw* mutant, while SA decreased MGG at a significant level only in unsterilized grains of the *tw2* mutant (**Figure 3** and [Table S3a](#)). Similar to 2,4-D, SA and TCA also inhibited root growth, but only in the allelic mutant *tw2* (**Figure 3b** and [Table S3b](#)). Neither compound showed any effect on the grain germination rate ([Table S3c](#)).

## 2.2. Fungi Spectrum in the Internal Grain Tissues of Barley *tw*-Type Mutants and Revertants

To reveal the possible differences in the fungal diversity in surface-sterilized mouldy germinating grains of tested barley genotypes, the spectrum of fungi species was investigated. Among the fungi that frequently reside in the internal tissues of barley grains, *B. sorokiniana* prevailed in all the tested plant genotypes (**Figure 4** and [Table S4](#)). In addition to the *WT*, *tw1* and *tw2* mutants, the fungi spectra were also studied in several revertants that arose during the phase of stabilization of *tw1* and *tw2* mutants. The revertant studies allowed for a broader understanding of the differences between *tw*-type mutants and the *WT*.




**Figure 4.** Spectra of fungi (%) in the internal grain tissues of barley *tw* mutants and the *WT*. The middle row—revertants from *tw*<sub>1</sub>; the lower row—revertants from *tw*<sub>2</sub>. N—revertants with normal spike and floral structure, C—revertants with normal floral structure but compactoid spikes. Grain was produced in 2013, and the analysis was performed in 2014. The asterisks represent significant differences (\*  $p < 0.05$ ; \*\*  $p < 0.01$ ; \*\*\*  $p < 0.001$ ) between the *tw*-type mutant and the *WT* (in the upper row) or between the revertant and the respective initial *tw* mutant (N1, N13 and C1 derived from *tw*<sub>1</sub>, N46, C6 and C7—from *tw*<sub>2</sub>).

In the grains of the *tw*<sub>1</sub> and *tw*<sub>2</sub> mutants, the *B. sorokiniana* frequency was lower than that in the *WT* strain, but the difference was only statistically significant in *tw*<sub>1</sub>. Interestingly, the level of *B. sorokiniana* was only significantly lower in grains of revertant N1 compared to *WT* ( $p < 0.05$ ) and remained the same as in grains of its parental mutant *tw*<sub>1</sub>. Such a tendency was not observed. A similar tendency was also observed in other revertants, except for the compactoid (C)-type revertants from *tw*<sub>2</sub>, but only in an insignificant manner. However, a decrease in the *Bipolaris* proportion occurred at the expense of the increasing *Fusarium* portion in the fungi spectra, and the observed effect was statistically significant (**Figure 4**). Comparable results were also obtained after analysing the fungi spectra in the internal grain tissues in our previous studies [3]. This finding provided a pretext for studying the effects of grain meals made from the different *tw*-type allelic mutants and the *WT* on *B. sorokiniana* growth.

### 2.3. The Impact of Meals from Grains of *tw*-Type Mutants on the Colony Growth of *Bipolaris sorokiniana*

The growth of *B. sorokiniana* colonies on MEA media supplemented with meals prepared from the grounded dry grains of allelic mutants *tw*, *tw*<sub>1</sub>, and *tw*<sub>2</sub> was compared with *B. sorokiniana* growth on MEA medium containing meals from the grains of *WT*. Additionally, the effects of the SA and TCA concentrations were investigated on such media (**Figure 5**). The meals from the allelic *tw* mutants significantly decreased the growth of *B. sorokiniana* colonies. In most cases, a further statistically significant decrease in *Bipolaris* colony growth occurred only after TCA but not SA addition. SA decreased the growth of *B. sorokiniana* in a concentration-dependent manner only in *tw*<sub>2</sub>. TCA also showed a strong inhibitory effect on *B. sorokiniana* growth on MEA media with meals from the *WT*. Even the lowest concentration of TCA, 0.05 mM, significantly decreased the growth of *Bipolaris* colonies, and a further increase in TCA concentration did not enhance its inhibitory effect on *B. sorokiniana* growth in *WT* media (**Figure 5**).



**Figure 5.** Effects of salicylic acid (SA) and *trans*-cinnamic acid (TCA) on the growth of *Bipolaris sorokiniana* after 7 days of growth on medium containing meal from *tw*. (a) The morphology of *B. sorokiniana* colonies grown on MEA media with different supplements. (b) Size of the colonies 7 days after inoculation. MEA, malt extract medium. Grain was produced in 2013, and the experiment was performed in 2014. The asterisks represent significant differences (\*  $p < 0.05$ ; \*\*  $p < 0.01$ ; \*\*\*  $p < 0.001$ ) between the control and the SA or TCA treatment. The numbers denote significant differences ( $^2 p < 0.01$ ) between the controls of the *tw*-type mutant and the WT.

### 3. Current Insights

The present study showed that more appreciable conclusions on the action of the modifying factors could be made when the MGG assay is performed in parallel with the root growth test. The inhibition of root elongation of germinating grains is one of the earliest and most distinct symptoms exhibited in response to auxins, especially 2,4-D [4], but auxin promotes lateral root formation [5], and this auxin feature is related to pathogen invasion [6]. In the present study, significant differences in 2,4-D-induced root growth inhibition among the different allelic *tw*-type mutants were revealed only in the lower range (10–50 mg L<sup>-1</sup>) of 2,4-D concentrations (Figure 1b). Moreover, significant root length inhibition with 2,4-D in the allelic mutants *tw* and *tw*<sub>2</sub> and the nonallelic mutant *twN18* was only observed in unsterilized germinating grains, whereas in sterilized grains, root growth inhibition in these mutants was absent (Figure 1). While differences in MGG frequency between sterilized and unsterilized grains are naturally expected, the dependence of root growth after 2,4-D treatment on grain sterilization status was quite an unexpected phenomenon. Hypochlorite-based agents, including commercial bleach, are routinely used for surface sterilization of various plant materials [7,8]. Hypochlorite was proposed to react with the seed surface, forming a chlorine cover that is not completely removed by rinsing, and subsequently can be converted into highly toxic chloramines that easily penetrate plant tissues [7]. Furthermore, various salts are known to antagonize the phytotoxicity of several herbicides, including 2,4-D [9,10]. After surface sterilization with commercial bleach, nonremovable chlorine compounds can antagonise exogenously applied 2,4-D and subsequently diminish the effect of root growth inhibition in comparison to that of unsterilized 2,4-D-treated grains. This observation highlights the importance of comparing sterilized and unsterilized grain conditions in studies of plant-auxin–pathogen relationships where the action of exogenous auxin will be analysed in germinating grains since hypochlorite-based sterilization itself can lead to underestimation of the 2,4-D effect on root growth.

### References

- Pinotti, L.; Ottoboni, M.; Giromini, C.; Dell'Orto, V.; Cheli, F. Mycotoxin contamination in the EU feed supply chain: A focus on cereal byproducts. *Toxins* 2016, 8, 45.
- Pascari, X.; Ramos, A.J.; Marín, S.; Sanchís, V. Mycotoxins and beer. Impact of beer production process on mycotoxin contamination. A review. *Food Res. Int.* 2018, 103, 121–129.
- Vaitkūnienė, V.; Varnaitė, A.; Balčiūnienė, L.; Rančelis, V.; Mačkinaitė, R.; Leistrumaitė, A. Two types of revertants from the same homeotic barley mutants tweaky spike. *Biologija* 2006, 2, 18–23.
- Scheitz, K.; Lüthen, H.; Schenck, D. Rapid auxin-induced root growth inhibition requires the TIR and AFB auxin receptors. *Planta* 2013, 238, 1171–1176.

5. Kazan, K.; Lyons, R. Intervention of phytohormone pathways by pathogen effectors. *Plant Cell* 2014, 26, 2285–2309.
6. Kidd, B.N.; Kadoo, N.Y.; Dombrecht, B.; Tekeoglu, M.; Gardiner, D.M.; Thatcher, L.F.; Aitken, E.A.; Schenk, P.M.; Mann  
ers, J.M.; Kazan, K. Auxin signaling and transport promote susceptibility to the root-infecting fungal pathogen *Fusarium*  
*oxysporum* in *Arabidopsis*. *Mol. Plant Microbe Interact.* 2011, 24, 733–748.
7. Miché, L.; Balandreau, J. Effects of rice seed surface sterilization with hypochlorite on inoculated *Burkholderia vietnami*  
*ensis*. *Appl. Environ. Microbiol.* 2001, 67, 3046–3052.
8. Lindsey, B.E.; Rivero, L.; Calhoun, C.S.; Grotewold, E.; Brkljacic, J. Standardized method for high-throughput sterilizati  
on of *Arabidopsis* seeds. *J. Vis. Exp.* 2017, 128, 56587.
9. Nalewaja, J.D.; Woznica, Z.; Matysiak, R. 2,4-D amine antagonism by salts. *Weed Technol.* 1991, 5, 873–880.
10. Nalewaja, J.D.; Matysiak, R. 2,4-D and salt combinations affect glyphosate phytotoxicity. *Weed Technol.* 1992, 6, 322–  
327.

---

Retrieved from <https://encyclopedia.pub/entry/history/show/36810>