

Climate Change Impacts on Sunflower Plants

Subjects: Area Studies

Contributor: Eloisa Agüera

Elevated concentrations of atmospheric carbon dioxide (CO₂) and increased temperatures, carbon and nitrogen metabolism will affect the plant's oxidative state in sunflower (*Helianthus annuus* L.) plants

Keywords: carbon metabolism ; growth

1. Effects of Elevated CO₂ and Elevated Temperatures on Sunflower Plants Growth

In general, elevated CO₂ levels, directly and indirectly affect plant growth and development, modifying numerous physiological processes. Elevated concentrations of CO₂ tend to increase plant growth and produce large quantities of biomass, especially C₃ plants, since they provide additional C (fertilization effect) ^[1]. Plant growth is determined by cell division and expansion. These processes are coordinated and controlled during organogenesis through a series of factors, including vegetable hormones, and they respond to environmental signals ^{[2][3][4]}. An elevated atmospheric CO₂ concentration level may positively influence cell division and expansion ^{[5][6]}. Increased cell expansion is associated with greater extensibility of the cell wall and increased activity of the enzymes that fluidify the wall, such as xyloglucan endotransglucosylase (XET) ^[7]. It has been found that in soy leaves and *Betula papyrifera*, which are grown in a CO₂-enriched environment, certain genes participating in the cell cycle (coding histones) or fluidifying the cell wall (coding expansins and XET) increase their expression ^{[8][9]}. It has been verified that a major supplement of carbon at elevated CO₂ concentrations may contribute to accelerating cell division and expansion in meristematic tissues and improves early plant growth and development ^[10]. Sunflowers grown at elevated CO₂ concentrations were shown to reveal improved growth, reflected in an increased specific leaf mass (SLM), which refers to the dry weight of young leaves (16 days) ^[11]. It is unclear whether or not this increased cell cycle activity resulting from the increased CO₂ is due to the fact that the plant has more photoassimilates for growth or whether it is because of the divergence produced in gene expression in response to the increased sugar levels ^[12]. However, in sunflower plants grown at elevated temperatures, a reduced growth has been observed, as reflected when determining the SLM and area of the leaf as well as the soluble protein content ^[13]. Elevated temperatures negatively affect cell division as well as cell expansion since temperature is one of the main stresses stimulating protein degradation and causing tissue senescence or death ^{[14][15]}. Elevated CO₂ stimulates the root and shoot growth of wheat, but this stimulation was found to reduce when plants were grown in combined elevated temperature and elevated CO₂ ^[16].

2. Elevated CO₂ Levels and Elevated Temperatures on Carbon Metabolism in Sunflower Plants

Elevated levels of CO₂ increase the photosynthetic rate; therefore, crop growth and productivity are increased ^[17]. It has been observed that an elevated concentration of CO₂ stimulates the photosynthetic fixation of CO₂, as well as stoma transpiration and conductance in young sunflower plant leaves ^[11]. Elevated levels of CO₂ concentration increase the photosynthesis rate in C₃ plants, since the Ribulose-1,5-bisphosphate carboxylase/oxygenase (Rubisco) enzyme involved in the fixation process of CO₂ and photorespiration, is not saturated in the environmental CO₂ concentration ^[18]. Therefore, an increase in atmospheric CO₂ would increase the leaf's level of internal CO₂, as well as the CO₂/O₂ ratio, affecting the Rubisco and thereby favoring the carboxylation reaction as compared with the oxygenation process. Elevated CO₂ concentrations may reduce the photorespiration process in C₃ plants and, therefore, the production of cellular hydrogen peroxide (H₂O₂) derived from the metabolism of glycolate ^{[19][20]}. On the other hand, it has been shown that the efficiency of photosystem I and II (PSI and PSII) increases at elevated levels of CO₂, producing more adenosine triphosphate (ATP) and reduced nicotinamide adenine dinucleotide phosphate (NADPH) ^{[21][22]}. In addition, increased efficiency in the use of light is observed as a result of the increased flow of electrons between the PSII and PSI under circumstances of high CO₂ ^[23]. Vicente et al. ^[24] revealed an increased gene and protein expression related to light reactions of photosynthesis.

This stimulating effect of photosynthesis caused by elevated levels of CO₂ may be temporary, given the acclimation of photosynthesis to elevated concentrations of CO₂, which initially stimulates the fixation of C but is followed by a slow decrease in the C fixation process [25]. Various studies have indicated that the acclimation of photosynthesis is due to factors such as reduced content of Rubisco [26], the inhibition of the assimilation of C due to the accumulation of non-structural carbohydrates that suppress the expression of genes related to photosynthesis [26][27], and a reduction in the concentration of nutrients, especially N in plant tissues, due to the inhibition of photoassimilates of NO₃⁻ [28][29][30]. In *Populus tremuloides* and *B. papyrifera* in the presence of elevated CO₂, net photosynthesis increased by 43–73% and the hexose ratio increased when compared with that of sucrose [31]. This was also observed in sunflower leaves [11]. When cucumber plants were grown at high concentrations of CO₂, an increase in the content of starch and soluble sugars was also observed in the leaf, as well as a decrease in the content of nitrogen [32][33]. However, the effect of the elevated CO₂ on the accumulation of hexose varied between species [34][35], as did the sensitivity of the distinct plant tissues [36].

The plant growth and yield depend upon the species specific temperature optimum [37]. An elevated temperature conditions the rate of enzymatic reactions and modifies the structure and activity of macromolecules [38]. In addition, it is known that elevated temperatures modify the composition and structure of cell membranes, increasing the fluidity of membrane lipids and decreasing electrostatic interactions between polar groups of the proteins within the aqueous phase of the membrane and producing a loss of ions [39]. Therefore, photosynthesis at elevated temperatures is modified, since the thylakoid membrane is altered along with the thylakoid shape and arrangement [40]. On the other hand, high temperatures also cause photoinhibition of the PSII through the effect on the oxygen emitter complex, which is destroyed by heat [41][42][43]. The decreased photosynthetic rate may also be due to the fact that elevated temperatures cause stomatal closure to prevent water loss, resulting in a decreased exchange of gases between the leaf and the atmosphere [44]. De la Mata et al. [13], attributed the lower net photosynthesis to elevated temperatures in primary sunflower leaves, compared with a control group, causing a reduction in photosynthetic pigments and partial stomatal closure. Greer and Weedon [45] observed that the average rates of photosynthesis of *Vitis vinifera* leaves decreased by 60% when temperatures increased from 25 to 45 °C. This reduction in photosynthesis was attributed to 15–30% stomatal closure. The photosynthetic rate is also determined by the capacity of carboxylation of Rubisco, which is highly dependent on temperature. Elevated temperatures decrease the state of activation of Rubisco due to the inactivation of the Rubisco activase enzyme, thereby affecting the carbamylation process of the Rubisco [46][47][48][49]. When Rubisco acts as carboxylase, products are frequently formed that prevent its activation, and these should be eliminated from the active site by the Rubisco activase [50][51]. Rubisco activase is relatively labile to heat [48][52]; therefore, its capacity to maintain the Rubisco's state of activation is expected to decrease with elevated temperatures. Plants expressing a more thermotolerant Rubisco activase have higher net photosynthesis at elevated temperatures [53][54]. On the other hand, as the temperature increases, the rate of photosynthesis decreases, with the rate of photorespiration increasing more rapidly [55]. There are two reasons for this. First, as temperatures increase, Rubisco's affinity for CO₂ decreases compared with that of the O₂. Thus, the oxygenation reaction of the Rubisco is more frequent [56][57]. Second, as the temperature increases, the O₂ solubility decreases more slowly than the CO₂ solubility [58]. Therefore, in warm environments, there is relatively more O₂ available to react with the Rubisco.

3. Elevated CO₂ Levels and Elevated Temperatures on Nitrogen Metabolism in Sunflower Plants

Nitrogen is the mineral with the greatest impact in terms of limiting the primary growth and productivity of plants in natural systems and in agriculture. In most soils, nitrogen tends to appear in the form of nitrate (NO₃⁻), since ammonium (NH₄⁺), including that which is added to the soil as fertilizer, is rapidly oxidized to NO₃⁻ by nitrifying bacteria. In plants, nitric nitrogen converts into ammonium nitrogen, a process known as assimilatory reduction in NO₃. The assimilation of NO₃⁻ is regulated by endogenous and/or exogenous factors, such as NO₃⁻, carbon compounds, and light. NH₄⁺ produced from the assimilatory reduction in NO₃⁻, combined with that resulting from other metabolic reactions, is added to the carbon compounds to synthesize nitrogenated compounds that the plant uses for its growth [59].

Stitt and Krapp [60] initially assumed that some plant species required a higher rate of NO₃⁻ assimilation to permit increased plant growth under conditions of elevated CO₂ concentrations. However, it was found that CO₂ enrichment inhibits the assimilation of NO₃⁻ in sunflowers [61] as well as in wheat plants, *Arabidopsis* [62], and field-grown wheat [28]. The assimilation of NO₃⁻ requires the reduced form of nicotinamide adenine dinucleotide (NADH) in order for the nitrate reductase (NR) to catalyze the formation of NO₂⁻ based on NO₃⁻. Photorespiration stimulates the release of malate from the chloroplasts and increases the availability of NADH in the cytosol, thereby increasing the NR activity [63], which permits the first step in NO₃⁻ assimilation [64]. Elevated CO₂ concentrations reduce photorespiration and thus, decrease the quantity of NADH available for the reduction in NO₃⁻. This may explain the decreased levels of NR activity observed in sunflower plants under conditions of elevated CO₂ [61]. However, six transporters from the Nar1 family are involved in the

translocation of NO_2^- from the cytosol to the chloroplast in *Chlamydomonas* some of these transport both NO_2^- as well as HCO_3^- [65]. Bloom et al. [66] revealed that HCO_3^- inhibits the entry of NO_2^- in isolated chloroplasts of wheat and peas, indicating that an analogous system is operating in higher plants. Therefore, a decrease in the affluence of NO_2^- to the chloroplast may result from higher CO_2 levels, which may also explain the reduced glutamine synthetase (GS) activity observed in sunflower plants grown under enriched CO_2 conditions [61].

In sunflower plants grown at elevated CO_2 levels (800 $\mu\text{L L}^{-1}$) and elevated nitrate availability (25 mM), the primary leaves reveal an increased growth, photosynthetic capacity, assimilation of nitrogen, and antioxidant defenses compared with plants grown at elevated CO_2 levels and limited nitrogen. This results in a delay in the leaf's senescence process, demonstrating that the induction of the senescence process is directly related to the C/N ratio of the leaf [67]. This C/N ratio should be balanced in order for the plant to be more productive. An elevated CO_2 increases this ratio in plants due to the decrease in nitrogen content in the leaf [61]. Sunflower plants that are biofertilized via inoculation with mycorrhizal fungi (*Rhizophagus irregularis*) and are grown in environments of elevated CO_2 , and reveal a decrease in the C/N ratio compared with plants grown at elevated CO_2 levels and without biofertilizers.

References

1. Nishimura, C.; Ohashi, Y.; Sato, S.; Kato, T.; Tabata, S.; Ueguchi, C. Histidine kinase homologs that act as cytokinin receptors possess overlapping functions in the regulation of shoot and root growth in Arabidopsis. *Plant Cell* 2004, 16, 1365–1377.
2. Tsukaya, H. Mechanism of leaf-shape determination. *Annu. Rev. Plant Biol.* 2006, 57, 477–496.
3. Riikonen, J.; Percy, K.E.; Kivimäenpää, M.; Kubiske, M.E.; Nelson, N.D.; Vapaavuori, E.; Karnosky, D.F. Leaf size and surface characteristics of *Betula papyrifera* exposed to elevated CO_2 and O_3 . *Environ. Pollut.* 2010, 158, 1029–1035.
4. Ranasinghe, S.; Taylor, G. Mechanism for increased leaf growth in elevated CO_2 . *J. Exp. Bot.* 1996, 47, 349–358.
5. Kinsman, E.; Lew, D.M.; Young, J.F.D.; Vilhar, B.; Ougham, H. Elevated CO_2 stimulates cells to divide in grass meristems: A differential effect in two natural populations of *Dactylis glomerata*. *Plant Cell Environ.* 1997, 20, 1309–1316.
6. Ferris, R.; Sabatti, M.; Miglietta, F.; Mills, R.; Taylor, G. Leaf area is stimulated in *Populus* by free air CO_2 enrichment (POPFACE), through increased cell expansion and production. *Plant Cell Environ.* 2001, 24, 305–315.
7. Ainsworth, E.A.; Rogers, A.; Vodkin, L.O.; Walter, A.; Schurr, U. The effects of elevated CO_2 concentration on soybean gene expression. An analysis of growing and mature leaves. *Plant Physiol.* 2006, 142, 135–147.
8. Kontunen-Soppela, S.; Parviainen, J.; Ruhanen, H.; Brosche, M.; Keinänen, M.; Thakur, R.C.; Kolehmainen, M.; Kanga-sjärvi, J.; Oksanen, E.; Karnosky, D.F. Gene expression responses of paper birch (*Betula papyrifera*) to elevated CO_2 and O_3 during leaf maturation and senescence. *Environ. Pollut.* 2010, 158, 959–968.
9. Wei, H.; Gou, J.; Yordanov, Y.; Zhang, H.; Thakur, R.; Jones, W.; Burton, A. Global transcriptomic profiling of aspen trees under elevated to identify potential molecular mechanisms responsible for enhanced radial growth. *J. Plant Res.* 2013, 126, 305–320.
10. Thilakarathne, C.L.; Tausz-Posch, S.; Cane, K.; Norton, R.M.; Fitzgerald, G.J.; Tausz, M.; Seneweera, S. Intraspecific variation in leaf growth of wheat (*Triticum aestivum*) under Australian Grain Free CO_2 Enrichment (AGFACE): Is it regulated through carbon and/or nitrogen supply? *Func. Plant Biol.* 2015, 42, 299–308.
11. De la Mata, L.; Cabello, P.; de la Haba, P.; Agüera, E. Growth under elevated atmospheric CO_2 concentration accelerates leaf senescence in sunflower (*Helianthus annuus* L.) plants. *J. Plant Physiol.* 2012, 169, 1392–1400.
12. Gamage, D.; Thompson, M.; Sutherland, M.; Hirotsu, N.; Makino, A.; Seneweera, S. New insights into the cellular mechanisms of plant growth at elevated atmospheric carbon dioxide concentrations. *Plant Cell Environ.* 2018, 41, 1233–1246.
13. De la Haba, P.; De la Mata, L.; Molina, E.; Agüera, E. High temperature promotes early senescence in primary leaves of sunflower (*Helianthus annuus* L.) plants. *Can. J. Plant Sci.* 2014, 94, 659–669.
14. Ferguson, D.L.; Guikema, J.A.; Paulsen, G.M. Ubiquitin pool modulation and protein degradation in wheat roots during high temperature stress. *Plant Physiol.* 1990, 92, 740–746.
15. Scheurwater, I.; Dunnebacke, M.; Eising, R.; Lambers, H. Respiratory costs and rate of protein turnover in the roots of a fast-growing (*Dactylis glomerata* L.) and a slow-growing (*Festuca ovina* L.) grass species. *J. Exp. Bot.* 2000, 51, 1089–1097.
16. Benlloch-González, M.; Bochicchio, R.; Berger, J.D.; Bramley, H.; Palta, J. High temperature reduces the positive effect of elevated CO_2 on wheat root system growth. *Fields Crops Res.* 2014, 165, 71–79.

17. Nowicka, B.; Ciura, J.; Szymanska, R.; Kruk, J. Improving photosynthesis, plant productivity and abiotic stress tolerance current trends and future perspectives. *J. Plant Physiol.* 2018, 231, 415–433.
18. Drake, B.G.; González-Meler, M.A.; Long, S.P. More efficient plants: A consequence of rising atmospheric CO₂? *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 1997, 48, 609–639.
19. Pritchard, S.G.; Ju, Z.; van Santen, E.; Qiu, J.; Weaver, D.B.; Prior, S.A.; Rogers, H.H. The influence of elevated CO₂ on the activities of antioxidative enzymes in two soybean genotypes. *Aust. J. Plant Physiol.* 2000, 27, 1061–1068.
20. Bauwe, H.; Hagemann, M.; Fernie, A.R. Photorespiration: Players, partners and origin. *Trends Plant Sci.* 2010, 15, 330–336.
21. Van Heerden, P.; Swanepoel, J.; Kruger, G. Modulation of photosynthesis by drought in two desert scrub species exhibiting C₃-mode CO₂ assimilation. *Environ. Ex Bot.* 2007, 61, 124–136.
22. Zhang, D.-Y.; Chen, G.-Y.; Gong, Z.-Y.; Chen, J.; Yong, Z.-H.; Zhu, J.-G.; Xu, D.-Q. Ribulose-1, 5-bisphosphate regeneration limitation in rice leaf photosynthetic acclimation to elevated CO₂. *Plant Sci.* 2008, 175, 348–355.
23. Sekhar, K.M.; Sreeharsha, R.V.; Reddy, A.R. Differential responses in photosynthesis, growth and biomass yields in two mulberry genotypes grown under elevated CO₂ atmosphere. *J. Photochem. Photobiol. B Biol.* 2015, 151, 172–179.
24. Vicente, R.; Pérez, P.; Martínez-Carrasco, R.; Usadel, B.; Kostadinova, S.; Morcuende, R. Quantitative RT–PCR platform to measure transcript levels of C and N metabolism-related genes in durum wheat: Transcript profiles in elevated and high temperature at different levels of N supply. *Plant Cell Physiol.* 2015, 56, 1556–1573.
25. Kant, S.; Seneweera, S.; Rodin, J.; Materne, M.; Burch, D.; Rothstein, S.J.; Spangenberg, G. Improving yield potential in crops under elevated CO₂: Integrating the photosynthetic and nitrogen utilization efficiencies. *Front. Plant Sci.* 2012, 3, 162.
26. Long, S.P.; Ainsworth, E.A.; Rogers, A.; Ort, D.R. Rising atmospheric carbon dioxide: Plants FACE the future. *Annu. Rev. Plant Biol.* 2004, 55, 591–628.
27. Kitaoka, S.; Matsuki, S.; Kitao, M.; Tobita, H.; Utsugi, H.; Maruyama, Y.; Koike, T. The photosynthetic response of four several deciduous broad-leaved tree seedlings grown under elevated CO₂ concentrations. *J. Agric. Meteorol.* 2016, 72, 3–49.
28. Bloom, A.J.; Burger, M.; Kimball, B.A.; Pinter, P.J. Nitrate assimilation is inhibited by elevated CO₂ in field-grown wheat. *Nat. Clim. Chang.* 2014, 4, 477–480.
29. Ruiz-Vera, U.M.; De Souza, A.P.; Long, S.P.; Ort, D.R. The role of sink strength and nitrogen availability in the down-regulation of photosynthetic capacity in field-grown *Nicotiana tabacum* L. at elevated CO₂ concentration. *Front. Plant Sci.* 2017, 8, 998.
30. Bahrami, H.; De Kok, L.J.; Armstrong, R.; Fitzgerald, G.J.; Bourgault, M.; Henty, S.; Tausz, M.; Tausz-Posch, S. The proportion of nitrate in leaf nitrogen, but not changes in root growth, are associated with decreased grain protein in wheat under elevated . *J. Plant. Physiol.* 2017, 216, 44–51.
31. Riikonen, J.; Kets, K.; Darbaj, J.; Oksanen, E.; Sober, A.; Vapaavuori, E.; Kubiske, M.E.; Nelson, N.; Karnosky, D.F. Carbon gain and bud physiology in *Populus tremuloides* and *Betula papyrifera* grown under long-term exposure to elevated concentrations of CO₂ and O₃. *Tree Physiol.* 2008, 28, 243–254.
32. Larios, B.; Agüera, E.; de la Haba, P.; Pérez-Vicente, R.; Maldonado, J.M. A short-term exposure of cucumber plants to rising atmospheric CO₂ increases leaf carbohydrate content and enhances nitrate reductase expression and activity. *Planta* 2001, 212, 305–312.
33. Agüera, E.; Ruano, D.; Cabello, P.; De la Haba, P. Impact of atmospheric CO₂ on growth, photosynthesis and nitrogen metabolism in cucumber (*Cucumis sativus* L.) plants. *J. Plant Physiol.* 2006, 163, 809–817.
34. Ainsworth, E.A.; Long, S.P. What have we learned from 15 years of free-air CO₂ enrichment (FACE)? A meta-analytic review of the responses of photosynthesis, canopy properties and plant production to rising CO₂. *New Phytol.* 2005, 165, 351–372.
35. Bae, H.; Sicher, R. Changes of soluble protein expression and leaf metabolite levels in *Arabidopsis thaliana* grown in elevated atmospheric carbon dioxide. *Field Crop Res.* 2004, 90, 61–73.
36. Araya, T.; Noguchi, K.; Terashima, I. Differences between sink and source leaves in carbohydrate repression of photosynthesis. In *Photosynthesis: Fundamental Aspects to Global Perspective*; van der Est, A., Bruce, D., Eds.; International Society of Photosynthesis: Montreal, QC, Canada, 2005; pp. 660–662.
37. Gray, S.B.; Brady, S.M. Plant developmental responses to climate change. *Develop. Biol.* 2016, 419, 64–77.
38. Arcus, V.L.; Prentice, E.J.; Hobbs, J.K.; Mulholland, A.J.; Van der Kamp, M.W.; Pudney, C.R.; Parker, E.J.; Schipper, L.A. On the temperature dependence of enzyme catalyzed rates. *Biochemistry* 2016, 55, 1681–1688.

39. Li, Q.; Zheng, Q.; Shen, W.; Cram, D.; Fowler, D.B.; Wei, Y.; Zou, J. Understanding the biochemical basis of temperature-induced lipid pathway adjustments in plants. *Plant Cell*. 2015, 27, 86–103.
40. Semenova, G. Structural reorganization of thylakoid systems in response to heat treatment. *Photosynthetica* 2004, 42, 521–527.
41. Aro, E.; Virgin, I.; Andersson, B. Photoinhibition of photosystem II. Inactivation, protein damage and turnover. *Biochim. Biophys. Acta* 1993, 1143, 113–134.
42. Murata, N.; Takahashi, S.; Nishiyama, Y.; Allakhverdiev, S.I. Photoinhibition of photosystem II under environmental stresses. *Biochim. Biophys. Acta* 2007, 1767, 414–421.
43. Takahashi, S.; Murata, N. How do environmental stresses accelerate photoinhibition? *Trends Plant Sci.* 2008, 13, 178–182.
44. Greer, D.H.; Weedon, M.M. Modelling photosynthetic responses to temperature of grapevine (*Vitis vinifera* cv. Semillon) leaves on vines grown in a hot climate. *Plant Cell Environ.* 2012, 35, 1050–1064.
45. Greer, D.H.; Weston, C. Heat stress affects flowering, berry growth, sugar accumulation and photosynthesis of *Vitis vinifera* cv. Semillon grapevines grown in a controlled environment. *Func. Plant Biol.* 2010, 37, 206–214.
46. Feller, U.; Crafts-Brandner, S.J.; Salvucci, M.E. Moderately high temperatures inhibit ribulose-1,5-bisphosphate carboxylase/oxygenase (rubisco) activase-mediated activation of rubisco. *Plant Physiol.* 1998, 116, 539–546.
47. Jiang, D.; Lu, Q.; Weng, X.; Zheng, B.; Xi, H. Regulation of rubisco carboxylation activity and photosynthetic rate by rubisco activase during leaf senescence in rice. *J. Zhejiang Univ. (Agric. Life Sci.)* 1999, 26, 119–124.
48. Salvucci, M.E.; Crafts-Brandner, S.J. Inhibition of photosynthesis by heat stress: The activation state of rubisco as a limiting factor in photosynthesis. *Physiol. Plant* 2004, 120, 179–186.
49. Demirevska-Kepova, K.; Holzer, R.; Simova-Stoilova, L.; Feller, U. Heat stress effects on ribulose-1, 5-bisphosphate carboxylase/oxygenase, rubisco binding protein and rubisco activase in wheat leaves. *Biol. Plant.* 2005, 49, 521–525.
50. Carmo-Silva, E.; Scales, J.C.; Madgwick, P.J.; Parry, M.A.J. Optimizing Rubisco and its regulation for greater resource use efficiency. *Plant Cell Environ.* 2015, 38, 1817–1832.
51. Bracher, A.; Whitney, S.M.; Hartl, F.U.; Hayer-Hartl, M. Biogenesis and metabolic maintenance of Rubisco. *Annu. Rev. Plant Biol.* 2017, 68, 29–60.
52. Crafts-Brandner, S.J.; Salvucci, M.E. Sensitivity of photosynthesis in a C4 plant, maize, to heat stress. *Plant Physiol.* 2002, 129, 1773–1780.
53. Itzhak Kurek, O.; Kai Chang, T.; Bertain, S.M.; Madrigal, A.; Liu, L.; Lassner, M.W.; Zhu, G. Enhanced thermostability of Arabidopsis Rubisco activase improves photosynthesis and growth rates under moderate heat stress. *Plant Cell* 2007, 19, 230–3241.
54. Kumar, A.; Verma, J.P. Does plant-microbe interaction confer stress tolerance in plants: A review? *Microbiol. Res.* 2018, 207, 41–52.
55. Long, S.P. Modification of the response of photosynthetic productivity to rising temperature by atmospheric CO₂ concentrations: Has its importance been underestimated? *Plant Cell Environ.* 1991, 4, 729–739.
56. Ku, S.B.; Edwards, G.E. Oxygen inhibition of photosynthesis: I. Temperature dependence and relation to O₂/CO₂ solubility ratio. *Plant Physiol.* 1977, 59, 986–990.
57. Jordan, D.B.; Ogren, W.L. The CO₂/O₂ specificity of ribulose 1,5-bisphosphate carboxylase/oxygenase. *Planta* 1984, 161, 308–313.
58. Ku, S.B.; Edwards, G.E. Oxygen inhibition of photosynthesis: II. Kinetic characteristics as affected by temperature. *Plant Physiol.* 1977, 59, 991–999.
59. Kishorekumar, R.; Bulle, M.; Wany, A.; Gupta, K.J. An overview of important enzymes involved in nitrogen assimilation of plants. In *Nitrogen Metabolism in Plants, Methods and Protocols*; Gupta, K.J., Ed.; Springer Science-Business Media, LLC, Part of Springer Nature: New York, NY, USA, 2020; pp. 1–13.
60. Stitt, M.; Krapp, A. The interaction between elevated carbon dioxide and nitrogen nutrition: The physiological and molecular background. *Plant Cell Environ.* 1999, 22, 583–621.
61. De La Mata, L.; De la Haba, P.; Alamillo, M.J.; Pineda, M.; Agüera, E. Elevated CO₂ concentrations alter nitrogen metabolism and accelerate senescence in sunflower (*Helianthus annuus* L.) plants. *Plant Soil Environ.* 2013, 59, 303–308.
62. Bloom, A.J.; Burger, M.; Rubio Asensio, J.S.; Cousins, A.B. Carbon dioxide enrichment inhibits nitrate assimilation in wheat and Arabidopsis. *Science* 2010, 328, 899–903.

63. Igamberdiev, A.U.; Bykova, N.V.; Lea, P.J.; Gardeström, P. The role of photorespiration in redox and energy balance of photosynthetic plant cells: A study with a barley mutant deficient in glycine decarboxylase. *Physiol. Plant.* 2001, 111, 427–438.
 64. Quesada, A.; Gómez-García, I.; Fernández, E. Involvement of chloroplast and mitochondria redox valves in nitrate assimilation. *Trends Plant Sci.* 2000, 5, 463–464.
 65. Mariscal, V.; Moulin, P.; Orsel, M.; Miller, A.J.; Fernández, E.; Galván, A. Differential regulation of the *Chlamydomonas* Nar1 gene family by carbon and nitrogen. *Protist* 2006, 157, 421–433.
 66. Bloom, A.J.; Smart, D.R.; Nguyen, D.T.; Searles, P.S. Nitrogen assimilation and growth of wheat under elevated carbon dioxide. *Proc. Natl. Acad. Sci. USA* 2002, 99, 1730–1735.
 67. Canales, F.J.; De la Haba, P.; Barrientos, E.; Aguera, E. Effect of CO₂ enrichment and increased nitrogen supply on the induction of sunflower (*Helianthus annuus* L.) primary leaf senescence. *Can. J. Plant. Sci.* 2016, 96, 1002–1013.
-

Retrieved from <https://encyclopedia.pub/entry/history/show/44358>