

Polar Lipids from Olives/Olive Oil

Subjects: [Food Science & Technology](#)

Contributor: Eliana Alves

Polar lipids are minor components of olives and olive oil and include a myriad of molecules such as phospholipids and glycolipids.

authentication

bioactive

by-product

glycolipid

lipidomics

mass spectrometry

phospholipid

traceability

1. Introduction

For millennia, olive oil has been an essential ingredient in the Mediterranean diet, as a food source of healthy fat. It is produced mostly by Spain, Italy, Greece and by other countries of Southern Europe and North Africa ^[1]. Nowadays, olive oil's economy has gained global importance, especially in gourmet cuisine, and its production has been extended to North and South Americas, Australia and Asia ^[1].

The increasing investment in the development of olive groves in these regions has been boosted by the benefits of olive oil's consumption which is directly related to its composition. Olive oil is mainly composed of triacylglycerols (Ca. 98%) ^[2], primarily consisting of monounsaturated fatty acids, acknowledged for improving several cardiovascular risk factors ^[3]. In addition to the primary compounds, high-quality olive oils, such as virgin olive oils (VOOs), possess a plethora of minor components in the remaining 2% of their composition ^[2]. Some of the minor components confer distinct features to olive oil in terms of sensorial attributes and health benefits ^{[4][5]}, and some components can be used for providing a chemical identity to olive oil ^[6].

Polar lipids are a group of minor components of olive oil ^[2]. The isolation, identification, and characterization of the minor components, such as polar lipids, might be essential to provide a molecular fingerprint for traceability and authenticity purposes ^[7]. The profiling of the major chemical components, such as triacylglycerols and total fatty acids, is insufficient to discriminate olives or olive oils, per se, and the simultaneous analysis of minor components is necessary ^[8]. VOOs are very susceptible to fraud and to tampering with other oils, as lower grade olive oils ^{[9][10]}. With recent analytical developments, new fast and sensitive methods have been claimed to evaluate olive oil's authenticity ^[11]. Therefore, it has become urgent to find foolproof analytical approaches and molecular markers to reveal a specific chemical identity for olives and olive oil and to detect adulterated olive oil ^[10]. Polar lipids have been suggested as promising molecular markers of identity ^{[12][13]}. Some research has been carried out towards their identification in olives and olive oil, mainly through mass spectrometry (MS)-based approaches, but there is still much to be done.

Another topic concerning olives' and olive oil's polar lipids is their positive impact on human nutrition and health, which has been little exploited [14][15]. Additionally, in recent years, polar lipids from olive-derived industrial by-products, such as olive seeds and olive pomace, have been studied as alternative sources of bioactive lipids. The new applications of polar lipids would favor the sustainable use of olive's industrial by-products and make them attractive from the biotechnological standpoint.

2. Identification of Polar Lipids from Olives, Olive Oil, and Their Industrial By-Products

The identification of polar lipids in olives and olive oil is a difficult task since they are minor components and include a broad range of lipid classes. Different analytical approaches have been used to unravel the polar lipidome of these matrices. The lipidomic workflows included lipid extraction, fractionation, analysis and quantification (Figure 1).




Figure 1. Schematic representation of the methodological approaches used for studying polar lipids from olives and olive oil. Abbreviations: HPLC, high-performance liquid chromatography; HPLC-MS, high-performance liquid

chromatography coupled to mass spectrometry; HPLC-MS/MS, high-performance liquid chromatography coupled to tandem mass spectrometry; HPLC-UV, high-performance liquid chromatography with ultraviolet detector; LLE, liquid/liquid extraction; NACE, non-aqueous capillary electrophoresis; NACE-MS, non-aqueous capillary electrophoresis coupled to mass spectrometry; NACE-MS/MS, non-aqueous capillary electrophoresis coupled to tandem mass spectrometry; NMR, nuclear magnetic resonance; SPE, solid-phase extraction.

Liquid/liquid extraction (LLE) has been used for extracting polar lipids from olives and olive oil. The most commonly used LLE methods were a modified Bligh and Dyer method [16], a modified Folch method [17] and a sequential LLE method developed by Galanos and Kapoulas [17][18][19]. Solid-phase extraction (SPE), using aminopropyl-bonded silica as sorbent, was recently used to obtain polar lipid-enriched fractions directly from olive oil [12]. There are other emerging extraction techniques that can be used for oil extraction from olives, such as ultrasound or microwave or CO₂-assisted techniques, but these approaches have not yet been reported for the analysis of polar lipids in olives or olive oil.

After extraction, the total lipid extract can be fractionated to obtain polar lipid-enriched fractions or specific polar lipid classes. Polar lipid-enriched fractions were obtained using SPE cartridges with different stationary phases (silica and diol-bonded silica) after olive oil's LLE [19].

³¹P nuclear magnetic resonance (NMR) spectroscopy [18] and non-aqueous capillary electrophoresis (NACE) coupled with MS [17] were used for the detection and characterization of the phospholipid classes of olive oil.

The separation of the polar lipid classes obtained from olive oil was carried out by high-performance liquid chromatography (HPLC) coupled to different detectors, as ultraviolet detectors (HPLC-UV) [20] or mass spectrometers (HPLC-MS) [12][16][19]. The structural characterization of the polar lipid molecules, namely the polar head and fatty acyl composition, has been achieved by using tandem MS (HPLC-MS/MS in [12][16][19] and NACE-MS/MS in [17]).

The analytical approaches used so far (**Table 1**) showed different results. In olive fruits, the polar lipidome has been studied in the oil extracted both from the pulp and the seed. Bianco et al. (1998) identified glycolipids in the olive pulp, namely digalactosyldiacylglycerols as DGDG(18:3/18:3) and DGDG(18:1/18:3) [20]. Montealegre et al. (2013) analyzed the glycerophospholipid profile of olive fruits from different Spanish cultivars and regions [17]. The glycerophospholipids identified in the olive pulp and in the seed included phosphatidic acid (PA), lyso-PA, phosphatidylethanolamine (PE), lyso-PE, phosphatidylcholine (PC), phosphatidylinositol (PI) and phosphatidylglycerol (PG) [17] (**Figure 2**).




Figure 2. Chemical structures of the classes of glycerophospholipids and glycolipids identified in olives and olive oil. Polar lipids include a broad range of molecules. Phospholipids are divided into two main classes depending on whether they contain glycerol (glycerophospholipids) or a sphingosyl (sphingophospholipids) backbone. Glycerophospholipids, besides the glycerol backbone, contain a polar phosphorus moiety. They derive mainly from *sn*-1,2-diacylglycerols and, thus, contain structures that are based on 3-*sn*-phosphatidic acid [21]. These lipids are grouped into classes based on the composition of their polar head group that is attached to the phosphate residue in *sn*-3 position. The polar head may be an amino acid, an amino-alcohol, a carbohydrate or another functional moiety. Each head group class is further differentiated into subclasses based on the *sn*-1 and *sn*-2 substituents on the glycerol backbone [21]. Glycolipids also include a wide variety of structures. These structures consist in acylglycerols (in the case of glycosylglycerides and sulfolipids) joined to a carbohydrate moiety by a glycosidic linkage at the *sn*-3 position [21]. Betaine lipids are ether-linked glycerolipids containing a betaine moiety. These lipids contain a polar group linked by an ether bond at the *sn*-3 position of the glycerol moiety, with the fatty acids esterified in the *sn*-1 and *sn*-2 positions [21]. 1,2-diacylglyceryl-3-O-4'-(*N,N,N*-trimethyl)-homoserine (DGTS) have been commonly found in lower plants, algae, fungi, and bacteria [22]. R, R1, and R2 represent fatty acyl chains.

Table 1. Summary of the polar lipid classes identified and quantified in olives and olive oils in different studies.

Reference	Sampling		Analysis		Polar Lipid Classes
	Type of Sample	Amount of Sample	Extraction	Method	
[20]	Olive fruit and olive oil from varieties Carolea and Ottobratica, both from Calabria region (Italy)	Olive fruit (250 g); olive oil (10 mL)	Glycosidic fraction in olive fruit: ethanol and "charcoal method"; glycosidic fraction in the aqueous phase of olive oil: ethyl acetate/dichloromethane (1:1 by volume) and water	HPLC-UV (μ -Bondapak C18 column)	DGDG
[16]	Tunisian commercial olive oil	Not said	Modified Bligh and Dyer method	HPLC-MS/MS (diol column)	PG (63%), PA (12%), PI (11%), PE (9%), PC (5%)
[18]	Greek virgin olive oil, refined olive oil and olive pomace oil from local cooperatives (7 regions and 5 cultivars)	100 g	According to Galanos and Kapoulas (1962)	^{31}P -NMR	PA, lyso-PA, lyso-PI, PI, PG (PG only in pomace oil), PC and PE (these two only in virgin olive oil).
[17]	Olive pulp and olive stone from Spanish Arbequina variety from three geographical regions (Córdoba, Jaén, and Toledo) and two Spanish varieties (Empeltre and Lechín de	Olive pulp or stone (2.5 g); olive oil (50 g)	PL from olive pulp and stone: modified Folch method; PL from olive oil: LLE according to Galanos and Kapoulas (1962)	NACE-ESI-MS and MS/MS	Olives (stone and pulp studied independently): PA (54–82%), PE (4–16%), PC (3–9%), lyso-PE (1.3–18%), PI (4.4–8%), PG (3.7–6.3%), and lyso-PA (0.1–0.2%).

Reference	Sampling		Analysis		Polar Lipid Classes
	Type of Sample	Amount of Sample	Extraction	Method	
	Sevilla) from the same region (Córdoba); commercial monovarietal extra virgin olive oil from Arbequina variety				Olive oil: PE (42%), PG (38%), PC (15%), lyso-PE (4.5%), and lyso-PA (0.2%)
[19]	Italian olive oil blend (Leccino, Frantoio and Picholine varieties) from a local mill of Emilia Romagna region (Italy)	100 g for LLE; 40 g for SPE	LLE according to Galanos and Kapoulas (1962) followed by SPE (diol and silica). PL eluted with methanol and chloroform/methanol/water (3:5:2 by volume)	HPLC-ESI-qTOF-MS (HILIC column)	Diol extracted veiled extra virgin olive oil (mg kg ⁻¹): lyso-PA (4.23), lyso-PC (1.21), PI (1.03), PC (0.90), PA (0.81), PG (0.07). Crystallized veiled virgin olive oil (mg kg ⁻¹): lyso-PA (1.15), lyso-PC (0.87), PC (0.74), PI (0.48), PA (0.14)
[12]	Portuguese commercial extra virgin and virgin olive oils	1 g	PL extracted by SPE (aminopropyl columns) and eluted with acetonitrile: ammonium hydroxide (95:5 by volume)	HPLC-ESI-ion trap-MS/MS (HILIC column)	PA, PE, PG, PC, PI, SQDG, SQMG, DGTS

3. Schwingshackl, L.; Hoffmann, G. Monounsaturated fatty acids and risk of cardiovascular disease: Synopsis of the evidence available from systematic reviews and meta-analyses. *Nutrients* 2012, 4, 1989–2007.

Legend: DGDG, digalactosyldiacylglycerol; DGTS, diacylglycerol-N,N,N-trimethylhomoserine; HILIC, hydrophilic interaction liquid chromatography; HPLC-ESI-MS/MS, hydrophilic interaction liquid chromatography coupled to electrospray ionization tandem mass spectrometry; HPLC, high performance liquid chromatography; HPLC-ESI-

4. Campestre, C.; Angelini, G.; Gasbarr, G.; Angerosa, F. The compounds responsible for the sensory profile in monovarietal virgin olive oils. *Molecules* 2017, 22, 1833.

51. Serrilli, M.; Ferraro, S.; Faludi, R.; Urbani, S.; Taticchi, A.; Marrazzi, F.; Salvaggio, R.; Montedone, G.; Perrini, C. **Polar lipids in olive oil: Antioxidant, pro- and anti-inflammatory, and neuroprotective activities according to their chemical structure.** *Inflammopharmacology* 2009, 17, 76–84. liquid/liquid extraction; MS/MS, tandem mass spectrometry; NACE-ESI-MS, non-aqueous capillary electrophoresis coupled to electro-spray ionization mass spectrometry; NMR, nuclear magnetic resonance; PA, phosphatidic acid; PC, phosphatidylcholine; PE, phosphatidylethanolamine; PG, phosphatidylglycerol; PI, phosphatidylinositol; PL, polar lipid; SPE, solid-phase extraction; SQDG, sulfoquinovosyldiacylglycerol; SQMG, sulfoquinovosylmonoacylglycerol.

7. Perri, E.; Benincasa, C.; Muzzalupo, I. Chapter 13. Olive oil traceability. In *Olive Germplasm-the Olive Cultivation Table Olive and Olive Oil Industry in Italy*; Muzzalupo, I., Ed.; InTechOpen: London, UK, 2012; pp. 265–286, ISBN 978-953-51-0883-2.

3. Potential Biotechnological Uses of Polar Lipids from Olives and Olive Oil's Industrial By-Products

8. Montealegre, C.; Marina Alegre, M.L.; García-Ruiz, C. Traceability markers to the botanical origin of olive oils and jointed table olives' producing industries generate several by-products, such as olive pomace and olive stones. These by-products can be recovered to create novel value-added products. In the case of polar lipids,

9. Aparicio, R.; Morales, M.T.; Aparicio-Ruiz, R.; Tena, N.; García-González, D.L. Authenticity of their concentration is tens to hundred times higher in olive pomace oil ^[18] and olive seed oil ^[23], comparatively to olive oil ^{[18][24]}. Thus, polar lipids from olive pomace and olive seeds have been regarded as potentially useful from the nutritional and biotechnological standpoints and have been suggested for several novel industrial applications.

10. Gallina Toschi, T.; Bendini, A.; Lozano-Sánchez, J.; Segura-Carretero, A.; Conte, L.

Olive pomace was proposed as the new promising lipid source for the sustainable production of animal feeds, namely functional fish feeds for aquaculture fish and as an ingredient for inclusion in animal feedstocks ^[25]. *Food Res. Int.* 2013, 54, 1205–1223.

Olive pomace after stoning has been extensively studied in mammal's species as feed integration for improving the 11. Bajoub, A.; Bendini, A.; Fernández-Gutiérrez, A.; Carrasco-Pancorbo, A. Olive oil authentication: A nutritional and nutraceutical properties of their meat as well as their milk and derived cheese ^{[26][27][28][29][30]}. Other comparative analysis of regulatory frameworks with especial emphasis on quality and authenticity studies carried out on fish species revealed that polar lipids from olive pomace oil ^[31] provide high nutritional value indices, and recent analytical techniques developed for their assessment. A review. *Crit. Rev. Food Sci. Nutr.* 2018, 58, 832–857.

12. Alves, S.A.E.; Melo, T.; Rey, F.; Moreira, J.A.S.; Domingues, P.; Deodrigues, M.R. Polar lipids from olive pomace oils as a useful tool for identifying decipies their unique fingerprints. *With Food Sci. Technol.* 2016, 74, 371–377.

13. Calvano, C.D.; De Ceglie, C.; D'Accolti, L.; Zamboni, C.G. MALDI-TOF mass spectrometry

Other residues resulting from table olives' production are the stones that contain the seeds. The economic potentialities of olive seeds and olive seed oil have been explored in the last few years, primarily by the industry ^[33]. *Food Chem.* 2012, 134, 1192–1198.

14. Karantonis, H.C.; Antonopoulou, S.; Demopoulos, C.A. Antithrombotic lipid minor constituents

Olive seed oil has 8–19% of phospholipids ^[23] and may have diverse technological uses in the soap, cosmetics, and pharmaceutical industries ^[23]. Phospholipids from olive seeds can also be used for lecithin production in the agri-food industry ^[23]. Food derived phospholipids have several biomedical applications, for instance, as emulsifiers in

15. Tsantila, N.; Karantonis, H.C.; Perrea, D.N.; Theocharis, S.E.; Iliopoulos, D.G.; Antonopoulou, S.; Demopoulos, C.A. Antithrombotic and antiatherosclerotic properties of olive oil and olive pomace

polar extracts in rabbits. *Med. Inflamm.* 2007, 2007, 11.

The potential biotechnological applications of the olive-derived by-products highlight the valuable alternatives that underlie the table olive's and olive oil's industries. However, more research is needed to characterize the polar lipids, its health benefits, and the cost benefit of being extracted from these by-products. 16. Boukhenina, S.; Sebba, K.; Chen, A.; Kafil, H.; Mayer, P.M. Identification of glycerophospholipids in rapeseed, olive, almond, and sunflower oils by LC-MS and LC-MS/MS. *Can. J. Chem.* 2004,

- 82, 1210–1215.
17. Montealegre, C.; Sanchez-Hernandez, L.; Crego, A.; Marina, M. Determination and characterization of glycerophospholipids in olive fruit and oil by nonaqueous capillary electrophoresis with electrospray-mass spectrometric detection. *J. Agric. Food Chem.* 2013, 61, 1823–1832.
 18. Hatzakis, E.; Koidis, A.; Boskou, D.; Dais, P. Determination of phospholipids in olive oil by ³¹P-NMR spectroscopy. *J. Agric. Food Chem.* 2008, 56, 6232–6240.
 19. Verardo, V.; Gómez-Caravaca, A.; Montealegre, C.; Segura-Carretero, A.; Caboni, M.; Fernández-Gutiérrez, A.; Bendini, A. Optimization of a solid phase extraction method and hydrophilic interaction liquid chromatography coupled to mass spectrometry for the determination of phospholipids in virgin olive oil. *Food Res. Int.* 2013, 54, 2083–2090.
 20. Bianco, A.; Mazzei, R.A.; Melchioni, C.; Scarpati, M.L.; Romeo, G.; Uccella, N. Microcomponents of olive oil. Part II: Digalactosyldiacylglycerols from *Olea europaea*. *Food Chem.* 1998, 62, 343–346.
 21. Scrimgeour, C.M.; Harwood, J.L. Chapter 1: Fatty acid and lipid structure. In *The Lipid Handbook*, 3rd ed.; Gunstone, F.D., Harwood, J.L., Dijkstra, A.J., Eds.; CRC Press: Boca Raton, FL, USA, 2007; pp. 1–36. ISBN 978-0849396885.
 22. Dembitsky, V.M. Betaine ether-linked glycerolipids: Chemistry and biology. *Prog. Lipid Res.* 1996, 35, 1–51.
 23. Moussaoui, R.; Labbaci, W.; Hemar, N.; Youyou, A.; Amir, Y. Physico-chemical characteristics of oils extracted from three compartments of the olive fruit (pulp, endocarp and seed) of variety chemlal cultivated in Kabylia (Algeria). *J. Food Agric. Environ.* 2008, 6, 52–55.
 24. Koidis, A.; Boskou, D. The contents of proteins and phospholipids in cloudy (veiled) virgin olive oils. *Eur. J. Lipid Sci. Technol.* 2006, 108, 323–328.
 25. Nasopoulou, C.; Zabetakis, I. Agricultural and aquacultural potential of olive pomace a review. *J. Agric. Sci.* 2013, 5, 116–127.
 26. Caputo, A.; Morone, G.; Di Napoli, M.A.; Rufrano, D.; Sabia, E.; Paladino, F.; Sepe, L.; Claps, S. Effect of destoned olive cake on the aromatic profile of cows' milk and dairy products: Comparison of two techniques for the headspace aroma profile analysis. *Ital. J. Agron.* 2015, 10, 15–20.
 27. Castellani, F.; Vitali, A.; Bernardi, N.; Marone, E.; Palazzo, F.; Grotta, L.; Martino, G. Dietary supplementation with dried olive pomace in dairy cows modifies the composition of fatty acids and the aromatic profile in milk and related cheese. *J. Dairy Sci.* 2017, 100, 8658–8669.
 28. Cibik, M.; Keles, G. Effect of stoned olive cake on milk yield and composition of dairy cows. *Revue Méd. Vét.* 2016, 167, 154–158.

29. Terramoccia, S.; Bartocci, S.; Taticchi, A.; Di Giovanni, S.; Pauselli, M.; Mourvaki, E.; Urbani, S.; Servili, M. Use of dried stoned olive pomace in the feeding of lactating buffaloes: Effect on the quantity and quality of the milk produced. *Asian-Australas. J. Anim. Sci.* 2013, 26, 971–980.
30. Vargas-Bello-Pérez, E.; Vera, R.R.; Aguilar, C.; Lira, R.; Peña, I.; Fernández, J. Feeding olive cake to ewes improves fatty acid profile of milk and cheese. *Anim. Feed Sci. Technol.* 2013, 184, 94–99.
31. Karantonis, H.C.; Tsantila, N.; Stamatakis, G.; Samiotaki, M.; Panayotou, G.; Antonopoulou, S.; Demopoulos, C.A. Bioactive polar lipids in olive oil, pomace and waste byproducts. *J. Food Biochem.* 2008, 32, 443–459.
32. Nasopoulou, C.; Smith, T.; Detopoulou, M.; Tsikrika, C.; Papaharisis, L.; Barkas, D.; Zabetakis, I. Structural elucidation of olive pomace fed sea bass (*Dicentrarchus labrax*) polar lipids with cardioprotective activities. *Food Chem.* 2014, 145, 1097–1105.
33. Are Olive Seeds the Next Superfood? Available online: (accessed on 7 January 2018).
34. Lodén, M. Role of topical emollients and moisturizers in the treatment of dry skin barrier disorders. *Am. J. Clin. Dermatol.* 2003, 4, 771–788.
35. Van Hoogevest, P.; Wendel, A. The use of natural and synthetic phospholipids as pharmaceutical excipients. *Eur. J. Lipid Sci. Technol.* 2014, 116, 1088–1107.

Retrieved from <https://encyclopedia.pub/entry/history/show/27218>