

# Ankyrin/Spectrin in Cardiac Conduction

Subjects: Cardiac & Cardiovascular Systems


Contributor: Thomas Hund

The cardiac conduction system is an extended network of excitable tissue tasked with generation and propagation of electrical impulses to signal coordinated contraction of the heart. The fidelity of this system depends on the proper spatio-temporal regulation of ion channels in myocytes throughout the conduction system. Importantly, inherited or acquired defects in a wide class of ion channels has been linked to dysfunction at various stages of the conduction system resulting in life-threatening cardiac arrhythmia. Members of the ankyrin and spectrin families have emerged as important nodes for normal expression and regulation of ion channels in myocytes throughout the conduction system. Human variants impacting ankyrin/spectrin function give rise to a broad constellation of cardiac arrhythmias. Furthermore, chronic neurohumoral and biomechanical stress promotes ankyrin/spectrin loss of function that likely contributes to conduction disturbances in the setting of acquired cardiac disease.

Keywords: pacemaking ; arrhythmia ; ankyrin ; spectrin

## 1. The Cardiac Conduction System in Health and Disease

Proper function of the cardiac conduction system depends on coordinated activity of a heterogeneous collection of excitable cells distributed throughout the heart. In turn, normal myocyte excitability depends on orchestrated function of a diverse assortment of ion channels (Figure 1). The myocyte maintains tight spatio-temporal control over ion channel function through organization of ion channel macromolecular complexes involving a host of adapter, cytoskeletal and regulatory molecules [1][2]. While defects in ion channels themselves are closely linked to cardiac conduction disturbances and arrhythmia, there is growing evidence that disruption of associated adapter/cytoskeletal proteins also underlies cardiac conduction system dysfunction.



**Figure 1.** Major ion channels underlying the action potential from different regions of the cardiac conduction system. (A) Schematic of major ion channels and representative action potentials from (A,B) sinoatrial node and (C,D) ventricular myocytes. Channels are color coded to correspond to phase of the AP where they make an important contribution. Abbreviations are as follows: Acetylcholine-activated K<sup>+</sup> current ( $I_{KACh}$ ); Inward rectifier K<sup>+</sup> current ( $I_{K1}$ ); L-type Ca<sup>2+</sup> current ( $I_{Ca}$ ); Na<sup>+</sup>/Ca<sup>2+</sup> exchanger ( $I_{NaCa}$ ); Na<sup>+</sup>/K<sup>+</sup> ATPase ( $I_{NaK}$ ); plateau K<sup>+</sup> current ( $I_{Kp}$ ); rapid ( $I_{Kr}$ ) and slow ( $I_{Ks}$ )

delayed rectifier  $K^+$  currents; sarcoplasmic reticulum (SR); SR  $Ca^{2+}$  release channel (RyR2); SR  $Ca^{2+}$  ATPase (SERCA); transient outward  $K^+$  current ( $I_{to}$ ); TWIK-related  $K^+$  channel 1 current ( $I_{trek}$ ); T-type  $Ca^{2+}$  current ( $I_{Ca,T}$ ); voltage-gated  $Na^+$  current ( $I_{Na}$ ).

### 1.1. Normal Physiology of the Cardiac Conduction System

The normal heartbeat begins with a spontaneous electrical impulse generated by the sinoatrial node (SAN), a specialized crescent-shaped tissue less than an inch in length located in the wall of the right atrium adjacent to the opening of the superior vena cava. From the SAN, activation spreads through the atria and to the atrioventricular node (AVN), which serves as a gateway for electrical activation between the atria and ventricles. The impulse then travels from the AVN to the ventricles through the His-Purkinje system comprised of the AV bundle, right and left bundle branches that run along the interventricular septum and finally Purkinje fibers that allow the action potential to spread rapidly in an organized fashion through the ventricular free walls. Each component of the cardiac conduction system is imbued with unique electrophysiological and structural features to support its specific role. For example, the SAN, unlike the majority of cardiac tissue, does not have a stable resting membrane potential, but instead generates spontaneous firing through cyclical depolarizing events. Changes in the firing rate of the SAN node contribute to changes in heart rate to meet changing physiologic demands of the body. The AVN also displays automaticity but at an intrinsic rate slower than the SAN allowing it to serve in a backup capacity for pacemaking in the event of SAN failure. The AVN supports very slow conduction, which helps insulate the ventricles in the case of rapid, arrhythmic activity in the atria. Purkinje myocytes support extremely rapid conduction and are bundled in insulated fibers to prevent premature or inappropriate ventricular activation. Purkinje-muscle junctions allow for coupling with slow and safe conduction to prevent propagation failure resulting from source-sink mismatch between the small, current generating volume of the Purkinje fiber and very large surrounding ventricular mass.

### 1.2. Cardiac Conduction System Dysfunction and Disease

Generation of the cardiac action potential within these various subregions of the heart and the electrical propagation between them is the result of the coordination of an array of  $Na^+$ ,  $K^+$  and  $Ca^{2+}$  ion channels. Mutations resulting in both gain- and loss-of-function changes in these ion transport proteins are not only sufficient to generate arrhythmic dysfunction in both rhythm and rate of the heart, but have been critical to understanding the function of for each of these players. Importantly, inherited and acquired defects in a host of ion channels have been identified in myocytes throughout the conduction system and are linked to various descriptions of cardiac arrhythmia, resulting in defects in cardiac automaticity and/or conduction [3]. Specific disruptions to heart rate include SAN dysfunction (also referred to as sinus node disease [SND] or sick sinus syndrome [SSS]) where the primary pacemaker is unable to generate appropriate heart rates for the physiologic demands of the body. This includes sinus bradycardia and tachycardia, sinus arrest, sinus-exit block, or chronotropic incompetence, leading to instances of exercise intolerance or syncope [4][5]. Development of such conditions can be the outcome of both genetic causes as well as acquired states, often being induced from electrical and structural remodeling associated with pathologic states such as atrial fibrillation, heart failure, ischemia, or simply aging. Acquired dysfunction may also be the result of drug intoxication from a variety of compounds, including  $\beta$ -blockers,  $Ca^{2+}$  channel blockers and digitalis [6]. Other rate disorders can arise from AV node block, which occurs when there is partial or complete block of impulse propagation through the AVN, resulting in delays between atrial and ventricular tissue in mild cases, or complete uncoupling in the most severe. Alternatively, a condition known as Wolff-Parkinson-White (WPW) syndrome, describes a state of early excitation of the ventricles through an accessory pathway that bypasses the slow conduction of the AV node, shortening the interval between atrial and ventricular excitation, resulting in tachycardia and impairment of proper filling of the ventricular chambers.


In addition to rate disturbances, cardiac conduction system dysfunction can result in abnormal rhythm arising from changes in myocyte excitability and/or intercellular communication that impact the ability of the electrical impulse to propagate through the heart. Disruption to conduction can lead to conduction disease, which includes left and right bundle branch block, causing asynchronous contraction of the ventricles. More severe consequences of conduction disorders are arrhythmias, including atrial flutter/fibrillation and ventricular tachycardia/fibrillation. Such arrhythmias can arise from disruption to normal states of conduction but are also the consequence of genetic or acquired remodeling of the cardiac action potential, including long QT syndrome, short QT syndrome, Brugada syndrome, arrhythmogenic cardiomyopathy, and catecholaminergic polymorphic ventricular tachycardia (CPVT). These conditions can provide pathologic substrates for creating arrhythmias by impacting the dynamics of cellular excitability or providing their own excitation decoupled from normal sinus rhythm. Despite appreciation for the association between disruption of ion channel function/expression and conduction disease, the mechanistic underpinnings, especially in acquired disease, are not fully understood. At the same time, there is growing appreciation that aside from primary defects in ion channels, dysfunction of associated adaptor and

cytoskeletal proteins give rise to arrhythmia. In particular, mounting data support a critical role for ankyrin and spectrin proteins in organizing ion channel macromolecular complexes in myocytes throughout the cardiac conduction system with strong link to inherited and acquired forms of disease. Moreover, these adapter complexes have been identified to regulate a range of ion channel targets, suggesting their potential influence to induce and/or restore proper balance of cardiac electrophysiology targets.

## 2. Ankyrins in the Cardiac Conduction System

Ankyrins are adapter proteins that link membrane proteins (e.g., ion channels and transporters, cell adhesion molecules, and signaling proteins) to the actin/spectrin-based cytoskeleton, ultimately serving to establish stable membrane expression for interacting proteins. The family is comprised of three members encoded by distinct genes: ankyrin-R (encoded by *ANK1*), ankyrin-B (encoded by *ANK2*), and ankyrin-G (encoded by *ANK3*). Ankyrin isoforms are expressed throughout the body, including kidney, lung, brain, skeletal muscle, and heart with ankyrin-R showing the most restricted expression, being found primarily in erythrocytes with secondary expression in muscle and neurons. Ankyrin-B and -G, on the other hand, are ubiquitously expressed, including in myocytes throughout the cardiac conduction system [7][8] and are the focus of our discussion here.


Canonical ankyrins are composed of four domains termed the membrane-binding domain (MBD), the spectrin-binding domain (SBD), the death domain (DD), and the C-terminal domain (CTD) (Figure 2) [9]. The MBD does not interact directly with the cell membrane but instead is primarily the site of interaction with membrane-bound proteins and is defined by a series of stacked  $\alpha$ -helices coupled by  $\beta$ -hairpin loops (24 ANK repeats) [10], which generate specific binding pockets for membrane associated proteins. Importantly, and particularly relevant for its role as a multimodal regulator of cellular excitability, the MBD allows for simultaneous interaction with a number of membrane proteins facilitating formation of large protein complexes. Despite significant sequence homology of the MBD across ankyrin isoforms, isoform-specific protein-protein interactions are achieved at least in part through distinct subcellular localization profiles. In cardiac myocytes for example, ankyrin-B localizes to T-tubules, the sarcoplasmic reticulum (SR), and lateral plasma membrane [11], while ankyrin-G is found primarily at the intercalated disc membrane where neighboring cells are electrically and mechanically coupled [12]. As its name implies, the SBD is the site for interaction with  $\beta$ -spectrin isoforms, linking ankyrin with the actin-based cytoskeleton to stabilize large protein complexes [13]. The last two domains (DD and CTD) are collectively referred to as the regulatory domain for their role in modulating ankyrin interactions. Residues within this domain are important for intermolecular interactions and localization of associated membrane proteins [14][15][16]. Moreover, the majority of disease-causing mutations in ankyrin genes are located within the regulatory domain, supporting the central role of ankyrins as hubs for macromolecular complexes [17]. While the many associated mutations in ankyrin have allowed identification of its interacting partners and functional impacts [8][17], it is critical to realize that acquired states of heart disease associate with reductions in ankyrin expression, suggesting that disruption of each of these interacting partners can be impacted by loss in ankyrin expression [18][19][20][21]. While this provides a strong rationale for understanding the pathologic remodeling of electrical activity in the diseased heart, it also illustrates the profound therapeutic potential that may exist in achieving ways of preserving ankyrin expression. Currently, it is unclear what mechanisms account for the observed loss in ankyrin expression with disease, whether it is a transcriptionally mediated response, or pathways affecting protein stability and/or localization. Undoubtedly, however, unraveling such mechanisms with the intent to preserve ankyrin expression in heart disease will represent a therapeutic approach with significant and wide reaching implications for many of the observed states of electrical disruptions. The following discussion of what these interacting partners are and their consequence on heart rate and rhythm will hopefully illustrate the profound role these cytoskeletal proteins serve in maintaining proper cardiac electrical activity.



**Figure 2.** Structure of canonical ankyrin and major interacting proteins organized by site of interaction for ankyrin-B and ankyrin-G.

## 2.1. Ankyrin-B

Ankyrin-B is expressed within the nodal cells, the cardiac conduction system, and atrial and ventricular myocytes, where it interacts with several channels and transporters important for  $\text{Ca}^{2+}$  handling in cardiomyocytes, including the  $\text{Na}^+/\text{Ca}^{2+}$  exchanger (NCX),  $\text{Na}^+/\text{K}^+$  ATPase (NKA) and inositol-1,4,5-triphosphate receptor ( $\text{IP}_3\text{R}$ ) [29][22] as well as the protein phosphatase, PP2A (Figure 2 and Figure 3) [23][24]. In SAN myocytes, ankyrin-B has also been found to associate with  $\text{Ca}_v1.3$ , a predominantly neuronal L-type  $\text{Ca}^{2+}$  channel that is an important determinant of spontaneous diastolic depolarization phase of the SAN action potential [25]. A human variant in ankyrin-B was first identified in a proband with prolongation of the QT interval on the electrocardiogram together with ventricular arrhythmias and sudden death, leading to the initial link to long QT syndrome (LQT type 4) [26]. It is now appreciated that loss of ankyrin-B function produces a complex phenotype (termed ‘ankyrin-B syndrome’) including SAN, atrial, and ventricular defects [25][27][28]. Relevant to the cardiac conduction system, loss of ankyrin-B leads to aberrant SAN myocyte excitability and pacemaking, displayed by severe bradycardia and rate variability, secondary to defects in  $\text{Ca}^{2+}$  cycling and loss of  $\text{Ca}_v1.3$  membrane targeting [25]. Ankyrin-B dysfunction also induces inappropriate afterdepolarizations, structural remodeling and arrhythmia in the atria with increased susceptibility to atrial fibrillation, ventricular fibrillation, [28][29] and has even been associated with WPW syndrome [30]. More recently, a loss-of-function mutation in ankyrin-B was identified in a patient with severe arrhythmogenic cardiomyopathy and sudden death [31]. The diversity of these phenotypes is likely the result of ankyrin-B mutations disrupting expression in subregions of the heart or subsets of interacting partners, while more complete disruption of ankyrin-B through heterologous knockdown in mice recapitulates a more comprehensive set of phenotypes [11][29][32][33][34].




**Figure 3.** Ankyrin-B/spectrin-based macromolecular complexes for regulation of pacemaking and membrane excitability in cardiac myocytes. Ankyrin-B (AnkB) associates primarily with  $\beta\text{I}$  and  $\alpha\text{I}$  or  $\alpha\text{II}$ -spectrin at the lateral membrane to target the  $\text{Na}^+/\text{Ca}^{2+}$  exchanger (NCX) and  $\text{Na}^+/\text{K}^+$  ATPase and, in sinoatrial node cells only, the voltage-gated  $\text{Ca}^{2+}$  channel  $\text{Ca}_v1.3$ . AnkB also associates with  $\alpha\text{II}$ - and  $\beta\text{II}$ -spectrin to regulate proteins at the cardiac dyad formed at the juxtaposition of the t-tubule (location of voltage-dependent  $\text{Ca}^{2+}$  channel  $\text{Ca}_v1.2$ ) and sarcoplasmic reticulum (SR) membranes. Importantly, this AnkB/spectrin-based complex is important for regulation of  $\text{Ca}^{2+}$  cycling by the protein phosphatase PP2A, in part through modulation of the SR  $\text{Ca}^{2+}$  release channel RyR2.

Beyond transgenic models investigating variants or knockdown, ankyrin-B has also been identified to experience protein loss in human ischemic and non-ischemic heart failure, leading to concomitant loss in binding partners like NKA [19]. The protease inhibitor calpain has been found to preserve ankyrin-B and NKA at the border zone in a mouse model of ischemia-reperfusion and reduce infarct size. Thus, defects in ankyrin-B function give rise to a diverse array of abnormal excitability throughout multiple locales of the cardiac conduction system with important implications for human disease.

## 2.2. Ankyrin-G

Ankyrin-G organizes macromolecular complexes to support conduction of the electrical impulse in neuronal and cardiac tissue, primarily through its association with voltage-gated  $\text{Na}^+$  channels ( $\text{Na}_v$ ) [35][36][37][38]. Specifically, ankyrin-G binds directly to a conserved motif found in DII-DIII linker of several  $\text{Na}_v$   $\alpha$ -subunits, including the predominant cardiac isoform  $\text{Na}_v1.5$ . Ankyrin-G and  $\text{Na}_v1.5$  colocalize primarily at the intercalated disc membrane important for electrical and mechanical communication between neighboring myocytes with secondary expression at t-tubules or lateral membrane (Figure 2 and Figure 4). A host of loss-of-function mutations have been identified in *SCN5A* (encodes  $\text{Na}_v1.5$ ) and linked to a wide range of cardiac conduction disorders, including sick sinus syndrome, cardiac conduction disease, atrial fibrillation, and Brugada syndrome [4][39][40][41][42]. Beyond direct defects in  $\text{Na}_v1.5$ , mutations in a large number of accessory proteins, including  $\beta$ -subunits, adapter, cytoskeletal and regulatory proteins, have been linked to cardiac conduction disorders and arrhythmia [43][44][45]. Relevant to ankyrins, a human mutation in the ankyrin-binding motif in  $\text{Na}_v1.5$  (E1053K) disrupts ankyrin-G/ $\text{Na}_v1.5$  interaction giving rise to loss of  $\text{Na}_v1.5$  membrane targeting and abnormal channel function. Specifically, HA-tagged E1053K  $\text{Na}_v1.5$  expressed in ventricular myocytes show minimal membrane surface expression compared to HA-tagged WT  $\text{Na}_v1.5$ , which express readily at intercalated disc and t-tubule domains. Studies in heterologous cells support that the trafficking defect is likely not due to alterations in  $\text{Na}_v1.5$  folding or stability. Interestingly, the E1053K mutation is associated with Brugada syndrome, characterized by potentially fatal ventricular arrhythmias with specific ECG abnormalities in the absence of structural heart disease [37]. Cardiac-specific deletion of ankyrin-G in mice gives rise to loss of  $\text{Na}_v1.5$  membrane expression and activity with pronounced bradycardia together with evidence of atrial and ventricular conduction slowing at baseline and increased susceptibility to AV block and ventricular arrhythmia following flecainide challenge [12].



**Figure 4.** Ankyrin-G/spectrin-based macromolecular complexes at the cardiomyocyte intercalated disc. Ankyrin-G (AnkG) is found primarily at the cardiomyocyte intercalated disc with  $\alpha_{II}$  and  $\beta_{IV}$ -spectrin in a macromolecular complex for regulation of the voltage-gated  $\text{Na}^+$  channel  $\text{Na}_v1.5$  by  $\text{Ca}^{2+}$ /calmodulin-dependent kinase II (CaMKII). A subpopulation of the protein phosphatase PP2A is also targeted by AnkG/spectrin for modulation of  $\text{Na}_v1.5$  activity (not pictured).  $\beta_{IV}$ -spectrin also associates with the two-pore domain  $\text{K}^+$  channel TREK-1 at the intercalated disc in an AnkG-independent manner.

Beyond direct interaction with  $\text{Na}_v1.5$ , ankyrin-G has been proposed to regulate cardiac conduction through association with mechanical adhesion proteins at the intercalated disc. While electrical coupling is supported by gap junctions comprised of connexin family members (Cx40, Cx45, and/or Cx43 in different layers of the conduction system), mechanical coupling and structural stabilization is provided by adherens junctions and desmosomes. Plakophilin, desmoplakin, and desmin are examples of desmosomal proteins that depend on ankyrin-G for proper expression and localization at the intercalated disc [46], which may explain development of dilated cardiomyopathy in cardiac-specific ankyrin-G knockout mice [12].



### 3. Spectrins in the Cardiac Conduction System

Spectrin is a critical component of the cytoskeleton in a broad distribution of metazoan cell types, including erythrocytes, neurons, beta cells, epithelial cells and cardiac myocytes [47][48]. Spectrin assembles as a heterotetramer of  $\alpha$ - and  $\beta$ -subunits to form an extended chain to provide support for fragile membranes and effectively link membrane proteins/lipids to the actin-based cytoskeleton. Mammals express 2  $\alpha$ - and 5  $\beta$ -subunits with  $\alpha_I$ ,  $\alpha_{II}$ ,  $\beta_I$ ,  $\beta_{II}$  and  $\beta_{IV}$  detected to varying degrees in cardiac myocytes. Canonical  $\alpha$ -spectrin is comprised of 20 triple-helical repeats (spectrin repeats) and a C-terminal calmodulin-related domain (Figure 2).  $\beta$ -spectrin consists of a conserved N-terminal actin-binding region, multiple spectrin repeats (17 for every isoform except for  $\beta_V$ -spectrin that has 30), and a C-terminal domain with multiple signaling motifs. Interaction with ankyrins occurs through a motif in  $\beta$ -spectrin repeat 15 that is highly conserved across isoforms. As discussed for ankyrin, isoform-specific macromolecular complexes are achieved at least in part through distinct subcellular localization profiles of ankyrins and spectrins. There is growing appreciation for the multifunctional nature of spectrin family members beyond providing structural support for the cell membrane. For example, spectrins support long-range cellular communication involving, in part, coordination of signaling nanodomains for ion channels [36][49][50][51][52][53][54][55][56]. More recently, it has been discovered that spectrins modulate gene expression to control remodeling of cell function in response to chronic stress stimuli, although the precise mechanisms remain to be determined [57][58][59]. Thus, spectrins have evolved to serve a host of unique challenges faced by metazoan cells.

Cardiomyocytes express several different spectrin isoforms with distinct subcellular localization profiles, similar to the situation for the different ankyrin isoforms.  $\alpha_{II}$ -spectrin is widely expressed at membrane domains throughout the myocyte, while  $\alpha_I$ -spectrin is mostly restricted to lateral membranes (Figure 3 and Figure 4). In contrast,  $\beta$ -spectrins appear to be more domain specific with  $\beta_I$ -spectrin found mostly at the lateral membrane,  $\beta_{II}$ -spectrin at the z-line in close proximity to sarcoplasmic reticulum (SR) and t-tubule membranes (secondary expression at intercalated disc), and  $\beta_{IV}$ -spectrin almost exclusively expressed at intercalated disc membrane (Figure 4).

Several reports have identified loss of function mutations in *SPTAN1* [60] and *SPTBN4* [61][62][63] in patients displaying a wide spectrum of neurodevelopmental phenotypes, central deafness, and motor dysfunction. While cardiac phenotypes have not been reported in these patients, studies in mice with similar loss of function result in both electrical and structural phenotypes associated with altered ion channel localization and expression (described below). It is possible that the severity of other neurological phenotypes overshadow cardiac complications in these patients. At the same time, a human mutation in *ANK2* has been shown to alter interaction with  $\beta_{II}$ -spectrin resulting in severe cardiac arrhythmia [64], supporting a role for spectrin proteins in human cardiac pathophysiology.

#### 3.1. $\alpha_{II}$ -Spectrin

$\alpha_{II}$ -spectrin (encoded by *SPTAN1*) is expressed in tissues throughout the body, including heart. Within the cardiomyocyte,  $\alpha_{II}$ -spectrin is found at lateral membranes, z-lines and the intercalated disc membrane [65]. Global  $\alpha_{II}$ -spectrin knockout in mice is embryonic lethal with pronounced malformation of multiple tissues including heart [66]. Cardiac-specific  $\alpha_{II}$ -spectrin knockout mice, on the other hand, are viable but show defects in cardiac electrical and mechanical function with evidence of conduction disturbances in the atria and ventricles [67].  $\alpha_{II}$ -spectrin deficient myocytes show a significant decrease in  $I_{Na}$  together with similar decrease in transient outward  $K^+$  current  $I_{to}$ . Interestingly, the cytoskeleton regulatory proteins Mena and VASP have been reported to interact with a specific  $\alpha_{II}$ -spectrin splice variant (SH3i) at z-line and intercalated disc [68]. Double deficiency of Mena and VASP leads to loss of structural integrity of these critical membrane domains and evidence of intra-atrial and intra-ventricular conduction slowing [68].

#### 3.2. $\beta_{II}$ -Spectrin

Similar to  $\alpha_{II}$ -spectrin,  $\beta_{II}$ -spectrin (encoded by *SPTBN1*) is widely distributed in tissues throughout the body and is the predominant  $\beta$ -spectrin isoform found in cardiomyocytes.  $\beta_{II}$ -spectrin associates with a host of critical ion channels, transporters and regulatory molecules through its interaction with ankyrin-B (discussed previously). Global deletion of  $\beta_{II}$ -spectrin in mice is embryonic lethal reflecting its wide distribution and important role in development [69]. Cardiac-specific  $\beta_{II}$ -spectrin knockout mice are viable with pronounced ventricular arrhythmia and even sudden death following acute catecholaminergic challenge [64]. Interestingly and relevant to the discussion of the cardiac conduction system, in addition to ventricular arrhythmias and remodeling,  $\beta_{II}$ -deficient mice show bradycardia at baseline with increased heart rate variability and episodes of AV block. At the myocyte level,  $\beta_{II}$ -spectrin deficiency induces abnormal intracellular  $Ca^{2+}$  homeostasis with increased frequency of inappropriate, spontaneous SR  $Ca^{2+}$  release events and action potential afterdepolarizations that serve as potential arrhythmia triggers. Altered myocyte  $Ca^{2+}$  handling and excitability is likely

related to loss in expression/membrane targeting of ankyrin-B and associated transporters NCX and NKA. Interestingly,  $\beta_{1I}$ -deficient myocytes also showed disorganized RyR2 subcellular localization and decreased RyR2 expression, although the mechanism remains unknown.

### 3.3. $\beta_{IV}$ -Spectrin

$\beta_{IV}$ -spectrin (encoded by *SPTBN4*) is highly expressed in brain and plays a critical role with ankyrin-G in  $\text{Na}_v$  clustering at axon initial segments and nodes of Ranvier [70][71][72]. Thus loss of  $\beta_{IV}$ -spectrin function is associated with defects in neuronal excitability in mice with prominent ataxia. In cardiomyocytes,  $\beta_{IV}$ -spectrin is expressed almost exclusively at the intercalated disc membrane [12][51]. As in neurons,  $\beta_{IV}$ -spectrin associates with ankyrin-G and  $\text{Na}_v$  (primarily  $\text{Na}_v1.5$ ) to orchestrate a platform for regulation of myocyte membrane excitability. While  $\beta_{IV}$ -spectrin is required for proper membrane targeting of  $\text{Na}_v$  in neurons, loss of  $\beta_{IV}$ -spectrin in myocytes does not affect membrane expression but rather disrupts channel regulation. Specifically,  $\beta_{IV}$ -spectrin binds directly to  $\text{Ca}^{2+}$ /calmodulin-dependent kinase II (CaMKII) and targets a kinase subpopulation with ankyrin-G/ $\text{Na}_v1.5$  at the intercalated disc. CaMKII phosphorylates  $\text{Na}_v1.5$  to modulate channel gating and the level of pathogenic so-called late current ( $I_{\text{Na},L}$ ). Loss of  $\beta_{IV}$ -spectrin/CaMKII interaction prevents CaMKII-dependent phosphorylation of  $\text{Na}_v1.5$  at a specific residue in the DI-DII linker (Ser571) and abrogates stress-induced potentiation of  $I_{\text{Na},L}$  [51][73][74]. Mutant mice expressing truncated  $\beta_{IV}$ -spectrin lacking the CaMKII binding domain ( $qv^{3J}$  mice) show normal excitability at baseline with resistance to sinus pause and premature ventricular contractions induced by acute adrenergic challenge.

More recently, it was discovered that  $\beta_{IV}$ -spectrin also regulates the cardiac conduction system through association with the two-pore domain  $\text{K}^+$  channel TREK-1 in cardiac myocytes. TREK-1 function has been well characterized in the nervous system with identified roles in nociception and depression phenotypes [75]. The channel is also expressed in heart across species, including mice and human [53][76][77][78][79]. Interestingly,  $\beta_{IV}$ -spectrin is required for normal TREK-1 membrane expression in SAN and ventricular myocytes. Furthermore, loss of  $\beta_{IV}$ -spectrin/TREK-1 interaction does not impact basal heart rate or conduction but promotes a dramatic increase in sinus pause in response to adrenergic stimulation. Consistent with this phenotype, cardiac-specific knockout of TREK-1 results in R-R prolongation, QTc prolongation, and increased incidence of sinus pause without major changes in baseline echocardiography features [74]. Notably, isolated SAN cells from these mice exhibit rapid spontaneous action potential firing, faster diastolic depolarization rate, depolarized maximum diastolic potential, and longer AP duration at 50% repolarization compared to SAN cells isolated from wildtype (WT) mice. TREK-1 has also been linked to defects in AV node conduction through its association with members of the Popeye domain containing (POPDC) family of cAMP effector proteins [80], implying a broader role for the channel in regulating function of the cardiac conduction system.

---

## References

1. Mohler, P.J.; Wehrens, X.H. Mechanisms of human arrhythmia syndromes: Abnormal cardiac macromolecular interactions. *Physiology* 2007, 22, 342–350.
2. Abriel, H.; Rougier, J.S.; Jalife, J. Ion channel macromolecular complexes in cardiomyocytes: Roles in sudden cardiac death. *Circ. Res.* 2015, 116, 1971–1988.
3. Park, D.S.; Fishman, G.I. The cardiac conduction system. *Circulation* 2011, 123, 904–915.
4. Benson, D.W.; Wang, D.W.; Dymment, M.; Knilans, T.K.; Fish, F.A.; Strieper, M.J.; Rhodes, T.H.; George, A.L., Jr. Congenital sick sinus syndrome caused by recessive mutations in the cardiac sodium channel gene (*SCN5A*). *J. Clin. Invest.* 2003, 112, 1019–1028.
5. Mangrum, J.M.; DiMarco, J.P. The evaluation and management of bradycardia. *N. Engl. J. Med.* 2000, 342, 703–709.
6. Mesirca, P.; Fedorov, V.V.; Hund, T.J.; Torrente, A.G.; Bidaud, I.; Mohler, P.J.; Mangoni, M.E. Pharmacologic approach to sinoatrial node dysfunction. *Annu. Rev. Pharmacol. Toxicol.* 2021, 61, 757–778.
7. Bennett, V.; Chen, L. Ankyrins and cellular targeting of diverse membrane proteins to physiological sites. *Curr. Opin. Cell Biol.* 2001, 13, 61–67.
8. Cunha, S.R.; Mohler, P.J. Cardiac ankyrins: Essential components for development and maintenance of excitable membrane domains in heart. *Cardiovasc. Res.* 2006, 71, 22–29.
9. Hashemi, S.M.; Hund, T.J.; Mohler, P.J. Cardiac ankyrins in health and disease. *J. Mol. Cell. Cardiol.* 2009, 47, 203–209.
10. Michaely, P.; Tomchick, D.R.; Machius, M.; Anderson, R.G. Crystal structure of a 12 ANK repeat stack from human ankyrinR. *EMBO J.* 2002, 21, 6387–6396.

11. Curran, J.; Mohler, P.J. Coordinating electrical activity of the heart: Ankyrin polypeptides in human cardiac disease. *Expert Opin. Ther. Targets*. 2011, 15, 789–801.
12. Makara, M.A.; Curran, J.; Little, S.C.; Musa, H.; Polina, I.; Smith, S.A.; Wright, P.J.; Unudurthi, S.D.; Snyder, J.; Bennett, V.; et al. Ankyrin-G coordinates intercalated disc signaling platform to regulate cardiac excitability in vivo. *Circ. Res.* 2014, 115, 929–938.
13. Bennett, V.; Stenbuck, P.J. Identification and partial purification of ankyrin, the high affinity membrane attachment site for human erythrocyte spectrin. *J. Biol. Chem.* 1979, 254, 2533–2541.
14. Hall, T.G.; Bennett, V. Regulatory domains of erythrocyte ankyrin. *J. Biol. Chem.* 1987, 262, 10537–10545.
15. Davis, L.H.; Davis, J.Q.; Bennett, V. Ankyrin regulation: An alternatively spliced segment of the regulatory domain functions as an intramolecular modulator. *J. Biol. Chem.* 1992, 267, 18966–18972.
16. Mohler, P.J.; Yoon, W.; Bennett, V. Ankyrin-B targets beta2-spectrin to an intracellular compartment in neonatal cardiomyocytes. *J. Biol. Chem.* 2004, 279, 40185–40193.
17. Smith, S.; Curran, J.; Hund, T.J.; Mohler, P.J. Defects in cytoskeletal signaling pathways, arrhythmia, and sudden cardiac death. *Front. Physiol.* 2012, 3, 122.
18. Hund, T.J.; Wright, P.J.; Dun, W.; Snyder, J.S.; Boyden, P.A.; Mohler, P.J. Regulation of the ankyrin-B-based targeting pathway following myocardial infarction. *Cardiovasc. Res.* 2009, 81, 742–749.
19. Kashef, F.; Li, J.; Wright, P.; Snyder, J.; Suliman, F.; Kilic, A.; Higgins, R.S.; Anderson, M.E.; Binkley, P.F.; Hund, T.J.; et al. Ankyrin-B protein in heart failure: Identification of a new component of metazoan cardioprotection. *J. Biol. Chem.* 2012, 287, 30268–30281.
20. Dun, W.; Lowe, J.S.; Wright, P.; Hund, T.J.; Mohler, P.J.; Boyden, P.A. Ankyrin-G participates in INa remodeling in myocytes from the border zones of infarcted canine heart. *PLoS ONE* 2013, 8, e78087.
21. Makara, M.A.; Curran, J.; Lubbers, E.R.; Murphy, N.P.; Little, S.C.; Musa, H.; Smith, S.A.; Unudurthi, S.D.; Rajaram, M.V.S.; Janssen, P.M.L.; et al. Novel mechanistic roles for Ankyrin-G in cardiac remodeling and heart failure. *JACC Basic Transl. Sci.* 2018, 3, 675–689.
22. Sucharski, H.C.; Dudley, E.K.; Keith, C.B.R.; El Refaey, M.; Koenig, S.N.; Mohler, P.J. Mechanisms and alterations of cardiac ion channels leading to disease: Role of Ankyrin-B in cardiac function. *Biomolecules* 2020, 10, 211.
23. Cunha, S.R.; Mohler, P.J. Obscurin targets ankyrin-B and protein phosphatase 2A to the cardiac M-line. *J. Biol. Chem.* 2008, 283, 31968–31980.
24. Bhasin, N.; Cunha, S.R.; Mduannayake, M.; Gigena, M.S.; Rogers, T.B.; Mohler, P.J. Molecular basis for PP2A regulatory subunit B56 alpha targeting in cardiomyocytes. *Am. J. Physiol. Heart Circ. Physiol.* 2007, 293, H109–H119.
25. Le Scouarnec, S.; Bhasin, N.; Vieyres, C.; Hund, T.J.; Cunha, S.R.; Koval, O.; Marionneau, C.; Chen, B.; Wu, Y.; Demolombe, S.; et al. Dysfunction in ankyrin-B-dependent ion channel and transporter targeting causes human sinus node disease. *Proc. Natl. Acad. Sci. USA* 2008, 105, 15617–15622.
26. Mohler, P.J.; Schott, J.J.; Gramolini, A.O.; Dilly, K.W.; Guatimosim, S.; duBell, W.H.; Song, L.S.; Haurogne, K.; Kyndt, F.; Ali, M.E.; et al. Ankyrin-B mutation causes type 4 long-QT cardiac arrhythmia and sudden cardiac death. *Nature* 2003, 421, 634–639.
27. Mohler, P.J.; Le Scouarnec, S.; Denjoy, I.; Lowe, J.S.; Guicheney, P.; Caron, L.; Driskell, I.M.; Schott, J.J.; Norris, K.; Lehenhardt, A.; et al. Defining the cellular phenotype of “ankyrin-B syndrome” variants: Human ANK2 variants associated with clinical phenotypes display a spectrum of activities in cardiomyocytes. *Circulation* 2007, 115, 432–441.
28. Cunha, S.R.; Hund, T.J.; Hashemi, S.; Voigt, N.; Li, N.; Wright, P.; Koval, O.; Li, J.; Gudmundsson, H.; Gumina, R.J.; et al. Defects in ankyrin-based membrane protein targeting pathways underlie atrial fibrillation. *Circulation* 2011, 124, 1212–1222.
29. Wolf, R.M.; Glynn, P.; Hashemi, S.; Zarei, K.; Mitchell, C.C.; Anderson, M.E.; Mohler, P.J.; Hund, T.J. Atrial fibrillation and sinus node dysfunction in human ankyrin-B syndrome: A computational analysis. *Am. J. Physiol. Heart Circ. Physiol.* 2013, 304, H1253–H1266.
30. Coban-Akdemir, Z.H.; Charnig, W.L.; Azamian, M.; Paine, I.S.; Punetha, J.; Grochowski, C.M.; Gambin, T.; Valdes, S.O.; Cannon, B.; Zapata, G.; et al. Wolff-Parkinson-White syndrome: De novo variants and evidence for mutational burden in genes associated with atrial fibrillation. *Am. J. Med. Genet. A* 2020, 182, 1387–1399.
31. Roberts, J.D.; Murphy, N.P.; Hamilton, R.M.; Lubbers, E.R.; James, C.A.; Kline, C.F.; Gollob, M.H.; Krahn, A.D.; Sturm, A.C.; Musa, H.; et al. Ankyrin-B dysfunction predisposes to arrhythmogenic cardiomyopathy and is amenable to therapy. *J. Clin. Invest.* 2019, 129, 3171–3184.



32. Skogestad, J.; Aronsen, J.M.; Tovsrud, N.; Wanichawan, P.; Hougen, K.; Stokke, M.K.; Carlson, C.R.; Sjaastad, I.; Sejersted, O.M.; Swift, F. Coupling of the Na<sup>+</sup>/K<sup>+</sup>-ATPase to Ankyrin B controls Na<sup>+</sup>/Ca<sup>2+</sup> exchanger activity in cardiomyocytes. *Cardiovasc. Res.* 2020, 116, 78–90.
33. Camors, E.; Mohler, P.J.; Bers, D.M.; Despa, S. Ankyrin-B reduction enhances Ca spark-mediated SR Ca release promoting cardiac myocyte arrhythmic activity. *J. Mol. Cell. Cardiol.* 2012, 52, 1240–1248.
34. Chu, L.; Greenstein, J.L.; Winslow, R.L. Na<sup>+</sup> microdomains and sparks: Role in cardiac excitation-contraction coupling and arrhythmias in ankyrin-B deficiency. *J. Mol. Cell. Cardiol.* 2019, 128, 145–157.
35. Zhou, D.; Lambert, S.; Malen, P.L.; Carpenter, S.; Boland, L.M.; Bennett, V. AnkyrinG is required for clustering of voltage-gated Na channels at axon initial segments and for normal action potential firing. *J. Cell. Biol.* 1998, 143, 1295–1304.
36. Jenkins, S.M.; Bennett, V. Ankyrin-G coordinates assembly of the spectrin-based membrane skeleton, voltage-gated sodium channels, and L1 CAMs at Purkinje neuron initial segments. *J. Cell. Biol.* 2001, 155, 739–746.
37. Mohler, P.J.; Rivolta, I.; Napolitano, C.; LeMaillet, G.; Lambert, S.; Priori, S.G.; Bennett, V. Nav1.5 E1053K mutation causing Brugada syndrome blocks binding to ankyrin-G and expression of Nav1.5 on the surface of cardiomyocytes. *Proc. Natl. Acad. Sci. USA* 2004, 101, 17533–17538.
38. Lowe, J.S.; Palygin, O.; Bhasin, N.; Hund, T.J.; Boyden, P.A.; Shibata, E.; Anderson, M.E.; Mohler, P.J. Voltage-gated Nav channel targeting in the heart requires an ankyrin-G dependent cellular pathway. *J. Cell. Biol.* 2008, 180, 173–186.
39. Abriel, H. Roles and regulation of the cardiac sodium channel Na v 1.5: Recent insights from experimental studies. *Cardiovasc. Res.* 2007, 76, 381–389.
40. Kapplinger, J.D.; Tester, D.J.; Alders, M.; Benito, B.; Berthet, M.; Brugada, J.; Brugada, P.; Fressart, V.; Guerchicoff, A.; Harris-Kerr, C.; et al. An international compendium of mutations in the SCN5A-encoded cardiac sodium channel in patients referred for Brugada syndrome genetic testing. *Heart Rhythm* 2010, 7, 33–46.
41. Calloe, K.; Refaat, M.M.; Grubb, S.; Wojciak, J.; Campagna, J.; Thomsen, N.M.; Nussbaum, R.L.; Scheinman, M.M.; Schmitt, N. Characterization and mechanisms of action of novel Nav1.5 channel mutations associated with Brugada syndrome. *Circ. Arrhythm. Electrophysiol.* 2013, 6, 177–184.
42. Ellinor, P.T.; Nam, E.G.; Shea, M.A.; Milan, D.J.; Ruskin, J.N.; MacRae, C.A. Cardiac sodium channel mutation in atrial fibrillation. *Heart Rhythm* 2008, 5, 99–105.
43. Olesen, M.S.; Jespersen, T.; Nielsen, J.B.; Liang, B.; Moller, D.V.; Hedley, P.; Christiansen, M.; Varro, A.; Olesen, S.P.; Haunso, S.; et al. Mutations in sodium channel beta-subunit SCN3B are associated with early-onset lone atrial fibrillation. *Cardiovasc. Res.* 2011, 89, 786–793.
44. Watanabe, H.; Darbar, D.; Kaiser, D.W.; Jiramongkolchai, K.; Chopra, S.; Donahue, B.S.; Kannankeril, P.J.; Roden, D.M. Mutations in sodium channel beta1- and beta2-subunits associated with atrial fibrillation. *Circ. Arrhythm. Electrophysiol.* 2009, 2, 268–275.
45. Rivaud, M.R.; Delmar, M.; Remme, C.A. Heritable arrhythmia syndromes associated with abnormal cardiac sodium channel function: Ionic and non-ionic mechanisms. *Cardiovasc. Res.* 2020, 116, 1557–1570.
46. Cerrone, M.; Delmar, M. Desmosomes and the sodium channel complex: Implications for arrhythmogenic cardiomyopathy and Brugada syndrome. *Trends Cardiovasc. Med.* 2014, 24, 184–190.
47. Bennett, V.; Baines, A.J. Spectrin and ankyrin-based pathways: Metazoan inventions for integrating cells into tissues. *Physiol. Rev.* 2001, 81, 1353–1392.
48. Unudurthi, S.D.; Greer-Short, A.; Patel, N.; Nassal, D.; Hund, T.J. Spectrin-based pathways underlying electrical and mechanical dysfunction in cardiac disease. *Expert Rev. Cardiovasc. Ther.* 2018, 16, 59–65.
49. Yang, Y.; Lacas-Gervais, S.; Morest, D.K.; Solimena, M.; Rasband, M.N. BetaIV spectrins are essential for membrane stability and the molecular organization of nodes of Ranvier. *J. Neurosci.* 2004, 24, 7230–7240.
50. Lacas-Gervais, S.; Guo, J.; Strenzke, N.; Scarfone, E.; Kolpe, M.; Jahkel, M.; De Camilli, P.; Moser, T.; Rasband, M.N.; Solimena, M. BetaIVSigma1 spectrin stabilizes the nodes of Ranvier and axon initial segments. *J. Cell. Biol.* 2004, 166, 983–990.
51. Hund, T.J.; Koval, O.M.; Li, J.; Wright, P.J.; Qian, L.; Snyder, J.S.; Gudmundsson, H.; Kline, C.F.; Davidson, N.P.; Cardona, N.; et al. A betaIV spectrin/CaMKII signaling complex is essential for membrane excitability in mice. *J. Clin. Invest.* 2010, 120, 3508–3519.
52. Bennett, V.; Stenbuck, P.J. The membrane attachment protein for spectrin is associated with band 3 in human erythrocyte membranes. *Nature* 1979, 280, 468–473.
53. Hund, T.J.; Snyder, J.S.; Wu, X.; Glynn, P.; Koval, O.M.; Onal, B.; Leymaster, N.D.; Unudurthi, S.D.; Curran, J.; Camardoglio, C.; et al. betaIV-Spectrin regulates TREK-1 membrane targeting in the heart. *Cardiovasc. Res.* 2014, 102, 166–175.

54. Kline, C.F.; Wright, P.J.; Koval, O.M.; Zmuda, E.J.; Johnson, B.L.; Anderson, M.E.; Hai, T.; Hund, T.J.; Mohler, P.J. betaIV-Spectrin and CaMKII facilitate Kir6.2 regulation in pancreatic beta cells. *Proc. Natl. Acad. Sci. USA* 2013, 110, 17576–17581.
55. Komada, M.; Soriano, P. BetaIV-spectrin regulates sodium channel clustering through ankyrin-G at axon initial segments and nodes of Ranvier. *J. Cell. Biol.* 2002, 156, 337–348.
56. Sikorski, A.F.; Sangerman, J.; Goodman, S.R.; Critz, S.D. Spectrin (betaSp1sigma1) is an essential component of synaptic transmission. *Brain Res.* 2000, 852, 161–166.
57. Patel, N.J.; Nassal, D.M.; Greer-Short, A.D.; Unudurthi, S.D.; Scandling, B.W.; Gratz, D.; Xu, X.; Kalyanasundaram, A.; Fedorov, V.V.; Accornero, F.; et al. betaIV-Spectrin/STAT3 complex regulates fibroblast phenotype, fibrosis, and cardiac function. *JCI Insight* 2019, 4, e131046.
58. Unudurthi, S.D.; Nassal, D.; Greer-Short, A.; Patel, N.; Howard, T.; Xu, X.; Onal, B.; Satroplus, T.; Hong, D.; Lane, C.; et al. betaIV-Spectrin regulates STAT3 targeting to tune cardiac response to pressure overload. *J. Clin. Investig.* 2018, 128, 5561–5572.
59. Lambert, M.W. Nuclear alpha spectrin: Critical roles in DNA interstrand cross-link repair and genomic stability. *Exp. Biol. Med.* 2016, 241, 1621–1638.
60. Syrbe, S.; Harms, F.L.; Parrini, E.; Montomoli, M.; Mutze, U.; Helbig, K.L.; Polster, T.; Albrecht, B.; Bernbeck, U.; van Binsbergen, E.; et al. Delineating SPTAN1 associated phenotypes: From isolated epilepsy to encephalopathy with progressive brain atrophy. *Brain* 2017, 140, 2322–2336.
61. Knierim, E.; Gill, E.; Seifert, F.; Morales-Gonzalez, S.; Unudurthi, S.D.; Hund, T.J.; Stenzel, W.; Schuelke, M. A recessive mutation in beta-IV-spectrin (SPTBN4) associates with congenital myopathy, neuropathy, and central deafness. *Hum. Genet.* 2017, 136, 903–910.
62. Wang, C.C.; Ortiz-Gonzalez, X.R.; Yum, S.W.; Gill, S.M.; White, A.; Kelter, E.; Seaver, L.H.; Lee, S.; Wiley, G.; Kelter, E.; et al. bIV spectrinopathies cause profound intellectual disability, congenital hypotonia, and motor axonal neuropathy. *Am. J. Hum. Genet.* 2018, 102, 1158–1168.
63. Hausler, M.G.; Begemann, M.; Lidov, H.G.; Kurth, I.; Darras, B.T.; Elbracht, M. A novel homozygous splice-site mutation in the SPTBN4 gene causes axonal neuropathy without intellectual disability. *Eur. J. Med. Genet.* 2020, 63, 103826.
64. Smith, S.A.; Sturm, A.C.; Curran, J.; Kline, C.F.; Little, S.C.; Bonilla, I.M.; Long, V.P.; Makara, M.; Polina, I.; Hughes, L.D.; et al. Dysfunction in the betaII spectrin-dependent cytoskeleton underlies human arrhythmia. *Circulation* 2015, 131, 695–708.
65. Baines, A.J.; Pinder, J.C. The spectrin-associated cytoskeleton in mammalian heart. *Front. Biosci.* 2005, 10, 3020–3033.
66. Stankewich, M.C.; Cianci, C.D.; Stabach, P.R.; Ji, L.; Nath, A.; Morrow, J.S. Cell organization, growth, and neural and cardiac development require alphaII-spectrin. *J. Cell. Sci.* 2011, 124, 3956–3966.
67. Lubbers, E.R.; Murphy, N.P.; Musa, H.; Huang, C.Y.; Gupta, R.; Price, M.V.; Han, M.; Daoud, E.; Gratz, D.; El Refaey, M.; et al. Defining new mechanistic roles for alphaII spectrin in cardiac function. *J. Biol. Chem.* 2019, 294, 9576–9591.
68. Benz, P.M.; Merkel, C.J.; Offner, K.; Abesser, M.; Ullrich, M.; Fischer, T.; Bayer, B.; Wagner, H.; Gambaryan, S.; Ursitti, J.A.; et al. Mena/VASP and alphaII-Spectrin complexes regulate cytoplasmic actin networks in cardiomyocytes and protect from conduction abnormalities and dilated cardiomyopathy. *Cell. Commun. Signal.* 2013, 11, 56.
69. Tang, Y.; Katuri, V.; Dillner, A.; Mishra, B.; Deng, C.X.; Mishra, L. Disruption of transforming growth factor-beta signaling in ELF beta-spectrin-deficient mice. *Science* 2003, 299, 574–577.
70. Berghs, S.; Aggujaro, D.; Dirkx, R., Jr.; Maksimova, E.; Stabach, P.; Hermel, J.M.; Zhang, J.P.; Philbrick, W.; Slepnev, V.; Ort, T.; et al. betaIV spectrin, a new spectrin localized at axon initial segments and nodes of Ranvier in the central and peripheral nervous system. *J. Cell. Biol.* 2000, 151, 985–1002.
71. Liu, C.H.; Stevens, S.R.; Teliska, L.H.; Stankewich, M.; Mohler, P.J.; Hund, T.J.; Rasband, M.N. Nodal beta spectrins are required to maintain Na(+) channel clustering and axon integrity. *eLife* 2020, 9, e52378.
72. Rasband, M.N. The axon initial segment and the maintenance of neuronal polarity. *Nat. Rev. Neurosci.* 2010, 11, 552–562.
73. Glynn, P.; Musa, H.; Wu, X.; Unudurthi, S.D.; Little, S.; Qian, L.; Wright, P.J.; Radwanski, P.B.; Gyorke, S.; Mohler, P.J.; et al. Voltage-gated sodium channel phosphorylation at Ser571 regulates late current, arrhythmia, and cardiac function in vivo. *Circulation* 2015, 132, 567–577.
74. Greer-Short, A.; Musa, H.; Alsina, K.M.; Ni, L.; Word, T.A.; Reynolds, J.O.; Gratz, D.; Lane, C.; El-Refay, M.; Unudurthi, S.; et al. Calmodulin kinase II regulates atrial myocyte late sodium current, calcium handling, and atrial arrhythmia. *H*

75. Honore, E. The neuronal background K2P channels: Focus on TREK1. *Nat. Rev. Neurosci.* 2007, 8, 251–261.
76. Unudurthi, S.D.; Wu, X.; Qian, L.; Amari, F.; Onal, B.; Li, N.; Makara, M.A.; Smith, S.A.; Snyder, J.; Fedorov, V.V.; et al. Two-pore K<sup>+</sup> channel TREK-1 regulates sinoatrial node membrane excitability. *J. Am. Heart Assoc.* 2016, 5, e002865.
77. Lugenbiel, P.; Wenz, F.; Syren, P.; Geschwill, P.; Govorov, K.; Seyler, C.; Frank, D.; Schweizer, P.A.; Franke, J.; Weis, T.; et al. TREK-1 (K2P2.1) K<sup>+</sup> channels are suppressed in patients with atrial fibrillation and heart failure and provide therapeutic targets for rhythm control. *Basic Res. Cardiol.* 2017, 112, 8.
78. Decher, N.; Ortiz-Bonnin, B.; Friedrich, C.; Schewe, M.; Kiper, A.K.; Rinne, S.; Seemann, G.; Peyronnet, R.; Zumhagen, S.; Bustos, D.; et al. Sodium permeable and "hypersensitive" TREK-1 channels cause ventricular tachycardia. *EMBO Mol. Med.* 2017, 9, 403–414.
79. Schmidt, C.; Wiedmann, F.; Kallenberger, S.M.; Ratte, A.; Schulte, J.S.; Scholz, B.; Muller, F.U.; Voigt, N.; Zafeiriou, M. P.; Ehrlich, J.R.; et al. Stretch-activated two-pore-domain (K2P) potassium channels in the heart: Focus on atrial fibrillation and heart failure. *Prog. Biophys. Mol. Biol.* 2017, 130, 233–243.
80. Rinne, S.; Ortiz-Bonnin, B.; Stallmeyer, B.; Kiper, A.K.; Fortmuller, L.; Schindler, R.F.R.; Herbort-Brand, U.; Kabir, N.S.; Dittmann, S.; Friedrich, C.; et al. POPDC2 a novel susceptibility gene for conduction disorders. *J. Mol. Cell. Cardiol.* 2020, 145, 74–83.