

Lavender Essential Oil

Subjects: **Plant Sciences**

Contributor: Inga Zinicovscaia

The producers of essential oils from the Republic of Moldova care about the quality of their products and at the same time, try to capitalize on the waste from processing. The purpose of the present study was to analyze the chemical composition of lavender (*Lavanda angustifolia* L.) essential oil and some by-products derived from its production (residual water, residual herbs), as well as to assess their “in vitro” antimicrobial activity. The gas chromatography-mass spectrometry analysis of essential oils produced by seven industrial manufacturers led to the identification of 41 constituents that meant 96.80–99.79% of the total. The main constituents are monoterpenes (84.08–92.55%), followed by sesquiterpenes (3.30–13.45%), and some aliphatic compounds (1.42–3.90%). The high-performance liquid chromatography analysis allowed the quantification of known triterpenes, ursolic, and oleanolic acids, in freshly dried lavender plants and in the residual by-products after hydrodistillation of the essential oil. The lavender essential oil showed good antibacterial activity against *Bacillus subtilis*, *Pseudomonas fluorescens*, *Xanthomonas campestris*, *Erwinia carotovora* at 300 µg/mL concentration, and *Erwinia amylovora*, *Candida utilis* at 150 µg/mL concentration, respectively.

Lavandula angustifolia L.

essential oil

by-products

terpenic compounds

chromatographic analyses

antimicrobial activity

statistical data analysis

1. Introduction

Lavandula angustifolia Mill. (syn. *Lavandula vera* DC, syn. *Lavandula officinalis* Chaix ex Vill., syn. *Lavandula spica* L.) is a perennial evergreen shrub of the family *Lamiaceae*, native to the Mediterranean region. Nowadays, this species is naturalized almost all over Europe, North Africa, United States, and Australia [1]. *L.angustifolia* (Lavander) is one of the most valuable medicinal and aromatic plants traditionally used to treat pain, parasitic infections, burns, insect bites, cramps, and muscle spasms [2]. In addition to its application in herbal treatment, lavender is also cultivated for the essential oils used in aromatherapy and the cosmetic, food, and flavour industries [3][4][5].

This is possible due to the presence of a set of biologically active substances, especially in essential oil, which possesses a multidirectional therapeutic activity being used in the treatment of gastrointestinal, cardiovascular, respiratory, and urinary infections [6]. Scientific studies reported anti-inflammatory [7], antioxidant [8][9], sedative [10], cytotoxic [11][12], analgesic [7], antimicrobial [6][13][14], and anticonvulsive [15] properties of *L. angustifolia* essential oil. Literature data reveal a huge variation in terms of *L. angustifolia* essential oil content, with values ranging between 0.5 and 6.25% in the case of essential oil obtained from fresh and dry inflorescences [16]. The main constituents

of *L. angustifolia* essential oil are linalool, linalyl acetate, 1,8-cineole, borneol, camphor, lavandulyl acetate, β -caryophyllene, β -ocimene, α -fenchone, terpinen-4-ol, caryophyllene oxide, limonene, pinenes, geranyl acetate, β -farnesene, santalene, lavandulol, camphene, geraniol, and α -terpineol [8][11][13][14][17][18][19][20][21][22][23][24][25][26]. The content of oxygenated monoterpenes prevails in *L. angustifolia* essential oil and varies between 36.33 and 92.90% [16].

The therapeutic effects of *L. angustifolia* are also determined by secondary metabolites such as oleanolic and ursolic acids, together with other pentacyclic triterpenes. [27][28]. It has been proven experimentally that both compounds in pure forms, as well as their synthetic derivatives, show multiple biological activities [29][30][31][32][33][34][35][36][37][38].

Some by-products, e.g., pomace or solid residues, that resulted after hydrodistillation of essential oil-producing plants could be considered as a source of biologically active compounds such as ursolic and oleanolic acids. In addition, residual distillation waters have various applications due to their aromatic and antimicrobial properties [39][40][41][42][43].

Antibiotic resistance is becoming one of the main problems of modern medicine since it substantially reduces the effectiveness of antibacterial treatments and is linked to increased patient mortality. As a result, known antibacterial preparations cease to be safe and effective against infections caused by resistant bacteria, leading to increasingly serious cases, including hospital-acquired complications. This requires the discovery of new classes of antibiotics or optimization and a combination of known compounds. However, microorganisms will likely evolve resistance in time and further research and development may be hard to sustain by the pharmaceutical companies. For this reason, studies are being conducted to identify effective remedies against multidrug-resistant strains. Preference is given to natural products among which are the essential oils [44], including lavender [45], or their combination with antibiotics [46]. Still, information about the antimicrobial activity of residual water and ethanolic extracts is very scanty and is mainly related to Lavender hydrosol, which is produced synthetically [47].

2. Discussion

2.1. Chemical Composition of Lavender Essential Oils

The essential oil manufactured by producer P1, destined for export, had the following physico-chemical properties: Density (20 °C)—0.8920 g/mL; refractive index (n_D^{20})—1.4660, and optical rotation (α_D^{20})— -7.0° .

The most multitudinous group of terpenic compounds are monoterpenes, which include C_{10} -hydrocarbones (8.72–15.32%) and their oxygenated derivatives (69.0–83.83%). The main constituents of this group which determine the quality and genuineness of lavender essential oil, according to the International Standard [48], are (%): 1,8-cineol (eucalyptol) (<1.0), (*E*)-ocimene (4.0–10.0), (*Z*)-ocimene (1.5–6.0), linalool (25.0–38.0), camphor (<0.5), terpin-1-en-4-ol (2.0–6.0), α -terpineol (<1.0), linalyl acetate (25.0–45.0), and lavandulyl acetate (>2.0) (Table 1 and Table 2).

Table 1. Phytochemical (terpenic and aliphatic compounds) composition of lavender essential oil of Moldovan origin.

No.	RT* (min)	Component	Producer, Content (%)						
			P1	P2	P3	P4	P5	P6	P7
1	4.416	α -Pinene	0.36	0.57	0.36	0.18	0.09	0.26	0.57
2	4.710	Camphene	0.34	0.47	0.30	0.09	0.09	0.26	0.63
3	5.179	Sabinene	0.14	-	-	-	-	-	-
4	5.240	1-Octen-3-ol		0.83	-	-	-	0.15	0.27
5	5.263	β -Pinene	0.51	-	0.34	0.39	0.38	0.22	0.57
6	5.398	Octan-3-one	0.28	0.51	0.31	0.25	0.39	0.12	0.21
7	5.489	β -Myrcene	1.06	1.50	0.96	0.80	0.89	0.62	0.89
8	5.577	Octan-3-ol	0.13	0.18	0.20	0.17	0.20	-	-
9	5.962	<i>n</i> -Hexyl acetate	0.59	1.27	0.31	0.42	0.55	0.52	1.17
10	6.284	<i>p</i> -Cymene	0.22	0.46	0.22	0.14	0.10	0.24	0.39
11	6.400	Limonene	0.52	1.79	0.79	0.45	0.55	1.17	1.93
12	6.455	1,8-Cineol (eucalyptol)	5.00	3.81	2.22	3.73	4.44	3.83	9.29
13	6.574	(<i>E</i>)-Ocimene	8.06	5.87	6.85	7.86	4.37	5.25	7.15
14	6.807	(<i>Z</i>)-Ocimene	3.74	3.45	2.59	2.55	1.86	1.76	2.16
15	7.087	γ -Terpinene	0.10	0.29	0.29	0.06	0.07	0.07	0.15
16	7.443	Linalool oxide	-	0.17	0.07	0.11	0.12	-	-
17	7.824	δ -Terpinene	0.27	-	-	-	-	-	-
18	7.825	α -Terpinolene	-	0.58	0.25	0.21	0.32	0.16	0.23
19	8.238	Linalool	23.54	27.98	29.06	25.57	40.68	33.29	26.19
20	8.392	Oct-1-en-3-yl acetate	0.56	0.82	0.60	0.71	0.63	0.39	0.58
21	9.308	Camphor	0.47	0.47	0.37	0.36	0.30	0.32	0.61
22	9.847	Borneol	1.92	2.15	1.68	1.28	1.65	1.41	2.40
23	10.00	(3 <i>E</i> ,5 <i>Z</i>)-Undeca-1,3,5-triene	0.17	-	-	-	-	-	-

No.	RT* (min)	Component	Producer, Content (%)						
			P1	P2	P3	P4	P5	P6	P7
24	10.15	Terpin-1-en-4-ol	1.30	4.65	5.98	0.94	1.67	1.41	1.03
25	10.41	Cryptone	0.29	0.29	-	-	-	0.30	0.33
26	10.50	α -Terpineol	2.42	3.31	2.02	1.42	7.95	1.49	1.61
27	11.49	Nerol	0.38	0.46	0.23	0.14	1.14	-	-
28	11.84	<i>p</i> -Cumic aldehyde	0.13	0.15	-	-	-	0.18	-
29	12.32	Linalyl acetate	26.55	20.26	28.65	32.25	16.68	33.30	28.10
30	13.01	Bornyl acetate	0.32	0.25	0.24	0.27	0.19	0.17	0.24
31	13.11	Lavandulyl acetate	4.88	2.84	2.36	4.83	4.78	2.56	3.07
32	14.98	Neryl acetate	0.78	0.91	0.39	0.33	1.53	0.31	0.37
33	15.47	Geranyl acetate	1.31	1.69	0.79	0.73	2.70	0.59	0.67
34	15.66	α -Zingiberene	0.15	-	-	- [48]	-	-	-
35	16.47	β -Caryophyllene	6.25	5.33	4.62	5.44	1.64	4.93	4.32
36	16.80	α -Bergamotene	0.27	0.28	0.19	0.20 [48]	0.05	0.16	-
37	17.31	(<i>E</i>)- β -Farnesene	4.86	2.59	3.65	3.94	1.23	2.46	2.45
38	17.97	β -Cubebene	1.12	0.82	1.03	0.69	0.17	-	-
39	18.75	γ -Cadinene	0.18	0.53	-	-	-	0.68	0.65
40	20.39	Caryophyllene oxide	0.45	0.69	0.19	0.29	0.21	0.35	0.28
41	21.69	Cadinol	0.17	0.57	-	-	-	0.18	-
Total content, (%)			99.80	98.79	98.17	96.80	97.62	99.11	98.51

Table 1. The content of components of lavender plant material (LPM).

Lavender Plant Material	Extract Yield (%)	Concentration (mg/g Extract)		Concentration (mg/100 g Lavender Plant Material, DW)	
		OA	UA	OA	UA
LPM 1	9.94	16.19	37.46	160.95	372.36
LPM 2	8.83	19.09	60.82	168.57	537.00
LPM 3	9.91	13.43	33.28	133.11	329.83
Monoterpenes					84.08–92.55
Monoterpene hydrocarbons					8.72–15.32

(1.92%), and α -terpineol (10.04%).

Class		Subclass		Content, (%)		A content and UA in their amount of OA
		Oxygenated monoterpenes		69.00–83.83		
		Sesquiterpenes		3.30–13.45		
		Sesquiterpene hydrocarbons		3.09–12.83		
		Oxygenated sesquiterpenes		0.19–1.26		
Aliphatic compounds				1.42–3.90		259.7 and
		Hydrocarbons		0.17		
Lavender by-Product (Solid Residue, SR)	Extract Yield (%)	Concentration (mg/g Extract)		Concentration (mg/100 g Dry Solid Residue)		
		OA	UA	OA	UA	
SR 1 [49]	3.88	29.21	80.82	113.47	313.95	
SR 2	3.68	39.37	135.56	144.98	499.15	
SR 3	4.15	27.48	87.90	114.07	364.89	

Phytopathogenic bacteria can cause various diseases of agricultural plants, especially the genera *Erwinia* and *Xanthomonas*. For example, *Erwinia amylovora*, the Gram-negative bacterium of the Enterobacteriaceae family, is the causative agent of fire blight, a devastating plant disease that affects a wide range of species of the family Rosaceae and is a major global threat to commercial apple and pear production. [\[50\]](#). Another species, *E. carotovora*, causes bacterial soft rot in economically important crops, such as potatoes, tomatoes, and cucumbers. In the case of potatoes, the soft rot of the stem and tubers occurs even after harvest, thus considerably reducing the yield [\[51\]](#). *Xanthomonas campestris* pv. *vesicatoria* is a biotrophic Gram-negative bacterium and is the agent that causes bacterial leaf scorch on tomatoes (*Solanum lycopersicum* L.) and peppers (*Capsicum annuum*), a disease that is present worldwide. Symptoms of bacterial infection include defoliation and chlorotic necrotic lesions on leaves, stems, fruits, and flowers, which subsequently lead to reduced fruit yield [\[52\]](#).

The species *Bacillus subtilis* and *Pseudomonas fluorescens* do not cause any disease to plants but were selected as reference bacteria from the Gram-positive and Gram-negative groups. They are also very suitable as test objects for evaluating the antibacterial activity of the lavender extract. *Candida utilis* and *Saccharomyces cerevisiae* are also non-pathogenic but were used as representatives of the yeast-fungus group for evaluating the antifungal activity of the extract.

It should be mentioned that there is a lack of information about any antimicrobial effects of lavender essential oil on *E. carotovora*, *E. amylovora*, and *C. utilis*.

The in vitro assessment of lavender essential oil of Moldovan origin showed good antibacterial activity against both non-pathogenic Gram-positive/Gram-negative bacteria (*B. subtilis* and *P. fluorescens*) at MBC of 300 µg/mL and good to high antifungal activity against phytopathogenic bacteria (*X. campestris*, *E. amylovora*, *E. carotovora*) and *C. utilis* fungi at MFC of 150–300 µg/mL (**Table 5**).

Table 5. The antimicrobial activity of lavender essential oil.

MBC and MFC, µg/mL						
Sample	<i>Bacillus subtilis</i>	<i>Pseudomonas fluorescens</i>	<i>Xantdomonas campestris</i>	<i>Erwinia amylovora</i>	<i>Erwinia carotovora</i>	<i>Candida utilis</i>
LEO	300	300	300	150	300	150

MBC: Minimal bactericidal concentration; MFC: Minimal fungicidal concentration. The two techniques employed for determining MBC and MFC are diffusion and the dilution methods have been developed to yield accurate measurements of antibacterial and antifungal activities and are routinely used in antimicrobial susceptibility testing.

According to the obtained results, the antibacterial activity was similar but the antifungal activity was slightly different, thus suggesting that the activity against different microorganisms could be caused by different components of the oil.

2.5. Statistical Data Analysis

Univariate as well as multivariate statistical data analysis (SDA) represent one of the most reliable methods that permit extracting useful information and inferring different hypotheses concerning the considered set of data. Given the great diversity of organic compounds which can be found in lavender essential oil, multivariate statistical data analysis was an appropriate method allowing to group samples, in this case, according to the lavender oil producer and based on the concentrations of organic compounds (R mode), or, to classify an experimentally determined organic compound based on the concentration in samples (Q mode) [53][54].

It is worth mentioning that, to avoid any errors induced by missing data, SDA was applied only in the cases of compounds with a non-negligible variation present in all the samples (Table 1), i.e., the compounds which permitted generating the box plots in Figure 1a,b.




Figure 1. Box plots representing the distribution of (a) 25 components of lavender oil and (b) total content of compounds present in all the samples (producers).

Univariate SDA was useful in establishing the extent to which the samples of lavender oil by the seven producers were similar. This information was obtained by analyzing the box plot shown in **Figure 1a**. It was observed that all the samples were quite similar. To confirm this, we used more univariate tests, such as one-way ANOVA, Tukey's pairwise test, Kruskal-Wallis test of equal medians, as well as Mann-Whitney U tests. All of them confirmed that between the lavender oil samples there are no statistically significant differences. For this reason, we have proceeded with multivariate SDA.

Within multivariate SDA, each sample (case) is characterized by independent parameters (variables), so that the final analysis can be performed in R mode (to study relations between samples based on variables) or Q mode (to study the interrelations between variables based on samples). As both methods were based on the same set of samples and variables, R and Q modes could be considered complementary, which significantly enhanced the analysis.

Depending on the situation, cases/variables can be grouped by a multitude of procedures among which covariance and correlation are frequently utilized.

In the case of lavender samples, the best results were obtained by the principal component analysis (PCA) applied in both R and Q modes. With respect to the other two SDA methods, cluster analysis and K mean clustering, PCA permitted evidencing the association of samples, i.e., seven producers of lavender oil in R mode, as well as 25 lavender oil compounds in Q mode. Moreover, in R mode, a tree diagram corresponding to the cluster analysis (Euclidean distances) is, concerning the number and structure of clusters, similar to PCA based on correlation. For this reason, we restrained our SDA to both R and Q mode PCA.

The results, represented by the principal component (PC) 2 vs. PC 1 bi-plots, are illustrated in **Figure 2a,b**, respectively. In both cases, the PCA was based on correlations between variables (organic compounds, R mode) or samples (lavender oil producer, Q mode). Moreover, the loadings of each variable or sample were represented by Factor 2 vs. Factor 1 bi-plots in the corresponding insets: Variables in **Figure 2a** and samples in **Figure 2b**.




Figure 2. The results of R (a) and Q (b) mode PCA. The insets illustrate the contribution of the corresponding principal component (PC) analysis.

Accordingly, the result of PCA in R mode is illustrated by the bi-plot in **Figure 1a**. The existence of at least three clusters can be remarked, two of which consist of only one member, i.e., producers P2 and P5, and a third one, grouping the rest of the producers. The bi-plot illustrating the contribution of each compound to the PC1 and PC2 showed a relatively balanced situation, as both Factors 1 and 2 had similar contributions to PC, consisting of 36.46 and 27.25%, respectively. It is worth mentioning that a similar result was obtained by considering the PC3 vs. PC2, which most probably could be explained by their contribution to the total variance, 25.25 and 17.57%, respectively. The corresponding screen plot in **Figure 3a** illustrated this finding.




Figure 3. The screen plots corresponding to R-mode (a) and Q-mode (b) PCA.

Complementary to the R-mode, a Q mode PC2 vs. PC1 bi-plot, shown in **Figure 2b**, consisted of three clusters, two of which contained a single organic compound, i.e., linalyl acetate and linalool, while the third one included all other 23 compounds. This result was in good agreement with the composition of the investigated samples, according to which, both linalyl acetate and linalool were characterized by the highest concentrations and variances.

On the contrary, Factor 2 vs. Factor 1 (**Figure 2b**, inset), except for Producers 2 (P2) and 5 (P5), were nearly coincident and negatively oriented along the first axis, which suggested an almost equivalent contribution to the total variance. This finding may explain the fact that PC1 contributed about 96% to the total variance, as shown in the corresponding screen plot (**Figure 3b**). In this regard, it is of interest to remark, as mentioned before, that P2 and P5 formed two different uni-component clusters (**Figure 2b**).

References

1. Basch, E.; Foppa, I.; Liebowitz, R.; Nelson, J.; Smith, M.; Sollars, D.; Ulbricht, C. Monograph from National Standard: Lavender (*Lavandula angustifolia* Miller). J. Herb. Pharmacother. 2004, 4, 63–78.

2. Denner, S.S. *Lavandula angustifolia* Miller. *Holist. Nurs. Pract.* 2009, 23, 57–64.
3. Da Porto, C.; Decorti, D.; Kikic, I. Flavour compounds of *Lavandula angustifolia* Mill. To use in food manufacturing. Comparison of three different extraction methods. *Food Chem.* 2009, 112, 1072–1078.
4. Erland, L.A.E.; Mahmoud, S.S. Lavender (*Lavandula angustifolia*) Oils. In *Essential Oils in Food Preservation, Flavor and Safety*; Preedy, V.E., Ed.; Academic Press: Amsterdam, The Netherlands; pp. 501–507.
5. Smigielski, K.; Sikora, M.; Majewska, M.; Raj, A. The application of essentials oils to natural and organic cosmetics. *Pol. J. Cosmet.* 2008, 11, 89–107.
6. Prusinowska, R.; Smigielski, K. Composition, biological activity and therapeutic effects of lavender (*Lavandula angustifolia* L.): A review. *Herba Pol.* 2014, 60, 56–66.
7. Hajhashemi, V.; Ghannadi, A.; Sharif, B. Anti-inflammatory and analgesic properties of the leaf extracts and essential oil of *Lavandula angustifolia* Mill. *J. Ethnopharm.* 2003, 89, 67–81.
8. Hamad, K.J.; Al-Shaheen, S.J.A.; Kaskoos, R.A.; Ahanad, J.; Jameel, M.; Mir, S.R. Essential oil composition and antioxidant activity of *Lavandula angustifolia* from Irak. *Int. Res. J. Pharm.* 2013, 4, 117–120.
9. Djenane, D.; Aider, M.; Yanguela, J.; Idir, L.; Gomez, D.; Roncales, P. Antioxidant and antibacterial effects of *Lavandula* and *Mentha* essential oils in minced beef inoculated with *E. coli* O157:H7 and *S. aureus* during storage at abuse refrigeration temperature. *Meat Sci.* 2012, 92, 667–674.
10. Adaszynska-Skwirzynska, M.; Swarcewicz, M.; Dobrowolska, A. The potential of use lavender from vegetable waste as effective antibacterial and sedative agents. *Med. Chem.* 2014, 4, 734–737.
11. Mantovani, A.L.L.; Vieira, G.P.G.; Cunha, W.R.; Groppo, M.; Santos, R.A.; Rodrigues, V.; Magalhaes, L.G.; Crotti, A.E.M. Chemical composition, antischistosomal and cytotoxic effects of the essential oil of *Lavandula angustifolia* grown in South-eastern Brazil. *Rev. Braz. Farmacogn.* 2013, 23, 877–884.
12. Shou-Dong, S.; Chang-Xu, C.; Ji-Shu, Q.; Ming-Hua, S. Study on antitumor effect of Lavender *angustifolia* extract. *Food Sci. Technol.* 2009, 2, 213–215.
13. Martucci, J.F.; Gende, L.B.; Neira, L.M.; Ruseckaite, R.A. Oregano and lavender essential oils as antioxidant and antimicrobial additives of biogenic gelatine films. *Ind. Crop. Prod.* 2015, 71, 205–213.
14. Tarek, N.; Hassan, H.M.; AbdelGhani, S.M.M.; Radwan, I.A.; Hammouda, O.; El-Gendy, A.O. Comparative chemical and antimicrobial study of nine essential oils obtained from medicinal plants growing in Egypt. *Beni-Seuf Univ. J. Appl. Sci.* 2014, 3, 149–156.

15. Yamada, K.; Mimaki, Y.; Sashida, Y. Anticonvulsive effects of inhaling lavender oil vapour. *Biol. Pharm. Bull.* 1994, 17, 359–360.
16. Aprotosoiaie, A.C.; Gille, E.; Trifan, A.; Luca, V.S.; Miron, A. Essential oil of *Lavandula* genus: A systematic review of their chemistry. *Phytochem. Rev.* 2017, 16, 761–799.
17. Smigielski, K.; Raj, A.; Krosowiak, K.; Grusca, R. Chemical composition of the essential oil of *Lavandula angustifolia* cultivated in Poland. *J. Essent. Oil-Bear. Plants* 2009, 12, 338–347.
18. Hassanpouraghdam, M.B.; Hassani, A.; Vojodi, L.; Hajisamadi, A.B.; Rostami, A. Essential oil constituents of *Lavandula angustifolia* Chaix from Northwest Iran. *Chemija* 2011, 22, 167–171.
19. Belhadj, M.M.; Kabouche, A.; Abaza, I.; Aburjai, T.; Tauzani, R.; Kabouche, Z. Chemotypes investigation of *Lavandula* essential oils growing at different North African soils. *J. Mater. Environ. Sci.* 2014, 5, 1896–1901.
20. Cong, Y.; Abulizi, P.; Zhi, L.; Wang, X. Chemical composition of the essential oil of *Lavandula angustifolia* from Xinjiang, China. *Chem. Nat. Compd.* 2008, 44, 810.
21. Renaud, E.N.C.; Denys, C.J.; Simon, J.E. Essential oil quantity and composition from 10 cultivars of organically grown lavender and lavandin. *J. Essent. Oil Res.* 2001, 13, 269–273.
22. Jianu, C.; Pop, G.; Gruia, A.T.; Horhat, F.G. Chemical composition and antimicrobial activity of essential oils of lavender (*Lavandula angustifolia*) and lavandin (*Lavandula x intermedia*) grown in western Romania. *Int. J. Agric. Biol.* 2013, 15, 772–776.
23. Zagorcheva, T.; Stanev, S.; Rusanov, K.; Atanassov, I. Comparative GC/MS analysis of lavender (*Lavandula angustifolia* Mill.) inflorescence and essential oil volatiles. *Agric. Sci. Technol.* 2013, 5, 459–462.
24. Chatzopolou, P.S.; Goliaris, A.H.; Katsiotis, T. Contribution to the analysis of the volatile constituents from some lavender and lavandin cultivars grown in Greece. *Sci. Pharm.* 2003, 71, 229–234.
25. Raina, A.P.; Negi, K.S. Comparative essential oil composition of *Lavandula* species from India. *J. Herb Spice Med. Plants* 2012, 18, 268–273.
26. Verma, R.S.; Rahman, L.U.; Chanotiya, C.S.; Verma, R.K.; Chauhan, A.; Singh, A.; Yadav, A.K. Essential oil composition of *Lavandula angustifolia* Mill. cultivated in the mid hills of Uttarakhand. India. *J. Serbian Chem. Soc.* 2010, 75, 343–348.
27. Jaeger, S.; Trojan, H.; Kopp, T.; Laszczyk, M.N.; Scheffler, A. Pentacyclic triterpene distribution in various plants-rich sources for a new group of multi-potent plant extracts. *Molecules* 2009, 14, 2016–2031.
28. Janicsak, G.; Veres, K.; Zoltan Kakasy, A.; Mathe, I. Study of the oleanolic and ursolic acid contents of some species of the Lamiaceae. *Biochem. Syst. Ecol.* 2006, 34, 392–396.

29. Duke, J.A. Handbook of Biologically Active Phytochemicals and Their Activities, 1st ed.; CRC Press: Boca Raton, FL, USA, 1992; p. 208.
30. Liu, J. Pharmacology of oleanolic acid and ursolic acid. *J. Ethnopharmacol.* 1995, 49, 57–68.
31. Wolska, K.; Grudniak, A.; Fiecek, B.; Krackiewicz-Dowjat, A.; Kurek, A. Antibacterial activity of oleanolic and ursolic acids and their derivatives. *Open Life Sci.* 2010, 5, 543–553.
32. Hsu, Y.L.; Kuo, P.L.; Lin, C.C. Proliferative inhibition, cell-cycle dysregulation, and induction of apoptosis by ursolic acid in human non-small cell lung cancer A549 cells. *Life Sci.* 2004, 75, 2303–2316.
33. Chiang, L.C.; Chiang, W.; Chang, M.Y.; Ng, L.T.; Lin, C.C. Antileukemic activity of selected natural products in Taiwan. *Am. J. Chin. Med.* 2003, 31, 37–46.
34. Zhang, W.; Hong, D.; Zhou, Y.; Zhang, Y.; Shen, Q.; Li, J.; Hu, L.; Li, J. Ursolic acid and its derivative inhibit protein tyrosine phosphatase 1B, enhancing insulin receptor phosphorylation and stimulating glucose uptake. *Biochim. Biophys. Acta* 2006, 1760, 1505–1512.
35. Tian, Z.; Lin, G.; Zheng, R.X.; Huang, F.; Yang, M.S.; Xiao, P.G. Anti-hepatoma activity and mechanism of ursolic acid and its derivatives isolated from *Aralia decaisneana*. *World J. Gastroenterol.* 2006, 12, 874–879.
36. Kim, Y.K.; Yoon, S.K.; Ryu, S.Y. Cytotoxic Triterpenes from Stem Bark of *Physocarpus intermedius*. *Planta Med.* 2000, 66, 485–486.
37. Kashiwada, Y.; Nagao, T.; Hashimoto, A.; Ikeshiro, Y.; Okabe, H.; Cosentino, L.M.; Lee, K.H. Anti-AIDS agents 38. Anti-HIV activity of 3-O-acyl ursolic acid derivatives. *J. Nat. Prod.* 2000, 63, 1619–1622.
38. Yan, S.; Huang, C.; Wu, S.; Yin, M. Oleanolic acid and ursolic acid induce apoptosis in four human liver cancer cell lines. *Toxicol. In Vitro* 2010, 24, 842–848.
39. Kowalski, R. Studies of selected plant raw materials as alternative sources of triterpenes of oleanolic and ursolic acid types. *J. Agric. Food Chem.* 2007, 55, 656–662.
40. Zheljazkov, V.D.; Astatkie, T. Effect of residual distillation water of 15 plants and three plant hormones on Scotch spearmint (*Mentha × gracilis* Sole). *Ind. Crop. Prod.* 2011, 33, 704–709.
41. Tiliacos, C.; Gaydou, E.M.; Bessiere, J.-M.; Agnel, R. Distilled lavandin (*Lavandula intermedia* Emeric ex. *Loise l*) wastes: A rich source of coumarin and herniarin. *J. Essent. Oil Res.* 2008, 20, 412–413.
42. Torras-Claveria, L.; Jauregui, O.; Bastida, J.; Codina, C.; Viladomat, F. Antioxidant Activity and Phenolic Composition of Lavandin (*Lavandula x intermedia* Emeric ex. *Loiseleur*). *J. Agric. Food Chem.* 2007, 55, 8436–8443.

43. Khadzhieva, P.; Aleksiev, K.; Topalova, I. Di-and triterpenoids in essential oil waste-from pinus, lavender, and salvia. *Int. Conf. Chem. Biotechnol. Biol. Act. Nat. Prod.* 1987, 5, 519–524.
44. Amorese, V.; Donadu, M.; Usai, D.; Sanna, A.; Milia, F.; Pisanu, F.; Molicotti, P.; Zanetti, S.; Doria, C. In vitro activity of essential oils against *Pseudomonas aeruginosa* isolated from infected hip implants. *J. Infect. Dev. Ctries.* 2018, 12, 996–1001.
45. Yap, P.S.X.; Krishnan, T.; Yiap, B.C.; Hu, C.P.; Chan, K.-G.; Lim, S.H.E. Membrane disruption and anti-quorum sensing effects of synergistic interaction between *Lavandula angustifolia* (lavender oil) in combination with antibiotic against plasmid-conferred multi-drug-resistant *Escherichia coli*. *J. Appl. Microbiol.* 2014, 116, 1119–1128.
46. Yap, P.S.X.; Lim, S.H.E.; Hu, C.P.; Yiap, B.C. Combination of essential oils and antibiotics reduce antibiotic resistance in plasmid-conferred multidrug resistant bacteria. *Phytomedicine* 2013, 20, 710–713.
47. Kunicka-Styczyńska, A.; Śmigielski, K.; Prusinowska, R.; Rajkowska, K.; Kuśmider, B.; Sikora, M. Preservative activity of lavender hydrosols in moisturizing body gels. *Lett. Appl. Microbiol.* 2015, 60, 27–32.
48. ISO 3515:2002 Oil of Lavender (*Lavandula Angustifolia* Mill.); ISO: Geneva, Switzerland, 2002.
49. Ivanov, I.; Petkova, N.; Tumbarski, Y.; Vrancheva, R.; Stoyanova, M. Lavender waste—promising source of triterpenoids and polyphenols with antioxidant and antimicrobial activity. *Ind. Technol.* 2018, 5, 26–32.
50. Piqué, N.; Miñana-Galbís, D.; Merino, S.M.; Tomás, J. Virulence Factors of *Erwinia amylovora*: A Review. *Int. J. Mol. Sci.* 2015, 16, 12836–12854.
51. Benada, M.; Boumaaza, B.; Boudalia, S.; Khaladi, O.; Guessas, B. Variability of aggressiveness and virulence of *Erwinia carotovora* subsp. *carotovorum* causing the soft rot on potato tubers in the western of Algeria. *Int. J. Plant Biol.* 2018, 9, 52–56.
52. Tamir-Ariel, D.; Navon, N.; Burdman, S. Identification of Genes in *Xanthomonas campestris* pv. *vesicatoria* induced during its interaction with tomato. *J. Bacteriol.* 2007, 189, 6359–6371.
53. Davies, J.C. *Statistics and Data Analysis in Geology*; Elsevier: Amsterdam, The Netherlands, 2002.
54. Shi, G. Cluster Analysis. In *Data Mining and Knowledge Discovery for Geoscientists*; Shi, G., Ed.; Elsevier: Amsterdam, The Netherlands, 2014.

Retrieved from <https://encyclopedia.pub/entry/history/show/33203>