

Trichoderma spp. and Cocoa Seed

Subjects: Plant Sciences

 Thiago

Submitted by: Vieira

Definition

Isolates of *Trichoderma* spp., a soil fungus, has been used to control diseases and promote plant growth, reducing the use of chemicals in the production of seedlings of different plant species. We evaluated the effect of some *Trichoderma* spp. isolates on seed treatment and seedling production of *Theobroma cacao*. Five isolates from the Amazon region were tested. In laboratory, the following variables were evaluated for seed treatments: germination, germination speed index, radicle and hypocotyl lengths, and fungi incidence. In nursery, the following forms of application were tested: via seeds; in the substrate at pre-planting; monthly in post-planting substrate, and also their combination. The following was evaluated: height, diameter, number of leaves, root length, leaf area, and shoot dry mass and root system. Inoculation with *Trichoderma* increased the length of the radicle and hypocotyl and showed no fungi in the seeds. In seedlings, some treatments increased height and plant root dry mass. The use of *Trichoderma* was beneficial for seeds and appeared favorable for *T. cacao* production.

1. Introduction

Recently, an increase in the use of biological agents in agriculture has been observed ^{[1][2]}, which consists of the application of microorganisms or their metabolites to protect seeds and promote germination and plant growth and management of different pathogens and pests ^{[3][4]}. Among the microorganisms that can be used, several isolates of *Trichoderma* spp. showed different mechanisms of action, such as antibiosis, mycoparasitism, competition, resistance induction, and growth promotion, in addition to solubilization of nutrients ^{[5][6][7][8]}. These properties provide productive and economic gains in the process of growing important crops.

Cocoa (*Theobroma cacao* L.) is a species of great economic and social importance globally. Worldwide, cocoa-producing areas occupy approximately 6.7 million hectares, with an estimated production of 4.7 million tons, generating a revenue of approximately 8.6 billion dollars per year ^[9]. In Brazil, this crop gained prominence with the expansion of chocolate consumption. Its cultivation represents a good example for composing agroforestry systems, as it can be cultivated under the shade of the thinned native forest ("cabruca" system) in association with other species, without the need of deforestation ^[10].

Pests and diseases affect cocoa crops strongly, constituting a major problem for producers ^[11]. The use of agrochemicals, applied from seed treatment and seedling preparation through cultivation to post harvest, has caused serious problems of environmental contamination, and therefore, products in favor of sustainable agriculture have been studied and developed.

The use of *Trichoderma* spp. on *T. cacao* has mainly been studied for disease control ^[12]. However, this fungus can also promote the development of seedlings and root systems in addition to delaying drought responses when plants are subjected to water stress ^[13]. Thus, research is important to identify *Trichoderma* isolates that are efficient and adapted to the environmental conditions of the producing region, which can be used for treating seeds and seedlings and in the development of cultivated plants.

The sanitary quality of seeds is essential for the production of healthy and vigorous seedlings ^[14] and for the production of quality, vigorous, and pest and disease-free seedlings with good adaptation and field growth ^[15]. Thus, the treatment of seeds and seedlings with *Trichoderma* spp. is an alternative that can generate good results in different plant species, as it is considered sustainable.

Our hypothesis was that the application of *Trichoderma* stimulates the germination of cacao seeds and the initial growth of seedlings, reducing the time the plants remain in the nursery, ensuring quality seedlings. In this article, we evaluate the effect of five *Trichoderma* isolates in different application modes on seed treatment and development of *T. cacao* seedlings.

2. *Trichoderma* Effect on *T. cacao* Seed Germination

There was a significant difference for the radicle and hypocotyl length and the incidence of fungi. In all treatments, germination was greater than 95% (**Table 1**). *Trichoderma* did not influence germination speed compared to control.

Table 1. Germination and germination speed index (SG) and radicle and hypocotyl lengths of *Theobroma cacao* seeds treated or not with different *Trichoderma* isolates. Incidence of fungi represent the % of seed contaminated with other fungi.

Treatment	Germination (%)	SG	Radicle (cm)	Hypocotyl (cm)	Incidence of Fungi (%)				
Control	97.0	ab	58.3	a	1.2	c	1.3	c	26.5
<i>Trichoderma</i> sp. Tc	98.0	a	61.7	a	4.3	a	2.6	a	0.0
<i>Trichoderma</i> sp. Tce	97.5	b	61.7	a	4.2	a	2.2	ab	0.0
<i>T. asperellum</i> Tam01	96.5	ab	61.9	a	4.1	a	2.3	ab	0.0
<i>T. asperellum</i> Tam02	98.0	a	54.4	a	3.2	b	1.6	bc	0.0
<i>T. asperellum</i> Tam03	95.0	b	52.6	a	3.1	b	2.4	a	0.0
CV(%)	1.3		7.7		9.8		15.8		

CV, coefficient of variation; means followed by the same letters in the columns do not differ from each other by the Tukey test ($p \leq 0.05$).

All *Trichoderma* isolates increased radicle length compared to the control (**Table 1, Figure 1**), ranging from 1.9 to 3.1 cm, corresponding to an increase from 158% to 258.3%, respectively. For the length of the hypocotyl, there was a significant difference except for the seeds treated with isolate Tam02, which did not differ from the control (**Table 1**). The increase in hypocotyl ranged from 0.9 to 1.3 cm, representing an increase from 69.2% to 100% for the Tce and Tc isolates, respectively.



Figure 1. Radicle and hypocotyl lengths of cocoa seedlings resulting from seeds treated or not with different *Trichoderma* spp. isolates.

The incidence of other fungi was observed only in the control treatment, with 26.5% of contaminated seeds (**Table 1**). They did not affect, however, the germination of seeds.

3. *Trichoderma* spp. in the Production of *T. cacao* Seedlings

A significant difference for the height and root dry mass of seedlings, at three and six months, was observed (**Table 2**).

Table 2. Height (cm), diameter of the collar (mm), number of leaves per plant, dry mass (DM) of aerial

shoot and root system (g), root length (cm), and leaf area (cm²) of *Theobroma cacao* seedlings submitted to different application modes and *Trichoderma* spp. isolates.

Treatments	Height		Collar Diameter		Number of Leaves		Aerial DM	Root DM	Root Length	Leaf Area
	3rd month	6th month	3rd month	6th month	3rd month	6th month				
Tc applied to seeds	26.0 ^{ns}	27.9 ^{ns}	5.4 ^{ns}	6.2 ^{ns}	7.7 ^{ns}	9.3 ^{ns}	2.8 ^{ns}	1.9 ^{ns}	49.1 ^{ns}	613.1 ^{ns}
Tc applied to pre-planting substrate	24.9 ^{ns}	29.6 ^{ns}	5.2 ^{ns}	6.4 ^{ns}	7.5 ^{ns}	10.6 ^{ns}	3.3 ^{ns}	2.0 ^{ns}	54.4 ^{ns}	437.7 ^{ns}
Tc applied monthly to post-planting substrate	25.7 ^{ns}	30.3 ^{ns}	5.7 ^{ns}	6.5 ^{ns}	7.8 ^{ns}	12.7 ^{ns}	4.1 ^{ns}	2.2 ^{ns}	47.0 ^{ns}	613.1 ^{ns}
Tc seeds + substrate + monthly	28.2 ^{ns}	32.3 ^{ns}	5.5 ^{ns}	6.3 ^{ns}	8.8 ^{ns}	11.2 ^{ns}	3.3 ^{ns}	2.1 ^{ns}	54.9 ^{ns}	448.7 ^{ns}
Tce applied to seeds	27.0 ^{ns}	30.4 ^{ns}	5.2 ^{ns}	6.4 ^{ns}	7.2 ^{ns}	10.3 ^{ns}	3.6 ^{ns}	2.3 ^{ns}	51.1 ^{ns}	551.8 ^{ns}
Tce applied to pre-planting substrate	26.5 ^{ns}	31.2 ^{ns}	5.2 ^{ns}	6.2 ^{ns}	8.0 ^{ns}	10.2 ^{ns}	3.0 ^{ns}	1.8 ^{ns}	51.7 ^{ns}	539.1 ^{ns}
Tce applied monthly to post-planting substrate	24.5 ^{ns}	30.6 ^{ns}	5.0 ^{ns}	6.2 ^{ns}	8.8 ^{ns}	11.1 ^{ns}	3.4 ^{ns}	2.0 ^{ns}	54.9 ^{ns}	438.6 ^{ns}
Tce seeds + substrate + monthly	26.0 ^{ns}	31.1 ^{ns}	5.4 ^{ns}	6.3 ^{ns}	9.0 ^{ns}	10.5 ^{ns}	3.6 ^{ns}	2.3 ^{ns}	52.8 ^{ns}	413.6 ^{ns}
Tam01 applied to seeds	26.9 ^{ns}	32.3 ^{ns}	5.4 ^{ns}	6.7 ^{ns}	8.5 ^{ns}	11.0 ^{ns}	3.6 ^{ns}	1.9 ^{ns}	52.3 ^{ns}	514.1 ^{ns}
Tam01 applied to pre-planting substrate	26.8 ^{ns}	29.8 ^{ns}	5.1 ^{ns}	6.0 ^{ns}	8.0 ^{ns}	10.6 ^{ns}	2.8 ^{ns}	1.9 ^{ns}	53.4 ^{ns}	380.6 ^{ns}
Tam01 applied monthly to post-planting substrate	30.0 [*]	34.2 [*]	5.0 ^{ns}	6.5 ^{ns}	8.7 ^{ns}	11.8 ^{ns}	3.8 ^{ns}	1.8 ^{ns}	50.0 ^{ns}	585.5 ^{ns}
Tam01 seeds + substrate + monthly	27.6 ^{ns}	31.5 ^{ns}	5.8 ^{ns}	6.6 ^{ns}	9.5 ^{ns}	11.1 ^{ns}	3.4 ^{ns}	2.8 [*]	55.2 ^{ns}	559.2 ^{ns}
Tam02 applied to seeds	27.9 ^{ns}	34.3 [*]	5.2 ^{ns}	6.3 ^{ns}	8.8 ^{ns}	11.2 ^{ns}	3.4 ^{ns}	1.7 ^{ns}	59.6 ^{ns}	529.9 ^{ns}
Tam02 applied to pre-planting substrate	25.7 ^{ns}	31.4 ^{ns}	5.2 ^{ns}	6.2 ^{ns}	8.6 ^{ns}	11.5 ^{ns}	3.9 ^{ns}	2.1 ^{ns}	61.9 ^{ns}	491.4 ^{ns}
Tam02 applied monthly to post-planting substrate	27.3 ^{ns}	29.4 ^{ns}	5.1 ^{ns}	6.1 ^{ns}	7.8 ^{ns}	9.2 ^{ns}	2.9 ^{ns}	1.6 ^{ns}	53.1 ^{ns}	442.2 ^{ns}
Tam02 seeds + substrate + monthly	28.8 ^{ns}	32.4 ^{ns}	5.5 ^{ns}	6.5 ^{ns}	8.8 ^{ns}	9.6 ^{ns}	3.5 ^{ns}	2.1 ^{ns}	56.1 ^{ns}	464.0 ^{ns}
Tam03 applied to seeds	25.8 ^{ns}	28.3 ^{ns}	5.3 ^{ns}	6.2 ^{ns}	7.8 ^{ns}	9.6 ^{ns}	3.2 ^{ns}	2.0 ^{ns}	43.6 ^{ns}	494.4 ^{ns}

Treatments	Height		Collar Diameter		Number of Leaves		Aerial DM	Root DM	Root Length	Leaf Area
	3rd month	6th month	3rd month	6th month	3rd month	6th month				
Tam03 applied to pre-planting substrate	26.5 ^{ns}	30.2 ^{ns}	5.3 ^{ns}	6.3 ^{ns}	7.2 ^{ns}	10.3 ^{ns}	2.8 ^{ns}	1.5 ^{ns}	53.2 ^{ns}	425.4 ^{ns}
Tam03 applied monthly to post-planting substrate	27.0 ^{ns}	29.9 ^{ns}	5.2 ^{ns}	6.0 ^{ns}	7.7 ^{ns}	10.7 ^{ns}	3.3 ^{ns}	1.7 ^{ns}	51.1 ^{ns}	422.8 ^{ns}
Tam03 seeds + substrate + monthly	26.8 ^{ns}	31.1 ^{ns}	5.5 ^{ns}	6.4 ^{ns}	7.8 ^{ns}	9.2 ^{ns}	3.1 ^{ns}	1.9 ^{ns}	56.4 ^{ns}	385.1 ^{ns}
Control	25.6	28.4	5.1	6.0	7.6	10.7	3.4	1.7	51.5	484.1
Coefficient of Variation (%)	9.9	10.5	10.4	11.4	18.6	22.1	28.0	31.9	23.2	20.7

* Significant; ns, not significant; by Dunnett test ($p < 0.05$). Tc, *Trichoderma* sp.; Tce, *Trichoderma* sp.; Tam01, *T. asperellum*; Tam02, *T. asperellum*; Tam03, *T. asperellum*.

Two *Trichoderma* isolates increased plant height compared to the control. They were *T. asperellum* Tam01, applied monthly to the substrate after planting in the two evaluation periods, and *T. asperellum* Tam02, applied to the seeds in the evaluation at six months only (**Table 2**, **Figure 2**). The increase in height provided by Tam01 achieved 4.4 cm at three months and 5.8 cm at six months, corresponding to an increase by 17.2% and 20.4%, respectively. For Tam02 applied to seeds, the increase in height was 5.9 cm (20.7%) when compared to the control.



Figure 2. Cocoa seedlings with and without application of *Trichoderma asperellum*, six months after planting.

Tam01 also showed an increase in root dry mass of plants in the triple combination of application modes (seeds + substrate pre-planting + monthly applications in the substrate post planting) (**Table 2**), with an increase of 1.1 g in relation to the control, corresponding to a 64% increment.

When comparing the treatments, there was a significant difference for the interaction between (*Trichoderma* isolates \times application modes) only for the plant height at six months after planting. In the mode of application via seeds, the height of seedlings treated with Tam02 was greater than those that

received the application of isolates Tc and Tam03. In the monthly applications in the post-planting substrate, a difference was observed only between Tam01 and Tam02, with mean height values of 34.2 cm and 29.4 cm, respectively (**Table 3**).

Table 3. Height of *Theobroma cacao* seedlings in relation to different modes of application modes of *Trichoderma* spp., six months after planting.

Modes of Application	Trichoderma Isolates									
	Tc		Tce		Tam01		Tam02		Tam03	
Seeds ¹	27.9	bB	30.3	aAB	32.3	abAB	34.3	aA	28.2	aB
Substrate ²	29.5	abA	31.2	aA	29.7	bA	31.3	abA	30.2	aA
Monthly Application ³	30.3	abAB	30.6	aAB	34.2	aA	29.4	bB	29.9	aAB
Seed + Subs. + Mon.Ap. ⁴	32.3	aA	31.0	aA	31.5	abA	32.4	abA	31.1	aA
CV (%) 10.5										

Means followed by the same lowercase letters in the columns and the same uppercase letters in the rows do not differ by Tukey's test ($p \leq 0.05$). ¹ = *Trichoderma* application to seeds; ² = application of *Trichoderma* in the substrate at pre-planting; ³ = monthly application of *Trichoderma* in the post-planting substrate; ⁴ = *Trichoderma* applied seeds + application in the substrate at pre-planting + monthly application in the substrate post planting.

4. Discussion

Biological agents applied to seeds and seedlings are important components of sustainable agriculture strategies. In this study, *Trichoderma* spp. natives of the Amazon region were used in cocoa, a naturally occurring species of this biome.

Although *Trichoderma* isolates did not increase germination parameters compared to the control, they positively affected seedlings growth. Data indicate a healthy condition of the seeds used, showing an average germination close to that of newly harvested fruits of this species, which can reach 100% [16]. This was likely due to storage lipids that act as carbon and energy reserves for embryos to germinate, favoring high germination rates [17]. On the opposite, the storage of seeds for long periods of time affects germination, as seeds do not tolerate dehydration or low temperatures [18].

Biological treatment positively influenced other variables related to cocoa seed germination, such as radicle and hypocotyl length. Data showed that the greatest increase in radicle length was 3.6-times higher than the control and twice the hypocotyl length. Indoleacetic acid (IAA) appears as a possible effector in this interaction. Root exudates are sources of the amino acid L-tryptophan, a precursor for IAA biosynthesis by the rhizosphere microbiota [19][20]. IAA regulates several stages of plant growth and development, such as cell elongation and division, tissue differentiation, and apical dominance [21]. In addition to auxins, cytokinin produced by microorganisms promote plant growth and cell division [22]. The effects observed on shoot growth may be hence related to the production active effectors translocated from the roots to the shoot and induced by auxin-producing microorganisms that thus stimulate the root system [23]. The production of auxins by *Trichoderma* spp. has already been shown [24][25].

There are several reports indicating a biostimulation effect induced by *Trichoderma* spp. Seeds of *Theobroma grandiflorum*, a relative of cocoa, treated with isolates of *T. asperellum* showed an increased length of the radicle in the seedlings but no effect on germination parameters [26]. Moreover, the incidence of fungi was observed only in untreated seeds, demonstrating the efficiency of *Trichoderma* in controlling fungi naturally occurring in cocoa seeds. Similarly, in *T. grandiflorum*, no

fungi were observed in seeds treated with the same *Trichoderma* isolates herein evaluated [26].

The absence of harmful fungi in seeds is important for developing of healthy seedlings and for their survival in the field. Among seeds colonizing fungi, species of *Aspergillus* and *Penicillium* have been reported as potential producers of mycotoxins in cocoa beans [27]. Seed treatments are important to reduce or eliminate microorganisms that can harm the seeds and affect the initial development of seedlings. Additionally, since *Trichoderma* spp. includes important biocontrol agents, biological treatments appear useful to protect the young plants from the attack of pathogens already present in the soil.

Identifying the ideal conditions for seed germination is an important factor to ensure uniform production of quality seedlings and for the establishment of a standard in the field production [28]. In the nursery assay, only two *Trichoderma* isolates applied in seeds and in the post-planting substrate influenced height (Tam01 and Tam02) and root dry mass (Tam01). Differences in isolates efficiency may be related to different mechanisms of action in plant growth promotion and biological control that may vary according to environmental conditions (substrate, nutrients, other microorganisms), species, or association with the host plant [29].

Growth promotion is not a general feature for all *Trichoderma* spp., as different effects on plants have been observed. In addition, promotion variability may be related to and influenced by environmental factors [30][31]. In a collection from Colombia, 60% of *Trichoderma* strains produced IAA derivatives in vitro, but only 18% showed plant growth effects on beans [32]. In addition to this, auxins and volatile organic compounds have been described as growth-promoting inducers in fungi, including *Trichoderma* [33]. Evaluation of *Trichoderma* spp. isolates in *Jatropha curcas* and *Ricinus communis* showed negative effects for some isolates on the diameter, number of leaves, and dry mass of the aerial part and root system of plants [34]. A promotion effect on root dry mass and height were instead reported in *Gochnatia polymorpha* plants treated with *Trichoderma harzianum* [29]. Seedlings of *Handroanthus serratifolius* (yellow ipe) treated with *Trichoderma* showed increased height, stem diameter, number of leaves, and root dry mass compared to the control [35]. Increased height of plants treated with *Trichoderma* was also found in *Khaya ivorensis* and *Eucalyptus camaldulensis*, with increased number of leaves and higher shoot and root dry mass, and *Pinus radiata* [36][37][38].

The 64% increase in root dry mass observed in the treatment with *Trichoderma* Tam01 isolate and applied in a triple combination (seeds + pre-planting substrate + monthly applications in post-planting substrate) is a promising result. The plant's investment in root growth is one of the main benefits attributed to microorganisms that produce IAA. For young plants, the growth and establishment of roots is essential, as the increase in root biomass promotes adherence to the substrate and greater absorption of water and nutrients. These effects increase nutrients accumulation and cell multiplication through the production of hormones and other metabolites, consequently increasing the chances of survival in the field [39][40][41][42][43][44][45][46].

The application of *Trichoderma* Tam01 isolate in the pre-planting substrate combined with applications in the post-planting substrate also positively influenced the development of *Euterpe oleracea* seedlings [2]. The use of *Trichoderma* in the pre-planting substrate increased the height, stem diameter, and number of leaves of *E. schomburgkii* seedlings [47]. *Trichoderma* spp. used in seeds or in combination with substrate application were also effective on seedlings of *Khaya ivorensis* [36]. In seedlings of *Parapiptadenia rigida*, the application of *Trichoderma* via seeds showed the greatest increase in plant height [48].

Trichoderma spp. have potential for application with several purposes, such as biological control of plant diseases, production of enzymes for industries, and other fundamental aspects, such as increasing the efficiency of nitrogen use, promoting the development of plants, as well as productivity and alleviating the impacts of salt stresses [49].

References

1. Jorge, D.M.; Souza, C.A.V. O papel da regulamentação dos produtos de origem biológica no avanço da agroecologia e da produção orgânica no Brasil. In *A Política Nacional de Agroecologia e Produção Orgânica no Brasil: Uma Trajetória de Luta Pelo Desenvolvimento Rural Sustentável*, 1st ed.; Sambuichi, R.H.R., Moura, I.F., Mattos, L.M., Ávila, M.L., Spínola, P.A.C., Silva, A.P.M., Eds.; IPEA: Brasília, Brasil, 2017; pp. 229–252.
2. Campos, B.F.; Araújo, A.J.C.; Felseburgh, C.A.; Vieira, T.A.; Lustosa, D.C. *Trichoderma* contributes to the germination and seedling development of Açai Palm. *Agriculture* 2020, 10, 456.
3. Andrade-Hoyos, P.; Silva-Rojas, H.V.; Romero-Arenas, O. Endophytic *Trichoderma* species isolated from *Persea americana* and *Cinnamomum verum* roots reduce symptoms caused by *Phytophthora cinnamomi* in avocado. *Plants* 2020, 9, 1220.
4. Silletti, S.; Di Stasio, E.; Van Oosten, M.J.; Ventorino, V.; Pepe, O.; Napolitano, M.; Marra, R.; Woo, S.L.; Cirillo, V.; Maggio, A. Biostimulant activity of *Azotobacter chroococcum* and *Trichoderma harzianum* in Durum wheat under water and nitrogen deficiency. *Agronomy* 2021, 11, 380.
5. Machado, D.F.M.; Parzianello, F.R.; Silva, A.C.F.; Antonioli, Z.I. *Trichoderma* no Brasil: O fungo e o bioagente. *Rev. Cienc. Agrar.* 2012, 35, 274–288.
6. Carillo, P.; Woo, S.L.; Comite, E.; El-Nakhel, C.; Roupheal, Y.; Fusco, G.M.; Borzacchiello, A.; Lanzuise, S.; Vinale, F. Application of *Trichoderma harzianum*, 6-Pentyl- α -pyrone and plant biopolymer formulations modulate plant metabolism and fruit quality of plum tomatoes. *Plants* 2020, 9, 771.
7. Halifu, S.; Deng, X.; Song, X.; Song, R.; Liang, X. Inhibitory mechanism of *Trichoderma virens* ZT05 on *Rhizoctonia solani*. *Plants* 2020, 9, 912.
8. Marra, R.; Coppola, M.; Pironi, A.; Grasso, F.; Lombardi, N.; d'Errico, G.; Sicari, A.; Bolletti Censi, S.; Woo, S.L.; Rao, R.; et al. The Application of *Trichoderma* strains or metabolites alters the olive leaf metabolome and the expression of defense-related genes. *J. Fungi* 2020, 6, 369.
9. Chitiva-Chitiva, L.C.; Ladino-Vargas, C.; Cuca-Suárez, L.E.; Prieto-Rodríguez, J.A.; Patiño-Ladino, O.J. Antifungal activity of chemical constituents from *Piper pesaesatum* C. DC. and derivatives against phytopathogen fungi of cocoa. *Molecules* 2021, 26, 3256.
10. Piasentin, F.B.; Saito, C.H. Os diferentes métodos de cultivo de cacau no sudeste da Bahia, Brasil: Aspectos históricos e percepções. *Bol. Mus. Para. Emílio Goeldi. Ciências Hum.* 2014, 9, 61–78.
11. Cilas, C.; Bastide, P. Challenges to cocoa production in the face of climate change and the spread of pests and diseases. *Agronomy* 2020, 10, 1232.
12. Rodrigues, G.S.; Magalhães, D.M.A.; Costa, A.M.; Luz, E.D.M.N. Antagonismo de *Trichoderma* spp. ao agente etiológico da murcha de *Ceratocystis* em cacauzeiro. *Summa Phytopathol.* 2018, 44, 72–78.
13. Bae, A.; Sicher, R.C.; Kim, M.S.; Kim, S.H.; Strem, M.D.; Melnick, R.L.; Bailey, B.A. The beneficial endophyte *Trichoderma hamatum* isolate DIS 219b promotes growth and delays the onset of the drought response in *Theobroma cacao*. *J. Exp. Bot.* 2009, 60, 3279–3295.
14. Lazarotto, M.; Muniz, M.F.B.; Beltrame, R.; Santos, A.F.; Mezzomo, R.P.G.; Blume, E. Qualidade fisiológica e tratamentos de sementes de *Cedrela fissilis* procedentes do sul do Brasil. *Rev. Árvore* 2013, 37, 201–210.
15. Cruz, C.A.F.; Paiva, H.N.; Gomes, K.C.O.; Guerrero, C.R.A. Efeito de diferentes níveis de saturação por bases no desenvolvimento e qualidade de mudas de ipê roxo (*Tabebuia impetiginosa* (Mart.) Standley). *Sci. For.* 2004, 66, 100–107.
16. Venial, L.R.; Alexandre, R.S.; Camata, H.; Lopes, J.C.; Zanotti, R.F.; Ferreira, A.; Aguilar, M.A.G. Biometria e armazenamento de sementes de genótipos de cacauzeiro. *Pesqui. Florest. Bras.* 2017, 37, 39–46.
17. Argout, X.; Salse, J.; Aury, J.-M.; Guiltinan, M.J.; Droc, G.; Gouzy, J.; Allegre, M.; Chaparro, C.; Legavre, T.; Maximova, S.N.; et al. The genome of *Theobroma cacao*. *Nat. Genet.* 2011, 43, 101–108.
18. Salles, B.P.A.; David, A.M.S.S.; Figueiredo, J.C.; Maia, V.M.; Prudêncio, J.R.S.; Pereira, K.K.G. Viabilidade de sementes de cacau e limitações no armazenamento. *Rev. Cienc. Agrar.* 2019, 42, 1010–1014.
19. Lum, M.R.; Hirsch, A.M. Rots and their symbiotic microbes: Strategies to obtain nitrogen and phosphorus in a nutrient limiting environment. *J. Plant Growth Regul.* 2003, 21, 368–382.
20. Khalid, A.; Arsha, M.; Zahir, Z.A. Screening plant growth-promoting rhizobacteria for improving growth and yield of wheat. *J. Appl. Microbiol.* 2004, 96, 473–480.
21. Gutiérrez-Luna, F.M.; López-Bucio, J.; Altamirano-Hernández, J.; Valencia-Cantero, E.; de la Cruz, E.R.; Macías-Rodríguez, L. Plant growth-promoting rhizobacteria modulate root-system architecture in *Arabidopsis thaliana* through volatile organic compound emission. *Symbiosis* 2010, 51, 75–83.
22. Germida, J.J.; Walley, F.L. Plant growth-promoting rhizobacteria alter rooting patterns and arbuscular mycorrhizal fungi colonization of field-grown spring wheat. *Biol. Fertil. Soils* 1996, 23, 113–120.
23. Orhan, E.; Esitken, A.; Ercisli, S.; Turan, M.; Sahin, F. Effects of plant growth promoting rhizobacteria (PGPR) on yield, growth and nutrient contents in organically growing raspberry. *Sci. Hortic.* 2006, 111, 38–43.
24. Oliveira, A.G.; Chagas Junior, A.F.; Santos, G.R.; Miller, L.O.; Chagas, L.F.B. Potencial de solubilização de fosfato e

- produção de AIA por *Trichoderma* spp. Rev. Verde De Agroecol. E Desenvol. Sustentável 2012, 7, 149-155.
25. Almança, M.A.K. Aspectos da interação Arroz-*Trichoderma* spp. em solos alagados. Ph.D. Thesis, Universidade Federal do Rio Grande do Sul, Porto Alegre, Brazil, 2008.
 26. Santos, M.F.; Costa, D.L.; Matos, J.C.N.; Silva, G.B.; Vieira, T.A.; Lustosa, D.C. Tratamento biológico de sementes de cupuaçu para o controle de fitopatógenos e promoção da germinação. Cad. Agroecol. 2018, 13, 1017.
 27. Sánchez-Hervás, M.; Gil, J.V.; Bisbal, F.; Ramón, D.; Martínez-Culebras, P.V. Mycobiota and mycotoxin producing fungi from cocoa beans. Int. J. Food Microbiol. 2008, 125, 336-340.
 28. Paixão, M.V.S.; Demuner, F.M.; Rodrigues, P.S.; Faria Junior, H.P.; Bozetti, M. Tratamentos pré germinativos na germinação de sementes de cacau. Int. J. Adv. Eng. Res. Sci. 2019, 6, 130-134.
 29. Machado, D.F.M.; Tavares, A.P.; Lopes, S.J.; Silva, A.C.F. *Trichoderma* spp. na emergência e crescimento de mudas de camarará (*Gochnatia polymorpha* (Less.) Cabrera)1. Rev. Árvore 2015, 39, 167-176.
 30. Lee, S.; Hung, R.; Yap, M.; Bennett, J.W. Age matters: The effects of volatile organic compounds emitted by *Trichoderma atroviride* on plant growth. Arch. Microbiol. 2015, 197, 723-727.
 31. Nieto-Jacobo, M.F.; Steyaert, J.M.; Salazar-Badillo, F.B.; Nguyen, D.V.; Rostás, M.; Braithwaite, M.; De Souza, J.T.; Jimenez-Bremont, J.F.; Ohkura, M.; Stewart, A.; et al. Environmental growth conditions of *Trichoderma* spp. affects Indole Acetic Acid derivatives, volatile organic compounds, and plant growth promotion. Front. Plant Sci. 2017, 9, 102.
 32. Hoyos-Carvajal, L.; Orduz, S.; Bissett, J. Growth stimulation in bean (*Phaseolus vulgaris* L.) by *Trichoderma*. Biol. Control 2009, 51, 409-416.
 33. Garnica-Vergara, A.; Barrera-Ortiz, S.; Muñoz-Parra, E.; Raya-González, J.; Méndez-Bravo, A.; Macias-Rodríguez, L.; Ruiz-Herrera, L.F.; López-Bucio, J. The volatile 6-pentyl-2H-pyran-2-one from *Trichoderma atroviride* regulates *Arabidopsis thaliana* root morphogenesis via auxin signaling and ETHYLENE INSENSITIVE 2 functioning. New Phytol. 2015, 209, 1496-1512.
 34. Martins, C.Y.S. Promoção de crescimento e colonização radicular por *Trichoderma* spp. em pinhão manso (*Jatropha curcas* L.) e mamoneira (*Ricinus communis* L.). Master's Thesis, Universidade Federal do Recôncavo da Bahia, Cruz das Almas, Brazil, 2010.
 35. Santos, M.F.; Santos, L.E.; Costa, D.L.; Vieira, T.A.; Lustosa, D.C. *Trichoderma* spp. on treatment of *Handroanthus serratifolius* seeds: Effect on seedling germination and development. Heliyon 2020, 6, e04044.
 36. Lustosa, D.C.; Araujo, A.J.C.; Campos, B.F.; Vieira, T.A. *Trichoderma* spp. and its effects on seeds physiological quality and seedlings development of African mahogany. Rev. Bras. De Cienc. Agrar. 2020, 15, 1-7.
 37. Azevedo, G.B.; Novaes, Q.S.; Azevedo, G.T.O.S.; Silva, H.F.; Rocha Sobrinho, G.G.; Novaes, A.B. Efeito de *Trichoderma* spp. no crescimento de mudas clonais de *Eucalyptus camaldulensis*. Sci. For. 2017, 45, 343-352.
 38. Chávez, D.; Pereira, G.; Machuca, A. Estimulación del crecimiento en plántulas de *Pinus radiata* utilizando hongos ectomicorrízicos y saprobios como biofertilizantes. Bosque 2014, 35, 57-63.
 39. Spaepen, S.; Vanderleyden, J.; Remans, R. Indole-3-acetic acid in microbial and microorganism-plant signaling. FEMS Microbiol. Rev. 2007, 31, 425-448.
 40. Farina, R. Diversidade de Bactérias Promotoras do Crescimento Vegetal Associadas a Cultura De Canola (*Brassica napus* L.) Cultivada no Município de Vacaria, Rio Grande do Sul. Ph.D. Thesis, Universidade Federal do Rio Grande do Sul, Porto Alegre, Brazil, 2012.
 41. Aguiar, A.R.; Aguiar, D.; Tedesco, S.B.; Silva, A.C.F. Efeito de metabólitos produzidos por *Trichoderma* spp. sobre o índice mitótico em células das pontas de raízes de *Allium cepa*. Biosci. J. 2015, 31, 934-940.
 42. Korasick, D.A.; Enders, T.A.; Strader, L.C. Auxin biosynthesis and storage forms. J. Exp. Bot. 2013, 64, 2541-2555.
 43. Carvalho Filho, M.R.; Mello, S.C.M.; Santos, R.P.; Menêzes, J.E. Avaliação de Isolados de *Trichoderma* na Promoção de Crescimento, Produção de Ácido Indolacético In Vitro e Colonização Endofítica de Mudas de Eucalipto; Embrapa Recursos Genéticos e Biotecnologia: Brasília, Brazil, 2008.
 44. Kapri, A.; Tewari, L. Potencial de solubilização de fosfato e atividade de fosfatase de *Trichoderma* spp. rizosférico. Braz. J. Microbiol. 2010, 41, 787-795.
 45. Ribas, P.P.; Rech, R.; Matsumura, A.T.S.; Sand, S.T.V.D. Potencial in vitro para solubilização de fosfato por *Trichoderma* spp. Rev. Bras. Biocienc. 2016, 14, 70-75. Available online: <http://www.ufrgs.br/seerbio/ojs/index.php/rbb/article/view/2967/1315> (accessed on 30 August 2021).
 46. Contreras-Cornejo, H.A.; Macías-Rodríguez, L.; Cortés-Penagos, C.; López-Bicio, J. *Trichoderma virens*, a plant beneficial fungus, enhances biomass production and promotes lateral root growth through an auxin-dependent mechanism in *Arabidopsis*. Plant Physiol. 2009, 149, 1579-1592.
 47. Santos, M.F.; Costa, D.L.; Vieira, T.A.; Lustosa, D.C. Effect of *Trichoderma* spp. fungus for production of seedlings in *Enterolobium Schomburgkii* (Benth.) Benth. Aust. J. Crop Sci. 2019, 10, 1706-1711.
 48. Junges, E.; Muniz, M.F.; Mezzomo, R.; Bastos, B.; Machado, R.T. *Trichoderma* spp. na produção de mudas de espécies florestais. Floresta Ambiente 2016, 23, 237-244.
 49. Bettiol, W.; Silva, J.C.; Castro, M.L.M.P. Uso atual e perspectivas do *Trichoderma* no Brasil. In *Trichoderma—Uso na Agricultura*; Meyer, M.C., Mazaro, S.M., Silva, J.C., Eds.; Embrapa: Brasília, Brazil, 2019; pp. 21-43.

Keywords

biocontrol;Theobroma cacao;biostimulant;germination;growth promotion

Retrieved from <https://encyclopedia.pub/16633>