

# Idiopathic Pulmonary Fibrosis

Subjects: [Immunology](#)

Contributor: Soo Jung Cho , Mihye Lee , Heather W. Stout-Delgado , Jong-Seok Moon

Idiopathic pulmonary fibrosis (IPF) is a chronic, progressive interstitial lung disease. Chronic lung inflammation is linked to the pathogenesis of IPF. DROSHA, a class 2 ribonuclease III enzyme, has an important role in the biogenesis of microRNA (miRNA). The function of miRNAs has been identified in the regulation of the target gene or protein related to inflammatory responses via degradation of mRNA or inhibition of translation. The absent-in-melanoma-2 (AIM2) inflammasome is critical for inflammatory responses against cytosolic double stranded DNA (dsDNA) from pathogen-associated molecular patterns (PAMPs) and self-DNA from danger-associated molecular patterns (DAMPs). The AIM2 inflammasome senses double strand DNA (dsDNA) and interacts with the adaptor apoptosis-associated speck-like protein containing a caspase recruitment domain (ASC), which recruits pro-caspase-1 and regulates the maturation and secretion of interleukin (IL)-1 $\beta$  and IL-18. A recent study showed that inflammasome activation contributes to lung inflammation and fibrogenesis during IPF. In the current review, we discuss recent advances in our understanding of the DROSHA–miRNA–AIM2 inflammasome axis in the pathogenesis of IPF.

DROSHA

miRNA

AIM2 inflammasome


IPF

## 1. Introduction

### 1.1. The Role of DROSHA in IPF

DROSHA is a nuclease of the RNase III family [1][2]. As a core nuclease in nucleus, DROSHA plays a role in the initiation of miRNA maturation (Figure 1). DROSHA produces precursor miRNA (pre-RNA) via cleavage of primary miRNA (pri-miRNA) transcripts that are transported to cytoplasm and affect the regulation of molecular targets [3]. The DROSHA protein is localized into the nucleus by phosphorylation and leads to ultimate miRNA processing function [4]. Treatment with MG-132, a selective proteasome inhibitor, also increases the level of DROSHA, which suggests that DROSHA is regulated by post-translational modification [5]. DGCR8 (DiGeorge syndrome critical region 8), a cofactor of “Microprocessor” was reported to stabilize DROSHA through protein-protein interactions [6]. Under stress conditions, p38MAPK phosphorylates DROSHA, which decreases the interaction of DROSHA with DGCR8 [7]. The next critical step in miRNA biogenesis is pre-miRNA processing by DROSHA, which generates pre-miRNA that defines the miRNA sequences embedded in long pre-miRNAs [8][9]. Recently, we have reported for the first time that the protein expression of DROSHA was increased in alveolar macrophages of patients with IPF and mice with bleomycin-induced pulmonary fibrosis [10]. Conversely, DGCR8, a cofactor of DROSHA, protein expression was comparable between IPF and the control lungs [10]. After pre-miRNAs are exported to cytoplasm by exportin 5 in a Ran-GTP-dependent manner [11], Dicer, which is a well conserved RNase III endoribonuclease,

performs the further process in cytoplasm as the maturation of miRNAs makes progress [12]. Interestingly, Dicer was noted to be downregulated in fibroblasts from IPF lung [13]. This suggests that DROSHA, DGCR8, and Dicer play independent roles in IPF pathogenesis.




**Figure 1.** The maturation of miRNA by drosha ribonuclease III (DROSHA).

## 1.2. The Role of miRNA in IPF

DROSHA is involved in maturation or processing and not the biogenesis of microRNAs (miRNAs), small non-coding RNA molecules that have 18–25 nucleotides [14][15]. The miRNAs are expressed in the various types of organs and cells [8][9][16]. The miRNAs could change the inflammatory responses via the regulation of gene expression by post-transcriptional modifications that cause degradation and inhibition of the translation of the molecular target [8][9][16]. Recent studies have shown that mature miRNAs could be potential biomarkers for diagnosis in many human diseases such as cancers, cardiovascular disease, and neurodegenerative disease [17][18][19][20]. In a study of IPF, the changes of miRNAs were demonstrated in the analysis of miRNA arrays containing probes for 450 miRNAs with isolated RNA from the lungs from patients with IPF or the control [21]. It is reported that approximately 10% of miRNAs are significantly changed in human IPF lungs (Figure 2). Pandit et al. showed that downregulation of let-7d in IPF and inhibition of let-7d caused increases in profibrotic effects including the production of N-cadherin-2, vimentin,  $\alpha$ -SMA, and HMGA1. In addition, Eliot et al. reported that let-7 downregulates ligand-independent estrogen receptor (ER)-mediated pulmonary fibrosis and this study suggested that let-7 dependent ER expression has an important role in male-predominant fibrotic lung disease [22]. Additionally, in a murine fibrosis model, a relative abundance of miRNA levels in bleomycin-treated lung was demonstrated [23]. Wei et al. showed that TGF- $\beta$ 1-induced miR-133a inhibits myofibroblast differentiation and pulmonary fibrosis by

downregulating  $\alpha$ -SMA, CTGF, and collagens [24]. The targets of miR-133a are TGF- $\beta$  receptor 1, CTGF, and collagen type 1-a1. Bahudhanapati et al. reported that miR-144-3p is upregulated in IPF fibroblasts compared with the control fibroblasts [25]. miR-144-3p controls the expression of relaxin/insulin-like family peptide receptor 1(RXFP1) in fibroblasts from lungs with IPF [26]. Recent studies have identified the interaction between long noncoding RNAs (lncRNAs) and miRNAs. Jiang et al. showed that inhibition of pulmonary fibrosis-regulatory lncRNA (PFRL) prevents pulmonary fibrosis by disrupting the miR-26a/smad2 loop [26]. In addition, another lncRNA, pulmonary fibrosis-associated lncRNA (PFAL) was found to be upregulated during lung fibrosis in vitro and in vivo fibrosis models [27]. Mechanistically, PFRL promoted lung fibroblast activation by acting as a competing endogenous RNA for miR-18a. Liu et al. analyzed the function and regulatory mechanism of miR-708-3p and showed that miR-708-3p is decreased in IPF lungs [28]. They reported that miR-708-3p directly controls a disintegrin and metalloproteinase 17 (ADAM17) [28]. IPF is the most common interstitial lung fibrosis that shares similar pathways with other fibrotic disease such as scleroderma [29]. Several microRNAs including, but not limited to miR-29, miR-21-5p, and miR-26a-5p play important roles in lung fibrosis and disease progression [30][31][32][33][34][35].



**Figure 2.** The various roles of miRNA in the pathogenesis of idiopathic pulmonary fibrosis (IPF).

What are the targets of these miRNAs in the pathogenesis of lung fibrosis? Lung epithelial cells and fibroblasts play a critical role in the development and progression of lung fibrosis [36]. Let-7d, miR-29b, miR-26a, and miR-98 mediates antifibrotic effect by regulating myofibroblast activation and differentiation [21][34][37][38]. Conversely, miR-

21, miR-424, and miR-145 are pro-fibrotic by suppressing negative regulators of TGF- $\beta$  signaling [39][40][41]. Another important pathway that plays a central role in IPF development are aberrant inflammatory responses. There are several miRNAs that have been related to pulmonary fibrosis, out of which only miR-29 has been shown to regulate the innate immune response in vivo [37]. The administration of miR-29 attenuates both inflammatory response and fibrosis in the bleomycin induced lung fibrosis model. Currently, miRNAs have a critical role as key gene regulators that control fibrosis and inflammation, which in turn are potential targets for IPF disease treatment (Figure 2).

## 2. The Activation of Inflammasomes in IPF

### 2.1. The Role of Inflammasome-Dependent Inflammation in IPF

Alveolar macrophages are resident innate immune cells in the airways [42][43][44][45]. Alveolar macrophages play a key role in antimicrobial phagocytosis and in the pathogenesis of fibrotic lung disease as a critical type of cell in the lung [42][43][44][45]. Inflammasomes are multiprotein complexes, expressed in innate immune cells such as macrophages. Inflammasomes make inflammatory responses by the recognition of pathogen-associated molecular patterns (PAMPs), which come from bacterial, virus, and fungi, and danger-associated molecular patterns (DAMPs) that come from injured or dead cells [46][47]. The types of inflammasomes are distinguished by the combination of specific sensor proteins including NOD-, LRR-, and pyrin domain-containing 1 (NLRP1), Nod-like receptor protein 3 (NLRP3), NLRC4, or PYHIN (pyrin and HIN domain-containing protein) family members absent in melanoma 2 (AIM2), with the adaptor protein, adaptor apoptosis-associated speck-like protein containing a caspase recruitment domain (ASC), and the effector protein, caspase-1 [48][49][50]. The activation of NLRP3 inflammasome by the inflammasome complex formation is induced by various bacteria and viral particles, whereas AIM2 inflammasome is activated by double strand DNA (dsDNA) originating from bacteria, virus, and abnormal hosts [51][52][53][54][55]. Specific inflammasome complexes are defined by their subunits, and the precise conglomeration of various subunits to create these complexes depends on specific biologic activators [56][57]. Since the sensor proteins in inflammasomes are activated by specific activators, the sensor proteins interact with ASC and caspase-1 [56][57]. This interaction promotes the ASC speck formation and the cleavage of caspase-1, which are required for inflammasome activation [56][57]. Finally, the cleaved caspase-1 p10 fragment induces maturation and secretion of pro-inflammatory cytokines IL-1 $\beta$  and IL-18 [56][57].

Recent studies suggest a role for NLRP3 inflammasome activation in lung inflammation and fibrosis [58][59][60][61]. Elevated inflammasome activation has been reported to be augmented in human IPF lungs. Specifically, IL-18 protein levels were highly expressed in alveolar macrophages and epithelial cells [62]. Lasithiotaki et al. showed that transcriptional levels of inflammasome components, NLRP3 and caspase-1, were elevated in IPF patients when compared to the controls [63]. In addition, NLRP3 inflammasome activation by ATP was significantly increased in alveolar macrophages from IPF patients when compared to the controls [63]. Other studies have shown that inflammasome activation is present in other fibrotic lung diseases such as systemic sclerosis [64]. Fibroblasts from systemic sclerosis showed increased expression of multiple genes that was associated with the inflammasome or downstream signaling molecules [64]. Several studies have demonstrated a role for the NLRP3

inflammasome and its regulated cytokines in lung fibrosis in experimental lung fibrosis [60][65][66]. Given the increased prevalence of IPF in older populations, Stout-Delgado et al. investigated the role of NLRP3 inflammasome activation in age-dependent pulmonary fibrosis. Aging was found to be associated with the increased production of mitochondrial reactive oxygen species, which led to NLRP3 inflammasome activation and IL-1 $\beta$  and IL-18 secretion [65][67]. The role of NLRP3 inflammasome has also been tested in other pulmonary fibrosis models including ventilator induced lung fibrosis, silica-induced fibrosis, and particulate matter-induced fibrosis [60][66][68]. Although the NLRP3 inflammasome is mainly expressed in innate immune cells, there are several reports of inflammasome independent function of NLRP3 in murine fibrosis models. Lv et al. demonstrated that mechanical stretch promotes the activation of endothelial-mesenchymal transition (EMT) and lung fibrosis via NLRP3-mediated signaling cascade [66]. Tian et al. showed that the NLRP3 inflammasome is activated in alveolar epithelial cells and that the NLRP3 inflammasome may regulate EMT through TGF- $\beta$ 1 [69]. While the study of the NLRP3 inflammasome has been explored in various lung fibrosis models, the role of AIM2 inflammasome has not been fully investigated. Terlizzi et al. demonstrated that the activation of the AIM2 inflammasome is increased in peripheral blood mononuclear cells (PBMCs) from patients with IPF. The AIM2 inflammasome activation contributes to the production and release of pro-fibrotic mediators [70]. Recently, our findings demonstrated that the expression and activation of AIM2 inflammasome expression and activation is enhanced in a lung fibrosis exacerbation model [71]. Therefore, recent studies suggest that the activation of NLRP3 or AIM2 inflammasome-dependent inflammation has a critical role in the progression of lung fibrosis.

## 2.2. The Mechanism of Inflammasome Activation in IPF

Several studies have shown the upstream mechanism by which the NLRP3 inflammasome is activated in animal fibrosis models. Vimentin, the most abundant intermediate filament in the cytoplasm, regulates the activation of NLRP3 inflammasome in the asbestos-induced lung injury and fibrosis model [72]. Inflammation, endothelial, and alveolar epithelial barrier permeability and fibrosis are attenuated in vimentin deficient mice after asbestos challenge, and vimentin-knockdown macrophages showed decreased active caspase-1 and IL-1 $\beta$  levels. Furthermore, in this study, the direct protein-protein interaction between NLRP3 and vimentin was demonstrated. Doster et al. showed that NLRP3 inflammasome activation is triggered by reactive oxygen species (ROS), which are generated by nicotinamide adenine dinucleotide phosphate (NADPH) oxidase (NOX) upon particle phagocytosis [73]. Inhibition of NADPH oxidase suppressed NLRP3 inflammasome activation and NLRP3 deficient mice had reduced inflammation and fibrosis after asbestos exposure. ROS can be repressed by autophagy [74]. Meng et al. demonstrated that the activation of autophagy suppressed angiotensin II-induced pulmonary fibrosis via the inhibition of ROS-derived NLRP3 inflammasome activation. Another study reported similar findings that autophagic dysfunction in alveolar epithelial cells leads to subsequent lung fibrosis in a silica nanoparticle-induced fibrosis model [75]. Metabolic stimulation can also stimulate NLRP3 inflammasome activation in lung fibrosis models. Xu et al. described that statin use is associated with interstitial lung abnormalities among smokers in chronic obstructive pulmonary disease (COPD) Gene, a well-defined large cohort of smokers [76]. In addition, they reported that statin administration increases bleomycin-induced lung inflammation and fibrosis via the NLRP3 inflammasome activation in a mouse model. While the upstream regulation of NLRP3 inflammasome activation has been well described in lung fibrosis models, the mechanism by which AIM2 inflammasome is activated in lung

fibrosis has not been fully elucidated. Recently, we have described that glucose transport 1 (GLUT1)-dependent glycolysis regulates the exacerbation of lung fibrosis via AIM2 inflammasome activation [71]. We reported that GLUT1 deficiency ameliorates *S. pneumoniae*-mediated exacerbation of lung fibrosis in a mouse model [71]. In this study, we found that GLUT1 contributes to AIM2 inflammasome activation in macrophages [71]. We also found that the deficiency of GLUT1 suppressed the activation of AIM2 inflammasome in lung tissues under *S. pneumoniae*-mediated exacerbation of lung fibrosis [71].

## 3. Drosha ribonuclease III (DROSHA)-Dependent AIM2 Inflammasome Activation in IPF

### 3.1. The Activation of DROSHA in Alveolar Macrophages during IPF

While the changes of selective miRNA, which regulate differentiation, proliferation, and interaction between cell and cell, were identified and investigated in IPF, the role of DROSHA in alveolar macrophages, which are a critical cell type for pulmonary inflammation in IPF pathogenesis, remains unclear. The amount of evidence supports that macrophages play critical roles that affect fibrotic responses [77]. Alveolar macrophages produce transforming growth factor-beta 1 (TGF- $\beta$ 1), which promotes collagen accumulation, in both humans and mice [78][79]. These results suggest that a potential mechanism by which alveolar macrophages contribute to IPF. In addition to their role in TGF- $\beta$ 1 production, macrophages lead to the secretion of various inflammatory cytokines including TNF- $\alpha$ , IL-1, IL-6, IL-8, IL-10, and IL-12 [88]. Additionally, macrophages produce various chemokines such as CXCL1, CXCL2, CXCL9, CXCL10, CXCL12, CCL5, CCL17, and CCL18 [80]. Additionally, the macrophage-derived production of lipid mediators including eicosanoids might contribute to fibrosis [81]. On the other hand, the function of lipid mediators has not been fully identified, further investigation is needed in patients with IPF and a mouse model of lung fibrosis. Macrophages also regulate ECM remodeling by the secretion of matrix metalloproteinases and by the rearrangement of collagen [82][83]. From a different aspect, macrophages could determine the metabolic fate of environmental cells, which might conduct the glycolytic reprogramming in fibroblast during fibrosis [84][85]. Additionally, macrophages produce vascular endothelial growth factor (VEGF), which is related to both pro-fibrotic responses or anti-fibrotic responses [86][87][88]. It depends on the pattern of expression of angiogenic factors and the target cell. Therefore, macrophages are critical for the development of pulmonary fibrosis.

In other cells, the function of DROSHA has been reported [7][12][89]. In smooth muscle cells, the roles of DROSHA are required for cell survival [89]. Additionally, DROSHA regulates stress-induced death by phosphorylation of its N terminus via p38 MAPK [7]. The translocation of DROSHA by phosphorylation into cytosol from the nucleus is the regulatory mechanism of stress-induced death [12]. In the function of T cells, DROSHA is critical for T cell compartment. Deficiency of DROSHA results in T lymphopenia, particularly in the CD8+ compartment [12]. This study suggests that DROSHA-dependent miRNA production is required for the homeostasis of mature T cells [78]. In our recent study, we showed an increase in the DROSHA protein expression levels in alveolar macrophages during IPF [10]. Immunohistochemistry staining indicated that the protein levels of DROSHA were significantly increased in lung tissues from patients with IPF compared to non-IPF patients [10]. In particular, the intensity and number of DROSHA-positive staining in CD68-positive alveolar macrophages were significantly increased in lung

tissues from patients with IPF [10]. These results showed that the high levels of DROSHA in alveolar macrophages contributed to pulmonary inflammation in patients with IPF [10]. Similarly, the protein levels of DROSHA were elevated in alveolar macrophages in a mouse model of bleomycin-induced pulmonary fibrosis [10]. In other words, the expression levels of DGCR8 were not changed in lung tissues between patients with IPF and non-IPF patients [10]. Our results demonstrated that the elevation of DROSHA in alveolar macrophages is critical for pulmonary inflammation in IPF pathogenesis [10].

### 3.2. The Role of DROSHA in AIM2 Inflammasome Activation

Among the various inflammasomes, the mechanism of AIM2 inflammasome activation remains unclear. Currently, the representative mechanism of AIM2 inflammasome activation has been identified by the recognition of dsDNA by AIM2 [51][52][53][54][55] or the increase of AIM2 expression levels [90]. These two mechanisms are required for the AIM2 inflammasome complex formation, which causes the activation of caspase-1 p10 via cleavage of pro-caspase-1.

In our recent study, we suggest that DROSHA is an upstream regulator of AIM2 inflammasome activation [10]. We showed that the deficiency of DROSHA suppresses the activation of caspase-1 by AIM2 inflammasome activation in alveolar macrophages [10]. The deficiency of DROSHA reduces the apoptosis-associated speck-like protein containing a CARD (ASC) speck formation in the AIM2 inflammasome complex formation during AIM2 inflammasome activation [10]. These results suggest that DROSHA is critical for the activation of caspase-1 and AIM2 inflammasome complex formation during AIM2 inflammasome activation [10]. As further investigation of the molecular mechanism for DROSHA-dependent AIM2 inflammasome activation is conducted, the understanding of the cellular pathway related to the regulation of DROSHA expression could be important for the discovery of a new mechanism for AIM2 inflammasome activation. Previous studies have shown that the expression of DROSHA is tightly controlled by various cellular pathways such as post-translational modifications, protein degradation pathways, and alternative splicing [80][81][82][83][84]. Link et al. showed that the alternative splicing of DROSHA causes the alteration of the subcellular localization of DROSHA between the cytoplasm and nucleus [4][91]. They suggest that the cytosolic DROSHA potentially alters the function depending on the interaction with partners such as substrate RNAs or regulatory mechanisms [92][93]. Additionally, Ye et al. showed that E3 ubiquitin-protein ligase Mdm2 directly interacts with DROSHA [94]. Its interaction induces the ubiquitination of DROSHA, which causes the degradation of DROSHA protein [95]. Therefore, further investigation for the molecular mechanisms that affect DROSHA expression via alternative splicing or ubiquitination of DROSHA under IPF pathogenesis in AIM2 inflammasome activation is needed.


### 3.3. miRNA as DAMPs in AIM2 Inflammasome Activation

AIM2 inflammasome is activated by the recognition of cytosolic self-dsDNA as well as foreign dsDNA from pathogens [91]. AIM2 has been shown to detect dsDNA in a sequence-independent manner [54][55]. Since miRNAs, which are produced by DROSHA, have a hairpin structure with double-stranded RNA (dsRNA), miRNAs could make hetero-triplex structures in the location of specific sequences of DNA through the interaction with dsDNA [3].

Recent reports suggest that sub-cellular localization is critical to miRNA function [96]. Among the miRNA compartment, the localization of miRNA in the cytoplasm is important for RNA granules, endomembranes, and the export of miRNA to extracellular space [96]. Since cytosolic miRNAs have double-strand structure as well as dsDNA, cytosolic miRNA could be an activator of the AIM2 inflammasome. In our recent study, we found that the transduction of miRNAs in a sequence-independent manner promoted the activation of the AIM2 inflammasome and the cleavage of caspase-1 in macrophages [10]. Moreover, the transduction of miRNAs increased the maturation and secretion of IL-1 $\beta$  and IL-18 [10]. Furthermore, transduction of miRNAs increased the protein complex formation of AIM2 inflammasome by ASC speck formation, which is important for AIM2 inflammasome activation [10]. These results suggest that the double-stranded structure of cytosolic miRNAs induces the AIM2 inflammasome activation via AIM2 inflammasome complex formation (Figure 3).

DAMPs could contribute to lung inflammation and fibrosis during IPF. There are various types of DAMPs that come from different origins. The simple type of DAMPs might be intracellular components including nucleic acids such as DNA or RNA and organelles that are released from dying and injured cells. DAMPs might be released by cells through the transportation via the processes using endosomes and membrane bound vesicles. Among the various types of DAMPs, certain DAMP can be produced by the characteristic alteration of intracellular proteins such as the transformation from collagens to collagen fragments. These kinds of DAMPs are recognized by innate immune receptors and the activated pro-inflammatory signaling pathway. The activation of these immune receptors could be protective or harmful events, which is determined by the type of DAMP and DAMP-related specific receptor. In IPF, the role of DAMPs has been investigated in IL-17A receptor (IL-17RA)-mediated NF- $\kappa$ B signaling and the role of high mobility group box 1 (HMGB1) during inflammation and fibrosis [94][97]. IL-17A stimulates proliferation and survival of airway smooth muscle cells via the activation of IL-17RA [98]. The stimulation of IL-17RA leads to changes to various cellular pathways including cell proliferation, myofibroblast differentiation, and the production of extracellular matrix (ECM) proteins [94]. The levels of HMGB1 in BAL were increased in patients with IPF compared the control [97]. In bleomycin-induced pulmonary fibrosis in mice, HMGB1 protein levels were elevated in bronchiolar epithelial cells at early stage and in alveolar epithelial and inflammatory cells at late stage [97]. Recent study showed that the levels of miRNAs were increased in bronchoalveolar lavage (BAL) from patients with IPF relative to the controls [96]. These findings showed that the miRNA in BAL could contribute to damage-associated molecular pattern (DAMP)-dependent chronic lung inflammation in the progression of IPF [96][97]. In our recent study, we showed that the intracellular excessive accumulation of miRNA promotes AIM2 inflammasome-dependent caspase-1 activation in alveolar macrophages [10]. Moreover, the intracellular excessive accumulation of miRNA increases the ASC speck formation that is required for AIM2 inflammasome complex formation. Our results suggest that miRNA could be a critical DAMP for AIM2 inflammasome activation in pulmonary inflammation during IPF [10]. Based on our observation, the high levels of miRNA in BAL or blood in patients with chronic lung diseases might be a prognostic biomarker for the prediction of IPF.





**Figure 3.** The role of miRNA-dependent AIM2 inflammasome activation in idiopathic pulmonary fibrosis (IPF).

### 3.4. Therapeutic Approach of AIM2 Inflammasome Activation

Recent studies have shown that potential inhibitors have been identified as the NLRP3 inflammasome inhibitor [99][100][101][102][103][104]. The effects of potential NLRP3 inflammasome inhibitors such as OLT1177, Tranilast, Oridonin, CY-09, and MCC950 were demonstrated in cells and a representative mouse model for high fat diet (HFD)-induced diabetes, acute arthritis, and Parkinson's disease [99][100][101][102][103][104]. Each potential NLRP3 inflammasome inhibitor reduced the activation of NLRP3 inflammasome through the alteration of upstream pathways for NLRP3 inflammasome activation including mitochondrial dysfunction and potassium efflux [99][100][101][102][103][104]. Unlike the NLRP3 inflammasome, the understanding of potential inhibitor for AIM2 inflammasome activation is still unclear. Recent studies showed that the potential molecules that could inhibit AIM2 including pyrin-only proteins (POPs), CARD-only proteins (COPs), and TRIM11 [105][106]. In particular, pyrin-only proteins 1 (POP1) and POP3 suppressed the activation of AIM2 inflammasome via the interference of PYD-PYD interaction [107][108]. Additionally, IFI16- $\beta$ , as a new inhibitor of AIM2 inflammasome activation, was investigated for new knowledge for the design and development of anti-inflammatory agents [109]. In human chronic skin disease, cathelicidin peptide LL-37

inhibits the activation of the AIM2 inflammasome by interfering cytosolic DNA in keratinocytes [110]. This result suggests that cathelicidin LL-37 might be a therapeutic target of the AIM2 inflammasome to ameliorate cutaneous inflammation in chronic skin disease.

In cancer, increased expression of AIM2 has been reported in nasopharyngeal carcinoma tumors [111][112], oral squamous cell carcinoma [113], and lung adenocarcinoma [114]. On the other hand, AIM2 has been shown to suppress the development of cancer such as colorectal cancer [115][116]. Reduced expression and instability of the AIM2 have been reported in cancer tissues from patients with colorectal cancer [116][117][118]. The reduction of AIM2 expression is linked to a poorer prognosis in colorectal cancer patients [119]. Furthermore, the reduction of AIM2 expression was found in prostate cancer [120]. In different types of tumor tissue, the variation of AIM2 expression may play a role in the pathogenesis of various cancers. Since the role and activation of AIM2 are different in each type of cancer, the drug discovery for the therapeutic target of the AIM2 inflammasome should be conducted carefully.

Currently, the selective chemical inhibitor for the AIM2 inflammasome has not been identified as a therapeutic agent in human diseases such as IPF. In future investigations, the finding and validation of the selective chemical inhibitor for the AIM2 inflammasome might be important for the treatment of IPF.

---

## References

1. Valery Filippov; Victor Solovyev; Maria Filippova; Sarjeet S. Gill; A novel type of RNase III family proteins in eukaryotes. *Gene* **2000**, *245*, 213-221, 10.1016/s0378-1119(99)00571-5.
2. Hongjiang Wu; Hong Xu; Loren J. Miraglia; Stanley T. Crooke; Human RNase III Is a 160-kDa Protein Involved in Preribosomal RNA Processing. *Journal of Biological Chemistry* **2000**, *275*, 36957-36965, 10.1074/jbc.m005494200.
3. Yoontae Lee; Chiyong Ahn; Jinju Han; Hyounjeong Choi; Jaekwang Kim; Jeongbin Yim; Junho Lee; Patrick Provost; Olof Rådmark; Sunyoung Kim; V. Narry Kim; The nuclear RNase III Drosha initiates microRNA processing. *Nature* **2003**, *425*, 415-419, 10.1038/nature01957.
4. Xiaoli Tang; Yingjie Zhang; Lynne Tucker; Bharat Ramratnam; Phosphorylation of the RNase III enzyme Drosha at Serine300 or Serine302 is required for its nuclear localization.. *Nucleic Acids Research* **2010**, *38*, 6610-9, 10.1093/nar/gkq547.
5. Xiaoli Tang; Sicheng Wen; Ng Zheng; Lynne Tucker; Lulu Cao; Dennis Pantazatos; Steven F. Moss; Bharat Ramratnam; Acetylation of Drosha on the N-Terminus Inhibits Its Degradation by Ubiquitination. *PLOS ONE* **2013**, *8*, e72503, 10.1371/journal.pone.0072503.
6. Jinju Han; Yoontae Lee; Kyu-Hyeon Yeom; Jin-Wu Nam; Inha Heo; Je-Keun Rhee; Sun Young Sohn; Yunje Cho; Byoung-Tak Zhang; V. Narry Kim; Molecular Basis for the Recognition of

- Primary microRNAs by the Drosha-DGCR8 Complex. *Cell* **2006**, *125*, 887-901, 10.1016/j.cell.2006.03.043.
7. Qian Yang; Wenming Li; Hua She; Juan Dou; Duc M. Duong; Yuhong Du; Shao-Hua Yang; Nicholas T. Seyfried; Haiyan Fu; Guodong Gao; Zixu Mao; Stress induces p38 MAPK-mediated phosphorylation and inhibition of Drosha-dependent cell survival.. *Molecular Cell* **2015**, *57*, 721-734, 10.1016/j.molcel.2015.01.004.
  8. Bonnie Bartel; MicroRNAs: genomics, biogenesis, mechanism, and function.. *Cell* **2004**, *116*, , null.
  9. Young Kook Kim; Boseon Kim; V. Narry Kim; Re-evaluation of the roles of DROSHA, Exportin 5, and DICER in microRNA biogenesis.. *Proceedings of the National Academy of Sciences* **2016**, *113*, E1881-9, 10.1073/pnas.1602532113.
  10. Soo Jung Cho; Kyoung Sook Hong; Ji Hun Jeong; Mihye Lee; Augustine M. K. Choi; Heather W. Stout-Delgado; Jong-Seok Moon; DROSHA-Dependent AIM2 Inflammasome Activation Contributes to Lung Inflammation during Idiopathic Pulmonary Fibrosis.. *Cells* **2019**, *8*, 938, 10.3390/cells8080938.
  11. Markus T. Bohnsack; Kevin Czaplinski; Dirk Görlich; Exportin 5 is a RanGTP-dependent dsRNA-binding protein that mediates nuclear export of pre-miRNAs.. *RNA* **2004**, *10*, , null.
  12. Mark M.W. Chong; Jeffrey P. Rasmussen; Alexander Y. Rudensky; Dan R. Littman; The RNaseIII enzyme Drosha is critical in T cells for preventing lethal inflammatory disease. *Journal of Experimental Medicine* **2008**, *205*, 2449-2449, 10.1084/jem.20071219090508c.
  13. Jeremy Herrera; Daniel J. Beisang; Mark Peterson; Colleen Forster; Adam Gilbertsen; Alexey Benyumov; Karen Smith; Christopher E. Korenczuk; Victor H. Barocas; Kacey Guenther; Ryan Hite; Lin Zhang; Craig A. Henke; Peter Bitterman; Dicer1 Deficiency in the Idiopathic Pulmonary Fibrosis Fibroblastic Focus Promotes Fibrosis by Suppressing MicroRNA Biogenesis.. *American Journal of Respiratory and Critical Care Medicine* **2018**, *198*, 486-496, 10.1164/rccm.201709-1823OC.
  14. Valentina Libri; Pascal Miesen; Ronald P. Van Rij; Amy H Buck; Regulation of microRNA biogenesis and turnover by animals and their viruses. *Cellular and Molecular Life Sciences* **2013**, *70*, 3525-44, 10.1007/s00018-012-1257-1.
  15. Neil R. Smalheiser; Regulation of mammalian microRNA processing and function by cellular signaling and subcellular localization. *Biochimica et Biophysica Acta (BBA) - Molecular Cell Research* **2008**, *1779*, 678-81, 10.1016/j.bbagrm.2008.03.009.
  16. Eric C. Lai; Micro RNAs are complementary to 3' UTR sequence motifs that mediate negative post-transcriptional regulation. *Nature Genetics* **2002**, *30*, 363-364, 10.1038/ng865.

17. Shan-Shan Zhou; Jing-Peng Jin; Ji-Qun Wang; Zhi-Guo Zhang; Jonathan H Freedman; Yang Zheng; Lu Cai; miRNAs in cardiovascular diseases: potential biomarkers, therapeutic targets and challenges. *Acta Pharmacologica Sinica* **2018**, *39*, 1073-1084, 10.1038/aps.2018.30.
18. Samantha Filipów; Łukasz Łaczmański; Blood Circulating miRNAs as Cancer Biomarkers for Diagnosis and Surgical Treatment Response.. *Frontiers in Genetics* **2019**, *10*, 169, 10.3389/fgen e.2019.00169.
19. Rimi Hamam; Dana Hamam; Khalid A Alsaleh; Moustapha Kassem; Waleed Zaher; MUSAAD Alfayez; Abdullah Aldahmash; Nehad M. Alajez; Circulating microRNAs in breast cancer: novel diagnostic and prognostic biomarkers.. *Cell Death & Disease* **2017**, *8*, e3045-e3045, 10.1038/cddi s.2017.440.
20. Emmette R. Hutchison; Eitan Okun; Mark P. Mattson; The Therapeutic Potential of microRNAs in Nervous System Damage, Degeneration, and Repair. *NeuroMolecular Medicine* **2009**, *11*, 153-61, 10.1007/s12017-009-8086-x.
21. Kusum Pandit; David L. Corcoran; Hanadie Yousef; Manohar Yarlagadda; Argyris Tzouveleakis; Kevin F. Gibson; Kazuhisa Konishi; Samuel A. Yousem; Mandal Singh; Daniel Handley; Thomas Richards; Moises Selman; Simon C. Watkins; Annie Pardo; Ahmi Ben-Yehudah; Demosthenes Bouros; Oliver Eickelberg; Prabir Ray; Panayiotis V. Benos; Naftali Kaminski; Inhibition and Role of let-7d in Idiopathic Pulmonary Fibrosis. *American Journal of Respiratory and Critical Care Medicine* **2010**, *182*, 220-229, 10.1164/rccm.200911-1698OC.
22. Sharon Elliot; Simone Periera-Simon; Xiaomei Xia; Paola Catanuto; Gustavo Rubio; Shahriar Shahzeidi; Fadi El Salem; Josh Shapiro; Karoline Briegel; Kenneth S. Korach; Marilyn K. Glassberg; MicroRNA let-7 Downregulates Ligand-Independent Estrogen Receptor-mediated Male-Predominant Pulmonary Fibrosis. *American Journal of Respiratory and Critical Care Medicine* **2019**, *200*, 1246-1257, 10.1164/rccm.201903-0508oc.
23. Ting Xie; Jiurong Liang; Rishu Guo; Ningshan Liu; Paul W. Noble; Dianhua Jiang; Comprehensive microRNA analysis in bleomycin-induced pulmonary fibrosis identifies multiple sites of molecular regulation.. *Physiological Genomics* **2011**, *43*, 479-87, 10.1152/physiolgenomics.00222.2010.
24. Peng Wei; Yan Xie; Peter W. Abel; Yapei Huang; Qin Ma; Linghai Li; Junfeng Hao; Dennis W. Wolff; Taotao Wei; Yaping Tu; Transforming growth factor (TGF)- $\beta$ 1-induced miR-133a inhibits myofibroblast differentiation and pulmonary fibrosis.. *Cell Death & Disease* **2019**, *10*, 670-17, 10.1038/s41419-019-1873-x.
25. Harinath Bahudhanapati; Jiangning Tan; Justin Dutta; Stephen B. Strock; John Sembrat; Diana Álvarez; Mauricio Rojas; Benedikt Jäger; Antje Prasse; Yingze Zhang; Daniel J. Kass; MicroRNA-144-3p targets relaxin/insulin-like family peptide receptor 1 (RXFP1) expression in lung fibroblasts from patients with idiopathic pulmonary fibrosis. *Journal of Biological Chemistry* **2019**, *294*, 5008-5022, 10.1074/jbc.ra118.004910.

26. Hua Jiang; Yingzhun Chen; Tong Yu; Xiaoguang Zhao; Huitong Shan; Jian Sun; Lu Zhang; Xuelian Li; Hongli Shan; HaiHai Liang; Inhibition of lncRNA PFRL prevents pulmonary fibrosis by disrupting the miR-26a/smad2 loop. *American Journal of Physiology-Lung Cellular and Molecular Physiology* **2018**, *315*, L563-L575, 10.1152/ajplung.00434.2017.
27. Li Xuelian; Tong Yu; Huitong Shan; Hua Jiang; Jian Sun; Xiaoguang Zhao; Wei Su; Lida Yang; Hongli Shan; HaiHai Liang; lncRNA PFAL promotes lung fibrosis through CTGF by competitively binding miR-18a. *The FASEB Journal* **2018**, *32*, 5285-5297, 10.1096/fj.201800055r.
28. Bo Liu; Rongrong Li; Jinjin Zhang; Chao Meng; Jie Zhang; Xiaodong Song; Changjun Lv; MicroRNA-708-3p as a potential therapeutic target via the ADAM17-GATA/STAT3 axis in idiopathic pulmonary fibrosis. *Experimental & Molecular Medicine* **2018**, *50*, e465-e465, 10.1038/emm.2017.311.
29. Gianluca Bagnato; William Neal Roberts; Jesse Roman; Sebastiano Gangemi; A systematic review of overlapping microRNA patterns in systemic sclerosis and idiopathic pulmonary fibrosis. *European Respiratory Review* **2017**, *26*, 160125, 10.1183/16000617.0125-2016.
30. Wajahat Khalil; Hong Xia; Vidya Bodempudi; Judy Kahm; Polla Hergert; Karen Smith; Mark Peterson; Matthew Parker; Jeremy Herrera; Peter Bitterman; Craig A. Henke; Pathologic Regulation of Collagen I by an Aberrant Protein Phosphatase 2A/Histone Deacetylase C4/MicroRNA-29 Signal Axis in Idiopathic Pulmonary Fibrosis Fibroblasts. *American Journal of Respiratory Cell and Molecular Biology* **2015**, *53*, 391-399, 10.1165/rcmb.2014-0150OC.
31. Matthew W. Parker; Daniel Rossi; Mark Peterson; Karen Smith; Kristina Sikström; Eric S. White; John E. Connett; Craig A. Henke; Ola Larsson; Peter Bitterman; Fibrotic extracellular matrix activates a profibrotic positive feedback loop.. *Journal of Clinical Investigation* **2014**, *124*, 1622-35, 10.1172/JCI71386.
32. Gabriela Díaz-Piña; Rosa María Ordoñez-Razo; Eduardo Montes; Ignacio Páramo; Carina Becerril; Alfonso Salgado; J. Alfredo Santibáñez-Salgado; Mariel Maldonado; Victor Ruiz; The Role of ADAR1 and ADAR2 in the Regulation of miRNA-21 in Idiopathic Pulmonary Fibrosis. *Lung* **2018**, *196*, 393-400, 10.1007/s00408-018-0115-9.
33. Gang Liu; Arnaud Friggeri; Jadranka Milosevic; Qiang Ding; Victor J. Thannickal; Naftali Kaminski; Edward Abraham; Yanping Yang; miR-21 mediates fibrogenic activation of pulmonary fibroblasts and lung fibrosis. *The Journal of Cell Biology* **2010**, *190*, , 10.1083/jcb1902oia3.
34. H Liang; Y Gu; T Li; Y Zhang; L Huangfu; M Hu; D Zhao; Y Chen; S Liu; Y Dong; X Li; Y Lu; B Yang; H Shan; Integrated analyses identify the involvement of microRNA-26a in epithelial-mesenchymal transition during idiopathic pulmonary fibrosis.. *Cell Death & Disease* **2014**, *5*, e1238-e1238, 10.1038/cddis.2014.207.
35. HaiHai Liang; Chaoqian Xu; Zhenwei Pan; Ying Zhang; Zhidan Xu; Yingzhun Chen; Tianyu Li; Xuelian Li; Ying Liu; Longtao Huangfu; Ying Lu; Zhihua Zhang; Baofeng Yang; Samuel Gitau;

- Yanjie Lu; Hongli Shan; Zhimin Du; The Antifibrotic Effects and Mechanisms of MicroRNA-26a Action in Idiopathic Pulmonary Fibrosis. *Molecular Therapy* **2014**, 22, 1122-1133, 10.1038/mt.2014.42.
36. Subbiah Rajasekaran; P. Rajaguru; P. S. Sudhakar Gandhi; MicroRNAs as potential targets for progressive pulmonary fibrosis. *Frontiers in Pharmacology* **2015**, 6, 473, 10.3389/fphar.2015.00254.
37. Toby Maher; Montgomery RI; Yu G; Latimer Pa; Stack C; Robinson K; Dalby Cm; Kaminski N; Van Rooij E; Faculty of 1000 evaluation for MicroRNA mimicry blocks pulmonary fibrosis.. *F1000 - Post-publication peer review of the biomedical literature* **2016**, 6, , 10.3410/f.718883296.793518552.
38. Shu-Yan Gao; Xue Zhou; You-Jie Li; Wei-Li Liu; Ping-Yu Wang; Min Pang; Shu-Yang Xie; Chang-Jun Lv; Arsenic trioxide prevents rat pulmonary fibrosis via miR-98 overexpression. *Life Sciences* **2014**, 114, 20-28, 10.1016/j.lfs.2014.07.037.
39. Mitsuhiro Yamada; Hiroshi Kubo; Chiharu Ota; Toru Takahashi; Yukiko Tando; Takaya Suzuki; Naoya Fujino; Tomonori Makiguchi; Kiyoshi Takagi; Takashi Suzuki; Masakazu Ichinose; The increase of microRNA-21 during lung fibrosis and its contribution to epithelial-mesenchymal transition in pulmonary epithelial cells. *Respiratory Research* **2013**, 14, 95-95, 10.1186/1465-9921-14-95.
40. Xiao Xiao; Chaoqun Huang; Chunling Zhao; Xuxu Gou; Lakmini K. Senavirathna; Myron Hinsdale; Pamela Lloyd; Lin Liu; Regulation of myofibroblast differentiation by miR-424 during epithelial-to-mesenchymal transition. *Archives of Biochemistry and Biophysics* **2015**, 566, 49-57, 10.1016/j.abb.2014.12.007.
41. Shanzhong Yang; Huachun Cui; N. Xie; Mert Icyuz; Sami Banerjee; Veena B. Antony; Edward Abraham; Victor J. Thannickal; Gang Liu; miR-145 regulates myofibroblast differentiation and lung fibrosis. *The FASEB Journal* **2013**, 27, 2382-2391, 10.1096/fj.12-219493.
42. P Bhattacharyya; D Acharya; S Roychowdhury; Role of matrix metalloproteinases in the pathophysiology of idiopathic pulmonary fibrosis. *Lung India* **2007**, 24, 61, 10.4103/0970-2113.44213.
43. Thomas A Wynn; Integrating mechanisms of pulmonary fibrosis. *The Journal of Experimental Medicine* **2011**, 208, 1339-1350, 10.1084/jem.20110551.
44. Byrne, A.J.; Maher, T.M.; Lloyd, C.M. Pulmonary Macrophages: A New Therapeutic Pathway in Fibrosing Lung Disease? *Trends Mol. Med.* 2016, 22, 303–316.
45. Erin Addis-Lieser; Jörg Köhl; Mónica G. Chiaramonte; Opposing regulatory roles of complement factor 5 in the development of bleomycin-induced pulmonary fibrosis.. *The Journal of Immunology* **2005**, 175, 1894-1902, 10.4049/jimmunol.175.3.1894.

46. Junya Masumoto; "The inflammasomes". *Japanese Journal of Clinical Immunology* **2011**, *34*, 346-354, 10.2177/jsci.34.346.
47. Petr Broz; Vishva M. Dixit; Inflammasomes: mechanism of assembly, regulation and signalling. *Nature Reviews Immunology* **2016**, *16*, 407-420, 10.1038/nri.2016.58.
48. Greta Guarda; Alexander So; Regulation of inflammasome activity. *Immunology* **2010**, *130*, 329-336, 10.1111/j.1365-2567.2010.03283.x.
49. T Fernandes-Alnemri; J Wu; Je-Wook Yu; P Datta; B Miller; W Jankowski; S Rosenberg; Jianke Zhang; E S Alnemri; The pyroptosome: a supramolecular assembly of ASC dimers mediating inflammatory cell death via caspase-1 activation. *Cell Death & Differentiation* **2007**, *14*, 1590-604, 10.1038/sj.cdd.4402194.
50. Jenny Ting; Martinon F; Burns K; Tschopp J; Faculty of 1000 evaluation for The inflammasome: a molecular platform triggering activation of inflammatory caspases and processing of proIL-beta.. *F1000 - Post-publication peer review of the biomedical literature* **2002**, *10*, , 10.3410/f.1008963.128907.
51. Mohamed Lamkanfi; Vishva M. Dixit; Inflammasomes and Their Roles in Health and Disease. *Annual Review of Cell and Developmental Biology* **2012**, *28*, 137-161, 10.1146/annurev-cellbio-101011-155745.
52. Eun-Kyeong Jo; Jin Kyung Kim; Ng-Min Shin; Chihiro Sasakawa; Molecular mechanisms regulating NLRP3 inflammasome activation. *Cellular & Molecular Immunology* **2015**, *13*, 148-159, 10.1038/cmi.2015.95.
53. Si Ming Man; Rajendra Karki; Thirumala-Devi Kanneganti; AIM2 inflammasome in infection, cancer, and autoimmunity: Role in DNA sensing, inflammation, and innate immunity.. *European Journal of Immunology* **2015**, *46*, 269-80, 10.1002/eji.201545839.
54. Veit K Hornung; Andrea Ablasser; Marie Charrel-Dennis; Franz Bauernfeind; Gabor L. Horvath; Daniel. R. Caffrey; Eicke Latz; Katherine A. Fitzgerald; AIM2 recognizes cytosolic dsDNA and forms a caspase-1-activating inflammasome with ASC. *Nature* **2009**, *458*, 514-8, 10.1038/nature07725.
55. Vijay A.K. Rathinam; Zhaozhao Jiang; Stephen N. Waggoner; Shruti Sharma; Leah E. Cole; Lisa Waggoner; Sivapriya Kailasan Vanaja; Brian G. Monks; Sandhya Ganesan; Eicke Latz; Veit Hornung; Stefanie N. Vogel; Eva Szomolanyi-Tsuda; Katherine A. Fitzgerald; The AIM2 inflammasome is essential for host defense against cytosolic bacteria and DNA viruses. *Nature Immunology* **2010**, *11*, 395-402, 10.1038/ni.1864.
56. Martina Proell; Motti Gerlic; Peter D. Mace; John C. Reed; Stefan J. Riedl; The CARD plays a critical role in ASC foci formation and inflammasome signalling. *Biochemical Journal* **2013**, *449*, 613-621, 10.1042/bj20121198.

57. Parimala R. Vajjhala; Ruth Mirams; Justine M. Hill; Multiple Binding Sites on the Pyrin Domain of ASC Protein Allow Self-association and Interaction with NLRP3 Protein\*. *Journal of Biological Chemistry* **2012**, *287*, 41732-41743, 10.1074/jbc.M112.381228.
58. Pamela Gasse; Nicolas Riteau; Sabine Charron; Sandra Girre; Lizette Fick; Virginie Petrilli; Jurg Tschopp; Vincent Lagente; Valerie F. J. Quesniaux; Bernhard Ryffel; Isabelle Couillin; Uric Acid Is a Danger Signal Activating NALP3 Inflammasome in Lung Injury Inflammation and Fibrosis. *American Journal of Respiratory and Critical Care Medicine* **2009**, *179*, 903-913, 10.1164/rccm.200808-1274oc.
59. Paméla Gasse; Caroline Mary; Isabelle Guenon; Nicolas Noulin; Sabine Charron; Silvia Schnyder-Candrian; Bruno Schnyder; Shizuo Akira; Valérie F.J. Quesniaux; Vincent Lagente; Bernhard Ryffel; Isabelle Couillin; IL-1R1/MyD88 signaling and the inflammasome are essential in pulmonary inflammation and fibrosis in mice. *Journal of Clinical Investigation* **2007**, *117*, 3786-3799, 10.1172/JCI32285.
60. Suzanne L. Cassel; Stephanie C. Eisenbarth; Shankar S. Iyer; Jeffrey J. Sadler; Oscar R. Colegio; Linda A. Tephly; A. Brent Carter; Paul B. Rothman; Richard A. Flavell; Fayyaz S. Sutterwala; The Nalp3 inflammasome is essential for the development of silicosis. *Proceedings of the National Academy of Sciences* **2008**, *105*, 9035-9040, 10.1073/pnas.0803933105.
61. Jianbo Wu; Zhibo Yan; David E. Schwartz; Jingui Yu; Asrar B. Malik; Guochang Hu; Activation of NLRP3 inflammasome in alveolar macrophages contributes to mechanical stretch-induced lung inflammation and injury.. *The Journal of Immunology* **2013**, *190*, 3590-9, 10.4049/jimmunol.1200860.
62. Yasuhiko Kitasato; Tomoaki Hoshino; Masaki Okamoto; Seiya Kato; Yoshiro Koda; Nobuhiko Nagata; Masaharu Kinoshita; Hideyuki Koga; -Young Yoon; Hironobu Asao; Hiroshi Ohmoto; Takeharu Koga; Toru Rikimaru; Hisamichi Aizawa; Enhanced Expression of Interleukin-18 and its Receptor in Idiopathic Pulmonary Fibrosis. *American Journal of Respiratory Cell and Molecular Biology* **2004**, *31*, 619-625, 10.1165/rcmb.2003-0306oc.
63. Ismini Lasithiotaki; Ioannis Giannarakis; Eliza Tsitoura; Katerina Samara; George A. Margaritopoulos; Christiana Choulaki; Eirini Vasarmidi; Nikolaos Tzanakis; Argyro Voloudaki; Prodromos Sidiropoulos; Nikolaos M. Siafakas; Katerina M. Antoniou; NLRP3 inflammasome expression in idiopathic pulmonary fibrosis and rheumatoid lung. *European Respiratory Journal* **2016**, *47*, 910-918, 10.1183/13993003.00564-2015.
64. Carol M. Artlett; Sihem Sassi-Gaha; Judy L. Rieger; Alina C. Boesteanu; Carol A. Feghali-Bostwick; Peter Katsikis; The inflammasome activating caspase 1 mediates fibrosis and myofibroblast differentiation in systemic sclerosis. *Arthritis & Rheumatism* **2011**, *63*, 3563-3574, 10.1002/art.30568.



65. Heather W. Stout-Delgado; Soo Jung Cho; Sarah G. Chu; Dana N. Mitzel; Julian Villalba; Souheil El-Chemaly; Stefan W. Ryter; Augustine M. K. Choi; Ivan O. Rosas; Age-Dependent Susceptibility to Pulmonary Fibrosis Is Associated with NLRP3 Inflammasome Activation. *American Journal of Respiratory Cell and Molecular Biology* **2016**, *55*, 252-263, 10.1165/rcmb.2015-0222OC.
66. Zhou Lv; Yan Wang; Yujian Liu; Yanfei Mao; Wenwen Dong; Zhong-Nuo Ding; Guang-Xun Meng; Lai Jiang; Xiaoyan Zhu; NLRP3 Inflammasome Activation Contributes to Mechanical Stretch-Induced Endothelial-Mesenchymal Transition and Pulmonary Fibrosis. *Critical Care Medicine* **2018**, *46*, e49-e58, 10.1097/ccm.0000000000002799.
67. W Bohn; [Modifications of the pulmonary hilus due to age].. *Virchows Archiv* **1950**, *318*, , null.
68. Runxiao Zheng; Lan Tao; Hui Jian; Yun Chang; Yan Cheng; Yanlin Feng; Haiyuan Zhang; NLRP3 inflammasome activation and lung fibrosis caused by airborne fine particulate matter. *Ecotoxicology and Environmental Safety* **2018**, *163*, 612-619, 10.1016/j.ecoenv.2018.07.076.
69. Rui Tian; Yong Zhu; Jiayi Yao; Xiaoxiao Meng; Jinfeng Wang; Hui Xie; Lina Zhao; NLRP3 participates in the regulation of EMT in bleomycin-induced pulmonary fibrosis. *Experimental Cell Research* **2017**, *357*, 328-334, 10.1016/j.yexcr.2017.05.028.
70. Michela Terlizzi; Antonio Molino; Chiara Colarusso; Chantal Donovan; Pasquale Imitazione; Pasquale Somma; Rita Aquino; Philip M. Hansbro; Aldo Pinto; Rosalinda Sorrentino; Activation of the Absent in Melanoma 2 Inflammasome in Peripheral Blood Mononuclear Cells From Idiopathic Pulmonary Fibrosis Patients Leads to the Release of Pro-Fibrotic Mediators. *Frontiers in Immunology* **2018**, *9*, , 10.3389/fimmu.2018.00670.
71. Soo Jung Cho; Jong-Seok Moon; Kiichi Nikahira; Ha Seon Yun; Rebecca Harris; Kyung Sook Hong; Huarong Huang; Augustine M K Choi; Heather Stout-Delgado; GLUT1-dependent glycolysis regulates exacerbation of fibrosis via AIM2 inflammasome activation. *Thorax* **2019**, *75*, 227-236, 10.1136/thoraxjnl-2019-213571.
72. Gimena Dos Santos; Micah R. Rogel; Margaret A. Baker; James R. Troken; Daniela Urich; Luisa Morales-Nebreda; Joseph A. Sennello; Mikhail Kutuzov; Albert Sitikov; Jennifer M. Davis; Anna P. Lam; Paul Cheresch; David Kamp; Dale K. Shumaker; G. R. Scott Budinger; Karen M. Ridge; Vimentin regulates activation of the NLRP3 inflammasome.. *Nature Communications* **2015**, *6*, 6574-6574, 10.1038/ncomms7574.
73. Catherine Dostert; Virginie Petrilli; Robin Van Bruggen; Chad Steele; Brooke T. Mossman; Jürg Tschopp; Innate Immune Activation Through Nalp3 Inflammasome Sensing of Asbestos and Silica. *Science* **2008**, *320*, 674-677, 10.1126/science.1156995.
74. Ying Meng; Miao-Xia Pan; Bo-Jun Zheng; Yan Chen; Wei Li; Qian-Jie Yang; Ze-Mao Zheng; Na-Na Sun; Yue Zhang; Xu Li; Autophagy Attenuates Angiotensin II-Induced Pulmonary Fibrosis by Inhibiting Redox Imbalance-Mediated NOD-Like Receptor Family Pyrin Domain Containing 3

- Inflammasome Activation. *Antioxidants & Redox Signaling* **2019**, *30*, 520-541, 10.1089/ars.2017.7261.
75. Xinyuan Zhao; Saisai Wei; Zhijian Li; Chen Lin; Zhenfeng Zhu; Desen Sun; Rongpan Bai; Jun Qian; Xiangwei Gao; Guangdi Chen; ZhengPing Xu; Autophagic flux blockage in alveolar epithelial cells is essential in silica nanoparticle-induced pulmonary fibrosis. *Cell Death & Disease* **2019**, *10*, 127, 10.1038/s41419-019-1340-8.
76. Xu, J.F.; Washko, G.R.; Nakahira, K.; Hatabu, H.; Patel, A.S.; Fernandez, I.E.; Nishino, M.; Okajima, Y.; Yamashiro, T.; Ross, J.C.; et al. Statins and pulmonary fibrosis: The potential role of NLRP3 inflammasome activation. *Am. J. Respir. Crit. Care Med.* 2012, *185*, 547–556.
77. Margaret L. Novak; Timothy J. Koh; Macrophage phenotypes during tissue repair.. *Journal of Leukocyte Biology* **2013**, *93*, 875-81, 10.1189/jlb.1012512.
78. Lisa R. Young; Peter M. Gulleman; Chelsi W. Short; Harikrishna Tanjore; Taylor Sherrill; Aidong Qi; Andrew P. McBride; Rinat Zaynagetdinov; John T. Benjamin; William E. Lawson; Sergey V. Novitskiy; Timothy S. Blackwell; Epithelial-macrophage interactions determine pulmonary fibrosis susceptibility in Hermansky-Pudlak syndrome.. *JCI Insight* **2016**, *1*, e88947, 10.1172/jci.insight.88947.
79. Z Toossi; C S Hirsch; B D Hamilton; C K Knuth; M A Friedlander; E A Rich; Decreased production of TGF-beta 1 by human alveolar macrophages compared with blood monocytes.. *The Journal of Immunology* **1996**, *156*, , null.
80. Guillermo Arango Duque; Albert Descoteaux; Macrophage Cytokines: Involvement in Immunity and Infectious Diseases. *Frontiers in Immunology* **2014**, *5*, , 10.3389/fimmu.2014.00491.
81. Huang, S.K.; Peters-Golden, M. Eicosanoid lipid mediators in fibrotic lung diseases: Ready for prime time? *Chest* 2008, *133*, 1442–1450.
82. MarYam G. Rohani; Ryan S. McMahan; Maria V. Razumova; Angie L. Hertz; Maryelise Cieslewicz; Suzie H. Pun; Michael Regnier; Ying Wang; Timothy P. Birkland; William C. Parks; MMP-10 Regulates Collagenolytic Activity of Alternatively Activated Resident Macrophages.. *Journal of Investigative Dermatology* **2015**, *135*, 2377-2384, 10.1038/jid.2015.167.
83. Daniel H. Madsen; Daniel Leonard; Andrius Masedunskas; Amanda Moyer; Henrik Jessen Jürgensen; Diane E. Peters; Panomwat Amornphimoltham; Arul Selvaraj; Susan S. Yamada; David A. Brenner; Sven Burgdorf; Lars Engelholm; Niels Behrendt; Kenn Holmbeck; Roberto Weigert; Thomas H. Bugge; M2-like macrophages are responsible for collagen degradation through a mannose receptor–mediated pathway. *The Journal of Cell Biology* **2013**, *202*, 951-966, 10.1083/jcb.201301081.
84. Subhra K. Biswas; Alberto Mantovani; Orchestration of Metabolism by Macrophages. *Cell Metabolism* **2012**, *15*, 432-437, 10.1016/j.cmet.2011.11.013.

85. Na Xie; Zheng Tan; Sami Banerjee; Huachun Cui; Jing Ge; Rui-Ming Liu; Karen Bernard; Victor J. Thannickal; Gang Liu; Glycolytic Reprogramming in Myofibroblast Differentiation and Lung Fibrosis. *American Journal of Respiratory and Critical Care Medicine* **2015**, *192*, 1462-1474, 10.1164/rccm.201504-0780oc.
86. Christian Stockmann; Yann M. Kerdiles; Marc Nomaksteinsky; Alexander Weidemann; Norihiko Takeda; Andrew Doedens; Antoni Xavier Torres-Collado; Luisa Iruela-Arispe; Victor Nizet; Randall S. Johnson; Loss of myeloid cell-derived vascular endothelial growth factor accelerates fibrosis. *Proceedings of the National Academy of Sciences* **2010**, *107*, 4329-4334, 10.1073/pnas.0912766107.
87. Shaney L. Barratt; Thomas Blythe; Caroline Jarrett; Khadija Ourradi; Golda Shelley-Fraser; Michael J. Day; Yan Qiu; Steve Harper; Toby M. Maher; Sebastian Oltean; Thomas J. Hames; Chris J Scotton; Gavin Iain Welsh; David O. Bates; Ann B. Millar; Differential Expression of VEGF-AxxxIsoforms Is Critical for Development of Pulmonary Fibrosis. *American Journal of Respiratory and Critical Care Medicine* **2017**, *196*, 479-493, 10.1164/rccm.201603-0568oc.
88. Shun Morizumi; Seidai Sato; Shuichi Abe; Hiroyasu Okazaki; Yanjuan Chen; Hisatsugu Goto; Yoshinori Aono; Hirohisa Ogawa; Masaki Hanibuchi; Hisanori Uehara; Yasuhiko Nishioka; Anti-fibrotic efficacy of nintedanib on pulmonary fibrosis via suppression of fibrocyte activity. *Diffuse Parenchymal Lung Disease* **2016**, *48*, , 10.1183/13993003.congress-2016.pa778.
89. Pei Fan; Zixuan Chen; Peng Tian; Wen Liu; Yan Jiao; Yi Xue; Anindya Bhattacharya; Jianmin Wu; Meifen Lu; Yuqi Guo; Yan Cui; Weikuan Gu; Weiwang Gu; Junming Yue; miRNA Biogenesis Enzyme Drosha Is Required for Vascular Smooth Muscle Cell Survival. *PLOS ONE* **2013**, *8*, e60888, 10.1371/journal.pone.0060888.
90. Jérôme Lugin; Fabio Martinon; The AIM2 inflammasome: Sensor of pathogens and cellular perturbations. *Immunological Reviews* **2017**, *281*, 99-114, 10.1111/imr.12618.
91. Anthony K.L. Leung; The Whereabouts of microRNA Actions: Cytoplasm and Beyond.. *Trends in Cell Biology* **2015**, *25*, 601-610, 10.1016/j.tcb.2015.07.005.
92. Steffen Link; Stefanie E. Grund; Sven Diederichs; Alternative splicing affects the subcellular localization of Drosha.. *Nucleic Acids Research* **2016**, *44*, 5330-43, 10.1093/nar/gkw400.
93. Lisheng Dai; Kevin Chen; Brenda Youngren; Julia Kulina; Acong Yang; Zhengyu Guo; Jin Li; Peng Yu; Shuo Gu; Cytoplasmic Drosha activity generated by alternative splicing.. *Nucleic Acids Research* **2016**, *44*, 10454-10466, 10.1093/nar/gkw668.
94. Jie Zhang; Dan Wang; Lei Wang; Shaohua Wang; Anja C. Roden; Hao Zhao; Xiujuan Li; Y. S. Prakash; Eric L. Matteson; Daniel J. Tschumperlin; Robert Vassallo; Profibrotic effect of IL-17A and elevated IL-17RA in idiopathic pulmonary fibrosis and rheumatoid arthritis-associated lung disease support a direct role for IL-17A/IL-17RA in human fibrotic interstitial lung disease.

- American Journal of Physiology-Lung Cellular and Molecular Physiology* **2019**, 316, L487-L497, 10.1152/ajplung.00301.2018.
95. Peiyong Ye; Yu Liu; Chong Chen; Fei Tang; Qi Wu; Xiang Wang; Chang-Gong Liu; Xiuping Liu; Runhua Liu; Yang Liu; Pan Zheng; An mTORC1-Mdm2-Drosha axis for miRNA biogenesis in response to glucose- and amino acid-deprivation.. *Molecular Cell* **2015**, 57, 708-720, 10.1016/j.molcel.2014.12.034.
96. Lichao Fan; Xiaoting Yu; Ziling Huang; Shaoqiang Zheng; Yongxin Zhou; Hanjing Lv; Yu Zeng; Jin-Fu Xu; Xuyou Zhu; Xianghua Yi; Analysis of Microarray-Identified Genes and MicroRNAs Associated with Idiopathic Pulmonary Fibrosis. *Mediators of Inflammation* **2017**, 2017, 1-9, 10.1155/2017/1804240.
97. Takashi Kida; Takahiro Seno; Hidetake Nagahara; Takuya Inoue; Amane Nakabayashi; Yuji Kukida; Kazuki Fujioka; Wataru Fujii; Makoto Wada; Masataka Kohno; Y. Kawahito; Roles of high-mobility group box 1 and thrombin in murine pulmonary fibrosis and the therapeutic potential of thrombomodulin. *American Journal of Physiology-Lung Cellular and Molecular Physiology* **2018**, 314, L473-L483, 10.1152/ajplung.00287.2017.
98. Ying Chang; Laila Al-Alwan; Paul-André Risse; Carolyn Baglole; James Martin; Qutayba Hamid; David H. Eidelman; Th17 Associated Cytokines Promote Human Airway Smooth Muscle Cell Proliferation. *B69. NOVEL INSIGHTS INTO AIRWAY INFLAMMATION AND REMODELING IN ASTHMA AND COPD* **2011**, null, , 10.1164/ajrccm-conference.2011.183.1\_meetingabstracts.a3601.
99. Rebecca C. Coll; Avril A. B. Robertson; Jae Jin Chae; Sarah C. Higgins; Raúl Muñoz-Planillo; Marco C. Inserra; Irina Vetter; Lara S. Dungan; Brian G. Monks; Andrea Stutz; Daniel E. Croker; Mark S. Butler; Moritz Haneklaus; Caroline E. Sutton; Gabriel Núñez; Eicke Latz; Daniel L. Kastner; Kingston H. G. Mills; Seth Lucian Masters; Kate Schroder; Matthew A. Cooper; Luke A. J. O'Neill; A small-molecule inhibitor of the NLRP3 inflammasome for the treatment of inflammatory diseases. *Nature Medicine* **2015**, 21, 248-255, 10.1038/nm.3806.
100. Yamuna Krishnan; Anand Saminathan; Gordon R; Albornoz Ea; Christie Dc; Langley; Kumar V; Mantovani S; Robertson Aab; Butler Ms; Rowe Db; O'Neill La; Kanthasamy Ag; Schroder K; Cooper Ma; Woodruff Tm; F1000Prime recommendation of Inflammasome inhibition prevents  $\alpha$ -synuclein pathology and dopaminergic neurodegeneration in mice.. *F1000 - Post-publication peer review of the biomedical literature* **2020**, 10, , 10.3410/f.734317494.793570817.
101. Joseph A. Duncan; Dan T Bergstralh; Yanhong Wang; Stephen B. Willingham; Zhengmao Ye; Albert G. Zimmermann; Jenny P.-Y. Ting; Cryopyrin/NALP3 binds ATP/dATP, is an ATPase, and requires ATP binding to mediate inflammatory signaling. *Proceedings of the National Academy of Sciences* **2007**, 104, 8041-8046, 10.1073/pnas.0611496104.

102. C Marchetti; B Swartzwelter; Mi Koenders; Ca Dinarello; La Joosten; OP0090 The human safe NLRP3 inflammasome inhibitor OLT1177 suppresses joint inflammation in murine models of experimental arthritis. *Annual European Congress of Rheumatology, 14–17 June, 2017* **2017**, null, 89, 10.1136/annrheumdis-2017-eular.2775.
103. Yi Huang; Hua Jiang; Yun Chen; Xiaqiong Wang; Yanqing Yang; Jinhui Tao; Xianming Deng; Gaolin Liang; Huafeng Zhang; Wei Jiang; Rongbin Zhou; Tranilast directly targets NLRP 3 to treat inflammasome-driven diseases. *EMBO Molecular Medicine* **2018**, 10, e8689, 10.15252/emmm.201708689.
104. HongBin He; Hua Jiang; Yun Chen; Jin Ye; Aoli Wang; Chao Wang; Qingsong Liu; Gaolin Liang; Xianming Deng; Wei Jiang; Rongbin Zhou; Oridonin is a covalent NLRP3 inhibitor with strong anti-inflammasome activity.. *Nature Communications* **2018**, 9, 2550, 10.1038/s41467-018-04947-6.
105. Magdalena Matusiak; Nina Van Opdenbosch; Mohamed Lamkanfi; CARD- and pyrin-only proteins regulating inflammasome activation and immunity. *Immunological Reviews* **2015**, 265, 217-230, 10.1111/imr.12282.
106. Tao Liu; Qin Tang; Kunpeng Liu; Weihong Xie; Xin Liu; Huishan Wang; Rong-Fu Wang; Jun Cui; TRIM11 Suppresses AIM2 Inflammasome by Degrading AIM2 via p62-Dependent Selective Autophagy. *Cell Reports* **2016**, 16, 1988-2002, 10.1016/j.celrep.2016.07.019.
107. Lucia De Almeida; Sonal Khare; Alexander V. Misharin; Rajul Patel; Rojo A. Ratsimandresy; Melissa C. Wallin; Harris Perlman; David R. Greaves; Hal M. Hoffman; Andrea Dorfleutner; Christian Stehlik; The PYRIN Domain-only Protein POP1 Inhibits Inflammasome Assembly and Ameliorates Inflammatory Disease.. *Immunity* **2015**, 43, 264-76, 10.1016/j.immuni.2015.07.018.
108. Sonal Khare; Rojo A. Ratsimandresy; Lucia De Almeida; Carla M. Cuda; Stephanie L. Rellick; Alexander V. Misharin; Melissa C. Wallin; Anu Gangopadhyay; Eleonora Forte; Eva Gottwein; Harris Perlman; John C. Reed; David R. Greaves; Andrea Dorfleutner; Christian Stehlik; The PYRIN domain–only protein POP3 inhibits ALR inflammasomes and regulates responses to infection with DNA viruses. *Nature Immunology* **2014**, 15, 343-353, 10.1038/ni.2829.
109. Pei-Hui Wang; Zi-Wei Ye; Jian-Jun Deng; Kam-Leung Siu; Wei-Wei Gao; Vidyanath Chaudhary; Yun Cheng; Sin-Yee Fung; Kit-San Yuen; Ting-Hin Ho; Ching-Ping Chan; Yan Zhang; Kin-Hang Kok; Wanling Yang; Dong-Yan Jin; Chi Ping Chan; Inhibition of AIM 2 inflammasome activation by a novel transcript isoform of IFI 16. *EMBO reports* **2018**, 19, e45737, 10.15252/embr.201845737.
110. Yvonne Dombrowski; Mark Peric; Sarah Koglin; Claudia Kammerbauer; Christine Göss; David Anz; Maren Simanski; Regine Gläser; Jürgen Harder; Veit Hornung; Richard L Gallo; Thomas Ruzicka; Robert Besch; Jürgen Schaubert; Christine Göß; Cytosolic DNA Triggers Inflammasome Activation in Keratinocytes in Psoriatic Lesions. *Science Translational Medicine* **2011**, 3, 82ra38-82ra38, 10.1126/scitranslmed.3002001.

111. Si Ming Man; Qifan Zhu; Liqin Zhu; Zhiping Liu; Rajendra Karki; Ankit Malik; Deepika Sharma; Liyuan Li; R. K. Subbarao Malireddi; Prajwal Gurung; Geoffrey Neale; Scott R. Olsen; Robert A. Carter; Daniel J. McGoldrick; Gang Wu; David Finkelstein; Peter Vogel; Richard J. Gilbertson; Thirumala-Devi Kanneganti; Critical Role for the DNA Sensor AIM2 in Stem Cell Proliferation and Cancer.. *Cell* **2015**, *162*, 45-58, 10.1016/j.cell.2015.06.001.
112. Justin E Wilson; Alex S Petrucelli; Liang Chen; Alicia Koblansky; Agnieszka D Truax; Yoshitaka Oyama; Arlin B Rogers; W June Brickey; Yuli Wang; Monika Schneider; Marcus Mühlbauer; Wei-Chun Chou; Brianne Barker; Christian Jobin; Nancy L Allbritton; Dale A Ramsden; Beckley K Davis; Jenny P.-Y. Ting; Inflammasome-independent role of AIM2 in suppressing colon tumorigenesis via DNA-PK and Akt. *Nature Medicine* **2015**, *21*, 906-913, 10.1038/nm.3908.
113. Katherine L Deyoung; Michael E Ray; Yan A Su; Sarah L Anzick; Ricky Johnstone; J. Trapani; Paul S Meltzer; Jeffrey M Trent; Cloning a novel member of the human interferon-inducible gene family associated with control of tumorigenicity in a model of human melanoma. *Oncogene* **1997**, *15*, 453-457, 10.1038/sj.onc.1201206.
114. Susanne Dihlmann; Sha Tao; Fabian Echterdiek; Esther Herpel; Lina Jansen; Jenny Chang-Claude; Hermann Brenner; Michael Hoffmeister; Matthias Kloor; Lack of Absent in Melanoma 2 (AIM2) expression in tumor cells is closely associated with poor survival in colorectal cancer patients. *International Journal of Cancer* **2014**, *135*, 2387-2396, 10.1002/ijc.28891.
115. M.L. Kochman; HNPCC-Associated Small Bowel Cancer: Clinical and Molecular Characteristics. *Yearbook of Gastroenterology* **2006**, *2006*, 81-82, 10.1016/s0739-5930(08)70298-8.
116. Stefan M. Woerner; Matthias Kloor; Yvette Schwitalle; Hanni Youmans; Magnus Von Knebel Doeberitz; Johannes Gebert; Susanne Dihlmann; The putative tumor suppressor AIM2 is frequently affected by different genetic alterations in microsatellite unstable colon cancers. *Genes, Chromosomes and Cancer* **2007**, *46*, 1080-1089, 10.1002/gcc.20493.
117. Tae-Min Kim; Peter W. Laird; Peter J. Park; The landscape of microsatellite instability in colorectal and endometrial cancer genomes.. *Cell* **2013**, *155*, 858-68, 10.1016/j.cell.2013.10.015.
118. Larissa Ponomareva; Hongzhu Liu; Xin Duan; Eric Dickerson; Hui Shen; Ravichandran Panchanathan; Divaker Choubey; AIM2, an IFN-Inducible Cytosolic DNA Sensor, in the Development of Benign Prostate Hyperplasia and Prostate Cancer. *Molecular Cancer Research* **2013**, *11*, 1193-1202, 10.1158/1541-7786.MCR-13-0145.
119. Li-Jie Wang; Chia-Wei Hsu; Chiu-Chin Chen; Ying Liang; Lih-Chyang Chen; David M. Ojcius; Ngan-Ming Tsang; Chuen Hsueh; Chih-Ching Wu; Yu-Sun Chang; Interactome-wide analysis identifies end-binding protein 1 as a crucial component for the speck-like particle formation of activated absence in melanoma 2 (AIM2) inflammasomes.. *Molecular & Cellular Proteomics* **2012**, *11*, 1230-44, 10.1074/mcp.M112.020594.

120. Chen, L.C.; Wang, L.J.; Tsang, N.M.; Ojcius, D.M.; Chen, C.C.; Ouyang, C.N.; Hsueh, C.; Liang, Y.; Chang, K.P.; Chen, C.C.; et al. Tumour inflammasome-derived IL-1beta recruits neutrophils and improves local recurrence-free survival in EBV-induced nasopharyngeal carcinoma. *EMBO Mol. Med.* 2012, 4, 1276–1293.
- 

Retrieved from <https://encyclopedia.pub/entry/history/show/6805>