

# Agonists for Peroxisome Proliferator-Activated Receptors

Subjects: **Cell Biology**

Contributor: hiroyuki miyachi

Nuclear receptors (NRs) are ligand-dependent transcription factors that modulate diverse aspects of development, reproduction, and energy homeostasis. This receptor superfamily includes receptors for vitamin D, steroid hormones, thyroid hormones and retinoids, as well as a large number of orphan receptors. NRs are composed of six functionally distinct regions (termed A to F). The N-terminal AB region is highly variable and contains a constitutionally active transactivation function-1 (AF-1) motif. The central C region (a DNA-binding region) is highly conserved among NRs and contains two zinc finger motifs that make contact with specific nucleotide sequences, termed hormone response elements. The C-terminal D, E and F regions are required for ligand binding and receptor dimerization. In most NRs, these regions also contain a second highly conserved transcriptional activation function-2 (AF-2) motif, which is important for ligand-dependent transcription.

peroxisome proliferator-activated receptor

PPAR agonist

structural biology

ligand superfamily concept

helix 12 holding induction concept


## 1. Overview

Progress in understanding peroxisome proliferator-activated receptor (PPAR) subtypes as nuclear receptors that have pleiotropic effects on biological responses has enabled the exploration of new subtype-selective PPAR ligands. Such ligands are useful chemical biology/pharmacological tools to investigate the functions of PPARs and are also candidate drugs for the treatment of PPAR-mediated diseases, such as metabolic syndrome, inflammation and cancer.

## 2. Nuclear Receptors

Nuclear receptors (NRs) are ligand-dependent transcription factors that modulate diverse aspects of development, reproduction, and energy homeostasis. This receptor superfamily includes receptors for vitamin D, steroid hormones, thyroid hormones and retinoids, as well as a large number of orphan receptors. NRs are composed of six functionally distinct regions (termed A to F) (**Figure 1A**). The N-terminal AB region is highly variable and contains a constitutionally active transactivation function-1 (AF-1) motif. The central C region (a DNA-binding region) is highly conserved among NRs and contains two zinc finger motifs that make contact with specific nucleotide sequences, termed hormone response elements. The C-terminal D, E and F regions are required for

ligand binding and receptor dimerization. In most NRs, these regions also contain a second highly conserved transcriptional activation function-2 (AF-2) motif, which is important for ligand-dependent transcription.



**Figure 1.** (A) Intact PPAR $\gamma$ –rosiglitazone complex structure. A to F regions are depicted. The position of H12 (helix 12) is indicated with a red dotted circle. (B) Schematic representation of ligand-dependent nuclear receptor activation (PPAR as an example). PPAR forms a heterodimer with another nuclear receptor, RXR in the nucleus.

In the basal state, NRs are functionally inactive because they are tightly bound with corepressors, such as NCoR and SMRT. Upon NR binding with an activating ligand, the corepressor dissociates and then coactivators, such as SRC-1 and NCoA, are recruited and the AF-2 helix located in the F region is stabilized to initiate transcription (Figure 1B) [1]. Over the past three decades, much attention has been focused on a subgroup of NRs, the peroxisome proliferator-activated receptors (PPARs).

### 3. Peroxisome Proliferator-Activated Receptors

PPARs are activated by endogenous unsaturated and saturated fatty acids and by synthetic ligands [2]. There are three PPAR subtypes: PPAR $\alpha$ , PPAR $\delta$ , and PPAR $\gamma$ . Each PPAR subtype is expressed in a tissue-specific manner. PPAR $\alpha$  is mostly expressed in tissues involved in lipid oxidation, such as liver and kidney. PPAR $\gamma$  is expressed in adipose tissue, macrophages and vascular smooth muscle, and also in tumors originating from various organs. PPAR $\delta$  is expressed in adipose tissue, skeletal muscle, heart, etc. [3].

Upon ligand binding, PPARs heterodimerize with another nuclear receptor, retinoid X receptor (RXR), in the nucleus, and the heterodimers regulate genes expression, such as carnitine palmitoyl acyl-CoA transferase 1A

(*CPT1A*), angiopoietin-like protein 4 (*ANGPTL4*) and adipocyte differentiation-related protein (*ADRP*) by binding to specific consensus DNA sequences, termed peroxisome proliferator responsive elements (PPREs) in the promoter regions of target genes. The structural basis of PPREs is a direct repeat of the hexameric AGGTCA recognition motif, separated by one nucleotide (termed DR1) [4].

## 4. Pleiotropic Effect of PPARs

PPAR subtypes play a key role in lipid, lipoprotein and glucose homeostasis. PPAR $\alpha$  regulates genes involved in fatty acid uptake,  $\beta$ -oxidation, and  $\omega$ -oxidation. It downregulates apolipoprotein C-III, which regulates triglyceride hydrolysis by lipoprotein lipase, and also regulates genes involved in reverse cholesterol transport, such as apolipoprotein A-I and apolipoprotein A-II [5]. PPAR $\delta$  activation regulates HDL cholesterol levels, and it influences glycemic control [6][7][8]. PPAR $\delta$  activation markedly improves glucose tolerance and insulin resistance [9]. PPAR $\gamma$  is a master regulator of adipocyte differentiation, but recent molecular studies have indicated that its activation is also linked to the expression of many important genes that affect energy metabolism, such as TNF- $\alpha$ , leptin, and adiponectin [10]. PPAR $\gamma$  also promotes cell cycle arrest by inhibiting cyclin-dependent kinase activity in several tumor cell lines [11].


Recent extensive biological studies clearly disclosed that PPARs function beyond metabolism. Each PPAR subtype plays major roles in a broad spectrum of biological processes, including cell proliferation and differentiation, fatty acid and eicosanoid signaling, bone formation, tissue repair and remodeling, insulin sensitivity [12].

The above examples demonstrate that PPARs are important pleiotropic NRs and attractive molecular targets for the treatment of various diseases. Therefore, we think it is important to develop potent and PPAR subtype-selective ligands as tools to investigate the detailed functions of individual PPARs. In this review, we summarize in historical order our structural development studies to create PPAR subtype-selective agonists. The discovery of many types of PPAR ligands indicates the validity of our strategy to create subtype-selective NR agonists.

## 5. Working Hypothesis of the NR Ligand Superfamily


For over twenty years, we have been engaged in NR ligand structure research, which has been based on our working hypothesis of the NR ligand superfamily [13]. The structural and functional features of the many different NRs are similar; therefore, we speculate that all NRs are derived from a single ancestral protein and have structurally evolved to fit various kinds of endogenous NR ligand. Similar evolution of NR ligands would have occurred from an ancestral ligand to form a superfamily of NR ligands, even though they now have diverse structures and functions. Based on this hypothesis, NR ligand structures are divided into two types, basic framework and branch. A common hydrophobic skeleton or a branch that fits into the ligand binding pocket of the basic ancestral protein are characteristic structural motifs that provide NR selectivity. PPAR subtype-selective agonists have certain unique structures associated with subtype selectivity. Examples include, thiazolidine-2,4-dione (TZD) and related structures, exemplified by pioglitazone, for PPAR $\gamma$  [Figure 2 (1)], the 2,2-dialkyl(usually

dimethyl)phenoxyacetic acid structure, exemplified by fenofibrate, for PPAR $\alpha$  [Figure 2 (2)], and the 2,2-unsubstituted phenoxyacetic acid structure, exemplified by GW-501516, for PPAR $\delta$  [Figure 2 (3)]. However, based on our working hypothesis, we predict that various kinds of subtype-selective, dual-, and pan-agonists can be created by starting with a common chemical framework as a template.



**Figure 2.** Structures of PPAR $\gamma$  agonist pioglitazone (1), PPAR $\alpha$  agonist fenofibrate (2), and PPAR $\delta$  agonist GW-501516 (3), and, below, our PPAR agonists.

Based on our hypothesis, we have successfully created various kinds of subtype-selective PPAR agonist, including KCL (PPAR  $\alpha$ -selective agonist) [14], APHM-19 (PPAR  $\alpha$ -selective agonist) [15], TIPP-401 (PPAR  $\alpha/\delta$  dual agonist) [16], APHM-13 (fluorescent PPAR  $\alpha/\delta$  dual agonist) [17], TIPP-204 (PPAR  $\delta$ -selective agonist) [18], TIPP-703 (PPAR pan agonist) [19], MO-4R (PPAR  $\gamma$ -selective agonist) [20], MEKT-21 (PPAR  $\gamma$ -selective partial agonist) [21], and MEKT-75 (PPAR  $\gamma$ -selective partial agonist) [22]. In addition, in collaboration with Dr. Oyama (Yamanashi University, Japan) and Dr. Shimizu (Tokyo University, Japan), we have solved many X-ray crystallographic structures of PPAR subtypes complexed with our ligands [23][24][25]. These structural biology studies have been integral to our medicinal chemistry research (Figure 3).



**Figure 3.** X-ray-crystallographic structures of our representative PPAR agonists complexed with each PPAR ligand binding domain (LBD).

## References

1. Chawta, A.; Repa, J.J.; Evans, R.M.; Mangelsdorf, D.J. Nuclear receptors and lipid physiology: Opening the x-files. *Science* 2001, 294, 1866–1870.
2. Banner, C.D.; Gottlicher, M.; Widmark, E.; Sjovall, J.; Rafter, J.J.; Gustafsson, J.A. A systematic analytical chemistry/cell assay approach to isolate activators of orphan nuclear receptors from biological extracts: Characterization of peroxisome proliferator-activated receptor activators in plasma. *J. Lipid Res.* 1993, 34, 1583–1591.
3. Wagner, K.D.; Wagner, N. Peroxisome proliferator-activated receptor  $\beta/\delta$  (PPAR $\beta/\delta$ ) acts as regulator of metabolism linked to multiple cellular functions. *Pharmacol. Ther.* 2010, 125, 423–435.
4. Keller, H.; Dreyer, C.; Medin, J.; Mahfoudi, A.; Ozato, K.; Wahli, W. Fatty acids and retinoids control lipid metabolism through activation of peroxisome proliferator-activated receptor-retinoid X receptor heterodimers. *Proc. Natl. Acad. Sci. USA* 1993, 90, 2160–2164.
5. Staels, B.; Auwerx, J. Role of PPAR in the pharmacological regulation of lipoprotein metabolism by fibrates and thiazolidinediones. *Curr. Pharm. Des.* 1997, 3, 1–14.
6. Lim, H.; Gupta, R.A.; Ma, W.G.; Paria, B.C.; Moller, D.E.; Morrow, J.D.; DuBois, R.N.; Trzaskos, J.M.; Dey, S.K. Cyclo-oxygenase-2-derived prostacyclin mediates embryo implantation in the mouse via PPAR $\delta$ . *Genes Dev.* 1999, 13, 1561–1574.

7. Sznaidman, M.L.; Haffner, C.D.; Maloney, P.R.; Fivush, A.; Chao, E.; Goreham, D.; Chao, E.; Goreham, D.; Sierra, M.L.; LeGrumec, C.; et al. Novel selective small molecule agonists for peroxisome proliferator-activated receptor  $\delta$  (PPAR $\delta$ ) synthesis and biological activity. *Bioorg. Med. Chem. Lett.* 2003, 13, 1517–1521.

8. Oliver, W.R., Jr.; Shenk, J.L.; Snaith, M.R.; Russell, C.S.; Plunket, K.D.; Bodkin, N.L.; Lewis, M.C.; Winegar, D.A.; Sznaidman, M.L.; Lambert, M.H.; et al. A selective peroxisome proliferator-activated receptor  $\delta$  agonist promotes reverse cholesterol transport. *Proc. Natl. Acad. Sci. USA* 2001, 98, 5306–5311.

9. Tanaka, T.; Yamamoto, J.; Iwasaki, S.; Asaba, H.; Hamura, H.; Ikeda, Y.; Watanabe, M.; Magoori, K.; Ionka, R.X.; Tachibana, K.; et al. Activation of peroxisome proliferator-activated receptor  $\delta$  induces fatty acid  $\beta$ -oxidation in skeletal muscle and attenuates metabolic syndrome. *Proc. Natl. Acad. Sci. USA* 2003, 100, 15924–15929.

10. Okuno, A.; Tamemoto, H.; Tobe, K.; Ueki, K.; Mori, Y.; Iwamoto, K.; Umesono, K.; Akanuma, Y.; Fujiwara, T.; Horikoshi, H.; et al. Troglitazone increases the number of small adipocytes without the change of white adipose tissue mass in obese Zucker rats. *J. Clin. Investig.* 1998, 101, 1354–1361.

11. Huang, J.-W.; Shiau, C.-W.; Yang, J.; Wang, D.-S.; Chiu, H.-C.; Chen, C.-S.; Chen, A.C.-Y. Development of small-molecule cyclin D1-ablative agents. *J. Med. Chem.* 2006, 49, 4684–4689.

12. Duszka, K.; Gregor, A.; Guillou, H.; König, J.; Wahli, W. Peroxisome Proliferator-Activated Receptors and Caloric Restriction—Common Pathways Affecting Metabolism, Health, and Longevity. *Cells* 2020, 9, 1708.

13. Hashimoto, Y.; Miyachi, H. Nuclear receptor antagonists designed based on the helix-folding inhibition hypothesis. *Bioorg. Med. Chem.* 2005, 13, 5080–5093.

14. Nomura, M.; Tanase, T.; Ide, T.; Tsunoda, M.; Suzuki, M.; Uchiki, H.; Murakami, K.; Miyachi, H. Design, Synthesis and Evaluation of Substituted Phenylpropanoic Acid Derivatives as Human Peroxisome Proliferator-Activated Receptor Activators; Discovery of Potent and Human PPAR $\alpha$  Subtype-Selective Activators. *J. Med. Chem.* 2003, 46, 3581–3599.

15. Oyama, T.; Kamata, S.; Ishii, I.; Miyachi, H. Crystal structures of the human peroxisome proliferator-activated receptor (PPAR) $\alpha$  ligand-binding domain in complexes with a series of phenylpropanoic acid derivatives generated by a ligand-exchange soaking method. *Bio. Pharm. Bull.* 2021. accepted for publication.

16. Kasuga, J.; Yamasaki, D.; Araya, Y.; Nakagawa, A.; Makishima, M.; Doi, T.; Hashimoto, Y.; Miyachi, H. Design, synthesis and evaluation of a novel series of  $\alpha$ -substituted phenylpropanoic acid derivatives as human peroxisome proliferator-activated receptor (PPAR)  $\alpha/\delta$  dual agonists for the treatment of metabolic syndrome. *Bioorg. Med. Chem.* 2006, 14, 8405–8414.

17. Araya, Y.; Kasuga, J.; Toyota, K.; Hirakawa, Y.; Oyama, T.; Makishima, M.; Morikawa, K.; Hashimoto, Y.; Miyachi, H. Structure-Based Design and Synthesis of Fluorescent PPAR $\alpha/\delta$  Co-agonist and Its Application as a Probe for Fluorescent Polarization Assay of PPAR $\delta$  Ligands. *Chem. Pharm. Bull.* 2008, 56, 1357–1359.

18. Kasuga, J.; Nakagome, I.; Aoyama, A.; Sako, K.; Ishizawa, M.; Ogura, M.; Makishima, M.; Hirono, S.; Hashimoto, Y.; Miyachi, H. Design, synthesis, and evaluation of potent, structurally novel peroxisome proliferator-activated receptor (PPAR)  $\delta$ -selective agonists. *Bioorg. Med. Chem.* 2007, 15, 5177–5190.

19. Kasuga, J.; Yamasaki, D.; Ogura, K.; Shimizu, M.; Sato, M.; Makishima, M.; Doi, T.; Hashimoto, Y.; Miyachi, H. SAR-oriented discovery of peroxisome proliferator-activated receptor pan agonist with a 4-adamantylphenyl group as a hydrophobic tail. *Bioorg. Med. Chem. Lett.* 2008, 18, 1110–1115.

20. Ohashi, M.; Oyama, T.; Nakagome, I.; Satoh, M.; Nishio, Y.; Nobusada, H.; Hirono, S.; Morikawa, K.; Hashimoto, Y.; Miyachi, H. Design, Synthesis, and Structural Analysis of Phenylpropanoic Acid-Type PPAR $\gamma$ -Selective Agonists: Discovery of Reversed Stereochemistry-Activity Relationship. *J. Med. Chem.* 2011, 54, 331–341.

21. Ohashi, M.; Oyama, T.; Putranto, E.W.; Waku, T.; Nobusada, H.; Kataoka, K.; Matsuno, K.; Yashiro, M.; Morikawa, K.; Huh, N.H.; et al. Design and synthesis of a series of  $\alpha$ -benzyl phenylpropanoic acid-type peroxisome proliferator-activated receptor (PPAR)  $\gamma$  partial agonists with improved aqueous solubility. *Bioorg. Med. Chem.* 2013, 21, 2319–2332.

22. Ohashi, M.; Oyama, T.; Miyachi, H. Different structures of the two peroxisome proliferator-activated receptor  $\gamma$  (PPAR $\gamma$ ) ligand-binding domains in homodimeric complex with partial agonist, but not full agonist. *Bioorg. Med. Chem. Lett.* 2015, 25, 2639–2644.

23. Oyama, T.; Toyota, K.; Waku, T.; Hirakawa, Y.; Nagasawa, N.; Kasuga, J.; Hashimoto, Y.; Miyachi, H.; Morikawa, K. Adaptability and selectivity of human peroxisome proliferator-activated receptor (PPAR) pan agonists revealed from crystal structures. *Acta Crystallogr. D Biol. Crystallogr.* 2009, 65, 786–795.

24. Kuwabara, N.; Oyama, T.; Tomioka, D.; Ohashi, M.; Yanagisawa, J.; Shimizu, T.; Miyachi, H. Peroxisome proliferator-activated receptors (PPARs) have multiple binding points that accommodate ligands in various conformations: Phenylpropanoic acid-type PPAR ligands bind to PPAR in different conformations, depending on the subtype. *J. Med. Chem.* 2012, 55, 893–902.

25. Ohashi, M.; Gamo, K.; Oyama, T.; Miyachi, H. Peroxisome proliferator-activated receptor  $\gamma$  (PPAR $\gamma$ ) has multiple binding points that accommodate ligands in various conformations: Structurally similar PPAR $\gamma$  partial agonists bind to PPAR $\gamma$  LBD in different conformations. *Bioorg. Med. Chem. Lett.* 2015, 25, 2758–2762.