

Forest Biomass

Subjects: Forestry

Contributors:  Abdelwahab Bessaad,  Nathalie Korboulewsky



Submitted by:  Abdelwahab Bessaad

Definition

Forest biomass is a sustainable source of renewable energy and a valuable alternative to finite fossil fuels. However, its overharvesting may lead to soil nutrient depletion and threaten future stand productivity, as well as affect the habitat for biodiversity.

1. Introduction

The European Union is resolved to reduce greenhouse gas emissions and promoting the development of new energy policies to reverse the trend of global warming associated with climate change ^{[1][2]}. These policies aim to develop new and sustainable energy sources and contribute to a reduction in the growing dependency of EU countries on fossil fuels. Amongst renewable resources, paramount importance has been given to the bioenergy because it has low negative environmental impacts. Biomass was one of humanity's earliest sources of energy, and it remains an important resource today and for the future to enhance security energy. Today, biomass accounts more than half of all the EU's renewable energy production and 10% of all energy sources ^[3]. This figure is likely to rise in the future because it is considered a promising alternative to fossil energy in order to shift towards sustainability. Woody biomass accounts for more than 60% of all EU domestic biomass ^[4]. Indeed, fuelwood seems to be a promising option in temperate forests and developed countries ^[5] by providing a local supply of renewable energy and by allowing forest owners to diversify their sources of income ^[6]. In most European countries, woody biomass demand is rising fast with many bioenergy strategies ^{[7][8]}, which have led to the adoption of new logging practices ^{[9][10][11]}. Different biomass harvesting methods are used in Europe; their selection depends on site conditions, silvicultural treatments, species composition, tree sizes, stand density, and the economic condition of each country. The main types of silvicultural operations, related to forest biomass extraction, are whole-tree harvesting in cleaning, thinning, and harvesting of woody debris and stumps before regeneration ^[12]. Nevertheless, the intensive harvesting of biomass for energy may cause additional pressure on forested ecosystems ^{[13][14][15][16][17][18][19]}. Recent interest in using woody debris and small-diameter material for biofuels is generating a renewed focus for studies on harvesting impacts on soils, biodiversity, and forestry practice sustainability. Indeed, this part of biomass, usually left in the forest in conventional harvesting systems, plays an essential role in nutrient cycles, maintenance of soil fertility, carbon sequestration, regeneration of communities, and biodiversity ^{[9][10][17][20]}.

Forest residues, downed wood, and deadwood are different terms used to define woody debris, including all nonliving biomass. Woody debris refers to fallen dead trees and the remains of twigs, branches, and small pieces of wood left on the ground in forests. It is generally divided into two main classes: (1) Fine Woody Debris (FWD) with a diameter less than, or equal to, 10 cm, and (2) Coarse Woody Debris (CWD) greater than 10 cm in diameter ^{[21][22]}. However, as there are no clearly defined criteria to separate these two classes, the diameter limit may vary depending on the countries or research purposes. For example, the limit between the two classes is 7.6 cm (3 inches) in some studies conducted in the United States ^{[23][24][25]}. In addition, the number of wood diameter classes can also vary according to the objectives of the studies; three to five classes are often used to better estimate the nutrient contents of woody debris and the mineral exports associated with their removal ^{[26][27][28]}.

Despite the relatively low proportion FWD compared to total harvested biomass, it contains a large amount of nutrients that are key elements for forest growth, and its overharvesting could compromise long-term forest site fertility ^{[5][16][29][30]}. Several studies have demonstrated that harvesting fine wood

affects the soil nutrient cycling and thus future site productivity, even with high rates of fertilizer application [18][31][32][33]. CWD is also an important component of forest ecosystems in that it provides habitats or micro-habitats, for saproxylic species, trophic substrate, soil texturing and structuring, and temporary carbon stock [10][20].

Studies in European forests have shown that there is a significant correlation between woody debris volume, harvest methods, and such forest parameters as management, site or stand attributes [34][35]. To prevent any reduction in the woody debris left in forests and negative effects of whole-tree harvesting, many countries have established guidelines for sustainable harvesting of forest biomass in order to limit nutrient exports mainly contained in leaves and fine wood [19][28][36][37]. The guidelines are generally based on volumes or on the number of wood pieces that should be left on the forest. However, the existing recommendations for sustainable biomass harvesting may vary between countries. For instance, Nordic countries and the United Kingdom first classify sites and stands according to their sensitivity to biomass extraction (forest type, soil fertility, wood production, soil compaction, and slope), and different restrictions are then imposed for the specific site types [38][39]. For example, Finnish guidelines recommend leaving up to 30% of woody debris, or the same amount of nutrients, on the forest floor [40], using ash recycling or fertilization [12][41]. In France, the recommendations for sustainable biomass harvesting have continuously been updated since 2006 [42] through several scientific studies and increasing technical expertise [28][36][42]. The latest recommendations take into account local conditions, in particular soil sensitivity to mineral exports [28]. It is highly recommended to leave at least 10% of fine wood with a diameter of less than, or equal to, 7 cm on the forest floor for non-sensitive soils, and to increase this value to 30% on moderately sensitive soils. In the case of highly sensitive or poor soils, harvesting fine wood is not recommended; all fine wood should be left on the forest floor. Furthermore, according to USDA guidelines for the Pacific North West, at least 4%–5% of total forest biomass should be retained on the operation site to avoid soil degradation and growth decline in the next rotation [37].

Overall, the national guidelines cover the same topics and often make similar general recommendations. To protect sensitive soils, most guidelines recommend restricting biomass removal on sites identified as sensitive; site sensitivity is often based on a combination of indicators, such as slope, soil depth, and soil moisture regime, all of which may be related to susceptibility to erosion, compaction, loss of fertility, and ecosystem resilience to harvesting. To protect biodiversity, most guidelines advise for example retaining at least 10 to 30% of the slash or tops, retaining biological legacies, such as large logs, snags, or wildlife trees, and restricting biomass harvesting near critical habitats and retaining appropriate vegetation and deadwood for habitats or along rivers [37][43][44]. On top of that, it is highly recommended to harvest during the leafless period to avoid exporting leaves from forests. However, when harvesting takes place within the leafed period on evergreen species, extracting crown biomass is recommended only after pre-drying operation [12][28][42]. Pre-drying felled trees is carried out on the forest before skidding operations. This operation has two major roles: first, it may allow the weakened leaves, twigs, and fine wood to fall off during the skidding. Second, it allows maintaining a certain amount of nutrients by leaching via rainfall, depending on weather conditions.

2. Study Area

The study area is located in the Centre-Val de Loire region, Central France (Lat 47.705141°; Long 1.647949°). Forests are mainly private (87%), composed of broadleaves species and cover a quarter of its total area [45]. They contain an important fuelwood resource but their soils are nutrient-poor and sensitive to the loss of fertility [46]. Half of the forested area of the region has long been managed in a coppice-with-standards (CWS) regime [47]. The standard layer is mostly dominated by deciduous oaks (*Quercus robur* L., *Q. petraea* (Matt.) Liebl. and in the coppice layer by hornbeam (*Carpinus betulus* L.), sweet chestnut (*Castanea sativa* Mill.) and other deciduous species.

Nine whole-tree harvesting sites in central France were sampled for three years 2017–2020 (Figure 1). Three sites are located in a public forest, and five are in different private forests. Some of the sites are located in protected areas of the Natura 2000 network, but none are concerned by a high-level protection

measures.




Figure 1. Location of the nine studied sites (S_1 to S_9) on the map of the Centre-Val de Loire region that comprises six departments: Cher (18), Eure-et-Loir (28), Indre (36), Indre-et-Loire (37), Loir-et-Cher (41), Loiret (45).

3. Aboveground Biomass and Harvest Intensity

The studied sites correspond to different stand structures, species composition, and management systems. Aboveground biomass estimates (Figure 2) showed a high variability between sites: standing biomass varied between 118 and 519 Mg ha^{-1} . In contrast, less variability was found between sites managed as coppice and CWS (S_1 , S_2 , S_5 , S_6 , S_7 , and S_8), the mean aboveground biomass was amounted to $174 \pm 56 \text{ Mg ha}^{-1}$. High forest sites (S_3 , S_4 , and S_9) contain obviously more biomass $383 \pm 127 \text{ Mg ha}^{-1}$. The proportion of wood [$d > 7 \text{ cm}$] in total biomass was greater in high forests ($83 \pm 5\%$) than in other stands ($69 \pm 7\%$). In contrast, the percentage of wood [$d \leq 7 \text{ cm}$] was lower in high forests $17 \pm 5\%$ compared to CWS sites $31 \pm 7\%$.




Figure 2. Estimates of standing biomass on the nine sites before harvest by diameter classes.

In general, the harvest intensity was highly variable and ranged between 13% and 100% (Table 2). On sites managed as coppice and CWS, more than 70% of aboveground biomass was harvested (S_2 , S_6 , S_7 , and S_8), 43% for S_5 and 33% for S_1 . However, the harvest intensity was lower in high forests (S_3 , S_4 , S_9) and ranged between 13% and 52%.

Table 2. Amounts (Mg ha^{-1}) and proportions (%) of woody debris left on sites presented for each wood diameter class.

Sites	Harvest Intensity (%)	FWD				CWD				Total Woody Debris	Total Woody Debris
		[d ≤ 4 cm]		[d = 4-7 cm]		Total [d ≤ 7 cm]		[d > 7 cm]			
		Mg ha ⁻¹	%	Mg ha ⁻¹	%	Mg ha ⁻¹	%	Mg ha ⁻¹	%	Mg ha ⁻¹	%
S_1	33	9.33	70	0.42	4	9.75	39	0	0	9.75	25
S_2	72	4.17	17	0.95	8	5.12	14	0.15	0	5.27	4
S_3	52	2.38	9	0.56	3	2.94	7	0.77	1	3.71	3
S_4	13	0.67	6	0.25	3	0.92	4	4.88	10	5.80	9
S_5	43	2.37	11	0	0	2.37	7	4.31	6	6.68	7
S_6	75	3.97	10	0	0	3.97	5	6.32	6	10.29	6
S_7	100	12.47	49	0.74	6	13.21	34	0	0	13.21	10
S_8	81	3.73	16	0.78	3	4.51	9	1.33	2	5.84	5
S_9	35	1.70	7	0	0	1.70	4	1.99	2	3.69	3

References

- European Union (EU). Directive (EU) 2018/2001 of the European Parliament and of the Council of 11 December 2018 on the promotion of the use of energy from renewable sources. Off. J. Eur. Union 2018, 5, 82–209. Available online: (accessed on 7 April 2020).
- European Commission (EC). Sustainable Bioeconomy for Europe: Strengthening the Connection between Economy, Society and the Environment; European Commission: Brussels, Belgium, 2018; p. 103.
- Scarlat, N.; Dallemand, J.-F.; Taylor, N.; Banja, M.; Sanchez, L.J.; Avraamides, M. Brief on Biomass for Energy in the European Union; European Commission: Brussels, Belgium, 2019; p. 8.
- European Commission. Report from the Commission to the European Parliament, the Council; The European Economic and Social Committee/Committee of the Regions: Brussels, Belgium, 2019; Available online: (accessed on 7 April 2020).
- Lattimore, B.; Smith, C.T.; Titus, B.D.; Stupak, I.; Egnell, G. Environmental factors in woodfuel production: Opportunities, risks, and criteria and indicators for sustainable practices. *Biomass Bioenergy* 2009, 33, 1321–1342.
- Manley, A.; Richardson, J. Silviculture and economic benefits of producing wood energy from conventional forestry systems and measures to mitigate negative impacts. *Biomass Bioenergy* 1995, 9, 89–105.
- Pelli, P.; Haapala, A.; Pykäläinen, J. Services in the forest-based bioeconomy—Analysis of European strategies. *Scand. J. For. Res.* 2017, 32, 559–567.
- Wolfslehner, B.; Linser, S.; Pülzl, H.; Bastrup-Birk, A.; Camia, A.; Marchetti, M. Forest bioeconomy—A new scope for sustainability indicators. *Sci. Policy* 2016, 4, 1–32.
- Verkerk, P.J.; Lindner, M.; Zanchi, G.; Zudin, S. Assessing impacts of intensified biomass removal on deadwood in European forests. *Ecol. Indic.* 2011, 11, 27–35.
- Bouget, C.; Lassauce, A.; Jonsell, M. Effects of fuelwood harvesting on biodiversity—A review focused on the situation in Europe. *Can. J. For. Res.* 2012, 42, 1421–1432.

11. Thiffault, E.; Hannam, K.D.; Pare, D.; Titus, B.D.; Hazlett, P.W.; Maynard, D.G.; Brais, S. Effects of forest biomass harvesting on soil productivity in boreal and temperate forests—A review. *Environ. Rev.* 2011, 19, 278–309.
12. Stupak, I.; Asikainen, A.; Röser, D.; Pasanen, K. Review of Recommendations for Forest Energy Harvesting and Wood Ash Recycling. In *Sustainable Use of Forest Biomass for Energy*; Springer: Dordrecht, The Netherlands, 2008; pp. 155–196.
13. Mälkönen, E. Effect of whole-tree harvesting on soil fertility. *Silva Fenn.* 1976, 10, 157–164.
14. Kaarakka, L.; Tamminen, P.; Saarsalmi, A.; Kukkola, M.; Helmisaari, H.-S.; Burton, A.J. Effects of repeated whole-tree harvesting on soil properties and tree growth in a Norway spruce (*Picea abies* (L.) Karst.) stand. *For. Ecol. Manag.* 2014, 313, 180–187.
15. Knust, C.; Schua, K.; Feger, K.H. Estimation of Nutrient Exports Resulting from Thinning and Intensive Biomass Extraction in Medium-Aged Spruce and Pine Stands in Saxony, Northeast Germany. *Forests* 2016, 7, 302.
16. Aherne, J.; Posch, M.; Forsius, M.; Lehtonen, A.; Harkonen, K. Impacts of forest biomass removal on soil nutrient status under climate change: A catchment-based modelling study for Finland. *Biogeochemistry* 2012, 107, 471–488.
17. Miettinen, J.; Ollikainen, M.; Nieminen, T.M.; Ukonmaanaho, L.; Laurén, A.; Hynynen, J.; Lehtonen, M.; Valsta, L. Whole-tree harvesting with stump removal versus stem-only harvesting in peatlands when water quality, biodiversity conservation and climate change mitigation matter. *For. Policy Econ.* 2014, 47, 25–35.
18. Walmsley, J.D.; Jones, D.L.; Reynolds, B.; Price, M.H.; Healey, J.R. Whole tree harvesting can reduce second rotation forest productivity. *For. Ecol. Manag.* 2009, 257, 1104–1111.
19. Achat, D.L.; Deleuze, C.; Landmann, G.; Pousse, N.; Ranger, J.; Augusto, L. Quantifying consequences of removing harvesting residues on forest soils and tree growth—A meta-analysis. *For. Ecol. Manag.* 2015, 348, 124–141.
20. Larrieu, L.; Cabanettes, A.; Gouix, N.; Burnel, L.; Bouget, C.; Deconchat, M. Post-harvesting dynamics of the deadwood profile: The case of lowland beech-oak coppice-with-standards set-aside stands in France. *Eur. J. For. Res.* 2019.
21. Nemeč, A.F.L.; Davis, G. Efficiency of Six Line Intersect Sampling Designs for Estimating Volume and Density of Coarse Woody Debris; Vancouver Forest Region: Nanaimo, BC, Canada, 2002.
22. Woodall, C.W.; Monleon, V.J. Sampling Protocol, Estimation, and Analysis Procedures for the Down Woody Materials Indicator of the FIA Program; US Department of Agriculture and Forest Service, Northern Research Station: Newtown Square, PA, USA, 2008; p. 68.
23. Brown, J.K. Handbook for Inventorying Downed Woody Material; US Department of Agriculture, Forest Service, Intermountain Forest and Range Experiment Station: Ogden, UT, USA, 1974; p. 24.
24. Waddell, K.L. Sampling coarse woody debris for multiple attributes in extensive resource inventories. *Ecol. Indic.* 2002, 1, 139–153.
25. Briedis, J.I.; Wilson, J.S.; Benjamin, J.G.; Wagner, R.G. Biomass retention following whole-tree, energy wood harvests in central Maine: Adherence to five state guidelines. *Biomass Bioenergy* 2011, 35, 3552–3560.
26. Phillips, T.; Watmough, S.A. A nutrient budget for a selection harvest: Implications for long-term sustainability. *Can. J. For. Res.* 2012, 42, 2064–2077.
27. Pyttel, P.L.; Kohn, M.; Bauhus, J. Effects of different harvesting intensities on the macro nutrient pools in aged oak coppice forests. *For. Ecol. Manag.* 2015, 349, 94–105.
28. Landmann, G.; Augusto, L.; Pousse, N.; Gosselin, M.; Cacot, E.; Deleuze, C.; Bilger, I.; Amm, A.; Bilot, N.; Boulanger, V.; et al. Recommendations pour une Récolte Durable de Biomasse Forestière pour l'Énergie—Focus sur les Menus Bois et les Souches; GIP-ECOFOR: Paris, France; ADEME: Angers, France, 2018; p. 52. Available online: (accessed on 23 March 2019).
29. Augusto, L.; Achat, D.L.; Bakker, M.R.; Bernier, F.; Bert, D.; Danjon, F.; Khelifa, R.; Meredieu, C.; Trichet, P. Biomass and nutrients in tree root systems—Sustainable harvesting of an intensively managed *Pinus pinaster* (Ait.) planted forest. *GCB Bioenergy* 2015, 7, 231–243.
30. Stevens, P.A.; Norris, D.A.; Williams, T.G.; Hughes, S.; Durrant, D.W.H.; Anderson, M.A.; Weatherley, N.S.; Hornung, M.; Woods, C. Nutrient losses after clearfelling in Beddgelert Forest: A comparison of the effects of conventional and whole-tree harvest on soil water chemistry. *For. Int. J. For. Res.* 1995, 68, 115–131.
31. Törmänen, T.; Kitunen, V.; Lindroos, A.-J.; Heikkinen, J.; Smolander, A. How do logging residues of different tree species affect soil N cycling after final felling? *For. Ecol. Manag.* 2018, 427, 182–189.
32. Nord-Larsen, T. Stand and site productivity response following whole-tree harvesting in early thinnings of Norway spruce (*Picea abies* (L.) Karst.). *Biomass Bioenergy* 2002, 23, 1–12.
33. Rocha, J.H.T.; Gonçalves, J.L.d.M.; Brandani, C.B.; Ferraz, A.d.V.; Franci, A.F.; Marques, E.R.G.; Arthur Junior, J.C.; Hubner, A. Forest residue removal decreases soil quality and affects wood productivity even with high rates of fertilizer application. *For. Ecol. Manag.* 2018, 430, 188–195.
34. Böhl, J.; Brändli, U.B. Deadwood volume assessment in the third Swiss National Forest Inventory: Methods and first results. *Eur. J. For. Res.* 2007, 126, 449–457.
35. Vítková, L.; Bače, R.; Kjučukov, P.; Svoboda, M. Deadwood management in Central European forests: Key considerations for practical implementation. *For. Ecol. Manag.* 2018, 429, 394–405.
36. Landmann, G.; Nivet, C. Projet RESOBIO—Gestion des Rémanents Forestiers: Préservation des Sols et de la

Biodiversité; Rapport Final; ADEME—Ministère de l'Agriculture, de l'Agroalimentaire et de la Forêt: Angers, France; GIP Ecofor: Paris, France, 2014; p. 243.

37. Titus, B.D.; Brown, K.; Helmisaari, H.-S.; Vanguelova, E.; Stupak, I.; Evans, A.; Clarke, N.; Guidi, C.; Bruckman, V.J.; Varnagiryte-Kabasinskiene, I. Sustainable forest biomass: A review of current residue harvesting guidelines. *Energy Sustain. Soc.* 2021, 11, 10.
38. Koistinen, A.; Aijala, O.; Mattson-Turku, G. *Uttag av Energived; Skogsbrukets Utvecklingscentral Tapio: Vammala, Finland, 2005.*
39. Nisbet, T.; Dutch, J.; Moffat, A.J. *Whole-Tree Harvesting: A Guide to Good Practice; Forestry Authority: Cambridge, UK, 1997.*
40. Aijala, O.; Kuusinen, M.; Halonen, M. *Energy Wood Harvest from Clearcuts: Guidelines; Forest Engineering Research Institute of Canada: Pointe-Claire, QC, Canada, 2005.*
41. Richardson, J.; Björheden, R.; Hakkila, P.; Lowe, A.; Smith, C. *Bioenergy from Sustainable Forestry: Guiding Principles and Practice; Springer Science & Business Media: New York, NY, USA, 2006; p. 71.*
42. Cacot, E.; Eisner, N.; Charnet, F.; Leon, P.; Rantien, C.; Ranger, J. *La Récolte Raisonnée des Rémanents en Forêt—Guide Pratique; ADEME: Angers, France; AFOCEL/INRAE/Union de la Coopération Forestière Française: Paris, France, 2006; p. 35.*
43. Evans, A.M.; Perschel, R.T.; Kittler, B.A. Overview of Forest Biomass Harvesting Guidelines. *J. Sustain. For.* 2013, 32, 89–107.
44. Bennett, N.; Bradley, M.; Broderick, S.; Brynn, D.; Bryan, R.; Campbell, R.; Dwyer, H.; Frost, E.; Hawthorn, B.; Ingerson, A.; et al. *Biomass Retention and Harvesting Guidelines for the Northeast; Forest Guild Biomass Working Group: Santa Fe, NM, USA, 2010; p. 17.*
45. Institut National de l'Information Géographique et Forestière (IGN). *Le Mémento Inventaire Forestier; Institut National de l'Information Géographique et Forestière: Paris, France, 2018; p. 29.*
46. Augusto, L.; Pousse, N.; Legout, A.; Seynave, I.; Jabiol, B.; Levillain, J. *Indicateurs de SENSibilité des Ecosystèmes Forestiers Soumis à une Récolte Accrue de Biomasse (INSENSE); ADEME: Angers, France, 2018; p. 262.*
47. Centre Régional de la Propriété Forestière (CRPF). *Schéma Régional de Gestion Sylvicole (SRGS); Centre Régional de la Propriété Forestière: Ile-de-France, France, 2007; p. 46.*

Keywords

whole-tree harvesting;biomass;fuelwood;fine woody debris;coarse woody debris

Retrieved from <https://encyclopedia.pub/12806>