Biodiesel from Castor Oil

Subjects: Energy & Fuels

Contributor: Carlos S. Osorio-González, Natali Gómez-Falcon, Fabiola Sandoval-Salas, Rahul Saini, Satinder K.

Brar, Antonio Avalos Ramírez

An attractive alternative to the use of fossil fuels is biodiesel, which can be obtained from a variety of feedstock through different transesterification systems such as ultrasound, microwave, biological, chemical, among others. The efficient and cost-effective biodiesel production depends on several parameters such as free fatty acid content in the feedstock, transesterification reaction efficiency, alcohol: oil ratio, catalysts type, and several parameters during the production process. However, biodiesel production from vegetable oils is under development, causing the final price of biodiesel to be higher than diesel derived from petroleum. An alternative to decrease the production costs will be the use of economical feedstocks and simple production processes. Castor oil is an excellent raw material in terms of price and quality, but especially this non-edible vegetable oil does not have any issues or compromise food security. Recently, the use of castor oil has attracted attention for producing and optimizing biodiesel production, due to the high content of ricinoleic fatty acid and the possibility to esterify with only methanol, which assures low production costs. Additionally, biodiesel from castor oil has different advantages over conventional diesel. Some of them are biodegradable, non-toxic, renewable, they can be used alone, low greenhouse gas emission, among others. This review discusses and analyzes different transesterification processes, technologies, as well as different technical aspects during biodiesel production using castor oil as a feedstock.

Biodiesel

Castor oil

Ricinus communis

1. Vegetable Oils to Produce Biodiesel

During biodiesel production it is necessary to consider the ecological and economic benefits during its production and its use as a biofuel. The biodiesel production cost depends greatly on the feedstock price. It is advisable to produce biodiesel in the same regions where the feedstock is located to decrease costs and environmental footprint related to feedstock transport [1].

The primary feedstock to produce biodiesel is raw vegetable oils and used cooking oils, as well as different animal grease. To produce biodiesel, several vegetable oils can be used such as rapeseed, soybean, cotton, peanut, corn, olive, sesame, safflower, and sunflower [2]. Some examples of tropical oily crops containing good quality oil that can be transesterified are *Raphanus sativus* (radish), *Jatropha curcas* (physic nut), *Cyperus esculentus* (tiger nut), *Simmondsia chinensis* (jojoba), *Gratissima persea* (avocado), *Lupinus albus* (white lupin), *Caryocar brasiliense* (pequi), the palm species *Acrocomia aculeata*, *Mauritia flexuosa*, *Elaeis oleifera*, *Syagrus coronata*, *Attalea speciose*, and *R. communis* (castor seeds) [3]. Table 1 shows some raw materials used in biodiesel production.

However, the most common feedstock is raw vegetable oils extracted from energy and non-energy crops such as soybean, rapeseed, sunflower, coconut, and palm oil, which are cultivated around the world $^{[\underline{4}]}$.

Table 1. Examples of oil-feedstock used in biodiesel production.

Feedstock Source (Oil Content, %)	Characteristics	Advantages	Disadvantages	Ref.
Castor beans (46–55)	Liquid at room temperature, light yellow color, and slightly pungent *US\$824/tonne of oil	Transesterification can be performed at room temperature Miscible in alcohol Non-edible Low acid value High flashpoint	Generation of toxic solid waste High viscosity Decrease fuel atomization	[<u>5]</u>
Jatropha (34–60)	Colorless after extraction (fresh) and pale yellow after standing time, liquid at room temperature **US\$250/tonne of oil	Biodiesel obtained is stable during storage Non-edible High cetane number, good oxidation stability, low viscosity	Engine corrosion due to free fatty acids Generation of toxic solid waste High cloud point Not suitable at low temperature High acid value	[Z] [<u>8</u>]
Soybean (12–22)	Fresh has a pale light color, and dark after storage, liquid at room temperature ***US\$746/tonne of oil	The yield of 98% crude biodiesel in refined oils High thermal stability Low viscosity	The high cost of production Biodiesel production in long-term is unsustainable Edible High acid value	[<u>9</u>] [<u>10]</u>
Sunflower (38–50)	Refined has a clear and vaguely yellowish-brown color Liquid at room temperature ***US\$689/tonne of oil	Low viscosity	Used to produce food and fiber Biodiesel production in long-term is unsustainable Edible High acid value	[<u>10]</u> [<u>11]</u>
Palm (18–40)	Semi-solid at room temperature, reddish and clear color, depending on extraction source (pulp or kernel) ***US\$535/tonne of oil	96% yield of crude biodiesel in refined oils Cheap feedstock Good oxidation stability Acceptable ratio of	High cloud point Conversion to biodiesel may not be sustainable long term Edible	[<u>9]</u> [<u>12</u>]

Feedstock Source (Oil Content, %)	Characteristics				Advantages saponification			vantages	Ref.	
					igh flashpoi					
	Depend	s on the c	ooking pro	cess	ow cloud po	int	High ra	atio of acid		
Oil		Saturate		ty Acid Comp Monounsa	Composition (wt%)			ated	Ref.	_
	C14:0	C16:0	C18:0	C18:1 cis-9	C18:1*	C18:2	C18:3	Others		
Castor	-	1.1	1.0	3.3	87.7	4.7	0.7	1.5	[20]	
Jatropha	-	12.80	6.20	39.94	-	45.40	-	<1.0	[<u>8</u>]	rted
Soybean	-	11.46	3.08	23.30	-	53.32	0.31	8.53	[20][21]	
Sunflower	0.08	8.03	3.26	29.27	-	59.32	-	0.04	[23]	nder
Palm	0	46.8	3.80	37.60	-	10.50	-	1.3	[24]	ition, otenti
Canola	-	3.90	1.10	64.40	-	20.4	9.60	0.6	[<u>25</u>]	nd [<u>18</u>

communis meets most of the properties considered desirable to produce useful feedstocks in biodiesel production, except the high viscosity of oil that may limit its use. They are composed mainly of triglycerides and slight amounts of free fatty acids. Table 2 shows the most abundant fatty acids in vegetable oils (palmitic oleic, and linoleic acids) from several sources. As mentioned above castor oil unlike other vegetable oils, has little variability in fatty acid food purposes and their use is restricted for this activity. Hence, castor oil for biotitel production shows several content as compared to other species oils. Ediction its ricinoleic acid, depending on its origin (ricinoleic acid) with hydroxyr groups that have a higher solubility in according to obtain methyl esters at low temperatures.

2. Castor Oil Biodiesel Production and Features

Castor oil is a light yellow and slightly pungent liquid, that is used in several industries around the world. Some of these industries are pharmaceutical, cosmetic, chemical, among others. However, in recent years this non-edible vegetable oil has been investigated to produce biofuels, specifically biodiesel. The above is due to the physicochemical, chemical, and physical characteristics of castor oil (Table 3) give to biodiesel when this is used as a feedstock.

Table 3. Main characteristics of raw castor oil.

Parameter	Zhang et al. [26]	Molefe et al. [27]	Kaur and Bhaskar ^[28]		
Acid value /(mg.g ⁻¹)	1	2.07	<4		
Saponification value /(mg.g ⁻¹)	180	175	178		

lodine value (g/100 g)	86	84	85
Refractive index (n20 D)	1.48	1.48	1.47
Relative density (g/cm ³)	0.956	0.961	0.965
Flashpoint (°C)	322	145	229
Specific heat (kJ/kg/K)	nd	0.089	0.089
Ricinic acid (%wt)	88	89.5	87-90
Oleic acid (%wt)	7	3	2-7
Linoleic acid (%wt)	5	4	1-5
Linolenic acid (%wt)	1	0.3	nd

nd: not determined

The process to produce biodiesel from oilseeds usually starts with oil extraction using a mechanical press or solvents. Figure 1 shows a general scheme for castor oil extraction. To perform mechanical oil extraction from castor beans it is necessary to consider the high oil viscosity. On the contrary, this is not necessary if the oil is extracted with solvents, but an evaporation step would be needed to recover the solvent. After extraction, different refining steps such as filtration, centrifugation, deodorization, discoloration, or winterization would be necessary to improve biodiesel quality.

References

- 1. Sadegh Azizzadeh Hajlari; Bahman Najafi; Sina F. Ardabili; Castor oil, a source for biodiesel production and its impact on the diesel engine performance. *Renewable Energy Focus* **2019**, *28*, 1-10, 10.1016/j.ref.2018.09.006.
- 2. Ragul Karthick Elango; Kiruthika Sathiasivan; Chandrasekaran Muthukumaran; Viruthagiri Thangavelu; Mathur Rajesh; Krishnamurthi Tamilarasan; Transesterification of castor oil for biodiesel production: Process optimization and characterization. *Microchemical Journal* **2019**, *145*, 1162-1168, 10.1016/j.microc.2018.12.039.
- 3. Nuria Sánchez; José María Encinar; Sergio Nogales-Delgado; Juan González; Biodiesel Production from Castor Oil by Two-Step Catalytic Transesterification: Optimization of the Process and Economic Assessment. *Catalysts* **2019**, *9*, 864, 10.3390/catal9100864.

- 4. Atiya Banerjee; Devyani Varshiney, Surentila Kumar, Payar Ciraludhary; Vinod Kumar Gupta; Biodiesel production from castor oil: ANN modeling and kinetic parameter estimation. *International Journal of Industrial Chemistry* **2017**, *8*, 253-262, 10.1007/s400 0-017-0122-3.
- 5. Bueno, A.V.; Pereira, M.P.B.; de Oliveira Pontes, J.V.; de Luna, F.M.T.; Cavalcante, C.L.;

 Periodinarical extraction Mechanical extraction Crusher

 Periodinarical extraction Mechanical extraction Crusher

 2017, 125, 559-566.

 Castor cake Crushed seeds
- 6. Hamed Bateni; Alireza Saraeian; Chad Able; Keikhosro Karimi; Biodiesel Purification and Upgrading Technologies. *Biofucial M Biorefinery Technologies* **2018**, 0, 5,7,11,00, 10.1007/978-3-03 0-00985-4_4.
- 7. A.E. Atabani; Arridina Susan Silitonga; H.C. Ong; T. M. I. Mahlia; H.H. Masjuki; Irfan Allium Badruddin; H. Fayaz; Non-edible vegetable oils: A critical evaluation of oil extraction, fatty acid compositions, biodiesel production, characteristics, engine performance and emissions production. *Renewable and Sustainable Energy Reviews* 2013, 18, 211-245, 10.1016/j.rser.2012. 10.013.
- 8. Khandual Sanghamitra; Ramirez Victorin Oramas; Rout Nutan Prasad; Comparative Yeild and Oil Quality of Toxic and Non-Toxic Mexican Jatropha curcas Grown in the Same Agroclimatic Conditions. *American Journal of Plant Sciences* **2013**, *5*, 230-234, 10.4236/ajps.2014.52030.
- 9. Titipong Issariyakul; Ajay K. Dalai; Biodiesel from vegetable oils. *Renewable and Sustainable Energy Reviews* **2014**, *31*, 446-471, 10.1016/j.rser.2013.11.001.
- 10. Ayten Sagiroglu; Hakki Mevlut Ozcan; Sebnem Selen Isbilir; Hatice Palüzar; Neslihan M. Figure 1. General scheme to produce biodiesel from castor oil (adapted from [29][30]). Toprakkiran; Alkali Catalysis of Different Vegetable Oils for Comparisons of Their Biodiesel
- Productivity. Journal of Sustainable Bioenergy Systems **2012**, 3, 79-85, 10.4236/jsbs.2013.31011. Once the biodiesel is obtained, this can be used directly in a neat form (100%) or blended with petrodiesel (25%, 150% at ad Pibzi; Danvird blenveto Medical Dosado that abeliance Gantia; blendo doseso Redes Macian teatestisk
- to the indicate feed stock soffes do included to the standard feed as the standard feed of th
- European Committee for Standardization for biodiesel (EN 14214). Jable 4 shows key parameters for biodiesel 12. Laura Cristina Barra Raiol; Fernando Kuss; André Guimarães Maciel E Silva; Bruno Cabral from castor oil and its comparison with the two standards mentioned above, as well as with petrodiesel Soares; Karla Débora Santana De Souza; Juliana Cristina Nogueira Colodo Colodo; José De
- characteristics. Castor oil biodiesel has different advantages in comparison with fossil diesel, the most significant Brito Lourenço Junior; Sandra Cristina De Avila; Nutrient intake and digestibility of the lipid are biodegradable, non-toxic, renewable, it can be used alone, low greenhouse gas emission (80% less carbon residue of biodiesel from palm oil in sheep. Revista Brasileira de Zootecnia 2012, 41, 2364-2368, dioxide, emissions and not sulfur content), and during combustion, it decreases the unburned and aromatic
- hydrocarbons. Nevertheless, one of the main disadvantages that castor oil biodiesel has is related to its high last companies in the standard and attended to its high last companies in the standard and attended to its high last companies in its last content and attended to its high last companies in its last content and attended to its high last content and attended to it
- 14e Bile tona Mile & Royale & Castor full publicums, Pails, & Bags | 18c. Buy Organic & Kosher Castor Oil
 - Drums, Pails, & Bags. Retrieved 2020-5-25

15. Jatropha Oil Table de Vatiropha Dilectude of Jatropha Dilectude a Dilectude de statropha Dilectude de Satropha Dilectude de Satr

1	Product	Kinematic Viscosity (mm²/s at 40 °C)	Density (kg/m³)	Acid Value (mgKOH/g)	Flash Point (°C)	Water Content (%)	Calorific Value (MJ/kg)	Cetane Number	Cloud Point (°C)	Pour Point (°C)	lodine Value (gl ₂ /100 g)	Ref.	arket	
1		23	960	13.12	262	nd	30.18	nd	3	-13	nd	[<u>32</u>]	lugo	
		14	926	nd	164	nd	37.90	nd	-23	nd	nd	[33]	erived 013-53	
1	*Biodiesel from	15	946	0.63	194	0.15	38.34	43	nd	-30	nd	[<u>34</u>]	ando	
7	castor oil	26	961	1.19	nd	0.31	nd	nd	nd	nd	80.5	[28]		
2		18	920	0.25	170	0.006	39.5	nd	nd	nd	nd	[<u>35</u>]	ustry. L38.	
2		14	923	nd	273.1	nd	37.34	50	nd	nd	83.40	[<u>36</u>]	by nass	
2	**Biodiesel according ASTM D 6751–06a	1.9–6.0	860– 900	≤0.50	≥130	≤0.50	nd	≥47	-3- -12	-15- 10	nd	[<u>37</u>]	a new	
2	**Biodiesel according EN 14214	3.5–5.0	860– 900	≤0.50	≥101	≤0.50	nd	≥51	nd	nd	nd	[<u>37</u>]	/j.rene olysis	
	***Fossil Diesel	2.5–4.5	820– 860	nd	68– 80	nd	≥45.56	≥46	-15-5	-35- 15	nd	[<u>38</u>]	oc.201	

- 24. Patrícia C. M. Da Rós; Larissa Freitas; Victor Haber Perez; Heizir F. De Castro; Enzymatic synthesis of biodiesel from palm oil assisted by microwave irradiation. *Bioprocess Engineering* **2012**, *36*, 443-451, 10.1007/s00449-012-0801-6.
- 25. A.C.F. Batista; A.T. Vieira; H.S. Rodrigues; T.A. Silva; R.M.N. Assunção; M.A. Beluomini; H.P. Rezende; Manuel Gonzalo Hernandez-Terrones; PRODUCTION AND PHYSICOCHEMICAL CHARACTERIZATION OF METHYLIC AND ETHYLIC BIODIESEL FROM CANOLA OIL / OBTENÇÃO E CARACTERIZAÇÃO DO BIODIESEL DE CANOLA PELAS ROTAS METÍLICA E ETÍLICA. Revista Brasileira de Engenharia de Biossistemas 2014, 8, 289, 10.18011/bioeng2014v *Wilmalp2896293.00)

- nd: not determined
- 27. Kasirajan Ramachandran; Pandian Sivakumar; Tamilarasan Suganya; Sahadevan Renganathan; As Production of biodicsal topoliniated wasted esetables of husing laboriorism discontinuation of the single point is a heterogeneous acid catalyst. *Bioresource Technology* **2011**, *102*, 7289-7293, 10.1016/j.biortec ENIL420111 European Committee for Standardization for biodiesel.
- 28. Rayneet Kaur. Thallada Bhaskar: Potential of castor plant (Ricinus communis) for production of Briodiesel Production around in word: is produced from different (Ricinus communis) for production of oils. Moreover, a Wide variety of methods to obtain this blotter exist, however, the most common is through transesterification. Biodiesel from castor oil offers environmental and technical benefits, therefore, it can be 2903 and a several ender white interesting the self-base furies of the identification. Biodiesel from castor oil offers environmental and technical benefits, therefore, it can be 2903 and a several ender white interesting the self-base furies of the identification. Biodiesel from castor oil offers environmental and technical benefits, therefore, it can be 2903 and a several ender the interesting the interesting the interesting of the interest
- 33. Aihua Zhang; Qin Wang; Yidan He; Pengying Lai; Yifu Miu; Zhihong Xiao; Preparation of Biodiesel Based on Alkaline Ionic Liquid [Bmim]OH Catalyzed Castor Oil. *IOP Conference Series: Materials Science and Engineering* **2020**, *729*, 012048, 10.1088/1757-899x/729/1/012048.
- 34. Masego Molefe; Diakanua Nkazi; Hembe E Mukaya; Method Selection for Biojet and Biogasoline Fuel Production from Castor Oil: A Review. *Energy & Fuels* **2019**, *33*, 5918-5932, 10.1021/acs.en ergyfuels.9b00384.
- 35. Jibrin Mohammed Danlami; Agus Arsad; Muhammad Abbas Ahmad Zaini; Characterization and process optimization of castor oil (Ricinus communis L.) extracted by the soxhlet method using polar and non-polar solvents. *Journal of the Taiwan Institute of Chemical Engineers* **2015**, *47*, 99-104, 10.1016/j.jtice.2014.10.012.
- 36. Kiany S.B. Cavalcante; Maria N.C. Penha; Karlene K.M. Mendonça; Hilton C. Louzeiro; Antonio C.S. Vasconcelos; Adeilton P. Maciel; Antonio G. De Souza; Fernando Silva; Optimization of

- transesterification of castor oil with ethanol using a central composite rotatable design (CCRD). *Fuel* **2010**, *89*, 1172-1176, 10.1016/j.fuel.2009.10.029.
- 37. Hemant Y Shrirame; N. L. Panwar; B. R. Bamniya; Bio Diesel from Castor Oil A Green Energy Option. *Low Carbon Economy* **2010**, *2*, 1-6, 10.4236/lce.2011.21001.
- 38. Dilip Kumar; Tapas Das; Balendu Shekher Giri; Bhawna Verma; Preparation and characterization of novel hybrid bio-support material immobilized from Pseudomonas cepacia lipase and its application to enhance biodiesel production. *Renewable Energy* **2020**, *147*, 11-24, 10.1016/j.rene ne.2019.08.110.

Retrieved from https://encyclopedia.pub/entry/history/show/7699