

HCFC1 Gene

Subjects: Genetics

Submitted by:  Dean Liu

(This entry belongs to Entry Collection "["MedlinePlus"](#)")

Definition

Host cell factor C1

1. Introduction

The *HCFC1* gene provides instructions for making a protein, called HCF-1, that helps regulate the activity of other genes. HCF-1 interacts with proteins called transcription factors, which attach (bind) to specific regions of DNA and help control the activity of particular genes.

A specific function of the HCF-1 protein is to control the activity of genes involved in the processing of vitamin B12 (also known as cobalamin), particularly the *MMACHC* gene. This gene plays a role in the conversion of vitamin B12 into one of two molecules, adenosylcobalamin (AdoCbl) or methylcobalamin (MeCbl). AdoCbl is required for the normal function of an enzyme known as methylmalonyl CoA mutase. This enzyme helps break down certain protein building blocks (amino acids), fats (lipids), and cholesterol. AdoCbl is called a cofactor because it helps methylmalonyl CoA mutase carry out its function. MeCbl is also a cofactor, but for an enzyme known as methionine synthase. This enzyme converts the amino acid homocysteine to another amino acid, methionine. The body uses methionine to make proteins and other important compounds.

HCF-1 helps regulate genes that are important in other cellular processes, such as progression of cells through the step-by-step process it takes to replicate themselves (called the cell cycle). This protein also plays a role in the distribution of cells in developing tissues and organs, including the brain.

2. Health Conditions Related to Genetic Changes

2.1. Methylmalonic Acidemia with Homocystinuria

At least six *HCFC1* gene mutations have been identified in individuals with methylmalonic acidemia with homocystinuria, cbIX type, one form of a disorder that causes developmental delay, eye defects, neurological problems, and blood abnormalities. Individuals with this form also have severe abnormalities in the development of the skull and face (craniofacial abnormalities). These mutations occur in regions of the protein that help it to interact with other proteins. It is thought that changes in these regions prevent HCF-1 from interacting with transcription factors, which disrupts normal gene activity. Impairment of *MMACHC* gene activity, in particular, prevents normal processing and transport of vitamin B12, impeding production of both AdoCbl and MeCbl. Because both of these cofactors are missing, the enzymes that require them (methylmalonyl CoA mutase and methionine synthase) do not function normally. As a result, certain amino acids, lipids, and cholesterol are not broken down and homocysteine cannot be converted to methionine. This dual defect results in a buildup of toxic compounds as well as homocysteine, and a decrease in the production of methionine within the body. This combination of imbalances leads to the signs and symptoms of methylmalonic acidemia with homocystinuria. Neurological and developmental problems are especially severe in individuals with cbIX type, in part due to disruption of the activity of other genes normally regulated by the HCF-1 protein.

2.2. Other Disorders

Mutations in the *HCFC1* gene have also been found in individuals with X-linked intellectual disability. These individuals have delayed development and other neurological problems but do not show other features of methylmalonic acidemia with homocystinuria, cbIX type. The *HCFC1* gene mutations lead to

production of an HCF-1 protein with reduced function. Partial reduction in this protein's function appears to disrupt normal brain development, leading to the features of X-linked disability, but does not severely impact vitamin B12 processing.

3. Other Names for This Gene

- CFF
- HCF
- HCF-1
- HCF1
- HFC1
- host cell factor 1
- MGC70925
- MRX3
- PPP1R89
- protein phosphatase 1, regulatory subunit 89
- VCAF
- VP16-accessory protein

References

1. Gérard M, Morin G, Bourillon A, Colson C, Mathieu S, Rabier D, Billette deVillemeur T, Ogier de Baulny H, Benoist JF. Multiple congenital anomalies in two boys with mutation in HCFC1 and cobalamin disorder. *Eur J Med Genet.* 2015Mar;58(3):148-53. doi: 10.1016/j.ejmg.2014.12.015.
2. Huang L, Jolly LA, Willis-Owen S, Gardner A, Kumar R, Douglas E, Shoubridge C, Wieczorek D, Tzschach A, Cohen M, Hackett A, Field M, Froyen G, Hu H, Haas SA, Ropers HH, Kalscheuer VM, Corbett MA, Gecz J. A noncoding, regulatory mutation implicates HCFC1 in nonsyndromic intellectual disability. *Am J Hum Genet.* 2012Oct 5;91(4):694-702. doi: 10.1016/j.ajhg.2012.08.011.
3. Jolly LA, Nguyen LS, Domingo D, Sun Y, Barry S, Hancarova M, Plevova P, Vlckova M, Havlovicova M, Kalscheuer VM, Graziano C, Pippucci T, Bonora E, Sedlacek Z, Gecz J. HCFC1 loss-of-function mutations disrupt neuronal and neural progenitor cells of the developing brain. *Hum Mol Genet.* 2015 Jun 15;24(12):3335-47. doi: 10.1093/hmg/ddv083.
4. Parker JB, Yin H, Vinckevicius A, Chakravarti D. Host cell factor-1 recruitment to E2F-bound and cell-cycle-control genes is mediated by THAP11 and ZNF143. *Cell Rep.* 2014 Nov 6;9(3):967-82. doi: 10.1016/j.celrep.2014.09.051.
5. Yu HC, Sloan JL, Scharer G, Brebner A, Quintana AM, Achilly NP, Manoli I, Coughlin CR 2nd, Geiger EA, Schneck U, Watkins D, Suormala T, Van Hove JL, Fowler B, Baumgartner MR, Rosenblatt DS, Venditti CP, Shaikh TH. An X-linked cobalamin disorder caused by mutations in transcriptional coregulator HCFC1. *Am J Hum Genet.* 2013 Sep 5;93(3):506-14. doi: 10.1016/j.ajhg.2013.07.022.
6. Zargar Z, Tyagi S. Role of host cell factor-1 in cell cycle regulation. *Transcription.* 2012 Jul-Aug;3(4):187-92. doi: 10.4161/trns.20711.

Keywords

genes
