

Drought/Salt Tolerance in Wheat

Subjects: Plant Sciences

Contributors:  Mario A. Pagnotta,  Ieva Urbanavičiūtė,  Luca Bonfiglioli

Submitted by:  Mario A. Pagnotta

Definition

Drought and salinity are major constraints to agriculture. Here, we present an overview of the global situation and the consequences of drought and salt stress connected to climatic changes.

1. Global Situation

Water and consequently an adequate hydric balance are essential for life, determining crop production in terms of quantity and quality. In the present century, the main crop production problems are related to hydric stress, including dwindling water resources and excessive salinity of irrigation water and soil. Moreover, it is expected that global warming and climate change would increase the frequency of drought events, leading to a severe threat to food security ^{[1][2][3]}.

In nature, plants are exposed to various abiotic factors and exhibit complex adaptation responses that depend on their degree of plasticity. In addition, they are facing environmental changes that increase the risk factors, such as rainfall distribution, loss of soil fertility and organic carbon, fluctuation in temperature and light, evaporative intensity, biological stresses, increasing pollution, and declining biodiversity ^[4]. Water is the most critical factor for the growth and development of living organisms, determining 82% of the variation in grain yield in areas receiving less than 400 mm of annual rainfall ^[5]. Land plants are anchored in the soil and rely on their root system to ensure water availability and nutrient uptake. About 30% of the Earth's land is arid or semi-arid, and this percentage is expected to increase due to climate change ^[6]. The prevision indicates an erratic rainfall distribution, an overall decrease in total annual rainfall, and a more pronounced occurrence of dry periods ^[7], in addition to an increase in temperatures with a consequent reduction in ecosystem health through species extinction or a reduction in biodiversity, due to migration and changes in behavioral patterns ^[8].

Among all stresses, drought and salinity (soil and water) are the main abiotic factors that reduce a crop's productivity, causing the most damage for grain production all over the world ^[9]. Other than grain production, water scarcity also affects the growth rate, leaf size, stem extension, root proliferation, susceptibility to disease, plant color, etc. ^{[10][11]}. Particularly in conventional farming, the rainfall water effectively decreases due to increased surface runoff, soil evaporation through unplanted and bare soil surfaces, and evaporation caused by the aeration of soil during intensive tillage ^[3]. Factors such as rainfall intensity, duration, exact time of rain events, and the soil properties affect the soil moisture status, especially the plant-available water (PAW), which can differ significantly with the same total amount of precipitation under different conditions. In addition, the water availability for plants decreases due to the high soluble salt concentrations in the soil, which causes absorption difficulties; i.e., water deficit or hydric stress. The main causes of soil salinization can be classified as natural or caused by human activity, named primary and secondary salinization, respectively (Figure 1) ^[12].




Figure 1. Primary and secondary soil salinity mechanisms (Reprinted with permission from ref. [12]. 2016, Elsevier).

Current estimations indicate that salt issues affect about 20% of global lands and almost 40% of irrigated lands. This phenomenon is increasing and 50% of total cultivated land worldwide will be salinized by 2050 [13][14][15]. Global warming is the increase in temperatures often associated with rainfall decrease, leading to more extreme and unpredictable events. As a conclusion, the need to have a system protecting agriculture from an erratic climate becomes important worldwide.

The global grain production is an important indicator of food security, but unfortunately it grows slower than the human population. This unfavorable situation is exacerbated by abiotic factors such as drought, high salinity, extreme temperatures, flooding, and heavy metals [16][17]. There is an urgent need to increase food, which cannot be done by simply raising the agricultural lands, which causes deforestation and consequently climate change problem. Increasing the production per unit area (i.e., yields) represents a potential solution. Yields can be improved by using genetics methods, appropriate crop rotation, adequate fertilization, and early planting. For example, a six-week delay in planting time resulted in yield reductions of about 42% in barley and 22–32% in wheat [18]. While, early planting, proper fertilization, and appropriate crop rotation were found to increase the water-use efficiency (WUE) [19].

Useful and adequate breeding programs, addressed to reduce crop yield lost under stress conditions, are necessary to face the consequences of climate change. Attention should be focused on the drought-related traits, which include leaf canopy temperature, cell osmotic adjustment, cell membrane stability, leaf water potential, stomatal resistance, leaf rolling index, and leaf waxiness [18]. In wheat, for example, the total aboveground part of the plant (biological yield) in a water-limited production area is directly related to the water supply [20]. Although biological yield offers little hope of modification through breeding, the harvest index, which reflects the ratio of grain to total biological yield, appears to be amenable to genetic improvement for enhanced drought tolerance.

2. Germplasm

Significant differences in drought and salinity stresses were reported between plant species [21]. Their growth rates were highly different under stress, indicating that germplasm is the main reserve of useful genes that could be used to improve cultivated crops. Figure 2 shows the differences in salt sensitivity among some plant species. Tall wheat grass (*Thinopyrum ponticum* (Podp.) Z.-W. Liu and R.-C. Wang; (synonymous *Agropyron elongatum* (Host) P. Beauv.) and saltbush (*Atriplex anicola* Paul G. Wilson) are extremely tolerant to salinity. *Arabidopsis thaliana* (L.) is more sensitive to salinity while alfalfa (*Medicago sativa* L.) is much more tolerant, compared to cereals. Among the cereals, rice (*Oryza sativa* L.) is the most sensitive while barley (*Hordeum vulgare* L.) is the most tolerant. Within the intermediate tolerant species, durum wheat (*Triticum turgidum* subsp. *durum* (Desf.) Husn) is more sensitive than bread wheat (*Triticum aestivum* L.) [21].




Figure 2. Salt tolerance differences among various species, expressed as NaCl concentration to inhibit dry matter production [21].

3. Morphological and Physiological Response

When plants suffer from hydric stress, significant changes in their morphology can be observed. Usually, hydric stresses affect plant size, which becomes smaller due to a decreasing number of leaves and decreasing area. However, the roots are lengthened searching for water, which results in an increase in the root-to-shoot ratio. This adverse effect of water scarcity on crop plants causes fresh and dry biomass losses. Early maturity or early flowering is another adaptation strategy where a shorter vegetative phase in wheat can help to avoid stress in further very sensitive flowering and post-anthesis grain filling stages [22].

Drought induces a plethora of negative physiological alterations, such as cell turgor loss, reduction in CO₂ assimilation, oxidative stress, and nutritional imbalance [23]. Nutritional imbalances due to drought stress occur by decreasing the water uptake and leaf transpiration, combined with alterations in nutrient uptake. Plants try to counteract these effects by activating drought resistance mechanisms, such as accumulation of salts and water, to improve the cell osmotic adjustment and stomata function. K⁺ and Cl⁻ ion regulation, as well water transport, have been important mechanism for plants under drought. K⁺ and Cl⁻ are involved in leaf cells osmotic adjustment, with the consequent water retention in cells; stomatal closure prevents water loss [24]. The presence of high ionic concentrations in soil, especially NaCl, is a great challenge for the plant's physiology. To balance the high sodium concentrations in the soil, plants cells must maintain high concentration of potassium and low concentration of sodium inside the cells.

The root system has been studied less than the aboveground morphology. During the juvenile phases, the root architecture is plastic and can adapt to several environmental conditions. It has been detected that roots can escape a high salinity zone, moving to less salty soil. Galvan-Ampudia et al. [25] identified halotropism as the plant's possibility to reduce their exposure to salinity by moving their root system to an adequate saline environment. The auxin distribution in the root tip affects the root response to salty environments, as summarized in Figure 3 [25]. Furthermore, changes in the PIN subcellular localization affect root auxin transport and can cause various deformities in root architecture, including the size, less lateral roots, root meristem collapse, and others [26].




Figure 3. Salt-activated phospholipase D (PLD) induces clathrin-mediated endocytosis of PIN2, which allows reorganizing the auxin and roots' growing direction (modified from Galvan-Ampudia et al. [25]).

Salinity inhibits the root length and reduces the number of roots, but also influences the root growth direction. Sun et al. [27] found that the root direction could change with NaCl increasing; the curved roots could be visible already with a concentration of 100 mM of NaCl in the media. Salt stress reduces gravitropism of root growth, altering the PIN-FORMED2 (PIN2) protein abundance and polar distribution and overly salt-sensitive (*sos1* and *sos2*) genes. Julkowska et al. [28] found that more than 100 genetic loci activated under salt stress are associated with root system architecture changes. Among these, *CYP79B2* is correlated with lateral root development under salt stress while *HKT1* reduce the lateral root development.

4. Mechanisms of Tolerance

Since plants cannot easily change habitats where conditions are more favorable, to survive and multiply, they evolutionarily adapted, gaining plasticity and specific responses. Generally, plant reactions to abiotic factors include morphological, physiological, biochemical responses, and various adaptation scenarios. In addition, a plant's response can be either common to many kinds of stresses or specific to each one. Stress-tolerance mechanisms are divided in two macro groups: mechanisms involved in osmotic-stress tolerance and mechanisms involved in ion-stress tolerance [29]. The stomatal adjustment belongs to the first class of mechanisms. Plants under drought and osmotic stresses in non-irrigated systems tend to close their stomata to minimize transpiration, which is reflected in a reduction in growth and production. Exclusion and compartmentalization of toxic ions, avoiding their high concentration in plant cells, are mechanisms of ion-stress tolerance [21]. Each type of physical stress is translated by the plant into a biochemical response, and then into a pathway of interconnected signals with the expression of specific genes that are involved in the stress-tolerance mechanisms. Table 1 lists some of the genes that are candidates for the pathway process and their defense mechanisms for osmotic and ion stresses [21].

Table 1. Plant salinity-tolerance mechanisms, ordered by processes and their relevance for the three components of salinity tolerance [21].

		Osmotic Stress	Ionic Stress		
Process	Candidate Genes	Osmotic Tolerance	Na ⁺ Excluding	Tissue Tolerance	References
Signaling	SOS3, SnRKs	Signaling regulation	Activation of ion antiporter	Regulation of vacuolar loading	[30]
Photosynthesis	ERA1, PP2C, AAPK, PKS3	Stomatal closure regulation	Protection of chloroplast from ion toxicity	Delay Na ⁺ toxicity effect in chloroplast	[31][32]
Accumulation of Na ⁺ in shoots	HKT, SOS1	-	Decreasing long distance transport of Na ⁺	Decreasing energy used on Na ⁺ exclusion	[33][34][35]
Accumulation of Na ⁺ in vacuoles	NHX, AVP	-	Increased sequestration of Na ⁺ into root vacuoles	Increased sequestration of Na ⁺ into leaf vacuoles	[33][36]
Accumulation of organic solutes	P5CS, OTS, MT1D, M6PR, S6PDH, IMT1	Increasing osmotic adjustment	Reduction of Na ⁺ accumulation	Accumulation of organic solutes in cytoplasm	[35][37]

References

1. Mann, M.E.; Gleick, P.H. Climate change and California drought in the 21st century. *Proc. Natl. Acad. Sci. USA* 2015, 112, 3858–3859.
2. Nam, W.H.; Hayes, M.J.; Svoboda, M.D.; Tadesse, T.; Wilhite, D.A. Drought hazard assessment in the context of climate change for South Korea. *Agric. Water Manag.* 2015, 160, 106–117.
3. Bot, A.; Benites, J. Drought-resistant soils: Optimization of Soil Moisture for Sustainable Plant Production. In *Proceedings of the Electronic Conference 'Drought-Resistant Soils. Optimization of Soil Moisture for Sustainable Plant Production.'* FAO Land and Water Development Division, 15 November–18 December 2004; Volume 11.
4. Cramer, W.; Guiot, J.; Fader, M.; Garrabou, J.; Gattuso, J.P.; Iglesias, A.; Lange, M.A.; Lionello, P.; Llasat, M.C.; Paz, S.; et al. Climate change and interconnected risks to sustainable development in the Mediterranean. *Nat. Clim. Chang.* 2018, 8, 972–980.
5. Lawlor, D.W.; Mitchell, R.A. Crop ecosystem responses to climatic change. In *Climate Change and Global Crop Productivity*; Nobel, P.S., Reddy, K.R., Hodges, H.F., Eds.; CABI Publishing: Wallingford, UK, 2001; pp. 57–80.

6. Huang, J.; Zhang, W.; Zuo, J.; Bi, J.; Shi, J.; Wang, X.; Chang, Z.; Huang, Z.; Yang, S.; Zhang, B.; et al. An overview of the semi-arid climate and environment research observatory over the loess plateau. *Adv. Atmos. Sci.* 2008, 25, 906–921.
7. Houghton, J.T.; Albritton, D.L.; Meira Filho, L.G.; Cubasch, U.; Dai, X.; Ding, Y.; Karl, T. Technical summary of working group 1. In *Intergovernmental Panel on Climate Change (IPCC), Climate Change 2001: The Scientific Basis*; Cambridge University Press: Cambridge, UK, 2001; pp. 3–45.
8. Rani, G.; Kaur, J.; Kumar, A.; Yogalakshmi, K.N. Ecosystem health and dynamics: An indicator of global climate change. In *Contemporary Environmental Issues and Challenges in Era of Climate Change*; Springer: Singapore, 2019; pp. 1–32.
9. Shrivastava, P.; Kumar, R. Soil salinity: A serious environmental issue and plant growth promoting bacteria as one of the tools for its alleviation. *Saudi J. Biol. Sci.* 2015, 22, 123–131.
10. Wegulo, S.; Giesler, L.; Harveson, R.; Jackson-Ziems, T.; Liu, B.; Korus, K. 2013 Crop Production Clinic Proceedings, Impacts of Drought on Disease Development and Management. *Pap. Plant Pathol.* 2013, 125–127.
11. Farooq, M.; Wahid, A.; Kobayashi, N.; Fujita, D.; Basra, S.M.A. Plant drought stress: Effects, mechanisms and management. *Agron. Sustain. Dev.* 2009, 153–188.
12. Daliakopoulos, I.N.; Tsanis, I.K.; Koutroulis, A.; Kourgialas, N.N.; Varouchakis, A.E.; Karatzas, G.P.; Ritsema, C.J. The threat of soil salinity: A European scale review. *Sci. Total Environ.* 2016, 573, 727–739.
13. Allen, M.R.; Ingram, W.J. Constraints on future changes in climate and the hydrologic cycle. *Nature* 2002, 419, 228–232.
14. Rozema, J.; Flowers, T.J. Crops for salinized world. *Science* 2008, 322, 1478–1480.
15. Konapala, G.; Mishra, A.K.; Wada, Y.; Mann, M.E. Climate change will affect global water availability through compounding changes in seasonal precipitation and evaporation. *Nat. Commun.* 2020, 11, 1–10.
16. Ali, S.; Rizwan, M.; Qayyum, M.F.; Ok, Y.S.; Ibrahim, M.; Riaz, M.; Arif, M.S.; Hafeez, F.; Al-Wabel, M.I.; Shahzad, A.N. Biochar soil amendment on alleviation of drought and salt stress in plants: A critical review. *Environ. Sci. Pollut. Res.* 2017, 24, 12700–12712.
17. Ahmad, P.; Prasad, M.N.V. *Abiotic Stress Responses in Plants: Metabolism, Productivity and Sustainability*; Springer Science + Business Media, LLC: New York, NY, USA, 2012.
18. Lawlor, D.W. Stress metabolism: Its implication in breeding programmes. In *Drought Tolerance in Winter Cereals*; Srivastava, E.J.P., Porceddu, E., Acevedo, S., Eds.; Varma. Icarda: Aleppo, Syria, 1987.
19. Hatfield, J.L.; Dold, C. Water-use efficiency: Advances and challenges in a changing climate. *Front. Plant Sci.* 2019, 10, 103.
20. Richards, R.A.; Passioura, J.B. Seminal Root Morphology and Water Use of Wheat I. *Environmental Effects I. Crop Sci.* 1981, 21, 249–252.
21. Munns, R.; Tester, M. Mechanisms of salinity tolerance. *Annu. Rev. Plant Biol.* 2008, 59, 651–681.
22. Shavrukov, Y.; Kurishbayev, A.; Jatayev, S.; Shvidchenko, V.; Zotova, L.; Koekemoer, F.; De Groot, S.; Soole, K.; Langridge, P. Early flowering as a drought escape mechanism in plants: How can it aid wheat production? *Front. Plant Sci.* 2017, 8, 1950.
23. Farooq, M.; Wahid, A.; Kobayashi, N.; Fujita, D.; Basra, S.M.A. Plant drought stress: Effects, mechanisms and management. In *Sustainable Agriculture*; Springer: Cham, Switzerland, 2009; pp. 153–188. ISBN 9789048126651.
24. Nieves-Cordones, M.; García-Sánchez, F.; Pérez-Pérez, J.G.; Colmenero-Flores, J.M.; Rubio, F.; Rosales, M.A. Coping With Water Shortage: An Update on the Role of K⁺, Cl⁻, and Water Membrane Transport Mechanisms on Drought Resistance. *Front. Plant Sci.* 2019, 10, 1619.
25. Galvan-Ampudia, C.S.; Julkowska, M.M.; Darwish, E.; Gandullo, J.; Korver, R.A.; Brunoud, G.; Haring, M.A.; Munnik, T.; Vernoux, T.; Testerink, C. Halotropism is a response of plant roots to avoid a saline environment. *Curr. Biol.* 2013, 23, 2044–2050.
26. Dai, M.; Zhang, C.; Kania, U.; Chen, F.; Xue, Q.; McCray, T.; Li, G.; Qin, G.; Wakeley, M.; Terzaghi, W.; et al. A PP6-type phosphatase holoenzyme directly regulates PIN phosphorylation and auxin efflux in Arabidopsis. *Plant Cell* 2012, 24, 2497–2514.
27. Sun, F.; Zhang, W.; Hu, H.; Li, B.; Wang, Y.; Zhao, Y.; Li, K.; Liu, M.; Li, X. Salt modulates gravity signaling pathway to regulate growth direction of primary roots in arabidopsis. *Plant Physiol.* 2008, 146, 178–188.
28. Julkowska, M.M.; Koevoets, I.T.; Mol, S.; Hoefsloot, H.; Feron, R.; Tester, M.A.; Keurentjes, J.J.B.; Korte, A.; Haring, M.A.; De Boer, G.J.; et al. Genetic components of root architecture remodeling in response to salt stress. *Plant Cell* 2017, 29, 3198–3213.
29. Mondini, L.; Pagnotta, M.A. *Drought and Salt Stress in Cereals*; Springer: Cham, Switzerland, 2015; pp. 1–31.
30. Sun, T.; Wang, Y.; Wang, M.; Li, T.; Zhou, Y.; Wang, X.; Wei, S.; He, G.; Yang, G. Identification and comprehensive analyses of the CBL and CIPK gene families in wheat (*Triticum aestivum* L.). *BMC Plant Biol.* 2015, 15, 1–17.
31. Bartels, D.; Sunkar, R. Drought and salt tolerance in plants. *CRC. Crit. Rev. Plant Sci.* 2005, 24, 23–58.
32. Zhu, J.K. Salt and drought stress signal transduction in plants. *Annu. Rev. Plant Biol.* 2002, 53, 247–273.
33. Faïc Xal Brini, F.; Hanin, M.; Mezghani, I.; Berkowitz, G.A.; Masmoudi, K. Overexpression of wheat Na⁺/H⁺ antiporter TNH1 and H⁺-pyrophosphatase TVP1 improve salt-and drought-stress tolerance in Arabidopsis thaliana plants. *J.*

Exp. Bot. 2007, 58, 301–308.

34. Ramezani, A.; Niazi, A.; Abolmoghadam, A.A.; Zamani Babgohari, M.; Deihimi, T.; Ebrahimi, M.; Akhtardanesh, H.; Ebrahimie, E. Quantitative expression analysis of TaSOS1 and TaSOS4 genes in cultivated and wild wheat plants under salt stress. *Mol. Biotechnol.* 2013, 53, 189–197.
35. Munns, R. Genes and salt tolerance: Bringing them together. *New Phytol.* 2005, 167, 645–663.
36. Zhang, Y.M.; Zhang, H.M.; Liu, Z.H.; Li, H.C.; Guo, X.L.; Li, G.L. The wheat NHX antiporter gene TaNHX2 confers salt tolerance in transgenic alfalfa by increasing the retention capacity of intracellular potassium. *Plant Mol. Biol.* 2015, 87, 317–327.
37. Tavakoli, M.; Poustini, K.; Alizadeh, H. Proline accumulation and related genes in wheat leaves under salinity stress. *J. Agric. Sci. Technol.* 2016, 18, 707–716.

Keywords

osmotic adjustment;ionic;redox homeostasis;transcription factors;salt-responsive genes;genetic diversity;germplasm;cross-transferability

Retrieved from <https://encyclopedia.pub/12199>