

DUX4-rearranged leukaemia

Subjects: **Oncology**

Contributor: Jacqueline A. Rehn , Matthew J. O'Connor , Deborah L. White , David T. Yeung

DUX4-rearrangement (*DUX4r*) is a recently discovered recurrent genomic lesion reported in 4-7% of B cell acute lymphoblastic leukaemia (B-ALL). The gene fusion most commonly links the hypervariable *IGH* gene to *DUX4* a gene located within the D4Z4 macrosatellite repeat on chromosome 4. *DUX4r* is cryptic to most standard diagnostic techniques and difficult to identify even with next-generation sequencing assays.

acute lymphoblastic leukaemia

DUX4

ERG

DUX4-IGH

DUX4-rearrangement

1. Introduction

B-cell acute lymphoblastic leukaemia (B-ALL) is a malignant disorder of the bone marrow resulting in over proliferation of immature B lymphoblasts. The disease can manifest at any age but the majority of patients are children, making B-ALL the most common childhood malignancy ^[1]. This heterogeneous disease is characterised by a variety of different genomic alterations including changes in chromosome number, chromosomal translocations and single nucleotide variants (SNV). Detection of the underlying genomic alterations assists clinical risk stratification and therapeutic triage. Cytogenetic analysis has proven adept at identifying several recurrent genomic alterations which result in diseases with distinct gene expression profiles (GEP) and defined prognosis. This includes high hyperdiploidy, hypodiploidy, and the translocations t(12;21) [*ETV6-RUNX1*], t(9;22) *BCR-ABL1*, t(1;19) *TCF3-PBX1* and alterations of chromosome 11q23 resulting in rearrangement of *KMT2A/MLL*. These alterations account for approximately 60% of pediatric B-ALL cases ^{[2][3]}. Remaining patients were historically classified as *B-other* and demonstrated highly variable prognosis and treatment response.

Molecular studies involving GEP and next-generation sequencing (NGS) have subsequently identified a number of additional recurrent molecular alterations not detectable with standard cytogenetics, several of which may be targetable by precision medicine approaches. This includes the newly recognized subtype of Philadelphia chromosome-like (Ph-like) ALL characterized by a gene expression profile similar to cases with a *BCR-ABL1* translocation, but instead carrying one of multiple kinase activating lesions. One example is rearrangement of the cytokine receptor gene *CRLF2* (*CRLF2r*), commonly with concurrence of Janus Kinase 2 (*JAK2*) mutations ^{[4][5]}, affecting 5–7% of children with B-ALL ^[3]. NGS has also recently identified a rearrangement of the homeodomain encoding the Double Homeobox 4 (*DUX4*) transcription factor with the immunoglobulin heavy chain (*IGH*) locus which results in a distinct genetic subtype.

As early as 2002 researchers identified a novel B-ALL subtype, with a distinct microarray GEP, not associated with any known recurrent genomic alterations, that appeared to confer a good prognosis. Interrogation of

overexpressed genes in these patients failed to uncover a causative lesion [6]. Follow-up studies involving copy number alteration (CNA) analysis revealed many of the patients with the distinct expression profile also demonstrated deletion of the *ERG* gene (ETS transcription factor *ERG*), a genomic alteration absent in almost all other subtypes. *ERG* deletion was consequently proposed as the driving lesion in this subtype [7][8]. Multiple studies have subsequently demonstrated that monoallelic deletion of *ERG* is observed in only a subset of patients demonstrating this GEP. Furthermore, *ERG* deletions were subclonal in several patients at diagnosis and either altered or absent at relapse [9][10][11].

In 2016, two independent studies identified rearrangement of the *DUX4* locus (*DUX4r*), most commonly partnered with *IGH*, present in patients with the previously detected GEP [12][13]. Transduction of the *DUX4* fusion transcript into NIH3T3 fibroblasts resulted in cellular transformations, demonstrating the oncogenic potential of this alteration [12]. Multiple studies have subsequently confirmed the unique GEP of *DUX4r* cases and identified the rearrangement in 4–7% of B-ALL patients [9][14][15][16][17][18][19], as well as within the NALM6 cell line [12]. Leukaemic cells carrying the *DUX4r* also display a unique methylation profile, associated with widespread hypomethylation [15][20][21], and express a specific non coding RNA signature [16][21]. Consequently, *DUX4r*, also reported in the literature as *DUX4/ERG*, has increasingly been accepted as a distinct molecular subtype in B-ALL. In this review, we present the current understanding of the molecular structure and biological effects of the *DUX4r*.

2. Description of the *DUX4* Rearrangement

The *DUX4* gene is present within each repeat of the D4Z4 tandem array located in the subtelomeric region of chromosome 4q [22], with an almost identical locus (>98% nucleotide identity) on 10q [23][24]. The D4Z4 array is polymorphic in length containing between 11–100 copies of the 3.3 kb repeat in healthy adults (Figure 1) [25]. In healthy tissue, transcription of *DUX4* is restricted to germline cells of the testes. Transcription has also been observed in induced pluripotent stem cells, suggesting a role for *DUX4* in germline development [26][27]. Expression of the full-length *DUX4* transcript is epigenetically silenced in somatic tissue [27]. Only the first exon of the spliced transcript contains a coding sequence for the protein which consists of two N-terminal homeodomains capable of DNA binding [28] and a C-terminal transactivation domain [29]. The *DUX4* protein is capable of binding to, and upregulating expression of, multiple genes as well as initiating expression from alternate promoters, producing non-canonical transcript isoforms [29].

Contraction of the D4Z4 region resulting in fewer than 10 repeats is associated with facioscapulohumeral muscular dystrophy (FSHD) [25][30], a genetically inherited disorder that initially manifests as progressive weakening of the facial, shoulder and upper arm muscles [31]. Partial deletion of the D4Z4 array is associated with hypomethylation and loss of repressive histone modifications that are believed to reduce chromatin packing of the subtelomeric region allowing *DUX4* expression [32][33]. Intriguingly, FSHD only manifests in patients who demonstrate D4Z4 contraction on chromosome 4q and not the homologous array on chromosome 10q. Furthermore, contraction of an alternative chromosome 4q allele (4qB) [24], does not result in disease. Sequencing efforts have subsequently revealed that the permissive 4qA allele carries a polymorphism in the region immediately distal to the final repeat of the array (Figure 1). This polymorphism creates a canonical polyadenylation signal in the 3'UTR of *DUX4* enabling

expression of a stable mRNA transcript [34]. The translated protein then binds multiple target genes resulting in widespread changes in gene expression that are ultimately cytotoxic [26].

DUX4 has also been implicated in several cancers involving rearrangements that produce chimeric proteins with altered transcriptional activity [35][36]. For example, a recurrent translocation between Capicua Transcriptional Repressor (*CIC*) and *DUX4* occurs in a proportion of patients with Ewing-like sarcoma. This chromosomal rearrangement produces an in-frame transcript containing the first 20 exons of *CIC* but replacing the terminal exons with the 3' portion of *DUX4*. Translation of the chimeric transcript produces a protein that retains the majority of *CIC*, including the N-terminal DNA binding domains, but replaces the C-terminus with the *DUX4* transactivation domain. As a result, the chimeric *CIC-DUX4* protein acts as an oncogenic transcriptional activator [37]. In B-ALL, translocation of *DUX4* results in a different chimeric protein, but one that again acts as an oncogenic transcriptional activator. In all but one reported case, the 5' coding sequence of *DUX4* is cryptically inserted into an alternate genomic location, resulting in expression of a chimeric transcript which retains sequence containing the N-terminus of *DUX4* but replacing the 3' coding sequence (Figure 2). While multiple potential fusion partners have been identified, including *ERG*, *DUX4* is most commonly inserted into the *IGH* locus [9][12][13][17].

Multiple rearrangements of *IGH* (*IGHr*) have been reported in B-ALL resulting in expression or overexpression of genes with oncogenic potential. The most common of these is a translocation between chromosome 14 and *CRLF2* of the pseudoautosomal region of chromosome X/Y resulting in increased expression of cytokine receptor-like factor 2 [4][38]. In the case of *DUX4*, most analyses report that the rearrangement involves a portion of the D4Z4 array on 4q or the homologous region on 10q, consisting of either a partial copy of *DUX4* or one complete and one partial D4Z4 repeat, being inserted into the *IGH* locus, placing them close to the *IGH* enhancer (E) [12]. As with other *IGHr*, the presence of the enhancer induces expression of the translocated gene [38]. Repeats containing *DUX4* can be inserted in either orientation resulting in expression from the positive or negative strand. In some cases, a more complex rearrangement involving sequences from a third genomic location have been reported (Figure 1C) [12][15]. Alternatively, Hi-C data performed on the NALM6 cell line suggest that a reciprocal translocation can occur in which the telomeric ends of 4/10q are exchanged with 14q [39].

IGH breakpoints are enriched in the 3.5 kb region preceding the *IGHM* constant allele and overlapping the *IGH D-J* junctions but can occur throughout the locus [9]. Breakpoint locations for the *DUX4* gene are harder to define given the repetitive nature of the D4Z4 array, but most commonly occur within the 5' region upstream of *DUX4* and within the 3' coding region of exon1. This results in a *DUX4* transcript which maintains the homeodomains encoded at the 5' end of the transcript fused to sequence, usually from *IGH-JH* or *IGH-DH* regions, but can also be another genomic location [9][12][13][40]. The resulting protein thus maintains its ability to bind *DUX4* targets but possesses a truncated C-terminus with inclusions of some amino acids encoded by the alternate locus. As the genomic breakpoints for this rearrangement are highly variable, the resultant length and amino acid sequence the C-terminal domain of the *DUX4*-fusion varies considerably between patients but consistently retains the DNA binding homeodomains (Figure 2) [9][12][39].




Figure 1. Potential chromosomal rearrangements involving *IGH* and *DUX4*. **(A)** Ideogram of chromosome 4 indicating location of the D4Z4 array and a depiction of the two alleles which vary in the sequence distal to the final repeat (repeat indicated by open triangles). This includes the permissive 4qA allele which can result in FSHD when contracted to fewer than 10 repeats. **(B)** Ideogram of chromosome 10 indicating location of the homologous D4Z4 array with 98% identical sequence. This chromosome is associated with a non-permissive allele which does not result in FSHD. **(C)** Schematic diagram of the repeat 3.3 kb repeat sequence indicating location and exons of the *DUX4* gene. **(D)** Ideogram of chromosome 14 and depiction of the *IGH* locus indicating constant (C_H), joining (J_H), diversity (D_H) and variable (V_H) alleles **(E)** Schematic diagram depicting possible rearrangements as a result of cryptic insertion of *DUX4* from either chromosome 4 or 10 into the *IGH* locus. *DUX4* can be inserted in either orientation, include only a partial or one complete and one partial copy of the repeat array and also be inserted with sequence from a third genomic location.

References

1. Inaba, H.; Greaves, M.; Mullighan, C.G. Acute lymphoblastic leukaemia. *Lancet* 2013, 381, 1943–1955.
2. Tasian, S.K.; Loh, M.L.; Hunger, S.P. Childhood acute lymphoblastic leukemia: Integrating genomics into therapy. *Cancer* 2015, 121, 3577–3590.

3. Hunger, S.P.; Mullighan, C.G. Redefining ALL classification: toward detecting high-risk ALL and implementing precision medicine. *Blood* 2015, 125, 3977–3987.
4. Harvey, R.C.; Mullighan, C.G.; Chen, I.-M.; Wharton, W.; Mikhail, F.M.; Carroll, A.J.; Kang, H.; Liu, W.; Dobbin, K.K.; Smith, M.A.; et al. Rearrangement of CRLF2 is associated with mutation of JAK kinases, alteration of IKZF1, Hispanic/Latino ethnicity, and a poor outcome in pediatric B-progenitor acute lymphoblastic leukemia. *Blood* 2010, 115, 5312–5321.
5. Mullighan, C.G.; Collins-Underwood, J.R.; Phillips, L.A.A.; Loudin, M.G.; Liu, W.; Zhang, J.; Ma, J.; Coustan-Smith, E.; Harvey, R.C.; Willman, C.L.; et al. Rearrangement of CRLF2 in B-progenitor-and Down syndrome-associated acute lymphoblastic leukemia. *Genet.* 2009, 41, 1243–1246. doi:10.1038/ng.469.
6. Yeoh, E.-J.; Ross, M.E.; Shurtleff, S.A.; Williams, W.K.; Patel, D.; Mahfouz, R.; Behm, F.G.; Raimondi, S.C.; Relling, M.V.; Patel, A.; et al. Classification, subtype discovery, and prediction of outcome in pediatric acute lymphoblastic leukemia by gene expression profiling. *Cancer Cell* 2002, 1, 133–143.
7. Mullighan, C.G.; Miller, C.B.; Su, X.; Radtke, I.; Dalton, J.; Song, G.; Zhou, X.; Pui, C.-H.; Shurtleff, S.A.; Downing, J.R. ERG Deletions Define a Novel Subtype of B-Progenitor Acute Lymphoblastic Leukemia. *Blood* 2007, 110, 691–691.
8. Harvey, R.C.; Mullighan, C.G.; Wang, X.; Dobbin, K.K.; Davidson, G.S.; Bedrick, E.J.; Chen, I.-M.; Atlas, S.R.; Kang, H.; Ar, K.; et al. Identification of novel cluster groups in pediatric high-risk B-precursor acute lymphoblastic leukemia with gene expression profiling: correlation with genome-wide DNA copy number alterations, clinical characteristics, and outcome. *Blood* 2010, 116, 4874–4884.
9. Zhang, J.; McCastlain, K.; Yoshihara, H.; Xu, B.; Chang, Y.; Churchman, M.L.; Wu, G.; Li, Y.; Wei, L.; Iacobucci, I.; et al. Deregulation of DUX4 and ERG in acute lymphoblastic leukemia. *Genet.* 2016, 48, 1481–1489.
10. Zaliouva, M.; Potuckova, E.; Hovorkova, L.; Musilova, A.; Winkowska, L.; Fiser, K.; Stuchly, J.; Mejstrikova, E.; Starkova, J.; Zuna, J.; et al. ERG deletions in childhood acute lymphoblastic leukemia with DUX4 rearrangements are mostly polyclonal, prognostically relevant and their detection rate strongly depends on screening method sensitivity. *Haematologica* 2019, 104, 1407–1416.
11. Potuckova, E.; Zuna, J.; Hovorkova, L.; Starkova, J.; Sary, J.; Trka, J.; Zaliouva, M. Intragenic ERG Deletions Do Not Explain the Biology of ERG-Related Acute Lymphoblastic Leukemia. *PLoS One* 2016, 11, e0160385.
12. Yasuda, T.; Tsuzuki, S.; Kawazu, M.; Hayakawa, F.; Kojima, S.; Ueno, T.; Imoto, N.; Kohsaka, S.; Kunita, A.; Doi, K.; et al. Recurrent DUX4 fusions in B cell acute lymphoblastic leukemia of adolescents and young adults. *Genet.* 2016, 48, 569–574.

13. Lilljebjörn, H.; Henningsson, R.; Hyrenius-Wittsten, A.; Olsson, L.; Orsmark-Pietras, C.; von Palffy, S.; Askmyr, M.; Rissler, M.; Schrappe, M.; Cario, G.; et al. Identification of ETV6-RUNX1-like and DUX4-rearranged subtypes in paediatric B-cell precursor acute lymphoblastic leukaemia. *Commun.* 2016, 7, 11790.
14. Zaliouva, M.; Stuchly, J.; Winkowska, L.; Musilova, A.; Fiser, K.; Slamova, M.; Starkova, J.; Vaskova, M.; Hrusak, O.; Sramkova, L.; et al. Genomic landscape of pediatric B-other acute lymphoblastic leukemia in a consecutive European cohort. *Haematologica* 2019, 104, 1396–1406.
15. Marincevic-Zuniga, Y.; Dahlberg, J.; Nilsson, S.; Raine, A.; Nystedt, S.; Lindqvist, C.M.; Berglund, E.C.; Abrahamsson, J.; Cavelier, L.; Forestier, E.; et al. Transcriptome sequencing in pediatric acute lymphoblastic leukemia identifies fusion genes associated with distinct DNA methylation profiles. *Hematol. Oncol.* 2017, 10, 148.
16. Vendramini, E.; Giordan, M.; Giarin, E.; Michielotto, B.; Fazio, G.; Cazzaniga, G.; Biondi, A.; Silvestri, D.; Valsecchi, M.G.; Muckenthaler, M.U.; et al. High expression of miR-125b-2 and SNORD116 noncoding RNA clusters characterize ERG-related B cell precursor acute lymphoblastic leukemia. *Oncotarget* 2017, 8, 42398–42413.
17. Gu, Z.; Churchman, M.L.; Roberts, K.G.; Moore, I.; Zhou, X.; Nakitandwe, J.; Hagiwara, K.; Pelletier, S.; Gingras, S.; Berns, H.; et al. PAX5-driven subtypes of B-progenitor acute lymphoblastic leukemia. *Genet.* 2019, 51, 296–307.
18. Li, J.F.; Dai, Y.T.; Lilljebjörn, H.; Shen, S.H.; Cui, B.W.; Bai, L.; Liu, Y.F.; Qian, M.X.; Kubota, Y.; Kiyoi, H.; et al. Transcriptional landscape of B cell precursor acute lymphoblastic leukemia based on an international study of 1,223 cases. *Natl. Acad. Sci. U. S. A.* 2018, 115, E11711–E11720. doi:10.1073/pnas.1814397115.
19. Liu, Y.-F.; Wang, B.-Y.; Zhang, W.-N.; Huang, J.-Y.; Li, B.-S.; Zhang, M.; Jiang, L.; Li, J.-F.; Wang, M.-J.; Dai, Y.-J.; et al. Genomic Profiling of Adult and Pediatric B-cell Acute Lymphoblastic Leukemia. *EBioMedicine* 2016, 8, 173–183.
20. Schroeder, M.P.; Bastian, L.; Eckert, C.; Gökbüget, N.; James, A.R.; Sanchez, J.O.; Schlee, C.; Isaakidis, K.; Häupl, B.; Baum, K.; et al. Integrated analysis of relapsed B-cell precursor Acute Lymphoblastic Leukemia identifies subtype-specific cytokine and metabolic signatures. *Rep.* 2019, 9, 4188.
21. James, A.R.; Schroeder, M.P.; Neumann, M.; Bastian, L.; Eckert, C.; Gökbüget, N.; Sanchez, J.O.; Schlee, C.; Isaakidis, K.; Schwartz, S.; et al. Long non-coding RNAs defining major subtypes of B cell precursor acute lymphoblastic leukemia. *Hematol. Oncol.* 2019, 12, 8.
22. Winokur, S.T.; Bengtsson, U.; Feddersen, J.; Mathews, K.D.; Weiffenbach, B.; Bailey, H.; Markovich, R.P.; Murray, J.C.; Wasmuth, J.J.; Altherr, M.R. The DNA rearrangement associated with facioscapulohumeral muscular dystrophy involves a heterochromatin-associated repetitive

element: implications for a role of chromatin structure in the pathogenesis of the disease.

Chromosome Res. 1994, 2, 225–234.

23. Cacurri, S.; Piazza, N.; Deidda, G.; Vigneti, E.; Galluzzi, G.; Colantoni, L.; Merico, B.; Ricci, E.; Felicetti, L. Sequence homology between 4qter and 10qter loci facilitates the instability of subtelomeric KpnI repeat units implicated in facioscapulohumeral muscular dystrophy. *J. Hum. Genet.* 1998, 63, 181–190.
24. van Geel, M.; Dickson, M.C.; Beck, A.F.; Bolland, D.J.; Frants, R.R.; van der Maarel, S.M.; de Jong, P.J.; Hewitt, J.E. Genomic analysis of human chromosome 10q and 4q telomeres suggests a common origin. *Genomics* 2002, 79, 210–217.
25. van Deutekom, J.C.; Wijmenga, C.; van Tienhoven, E.A.; Gruter, A.M.; Hewitt, J.E.; Padberg, G.W.; van Ommen, G.J.; Hofker, M.H.; Frants, R.R. FSHD associated DNA rearrangements are due to deletions of integral copies of a 3.2 kb tandemly repeated unit. *Mol. Genet.* 1993, 2, 2037–2042.
26. Geng, L.N.; Yao, Z.; Snider, L.; Fong, A.P.; Cech, J.N.; Young, J.M.; van der Maarel, S.M.; Ruzzo, W.L.; Gentleman, R.C.; Tawil, R.; et al. DUX4 activates germline genes, retroelements, and immune mediators: implications for facioscapulohumeral dystrophy. *Cell* 2012, 22, 38–51.
27. Snider, L.; Geng, L.N.; Lemmers, R.J.L.F.; Kyba, M.; Ware, C.B.; Nelson, A.M.; Tawil, R.; Filippova, G.N.; van der Maarel, S.M.; Tapscott, S.J.; et al. Facioscapulohumeral dystrophy: Incomplete suppression of a retrotransposed gene. *PLoS Genet.* 2010, 6, e1001181.
28. Gabriëls, J.; Beckers, M.C.; Ding, H.; De Vriese, A.; Plaisance, S.; van der Maarel, S.M.; Padberg, G.W.; Frants, R.R.; Hewitt, J.E.; Collen, D.; et al. Nucleotide sequence of the partially deleted D4Z4 locus in a patient with FSHD identifies a putative gene within each 3.3 kb element. *Gene* 1999, 236, 25–32.
29. Choi, S.H.; Gearhart, M.D.; Cui, Z.; Bosnakovski, D.; Kim, M.; Schennum, N.; Kyba, M. DUX4 recruits p300/CBP through its C-terminus and induces global H3K27 acetylation changes. *Nucleic Acids Res.* 2016, 44, 5161–5173.
30. Wijmenga, C.; Hewitt, J.E.; Sandkuijl, L.A.; Clark, L.N.; Wright, T.J.; Dauwerse, H.G.; Gruter, A.M.; Hofker, M.H.; Moerer, P.; Williamson, R. Chromosome 4q DNA rearrangements associated with facioscapulohumeral muscular dystrophy. *Genet.* 1992, 2, 26–30.
31. Pandya, S.; King, W.M.; Tawil, R. Facioscapulohumeral dystrophy. *Ther.* 2008, 88, 105–113.
32. Zeng, W.; de Greef, J.C.; Chen, Y.-Y.; Chien, R.; Kong, X.; Gregson, H.C.; Winokur, S.T.; Pyle, A.; Robertson, K.D.; Schmiesing, J.A.; et al. Specific loss of histone H3 lysine 9 trimethylation and HP1gamma/cohesin binding at D4Z4 repeats is associated with facioscapulohumeral dystrophy (FSHD). *PLoS Genet.* 2009, 5, e1000559.

33. van Overveld, P.G.M.; Lemmers, R.J.F.L.; Sandkuijl, L.A.; Enthoven, L.; Winokur, S.T.; Bakels, F.; Padberg, G.W.; van Ommen, G.-J.B.; Frants, R.R.; van der Maarel, S.M. Hypomethylation of D4Z4 in 4q-linked and non-4q-linked facioscapulohumeral muscular dystrophy. *Genet.* 2003, 35, 315–317.
34. Lemmers, R.J.L.F.; van der Vliet, P.J.; Klooster, R.; Sacconi, S.; Camaño, P.; Dauwerse, J.G.; Snider, L.; Straasheijm, K.R.; van Ommen, G.J.; Padberg, G.W.; et al. A unifying genetic model for facioscapulohumeral muscular dystrophy. *Science* 2010, 329, 1650–1653.
35. Dmitriev, P.; Kairov, U.; Robert, T.; Barat, A.; Lazar, V.; Carnac, G.; Laoudj-Chenivresse, D.; Vassetzky, Y.S. Cancer-related genes in the transcription signature of facioscapulohumeral dystrophy myoblasts and myotubes. *Cell. Mol. Med.* 2014, 18, 208–217.
36. Dib, C.; Zakharova, V.; Popova, E.; Kiseleva, E.; Chernyak, B.; Lipinski, M.; Vassetzky, Y.S. DUX4 Pathological Expression: Causes and Consequences in Cancer. *Trends Cancer Res.* 2019, 5, 268–271.
37. Kawamura-Saito, M.; Yamazaki, Y.; Kaneko, K.; Kawaguchi, N.; Kanda, H.; Mukai, H.; Gotoh, T.; Motoi, T.; Fukayama, M.; Aburatani, H.; et al. Fusion between CIC and DUX4 up-regulates PEA3 family genes in Ewing-like sarcomas with t(4;19)(q35;q13) translocation. *Mol. Genet.* 2006, 15, 2125–2137.
38. Dyer, M.J.S.; Akasaka, T.; Capasso, M.; Dusanj, P.; Lee, Y.F.; Karran, E.L.; Nagel, I.; Vater, I.; Cario, G.; Siebert, R. Immunoglobulin heavy chain locus chromosomal translocations in B-cell precursor acute lymphoblastic leukemia: Rare clinical curios or potent genetic drivers? *Blood* 2010, 115, 1490–1499. doi:1182/blood-2009-09-235986.
39. Tian, L.; Shao, Y.; Nance, S.; Dang, J.; Xu, B.; Ma, X.; Li, Y.; Ju, B.; Dong, L.; Newman, S.; et al. Long-read sequencing unveils IGH-DUX4 translocation into the silenced IGH allele in B-cell acute lymphoblastic leukemia. *Commun.* 2019, 10, 2789.
40. Zur Stadt, U.; Alawi, M.; Adao, M.; Indenbirken, D.; Escherich, G.; Horstmann, M.A. Characterization of novel, recurrent genomic rearrangements as sensitive MRD targets in childhood B-cell precursor ALL. *Blood Cancer J.* 2019, 9, 96.

Retrieved from <https://encyclopedia.pub/entry/history/show/6193>