

Trace Amine-Associated Receptor 1

Subjects: **Pathology**

Contributor: Udo Jeschke

Trace amine-associated receptor 1 (TAAR1) is a G α s- protein coupled receptor that plays an important role in the regulation of the immune system and neurotransmission in the CNS. In ovarian cancer cell lines, stimulation of TAAR1 via 3-iodothyronamine (T1AM) reduces cell viability and induces cell death and DNA damage.

TAAR1

ovarian cancer

prognostic factor

immunohistochemistry

1. Introduction

With more than 7000 new cases in Germany in 2016, ovarian cancer (OC) is representing 3.1% of cancer cases and 5.2% of cancer deaths in females ^[1]. Due to its high potential for cancer dissemination into the peritoneum, frequent late symptoms and detection, heterogeneity, and acquired and intrinsic chemoresistance ^{[2][3]}, epithelial ovarian cancer shows relative 5 years survival rates of less than 50% ^[4] with only slight improvement and a wide range of 5 years survival rates in geographic distribution ^[5]. OC presents in a histologically diverse way, with epithelial type being the most frequent, accounting for more than 90% of all primary ovarian tumors ^[6]. Other histological subtypes include sex cord-stromal (5–6%), germ cell (2–3%), and rare miscellaneous entities ^[6]. This article focuses on epithelial tumors, which can be subdivided into high-grade serous carcinomas (70–80% of cases), low-grade serous carcinomas, endometrioid (<5%), clear cell (3%), and mucinous cancers (<3%) ^[7].

Therapeutic management of OC lacks effective screening methods. Clinical prognosticators such as disease stage at diagnosis, postsurgical residual disease, histological subtype, grading, mutation in the breast cancer gene (BRCA), general state of health, age, guideline-based therapy, and volume of ascites are of value ^{[1][3]}. Platinum resistance after chemotherapy can worsen the prognosis of ovarian cancer patients ^{[8][9]}. Attempts of establishing new prognostic factors have been published in multiple studies ^{[10][11][12]}.

Ovarian cancer is regulated by thyroid hormones and its derivatives ^{[13][14]}. Thyroid hormones also have a pro-angiogenic role in various cancer types, making its receptors potential targets for therapeutic treatment ^[15]. Similar to ovarian cancer, hypothyroidism occurs predominantly in aging women ^{[16][17]}. Hyperthyroidism is prevalent in women from the 3rd to 5th decade of life with an 8:1 female:male ratio ^[18].

Thyronamines, such as 3-iodothyronamine (T₁AM), derive from thyroid hormones ^[19] via degradation by the enzyme ornithine decarboxylase (ODC) and intestinal deiodinases ^[20]. In comparison with the mostly epigenetically acting classical thyroid hormones, decarboxylated thyroid hormones can function rapidly (e.g., via fast change of heart rate and body temperature) ^[19]. T₁AM is described to inhibit cell growth and viability, induce cell death, and

lead to DNA damage in ovarian cancer cells [13]. Furthermore, T₁AM and its metabolites, T₀AM and thyroacetic acid (TA₁), can influence pancreatic islets, brain, heart, and other tissues through the G_{αs}-protein coupled receptor trace amine-associated receptor 1 (TAAR1) [20]. Other agonists of TAAR1 are trace amines, which are closely associated metabolically with the dopamine, serotonin, and noradrenaline systems [21]. The degradation and synthesis of both trace amines and thyronamines proceed via enzymes working through decarboxylation. L-thyroxin (T₄) is degraded to thyronamines (T₁AM) via ornithine decarboxylase (ODC) [20] while trace amines are synthesized via L-amino acid decarboxylase [21]. TAAR1 is widely expressed including placenta, brain, spinal cord, immune cells such as leukocytes, macrophages and dendritic cells, breast cancer tissue, D-cells in stomach, and pancreatic β cells [22][23][24][25][26]. Activation of TAAR1 leads to a G_{αs}-protein mediated increase in intracellular cAMP levels [27][28].

2. Differences of TAAR1 Expression in Histological Subtypes of Ovarian Cancer

TAAR1 was detected in membrane as well as in cytoplasm. Membrane TAAR1 staining could be gained in $n = 134$ cases and presented in endometrioid tumors ($n = 16$) and serous carcinoma ($n = 98$), with a median IRS of 3, in clear cell ($n = 9$) and mucinous tumor ($n = 11$) with a median IRS of 1 ($p = 0.003$), as shown in **Figure 1**. In $n = 22$ cases, either staining was not successful, or no ovarian cancer tissue was hit.




Figure 1. Correlation of membrane TAAR1 expression with histological subtype ($p = 0.003$) (a) boxplot of membrane TAAR1 expression and histological subtype (b) serous carcinoma ($n = 98$) with a membrane TAAR1 IRS of 4, magnification $\times 10$ and $\times 25$ in the inset (c) clear cell carcinoma ($n = 9$) with a membrane TAAR1 IRS of 1, magnification $\times 10$ and $\times 25$ in the inset (d) endometrioid carcinoma ($n = 16$) with a membrane TAAR1 IRS of 4, magnification $\times 10$ and $\times 25$ in the inset (e) mucinous carcinoma ($n = 11$) with a membrane TAAR1 IRS of 0, magnification $\times 10$ and $\times 25$ in the inset; $* p < 0.05$ was considered statistically significant.

Likewise, correlation of cytoplasmic TAAR1 expression and histological subtype differs significantly. A total of $n = 128$ cases could be observed. While endometrioid tumor ($n = 19$) showed a median IRS of 4, clear cell carcinoma ($n = 10$) showed a median IRS of 3.5, serous carcinoma a median IRS of 3 ($n = 89$), and mucinous carcinoma ($n = 10$) a median IRS of 2.5 ($p = 0.009$), as shown in **Figure 2**. On $n = 28$ slides, staining was not successful, or no ovarian cancer tissue was gained.




Figure 2. Correlation of cytoplasmic TAAR1 expression with histological subtype ($p = 0.009$) (a) boxplot of cytoplasmic TAAR1 expression and histological subtype (b) serous carcinoma ($n = 89$) with a cytoplasmic TAAR1 IRS of 3, magnification $\times 10$ and $\times 25$ in the inset (c) clear cell carcinoma ($n = 10$) with a cytoplasmic TAAR1 IRS of 3, magnification $\times 10$ and $\times 25$ in the inset (d) endometrioid carcinoma ($n = 19$) with a cytoplasmic TAAR1 IRS of 4, magnification $\times 10$ and $\times 25$ in the inset (e) mucinous carcinoma ($n = 10$) with a cytoplasmic TAAR1 IRS of 2, magnification $\times 10$ and $\times 25$ in the inset; $* p < 0.05$ was considered statistically significant.

3. TAAR1 Expression in High-Grade and Low-Grade Serous Ovarian Cancer

$n = 110$ tissue sections from patients diagnosed with serous ovarian carcinoma were stained (high-grade, $n = 80$; low-grade, $n = 24$; unknown, $n = 6$). Due to missed hits of ovarian cancer tissue and/or failed staining in $n = 17$ cases, $n = 93$ slides were examined. Correlation of membrane TAAR1 expression, with grading of serous carcinoma, showed that TAAR1 is expressed significantly higher in low-grade serous carcinoma (median IRS of 4; $n = 22$) compared to high-grade serous carcinoma (median IRS of 3; $n = 71$) ($p = 0.028$) (Figure 3).




Figure 3. Correlation of membrane TAAR1 expression with grading in serous carcinoma ($p = 0.028$) (a) boxplot of membrane TAAR1 expression and grading in serous carcinoma (b) low grade serous carcinoma ($n = 22$) with a membrane TAAR1 IRS of 4, magnification $\times 10$ and $\times 25$ in the inset (c) high-grade serous carcinoma ($n = 71$) with a membrane TAAR1 IRS of 2, magnification $\times 10$ and $\times 25$ in the inset; * $p < 0.05$ was considered statistically significant.

References

1. Leitlinienprogramm Onkologie (Deutsche Krebsgesellschaft, Deutsche Krebshilfe, AWMF): S3-Leitlinie Diagnostik, Therapie und Nachsorge Maligner Ovarialtumoren, Langversion 4.0, 2020, AWMF-Registernummer: 032/035OL. Available online: <https://www.leitlinienprogramm-onkologie.de/leitlinien/ovarialkarzinom/> (accessed on 1 May 2021).
2. Thibault, B.; Castells, M.; Delord, J.P.; Couderc, B. Ovarian cancer microenvironment: Implications for cancer dissemination and chemoresistance acquisition. *Cancer Metastasis Rev.* 2014, 33, 17–39.
3. Davidson, B.; Trope, C.G. Ovarian cancer: Diagnostic, biological and prognostic aspects. *Womens Health* 2014, 10, 519–533.

4. American Cancer Society. Survival Rates for Ovarian Cancer. Available online: <https://www.cancer.org/cancer/ovarian-cancer/detection-diagnosis-staging/survival-rates.html> (accessed on 3 August 2021).
5. Allemani, C.; Weir, H.K.; Carreira, H.; Harewood, R.; Spika, D.; Wang, X.S.; Bannon, F.; Ahn, J.V.; Johnson, C.J.; Bonaventure, A.; et al. Global surveillance of cancer survival 1995-2009: Analysis of individual data for 25,676,887 patients from 279 population-based registries in 67 countries (CONCORD-2). *Lancet* 2015, 385, 977–1010.
6. Reid, B.M.; Permeth, J.B.; Sellers, T.A. Epidemiology of ovarian cancer: A review. *Cancer Biol. Med.* 2017, 14, 9–32.
7. Sundar, S.; Neal, R.D.; Kehoe, S. Diagnosis of ovarian cancer. *BMJ Br. Med. J.* 2015, 351, h4443.
8. Wilson, M.K.; Pujade-Lauraine, E.; Aoki, D.; Mirza, M.R.; Lorusso, D.; Oza, A.M.; du Bois, A.; Vergote, I.; Reuss, A.; Bacon, M.; et al. Fifth Ovarian Cancer Consensus Conference of the Gynecologic Cancer InterGroup: Recurrent disease. *Ann. Oncol.* 2017, 28, 727–732.
9. Trillsch, F.; Mahner, S.; Czogalla, B.; Rottmann, M.; Chekerov, R.; Braicu, E.I.; Oskay-Öczelik, G.; Wimberger, P.; Richter, R.; Sehouli, J. Primary platinum resistance and its prognostic impact in patients with recurrent ovarian cancer: An analysis of three prospective trials from the NOGGO study group. *J. Gynecol. Oncol.* 2021, 32, e37.
10. Salminen, L.; Braicu, E.I.; Lääperi, M.; Jylhä, A.; Oksa, S.; Hietanen, S.; Sehouli, J.; Kulbe, H.; Bois, A.D.; Mahner, S.; et al. A Novel Two-Lipid Signature Is a Strong and Independent Prognostic Factor in Ovarian Cancer. *Cancers* 2021, 13, 1764.
11. Fraungruber, P.; Kaltofen, T.; Heublein, S.; Kuhn, C.; Mayr, D.; Burges, A.; Mahner, S.; Rathert, P.; Jeschke, U.; Trillsch, F. G Protein-Coupled Estrogen Receptor Correlates With Dkk2 Expression and Has Prognostic Impact in Ovarian Cancer Patients. *Front. Endocrinol.* 2021, 12, 564002.
12. Czogalla, B.; Partenheimer, A.; Jeschke, U.; von Schönfeldt, V.; Mayr, D.; Mahner, S.; Burges, A.; Simoni, M.; Melli, B.; Benevelli, R.; et al. β -arrestin 2 Is a Prognostic Factor for Survival of Ovarian Cancer Patients Upregulating Cell Proliferation. *Front. Endocrinol.* 2020, 11, 554733.
13. Shinderman-Maman, E.; Cohen, K.; Moskovich, D.; Hercbergs, A.; Werner, H.; Davis, P.J.; Ellis, M.; Ashur-Fabian, O. Thyroid hormones derivatives reduce proliferation and induce cell death and DNA damage in ovarian cancer. *Sci. Rep.* 2017, 7, 16475.
14. Shinderman-Maman, E.; Weingarten, C.; Moskovich, D.; Werner, H.; Hercbergs, A.; Davis, P.J.; Ellis, M.; Ashur-Fabian, O. Molecular insights into the transcriptional regulatory role of thyroid hormones in ovarian cancer. *Mol. Carcinog.* 2018, 57, 97–105.
15. Mousa, S.A.; Lin, H.Y.; Tang, H.Y.; Hercbergs, A.; Luidens, M.K.; Davis, P.J. Modulation of angiogenesis by thyroid hormone and hormone analogues: Implications for cancer management. *Angiogenesis* 2014, 17, 463–469.

16. Giri, S.K.; Nayak, B. Management of Ovarian Cancer in Elderly. *Rev. Recent Clin. Trials* 2015, 10, 270–275.
17. Hollowell, J.G.; Staehling, N.W.; Flanders, W.D.; Hannon, W.H.; Gunter, E.W.; Spencer, C.A.; Braverman, L.E. Serum TSH, T4, and Thyroid Antibodies in the United States Population (1988 to 1994): National Health and Nutrition Examination Survey (NHANES III). *J. Clin. Endocrinol. Metab.* 2002, 87, 489–499.
18. Taylor, P.N.; Albrecht, D.; Scholz, A.; Gutierrez-Buey, G.; Lazarus, J.H.; Dayan, C.M.; Okosieme, O.E. Global epidemiology of hyperthyroidism and hypothyroidism. *Nat. Rev. Endocrinol.* 2018, 14, 301–316.
19. Scanlan, T.S.; Suchland, K.L.; Hart, M.E.; Chiellini, G.; Huang, Y.; Kruzich, P.J.; Frascarelli, S.; Crossley, D.A.; Bunzow, J.R.; Ronca-Testoni, S.; et al. 3-Iodothyronamine is an endogenous and rapid-acting derivative of thyroid hormone. *Nat. Med.* 2004, 10, 638–642.
20. Hoefig, C.S.; Zucchi, R.; Kohrle, J. Thyronamines and Derivatives: Physiological Relevance, Pharmacological Actions, and Future Research Directions. *Thyroid* 2016, 26, 1656–1673.
21. Burchett, S.A.; Hicks, T.P. The mysterious trace amines: Protean neuromodulators of synaptic transmission in mammalian brain. *Prog. Neurobiol.* 2006, 79, 223–246.
22. Gozal, E.A.; O'Neill, B.E.; Sawchuk, M.A.; Zhu, H.; Halder, M.; Chou, C.C.; Hochman, S. Anatomical and functional evidence for trace amines as unique modulators of locomotor function in the mammalian spinal cord. *Front. Neural. Circuits* 2014, 8, 134.
23. Babusyte, A.; Kotthoff, M.; Fiedler, J.; Krautwurst, D. Biogenic amines activate blood leukocytes via trace amine-associated receptors TAAR1 and TAAR2. *J. Leukoc. Biol.* 2013, 93, 387–394.
24. Vattai, A.; Akyol, E.; Kuhn, C.; Hofmann, S.; Heidegger, H.; von Koch, F.; Hermelink, K.; Wuerstlein, R.; Harbeck, N.; Mayr, D.; et al. Increased trace amine-associated receptor 1 (TAAR1) expression is associated with a positive survival rate in patients with breast cancer. *J. Cancer Res. Clin. Oncol.* 2017, 143, 1637–1647.
25. Adriaenssens, A.; Lam, B.Y.; Billing, L.; Skeffington, K.; Sewing, S.; Reimann, F.; Gribble, F. A Transcriptome-Led Exploration of Molecular Mechanisms Regulating Somatostatin-Producing D-Cells in the Gastric Epithelium. *Endocrinology* 2015, 156, 3924–3936.
26. Regard, J.B.; Kataoka, H.; Cano, D.A.; Camerer, E.; Yin, L.; Zheng, Y.W.; Scanlan, T.S.; Hebrok, M.; Coughlin, S.R. Probing cell type-specific functions of Gi in vivo identifies GPCR regulators of insulin secretion. *J. Clin. Investig.* 2007, 117, 4034–4043.
27. Borowsky, B.; Adham, N.; Jones, K.A.; Raddatz, R.; Artymyshyn, R.; Ogozalek, K.L.; Durkin, M.M.; Lakhani, P.P.; Bonini, J.A.; Pathirana, S.; et al. Trace amines: Identification of a family of mammalian G protein-coupled receptors. *Proc. Natl. Acad. Sci. USA* 2001, 98, 8966–8971.

28. Bunzow, J.R.; Sonders, M.S.; Arttamangkul, S.; Harrison, L.M.; Zhang, G.; Quigley, D.I.; Darland, T.; Suchland, K.L.; Pasumamula, S.; Kennedy, J.L.; et al. Amphetamine, 3,4-methylenedioxymethamphetamine, lysergic acid diethylamide, and metabolites of the catecholamine neurotransmitters are agonists of a rat trace amine receptor. *Mol. Pharmacol.* 2001, 60, 1181–1188.
-

Retrieved from <https://encyclopedia.pub/entry/history/show/31466>