

# Tea Tree Oil Induced Resistance

Subjects: **Plant Sciences**

Contributor: Ronaldo J. D. Dalio , Heros J. Maximo , Rafaela Roma-Almeida , Janaína N. Barretta , Eric M. José , Agnelo J. Vitti , Daphna Blachinsky , Moshe Reuveni , Sérgio F. Pascholati

The essential tea tree oil (TTO) derived from *Melaleuca alternifolia* plant is widely used as a biopesticide to protect crops from several plant-pathogens. TTO was approved by the European Union (EU) and included in the positive list of the EU, in Annex I of Directive 91/414/EEC for registration of Plant Protection Products. TTO is classified as a low risk substance in Europe. This oil is in large use in cosmetics and in medicine.

[Melaleuca alternifolia](#)

[Tea Tree Oil](#)

[Plant Protection](#)

[Plant Microbe Interactions](#)

[Fusarium oxysporum](#)

[Xanthomonas campestris](#)

[Systemic Acquired Resistance \(SAR\)](#)

[Induced Systemic Resistance \(ISR\)](#)

[Priming](#)

[Agriculture](#)

[Agriculture](#)

## 1. Introduction

Plants are exposed to various pathogens in their environment and have developed immune systems with multiple defense layers to prevent infections [1][2]. However, often pathogens overcome these resistance barriers, infect plants and cause disease. Pathogens that cause diseases on economically important crop plants incur huge losses to the agriculture industry and important ecosystems [3][4].

Chemical pesticides have often been used to control diseases in plants, but this conduct is associated to negative environmental impacts, potential human exposure to pesticides, and deposition of residues on the plant organs. However, the effectiveness of synthetic pesticides has been reduced by the frequent development of resistance by the pathogens. Hence there is a great demand for safer, alternative and effective agents [5].

Bananas, the world's most important fruit in terms of production, volume and trade [6] and among the world's top 10 staple foods, is seriously threatened by Fusarium wilt (FW) caused by the fungus *Fusarium oxysporum* f. sp. *cubense* (Foc). The fungus penetrates the plant through the root system and may causes plant death. After the death of the mother plant, the fungus continues infecting the daughter plant and remains active on the clump for years [7][8][9].

The tomato production is yearly affected by disease caused by *Xanthomonas* spp. [10]. The disease is characterized by necrotic lesions on the leaves, stems, petals, flowers, and fruit [11]. During the initial stages of symptom development, circular water-soaked lesions appear, which later dry and turn dark brown to black with a wet to greasy appearance [12].

Both microorganisms are important because of the losses they can cause in crop production. No chemicals are known to control Fusarium wilt [8] and they are very limited for bacterial diseases.

An alternative procedure to protect plants against disease is to activate their inherent defense mechanisms by specific biotic or abiotic elicitors. Plants can be induced to switch on defense reactions to a broad range of pathogens as a result of prior exposure to pathogens or to various chemicals or physical stresses. Induced resistance is expressed locally, at the site of the infection or systemically, at sites located far from the initial infection. Upon recognition of the initial stimulus by the plant, a signal transduction pathway is set in motion, that includes intra and intercellular signals, and results in the activation of defense mechanisms, mostly by the expression of new genes [13][14].

During the interactions with microorganisms and molecular patterns from the environment, plants produce several hormones that act as signals that trigger the production of antimicrobial compounds and activate defense in general [15][16]. Upon recognition, plants produce salicylic acid, jasmonate and/or ethylene as major defense signals [17]. Salicylic acid (SA), for example, plays an important role in defense in biotrophic interactions. SA biosynthesis is induced in the infection site where it contributes to reactive oxygen species (ROS) production and cell death in a positive feedback [18][19][20][21]. Jasmonate (JA) and ethylene (ET), on the other hand, are generally associated to the regulation of defenses against herbivores and necrotrophic pathogens, although exceptions can be seen [22][23]. The activation of both jasmonic acid and ethylene pathways is frequently associated with induced systemic resistance (ISR) [20]. Thus, the genes for the three defense related pathways can be considered as molecular markers to study induced resistance in plants.

Frequently, SAR or ISR are accompanied by the priming effect. Priming is a phenomenon in which a plant activates their defense system by recognizing a molecular pattern from the environment in the absence of a pathogen and no development of disease, the defense system is deactivated and the plant turns back to homeostasis [14]. However, if the plant is subsequently challenged by a pathogen or abiotic stress, through epigenetics, it can activate their defense system more rapidly and robustly [24][25][26]. Inducing priming is preferable rather than the classic resistance induction where the plants may allocate too much energy on defense and that may affect growth and production. [24][25].

The global search for plant-protection solutions, that are both environmentally safe and efficacious, is an important aspect of sustainable agriculture. This is driven by the need to supply food to the ever-growing world population, and the call for chemical load reduction.

## 2. The Essential Tea Tree Oil (TTO)

The essential tea tree oil (TTO) extracted from *Melaleuca alternifolia* plant, contains many components, mostly terpenes and their alcohols, was found to be an effective antiseptic, bactericide [27][28][29] and more recently as an effective fungicide [30][31][32]. Based on TTO, as an active ingredient, a natural fungicide Timorex Gold® (22.3 EC W/V), was prepared in order to enable the use of TTO on plant tissue [33]. In numerous crops, including bananas

and fruit trees, this product was found effective against a broad range of plant-pathogenic fungi [32][33][34][35]. The high effectiveness of TTO raised questions regarding its ability to perform as a resistance inducer. TTO is highly applied in banana and tomato plants worldwide, however very little is known about its mode of action in these hosts physiology.

Thus, this study was undertaken to examine the induction of systemic resistance in field-grown banana plants to *Fusarium* wilt and greenhouse-grown tomato plants to bacterial disease. Symptoms, peroxidase and  $\beta$ -1,3-glucanase activities, and the expression of different marker genes related to SAR and ISR were evaluated in response to early treatment with the TTO.

## 3. Conclusions

TTO is able to prime tomato plants to have a strong defense reaction to subsequent challenges, such as mechanical wounding. These results also demonstrate that TTO provides protection to the plant independently of the fungicide effect.

Data in this entry paper also show that TTO can be consider as an efficient resistance inducer, since it has enhanced the expression of marker genes in non-symptomatic banana plants for both SAR and ISR at the three main defense related pathways—salicylic acid, jasmonic acid and ethylene. To the best of our knowledge, this is the first report on the effect of TTO as a resistance inducer, which opens new possibilities for this product to be used in strategies to control pathogens by decreasing the number of traditional defensive applications and inhibiting the spread of both *Fusarium oxysporum* in banana and *Xanthomonas campestris* in tomato as well as other type of pathogens in different crops.

## References

1. Janeway, C.A.; Medzhitov, R. Innate immune recognition. *Annu. Rev. Immunol.* 2002, 20, 197–216.
2. Jones, J.D.G.; Dangl, J.L. The plant immune system. *Nature* 2006, 444, 323–329.
3. Agrios, G. *Plant Pathology*, 5th ed.; Academic Press: Cambridge, MA, USA, 2004. [Google Scholar]
4. Dalio, R.J.D.; Magalhães, D.M.; Rodrigues, C.M.; Arena, G.D.; Oliveira, T.S.; Souza-Neto, R.R.; Picchi, S.C.; Martins, P.M.M.; Santos, P.J.C.; Maximo, H.J.; et al. PAMPs, PRRs, effectors and R-genes associated with citrus–pathogen interactions. *Ann. Bot.* 2017, 119, 749–774.
5. Liu, C.; Zhao, C.; Pan, H.-H.; Kang, J.; Yu, X.-T.; Wang, H.-Q.; Li, B.-M.; Xie, Y.-Z.; Chen, R.-Y. Chemical constituents from *Inonotus obliquus* and their biological activities. *J. Nat. Prod.* 2013, 77, 35–41.

6. Food and Agriculture Organization of the United Nations Rome. Available online: <http://www.fao.org/3/ca5625en/ca5625en.pdf> (accessed on 2 September 2020).
7. O'Donnell, K.; Gueidan, C.; Sink, S.; Johnston, P.R.; Crous, P.W.; Glenn, A.E.; Riley, R.; Zitomer, N.C.; Colyer, P.; Waalwijk, C.; et al. A two-locus DNA sequence database for typing plant and human pathogens within the *Fusarium oxysporum* species complex. *Fungal Genet. Biol.* 2009, 46, 936–948.
8. Ploetz, R.C. Management of *Fusarium* wilt of banana: A review with special reference to tropical race 4. *Crop Prot.* 2015, 73, 7–15.
9. Ploetz, R.C.; Kema, G.H.J.; Ma, L.-J. Impact of diseases on export and smallholder production of banana. *Annu. Rev. Phytopathol.* 2015, 53, 269–288.
10. Obradovic, A.; Jones, J.B.; Momol, M.T.; Balogh, B.; Olson, S.M. Management of tomato bacterial spot in the field by foliar applications of bacteriophages and SAR inducers. *Plant Dis.* 2004, 88, 736–740.
11. Jones, J.B.; Jones, J.P.; Stall, R.E.; Zitter, T.A. *Compendium of Tomato Diseases*; APS Press: St. Paul, MN, USA, 1991.
12. Vallad, G.E.; Subbarao, K.V. Colonization of resistant and susceptible lettuce cultivars by a green fluorescent protein-tagged isolate of *Verticillium dahliae*. *Phytopathology* 2008, 98, 871–885.
13. Metraux, J.-P.; Nawrath, C.; Genoud, T. Systemic acquired resistance. *Euphytica* 2002, 124, 237–243.
14. Choudhary, D.K.; Prakash, A.; Johri, B.N. Induced systemic resistance (ISR) in plants: Mechanism of action. *Indian J. Microbiol.* 2007, 47, 289–297.
15. Verma, V.; Ravindran, P.; Kumar, P.P. Plant hormone-mediated regulation of stress responses. *BMC Plant Biol.* 2016, 16, 86.
16. Verhage, A.; van Wees, S.C.; Pieterse, C.M. Plant immunity: It's the hormones talking, but what do they say? *Plant Physiol.* 2010, 154, 536–540.
17. Robert-Seilantianz, A.; Grant, M.; Jones, J.D. Hormone crosstalk in plant disease and defense: More than just jasmonate-salicylate antagonism. *Annu. Rev. Phytopathol.* 2011, 49, 317–343.
18. Fu, Z.Q.; Dong, X. Systemic acquired resistance: Turning local infection into global defense. *Annu. Rev. Plant Biol.* 2013, 64, 839–863.
19. Park, S.-W.; Kaimoyo, E.; Kumar, D.; Mosher, S.; Klessig, D.F. Methyl salicylate is a critical mobile signal for plant systemic acquired resistance. *Science* 2007, 318, 113–116.
20. Pascholati, S.F.; Blumer, S.; Rezende, D.C.; Brand, S.C. Systemic acquired resistance (SAR) x induced systemic resistance (ISR). In NEFIT. *Indução de Resistência—Novos Conceitos e*

Aplicações; UFLA: Lavras, Brazil, 2010; pp. 29–40.

21. Shah, J.; Zeier, J. Long-distance communication and signal amplification in systemic acquired resistance. *Front. Plant Sci.* 2013, 4, 30.

22. Zhu, Z. Molecular basis for jasmonate and ethylene signal interactions in *Arabidopsis*. *J. Exp. Bot.* 2014, 65, 5743–5748.

23. Bari, R.; Jones, J.D. Role of plant hormones in plant defence responses. *Plant Mol. Biol.* 2009, 69, 473–488.

24. Dalio, R.J.D.; Fleischmann, F.; Humez, M.; Osswald, W. Phosphite Protects *Fagus sylvatica* seedlings towards *Phytophthora plurivora* via local toxicity, priming and facilitation of pathogen recognition. *PLoS ONE* 2014, 9, e87860.

25. Havko, N.E.; Major, I.T.; Jewell, J.B.; Attaran, E.; Browse, J.; Howe, G.A. Control of carbon assimilation and partitioning by jasmonate: An accounting of growth–defense tradeoffs. *Plants* 2016, 5, 7.

26. Espinas, N.A.; Saze, H.; Saijo, Y. Epigenetic control of defense signaling and priming in plants. *Front. Plant Sci.* 2016, 7, 1201.

27. Carson, C.; Riley, T.V. Antimicrobial activity of the essential oil of *Melaleuca alternifolia*. *Lett. Appl. Microbiol.* 1993, 16, 49–55.

28. Carson, C.; Hammer, K.A.; Riley, T.V. *Melaleuca alternifolia* (tea tree) oil: A review of antimicrobial and other medicinal properties. *Clin. Microbiol. Rev.* 2006, 19, 50–62.

29. Cox, S.D.; Mann, C.M.; Markham, J.L.; Gustafson, J.E.; Warmington, J.R.; Wyllie, S.G. Determining the antimicrobial actions of tea tree oil. *Molecules* 2001, 6, 87–91.

30. Shao, X.; Cheng, S.; Wang, H.; Yu, D.; Mungai, C. The possible mechanism of antifungal action of tea tree oil on *Botrytis cinerea*. *J. Appl. Microbiol.* 2013, 114, 1642–1649.

31. Shao, X.; Wang, H.; Xu, F.; Cheng, S. Effects and possible mechanisms of tea tree oil vapor treatment on the main disease in postharvest strawberry fruit. *Postharvest Biol. Technol.* 2013, 77, 94–101.

32. Reuveni, M.; Barbier, M.; Viti, A.J. Essential tea tree oil as a tool to combat black Sigatoka in banana. *Outlooks Pest Manag.* 2020.

33. Reuveni, M.; Neifeld, D.; Dayan, D.; Kotzer, Y. BM-608—A novel organic product based on essential tea tree oil for the control of fungal diseases in tomato. *Acta Hortic.* 2009, 808, 129–132.

34. Martillo, E.E.; Reuveni, M. A new potent bio-fungicide for the control of Banana Black Sigatoka. *Phytopathology* 2009, 99, S80.

35. Vardi, Y.; Reuveni, M. Antifungal activity of a new broad spectrum biocide in the controlling of plant diseases. *Phytopathology* 2009, 99, 5134.

---

Retrieved from <https://encyclopedia.pub/entry/history/show/6707>