Heterologous Synthesis of Xanthophyll in Model Microorganisms

Subjects: Biotechnology & Applied Microbiology

Contributor: Nan Wang, Huakang Peng, Caifeng Yang, Wenfang Guo, Mengqi Wang, Gangqiang Li, Dehu Liu

Xanthophyll is an oxidated version of carotenoid. It presents significant value to the pharmaceutical, food, and cosmetic industries due to its specific antioxidant activity and variety of colors. Chemical processing and conventional extraction from natural organisms are still the main sources of xanthophyll.

Keywords: xanthophyll ; biosynthetic pathway ; model microorganisms

1. Introduction

Carotenoids are a class of secondary metabolites with a tetraterpene structure that are widely found in plants, algae, yeast, archaea, and some bacteria species [1][2][3][4][5]. Carotenoids are classified into two categories depending on whether they contain oxygen (**Figure 1**): carotene and xanthophyll $^{[6][2]}$. The structure and function of xanthophyll present abundant diversity attributed to the oxygen group, which exists in various forms, such as hydroxyl-, keto-, expo-xanthophyll, and expo-xanthophyll derivatives $^{[8]}$. So far, more than 600 xanthophylls (including isomers) have been found or identified among approximately 1000 carotenoids (<u>https://coconut.naturalproducts.net/</u> (accessed on 3 March 2023).



Figure 1. Xanthophyll biosynthesis pathway. Abbreviations: MVA, mevalonate; MEP, 2-C-methyl-D -erythritol-4phosphate; IPP, isoprene diphosphate; DMAPP, dimethylallyl diphosphate; GPP, geranyl diphosphate; FPP, farnesyl diphosphate; GGPP, geranylgeranyl diphosphate; IDI, IPP isomerase; FPPS, farnesyl diphosphate synthase; CrtE, GGPP synthase; CrtB, phytoene synthase; CrtI, phytoene desaturase; ε -LCY, lycopene ε -cyclase; β -LCY, lycopene β -cyclase; CYP97A, α -carotene β -ring hydroxylase; CYP97C, α -carotene ε -ring hydroxylase; CrtZ, β -carotene hydrolase; CrtW, β carotene ketolase (from bacteria); BKT, β -carotene ketolase (from algae); ZEP, zeaxanthin epoxidase; CCS, capsanthin/capsorubin synthase; NSY, neoxanthin synthase. The dotted line indicates that the enzyme catalyzing the reaction has not been identified.

Xanthophyll has been used commercially in a wide variety of industries, such as food, feed, pharmaceuticals, and cosmetics ^{[8][9][10]}. Currently, chemical processing and conventional extraction from natural organisms are the mainly source of xanthophyll. Despite the low cost and high yield of chemical synthesis methods, the formation of unnatural chiral isomer reduces the bioavailability of the product. In addition, petrochemical-refined xanthophyll is not allowed to be used in food and pharmaceutical products. The yield of xanthophyll extracted from plants or algae is too low to meet industrial needs ^{[11][12]}. In recent decades, metabolic engineering based on model microbes for the production of nature bioactivity compounds has experienced extensive development ^{[13][14][15]}, which brings a promising potential for the industrial synthesis of natural xanthophyll.

The elucidation of the natural synthetic pathways and key enzymes of several important xanthophylls has laid the foundation for xanthophyll synthesis by the metabolic engineering of model microorganisms ^{[16][17][18][19][20]}. The natural

synthetic pathway of xanthophyll is a cascade reaction that begins with the oxidative modification of the α - or β -carotenoid terminal group (**Figure 1**). The hydroxylation of α - and β -carotene terminal groups enables the production of lutein and zeaxanthin, respectively ^{[9][21][22][23]}. Lutein and zeaxanthin, two yellow hydroxyl-xanthophylls, are used to treat aging-related macular degeneration because they are the only and necessary carotenoids for the macular function of the human eye ^{[2][21]}. The β -carotene and zeaxanthin can be ketonylated to produce canthaxanthin and astaxanthin, two orange or red keto-xanthophylls ^{[24][25]}. They are used as feed additives in aquaculture and poultry farming to give fish or egg yolks a red color ^{[1][11][26]}. Astaxanthin is also widely used in nutraceuticals, pharmaceuticals, and cosmetics due to its excellent antioxidant activity ^{[11][27]}. Epoxy-xanthophyll, antheraxanthin, and violaxanthin are derived from another branch of the metabolic pathway of zeaxanthin, capsorubin, neoxanthin, etc. ^{[29][30][31][32][33][34][35][36]}. These different modifications allow xanthophyll to show a broader color variation and stronger antioxidant activity compared to carotene ^{[24][27][38][39][40]}.

2. General Strategy for the Xanthophyll Production by Metabolic Engineering of Model Microorganisms

Briefly, the microbial metabolic engineering of xanthophyll involves the docking of the exogenous pathway of xanthophyll synthesis with the endogenous core metabolism of the microbial host, allowing the flows of carbon and energy to the synthesis of xanthophyll.

In model microorganisms, glucose or other carbon sources undergo glycolysis to produce acetyl coenzyme A (acetyl-CoA) and pyrurate. The acetyl-CoA and pyrurate then pass through the ergosterol synthesis pathway of the host cell, MVA (in plants, yeast, and algae) or MEP (in plants and bacteria) ^{[16][17]}, to produce geranylgeranyl diphosphate (GGPP), an intermediate metabolite of the ergosterol pathway. GGPP is the connect point between core microbial metabolism and exogenous carotinoid synthesis pathways ^[41]. The subsequent synthesis of carotene and xanthophyll requires various heterologous carotenogenic enzymes and cofactors to convert GGPP further ^{[18][19][20]}.

Experimental manipulation is used to introduce the genes of carotegenesis enzymes and cofactors into the microorganism and utilize the transcriptional, translational, and metabolic regulatory functions of the microbial host to control the expression of these genes. The key genetic elements involved in this process include highly active key enzyme genes, cofactor genes matching the key enzymes, various promoters and terminators, etc. The key enzyme genes and cofactor genes will be described in detail in Section 3, and the structures of expression cassettes consisting of the most effective promoters, genes, and terminators are listed in **Table 1**.

3. Progress in Metabolic Engineering Synthesis of Various Xanthophylls

3.1. Hydroxy-Xanthophyll

Hydroxy-xanthophyll mainly refers to zeaxanthin, lutein, β -cryptoxanthin, zeinoxanthin, etc. The β -cryptoxanthin and zeinoxanthin are the intermediate metabolites of zeaxanthin and lutein ^{[9][21][23]}, respectively, and the metabolic engineering of model microorganisms taking these intermediate metabolites as the end products has not yet been reported, so this section focuses on the engineering biosynthesis of zeaxanthin and lutein.

Zeaxanthin is the first xanthophyll in the β -carotene branch of the xanthophyll synthetic pathway and is the precursor of various high-value xanthophyll, such as astaxanthin, violaxanthin etc. (**Figure 1**). Unfortunately, although the production of zeaxanthin in various microbial engineering hosts has shown an increasing trend in recent years, it is still too low, with a maximum yield of only 18.7 mg/g^[42], and it is the speed-limiting step of the engineering synthesis of downstream xanthophyll.

The reported model microorganisms for zeaxanthin synthesis include *E. coli*, *S. cerevisiae*, and *Y. lipolytica*. Zeaxanthin engineering synthesis is generally achieved by first introducing the exogenous β -carotene pathway into the host to construct a carotene chassis strain; the heterologous CrtZ is then expressed in this chassis strain to hydroxylate the β -ring at each end of β -carotene to produce zeaxanthin. The CrtZ gene has been cloned from a variety of algae and bacteria and applied to the metabolic engineering synthesis of zeaxanthin. In *E. coli*, compared with those from *Pantoea agglomerans* and *Haematococcus pluvialis*, the CrtZ of *Pantoea ananatis* has the highest β -carotene hydroxylation activity [42][43][44][45]. In *Y. lipolytica*, the catalytic activity of CrtZ from *Brevundimonas Vesicularis*, *H. lacustris* and *P. ananatis* was compared ^[46], and *P. ananatis* CrtZ had the highest activity, as in *E. coli*. In *S. cerevisiae*, the synthesis of zeaxanthin was mainly used as a reporting approach to study the feasibility of experimental methods such as multi gene co-transformation ^{[47][48]}; research aimed at improving the yield of zeaxanthin is rare. The discovery, optimization, and heterologous functional expression of highly active CrtZ are critical and urgent problems for zeaxanthin synthesis.

In addition to the above CrtZs, CrtZ proteins sequences of other species obtained by gene prediction or protein sequence alignment can be found on the NCBI GenBank database published in recent years, such as *Massilia varians* (GenBank: BDT58294.1), *Xanthomonadaceae bacterium* (GenBank: RZA32178.1), *Oxalobacteraceae bacterium* (GenBank: USX26554.1), etc. Their identities of protein sequences aligning against the CrtZ of *P. ananatis* are generally in the range

of 60–80%, which is not lower than the identities between the above reported CrtZs. The potential of these putative CrtZ applications for the metabolic engineering of zeaxanthin also deserves further exploration.

Another important representative of hydroxyl-xanthophyll is lutein. The metabolic engineering of lutein is more complex and difficult than zeaxanthin due to the asymmetric cyclization of lycopene. α - carotene is the direct precursor of lutein, and the β -ring and ϵ -ring on the two ends of α - carotene are catalyzed by lycopene β -cyclase and ϵ -cyclase, respectively (**Figure 1**). The hydroxylation of the β -ring and ϵ -ring are catalyzed by two cytochrome P450 enzymes, CYP97A and CYP97C, respectively. These hydroxylation reactions require ferredoxin-NADP+ reductase (FNR), with ferredoxin redox partner, NADPH, and flavin adenine dinucleotide (FAD) as cofactors [49][50][51].

Studies have shown that the heterologously expressed activity of β -cyclase is higher than that of ϵ -cyclase, leading to a preference for lycopene for the formation of β -carotene with two β -rings ^{[52][53][54]}. Therefore, in addition to the hydroxylation, the asymmetric cyclization of lycopene is also a hindrance to lutein production in model microorganisms. The reported engineering microorganisms producing lutein are only *E. coli* and *S. cerevisiae*, and the yield is very low. In *E. coli*, through the screening of ϵ -cyclase from three sources (*Lactuca sativa, Tagetes erecta,* and *Marchantia polymorpha*) and CYP97C from nine sources (*Chlamydomonas reinhardtii, H. pluvialis, Brassica napus, Chenopodium quinoa, Oryza sativa, L. sativa, Nicotiana tabacum, Helianthus annuus,* and *M. polymorpha*), only the enzymes from *M. polymorpha* have lutein biosynthetic activity ^{[52][55]}. In *S. cerevisiae*, the synthesis of lutein can be achieved by the co-expression of β -cyclase from *Xanthophyllomyces dendrorhous*, ϵ -cyclase from *Tagetes erecta*, and CYP97A and CYP97C from *Arabidopsis thaliana* ^{[53][54]}.

3.2. Keto-Xanthophyll

Among all keto-xanthophylls (including astaxanthin, canthaxanthin, echinenone, etc.), astaxanthin has received the most attention due to its strongest antioxidant activity; this promotes the most in-depth research and the highest fermentation yield, and has shown great industrialization potential. Under controlled bioreactor fermentation, the maximum yield can reach tens of mg/g DCW ^[56], which could be comparable to a yield of approximately 40 mg/g DCW of the industrial algae natural-producer *H. pluvialis* ^[57].

The most commonly used model microorganisms for the production of astaxanthin include *E. coli, S. cerevisiae*, and *Y. lipolytica* ^{[24][58]}. *Kluyveromyces maximus* ^{[59][60]}, *P. pastoris* ^{[61][62]}, *Corynebacterium glutamicum* ^{[63][64]} etc., as hosts have also been reported.

The first step in astaxanthin synthesis by metabolic engineering is the heterologous functional expression of CrtW and CrtZ. Since both enzymes involve two substrates, there is an issue of substrate preference. Therefore, the yield of astaxanthin is closely related to the enzymatic activity, the substrate preference, and the combination of the two enzymes. Accordingly, the first step of almost all metabolic engineering research on the heterologous production of astaxanthin is the screening and adaptability analysis of these two enzymes. In E. coli, the combination of CrtW from Brevundimonas sp. SD212 and CrtZ from Pantoea sp. (including P. agglomerans and P. ananatis), or the CrtZ from Paracoccus sp. PC1 as a supplement to CrtZ from Pantoea sp., produces the highest astaxanthin yield [65][66][67][68][69][70][71]. The CrtZs from Pantoea sp. and Paracoccus sp. PC1 have different substrate preferences; a high conversion efficiency from β -carotene and canthaxanthin to astaxanthin can be achieved when the two enzymes have a combinatorial expression at a specific copy number ratio [65]. In S. cerevisiae, the higher astaxanthin production comes from the combination of the mutant BKT (H165R/V264D/F298Y/M1T/N188D/L271R) and CrtZ (L288R) of H. pluvialis, or the combination of the CrtW of B. vesicularis and CrtZ of Agrobacterium aurantiacum [72][73][74][75][76][77][78][79]. In Y. lipolytica, Wang [25] and Ma [80] et al. reported that the combination of the CrtZ from Paracoccus sp. and the CrtW from H. pluvialis yielded a higher astaxanthin production. Furthermore, the result of Wang et al. also showed that astaxanthin cannot be detected with the co-expression of CrtW and CrtZ from H. pluvialis in Y. lipolytica, On the contrary, Zhu [63] and Tramontin [81] reported that a higher astaxanthin production was produced by these CrtW and CrtZ from H. pluvialis. This contradiction needs to be further verified. In general, no matter the type of host, the bacteria CrtW and CrtZ usually have higher activity, followed by algae enzymes, and X. dendrorhous and plants (such as Adonis aestivalis) have very low or almost no activity [78][82].

Almost all reports indicate that CrtZ is a rate-limiting enzyme when the astaxanthin synthetic pathway is introduced in a model microbial host, which means the conversion from β -carotene to zeaxanthin, or from canthaxanthin and echinenone to astaxanthin, which is a key step for limiting the production of astaxanthin. Thus, in addition to the CrtZs listed in **Table 1**, the functional identification of other putative CrtZs mentioned in 2.1 may also be of significance for the synthesis of astaxanthin.

Canthaxanthin is another keto-xanthophyll that is widely used commercially. At present, only *E. coli* ^[83] and *S. cerevisiae* ^[84] have been reported as engineering microorganisms that take canthaxanthin as the target product. Theoretically, the synthesis of canthaxanthin does not require CrtZ, and its production should be higher than astaxanthin. Indeed, comparing the experimental results of Ye et al. ^{[76][84]}, the yield of canthaxanthin (approximately 10–15 mg/g) in *S. cerevisiae* was higher than that of axtathophyll (5.7 mg/g) when almost the same strategies and shake flask culture conditions were used. This further demonstrates the rate-limiting effect of CrtZ.

3.3. Epoxy-Xanthophyll

Epoxy-xanthophyll mainly includes violaxanthin and antheraxanthin. Antheraxhanthin was mentioned mainly as the intermediate metabolite of violaxanthin, so the researchers only focused on the engfigureineering biosynthesis of violaxanthin in this section. The biosynthesis of violaxanthin by metabolic engineering is achieved by prolonging the pathway of zeaxanthin with the introduction of ZEP into a zeaxanthin-producing microbial host. Although the cloning and heterologous expression of ZEP have been studied for nearly 30 years [85][86][87][88], as far as the metabolic engineering of violaxanthin is concerned, it has only been described in detail in recent years.

ZEP belongs to FAD-dependent monooxygenase and catalyzes the epoxidation of zeaxanthin in the presence of NADPH, FAD, ferredoxin (FD), and ferredoxin-NADPH oxidoreductase (FNR) [89][90]. Therefore, the synthesis of violaxanthin in an engineering host requires the reducing power and the redox pair ether from the endogenous metabolism of the host or the functional heterologous expression of relative genes. In general, the reducing power of host cells cannot meet the needs of an efficient synthesis of violaxanthin, and not all redox pairs can transfer the electron to ZEP due to the specificity of ZEP for the redox partner. Therefore, the first challenge of violaxanthin metabolic engineering is the heterologous expression of highly active ZEP and its matching redox partner. The reported microorganism hosts for violaxanthin synthesis include E. coli and S. cerevisiae. In E. coli, the activity of ZEPs from seven higher-plants (Capsicum annuum, A. thaliana, etc.), one liverwort (M. polymorpha), and one algae (Phaeodactylum tricornutum) was compared [91]. The C. annuum ZEP had the highest activity, whereas that of Prunus ameniaca, Zea mays, M. polymorpha, and P. tricornutum showed extremely low activity in E. coli. Although the NADPH and electron transport system of E. coli can make ZEP active, the yield of violaxanthin is low. Exogenous redox partners, spinach ferredoxin and ferredoxin oxidoreductase, can improve the ZEP activity, whereas Nostoc sp. severely decreases its activity. The Bacillus subtilis glucose dehydrogenase as the NADPH-regenerating enzyme can increase the synthesis of violaxanthin in E. coli. Furthermore, the E. coli strains and ribosome-binding site (RBS) sequences also impacted the yield of violaxanthin. The best E. coli strain, JM101, had a yield of 231 µg/g of violaxanthin; however, the ratio of violaxanthin to total carotenoids was only 1.4-21%, which was still low. In S. cerevisiae, compared with the ZEPs of A. thaliana and Solanum lycopersicum, the Haematococcus lacustris ZEP showed the highest activity [92]. The redox partner from A. thaliana can improve the violaxanthin yield, whereas the yeast mitochondrial ferredoxin-like protein and its reductase has no effect on the yield. In the best violaxanthin-producing S. cerevisiae, the final yield of violaxanthin reached 7.3mg/g, corresponding to 58.4% of total carotenoids.

From the above, in *E. coli* and in *S. cerevisiae*, a considerable amount of upstream carotene remained, indicating that the catalytic efficiency of the heterologously expressed ZEP was low. The characterization and screening of ZEP and the matching redox partner from different species in various microbial hosts need to be further explored.

3.4. Epoxy-Xanthophyll Derivative

Epoxy-xanthophyll derivatives include capsanthin, capsorubin, neoxanthin, fucoxanthin, diadinoxanthin, etc. Capsanthin, capsorubin, and neoxanthin are usually considered as the last metabolites of the carotenoid synthesis pathway in higher plants. The epoxy-xanthophyll derivatives have more complex terminal groups, and they may have multiple terminal groups, such as hydroxyl, ketone, epoxy, allenic, cyclopentane, cyclohexane, etc. (Figure 1).

Except for neoxanthin synthase (NSY) and the capsanthin/capsorubin synthase (CCS), the synthases for other epoxyxanthophyll derivatives have not yet been identified or characterized. The NSY catalyze the opening of cyclohexenyl 5–6 epoxides at one end of violaxanthin to form an allenic group through a transient carbocation ^[31]. Although the NSY from *Lycopersicum esculentum* ^[31], *S. tuberosum* ^[93], *Arabidopsis* ^[32], and *Chinese Kale* ^[33] has been cloned, and the activity of *L. esculentum* NSY ^[94] conversing from violaxanthin to neoxanthin has been verified in *E. coli* (the substrate violaxanthin was added to the system and the yield of neoxhanthin was not mentioned), to date, neoxanthin production by microbial metabolic engineering has not been reported in detail.

Table 1. Non-native xanthophyll-producing microorganisms with the highest yield or titer ^a.

Xanthophyll	Engineering Microbial Hosts	Key Enzymes	Natural Origin Species	Key Expression Cassettes ^b	Methods or Principles of Host Transformation	Key Strategies	Yield (mg/g DCW)	Ti (n
	E. coli	ε-LCY	M. polymorpha	P _{tac} -IDI- CrtE-CrtB- CrtI-		Selection of ε - LCY and CYP97C from different species, decreasing the activity of β - LCY and increasing the copy number of ε -LCY gene Selection of ε - LCY form different species, regulation of ratios of CYP97A3 and RENR1/ED3.		
		β-LCY li	M. polymorpha	MpLCYb- MpLCYe- CrtZ-T _{rrnB}	Electroporation			2.
		CYP97C	M. polymorpha	and P _{T7} - <i>MpCYP97C-</i> T _{T7} - P _{T7} -				
		CrtZ	P. ananatis	MpLCYe- T _{T7}				
	S. cerevisiae	ε-LCY	A. thaliana	P _{TER1} - tHMG1- T _{CYC1} , P _{PGK1} - CrtE03M-	Se LC diff spo reg rati CY Chemical RF transformation and hie dyn reg bas ten			
		CrtYB	X. dendrorhous	T _{ADH1} , P _{PGK1} - <i>CrtYB11M</i> - T _{ADH1} , P _{TEF} - <i>CrtI</i> -T _{CYC1} , P _{ACT1} -				
		СҮР97А3	A. thaliana	Gal4M9- T _{ADH1} , P _{GAL1} - <i>CrtYB</i> - T _{CYC1} , T _{CYC1} -				
		Lut1	A. thaliana	CYP97A3- P _{GAL1} - P _{GAL10} - <i>LUT1-</i> T _{ADH1} - T _{PGK1} -FD3-		and hierarchical dynamic regulation based on the temperature-	4.53	1
		RFNR1	A. thaliana	P _{GAL2} - P _{GAL7} - <i>RFNR1</i> - T _{TPS1} , P _{TEF1} - <i>PM^{SeV-C}-At</i> -		responsive promoter		
		FD3	A.s thaliana	LCYE- T _{CYC1} , P _{GAL1} - CYP97A3- T _{CYC1}				

Xanthophyll	Engineering Microbial Hosts	Key Enzymes	Natural Origin Species	Key Expression Cassettes ^b	Methods or Principles of Host Transformation	Key Strategies	Yield (mg/g DCW)	Ti (n
zeaxanthin	E. coli	CrtZ	P. ananatis	P _{T5} - CrtEIBipi- TTR, P37- CrtY- 2CrtZ- T _{rmB} , pZSP _{IA44} - MevT _{TIGR} - MevB _{TIGR} IS-2	Electroporation	Introduction and dynamic control of the MVA pathway of <i>S. cerevisiae</i> to increase the precursors supply and prevent the accumulation of toxic metabolites	18.7	5
	S. cerevisiae	CrtZ	P. ananatis	P _{PDC1} -CrtE- T _{PDC1} , P _{TP11} -CrtB- T _{TP11} , P _{GPM1} -CrtI- T _{GPM1} , P _{GPD} -CrtY- T _{GPD} , P _{FBA1} -CrtZ- T _{FBA1}	Chemical transformation	Zeaxanthin as a reporter gene for identification of promoter strength		0.
	Y. Lipolytica	CrtZ	P. ananatis	P _{TEF1N} - <i>CrtE</i> -T _{xpr2} , P _{TEF1N} - <i>CrtB</i> -T _{xpr2} , P _{TEF1N} - <i>CrtI</i> - T _{xpr2} , P _{TEF1N} - <i>CarRP</i> - T _{xpr2} , P _{TEF1N} - <i>CrtZ</i> -T _{xpr2}	Frozen-EZ Yeast Transformation II Kit	High-copy- number integration of CrtZ gene into ribosomal DNA region		2: m in m

Xanthophyll	Engineering Microbial Hosts	Key Enzymes	Natural Origin Species	Key Expression Cassettes ^b	Methods or Principles of Host Transformation	Key Strategies	Yield (mg/g DCW)	Ti (n
astaxanthin	E. coli	CrtZ CrtW	P. ananatis Brevundimonas sp. SD212	P _{TM2} - CrtEBIA, P _{T7} -RLZ- CrtZ-RLW- CrtW		Screening and regulation of promoters and RBSs	15.1	6:
	S.	CrtZ	A. aurantiacum	A high β- carotene producing strain with	Homologous	Selection and optimization of combinations of CrtW and CrtZ	6.05	
cerevisiae	CrtW	Alcaligenes sp.	T _{ADH1} , P _{TDH3} - <i>CrtW</i> - T _{TDH2}		from different species			
	Y. Lipolytica	CrtZ	H. pluvialis	P _{TEF} -carRP- T _{XPR2} , P _{TEF} - <i>thmgR</i> - T _{XPR2} , P _{TEF} - GGS1-		Selection of CrtW and CrtZ		
Y. Lipolytic P. pastoris		ВКТ	H. pluvialis	T _{XPR2} , P _{TEF} - <i>carB</i> -T _{XPR2} , P _{TEF} - <i>CrtW</i> . linker-RIDD ⁻ T _{XPR2} -P _{TER} - <i>CrtZ</i> -linker- RIAD ⁻ T _{XPR2}	Chemical from differen transformation species and fine-tuning th transcription	from different species and fine-tuning their transcription	17.5	
	P. pastoris	CrtZ P. pastoris		P _{AOX1} -CrtI- T _{CYC1} , P _{AOX1} -CrtE- CrtZ-T _{CYC1} ,	CRISPR/Cas9	Astaxanthin as a reporter gene for marker-less integration of multigene		A 2.
		CrtW		P _{AOX1} - <i>CrtYB-CrtW</i> - T _{CYC1}		pathways into <i>Pichia pastoris</i> via CRISPR/Cas9		۲.

Xanthophyll	Engineering Microbial Hosts	Key Enzymes	Natural Origin Species	Key Expression Cassettes ^b	Methods or Principles of Host Transformation	Key Strategies	Yield (mg/g DCW)	Ti (n
		CrtZ	H. pluvialis	P _{KILac4} -CrtZ- T _{KILac4} , P _{ScGapDH} - CrtE-T _{ScGap} , P _{ScPGK} - CrtZ- T _{ScPGK} , P _{KIGapDH} - kanMX-	Homologous	Increasing the copy number of Hpchyb and BKT genes and	3.125 in YPL	
	n. marxianus	ВКТ	C. reinhardtii	P _{ICL} -Crtl- T _{35S} , P _{KIPGK} -BKT- T _{SCPGK} , P _{KIADH1} - CrtYB- T _{SCGap} , P _{SCADH1} - tHMG- T _{SCADH1}	recombination	modifying the Hpchyb by site- directed mutagenesis	5.701 in YPG medium	
	C. glutamicum	CrtZ	F. pelagi	P _{tuf} - <i>CrtZ-</i> linker- <i>CrtW</i>	electroporation with xenogeneic plasmid DNA	Fusion expression of CrtZ and CrtW, increasing the expression of upstream enzymes,	3.1	
		CrtW	B. aurantiaca			mediated medium composition		

Xanthophyll	Engineering Microbial Hosts	Key Enzymes	Natural Origin Species	Key Expression Cassettes ^b	Methods or Principles of Host Transformation	Key Strategies	Yield (mg/g DCW)	Ti (n
	E. coli	вкт	Anabaena variabilis	P _{Trc} -CrtW	Electroporation	Overexpression of host genes increases the carbon flux into the canthaxanthin biosynthetic pathway	Approximately 10.65	24
canthaxanthin	S. cerevisiae	OBKTM29 (mutant BKT)	H. pluvialis	P _{GAL1} - <i>mBKT</i> - T _{CYC1} - P _{GAL10} - <i>CrtE03</i> - T _{ADH2} , T _{CYC1} - <i>PM^{SeV-C}</i> - <i>mBKT</i> - P _{GAL1} -	Homologous recombination and CRISPR/cas9	Subcellular re- localization of OBKTM29 and its copy number adjustment both in the cytoplasm and on the periplasmic membrane, pleiotropic drug resistance (PDR) regulator overexpression	approximately 20–30	1(

Xanthophyll	Engineering Microbial Hosts	Key Enzymes	Natural Origin Species	Key Expression Cassettes ^b	Methods or Principles of Host Transformation	Key Strategies	Yield (mg/g DCW)	Ti (n
violaxanthin	E. coli	CrtZ	P. ananatis	P _{lac} -CrtE- CrtY-CrtI- CrtB-CrtZ, P _{lac} -ZEP, P _{T7} -gdh	Ξ- /- tZ, 2,	Selection of ZEP from different species and		
		ZEP	C. annuum			optimization of <i>E. coli</i> strain, expression vector, and	0.231	
		glucose dehydrogenase (gdh)	B. subtilis			ribosome- binding site (RBS) sequence		
	S. cerevisiae	CrtZ	P. ananatis	Р _{тDH3} - <i>CrtYB</i> - Т _{СYC1} , Р _{TDH3} -CrtI-	Select CrtZ, Z F- redox from d E- specie Modified their tr z- homologous varian increa gene d numbe upstre carote genes	Selection of CrtZ, ZEP and redox partner		
		ZEP	H. lacustris	P _{TDH3} -CrtE- T _{CYC1} , P _{TEF1} -CrtZ-		species and their truncated variants,	7.3	
		RFNR1	A. thaliana	linker- <i>trZEP</i> - T _{ADH1} , P _{TEF1} - <i>trRFNR1</i> -		increasing gene copy number of upstream		
		FD3	A. thaliana	T _{ADH1} , P _{PGK1} - <i>trFD3</i> -T _{CYC1}		carotenogenic genes		
		CrtZ	P. ananatis	P _{lac} -HpIDI- CrtE-CrtY-		A particularly		
capsanthin	E. coli	ZEP	C. annuum	Crtl-CrtB- CrtZ, P _{tac/T7} -		high expression of	С	0.
		CCS	C. annuum	CaZEP				

^a Under shake flask culture conditions. Under the conditions of fermentation tank, it is difficult to compare the yield or titer due to different fermentation types, medium, and fermenting duration. In order to relatively accurately reflect the ability of engineered cells to synthesize xanthophyll, the highest yields or titers here are for shake flask culture conditions. ^b Capital P and T indicate promoters and terminators, respectively.

Capsanthin is the only epoxy-xanthophyll derivative whose synthesis has been achieved by microbial metabolic engineering. Capsanthin and capsorubin are synthesized by CCS catalysis from antheraxanthin and violaxanthin, respectively (**Figure 1**). Although there are at least ten protein sequences from different species identified as CCSs in NCBI GenBank, only CCSs from *Capsicum annuum* ^[95] and *Tiger lily* ^[30] have been cloned and verified for their functions so far.

CCS is a multifunctional enzyme with capsanthin/capsorubin synthesis and β -carotene synthesis, and its activity requires the simultaneous existence of FAD and NADPH. In addition, it was also found that CCS catalyzes the reverse reaction in the presence of NAD⁺; that is, the reverse conversion from capsorubin to violaxanthin ^{[29][96][97][98]}. In 2021, by introducing ZEP and CCS from *C. annuum* into β -carotene-producing *E. coli*, combined with greatly enhancing the expression of CCS, the production of capsanthin was realized with a titer of 0.5 mg/L ^[99]. The high-level expression of CCS is necessary for the production of capsanthin; otherwise, only lycopene cyclization activity can be detected. Moreover, in the best strains, there are still considerable residues of precursors (zeaxanthin, antheraxanthin, and violaxanthin), which proves that CCS as well as ZEP are the rate-limiting enzymes of the capsanthin synthetic pathway in the engineering of *E. coli*. In addition, no capsorubin was detected in this study, which was consistent with the phenotype of the capsanthin accounting for a large amount in pepper, whereas the capsorubin was only a minor product [37]. Therefore, it can be speculated that CCS may have a preference for violaxanthin over antheraxanthin. Any metabolic engineering of capsanthin or capsorubin in yeast has not yet been reported.

From the above, it is known that the metabolic engineering of epoxy-xanthophyll derivatives is still at the beginning stage. The cloning and characterization of key enzymes, substrate preference, and cofactor specificity need to be further investigated. From the catalytic properties of CCS, it can be speculated that the key enzymes, such as the NSY, fucoxanthin, or diadinoxanthin synthetic enzymes, may also involve multiple substrates or multiple products, which implies that the metabolic engineering of epoxy-xanthophyll derivatives is more difficult to control.

References

- 1. Maoka, T. Carotenoids as natural functional pigments. J. Nat. Med. 2020, 74, 1-16.
- Aziz, E.; Batool, R.; Akhtar, W.; Rehman, S.; Shahzad, T.; Malik, A.; Shariati, M.A.; Laishevtcev, A.; Plygun, S.; Heydari, M.; et al. Xanthophyll: Health benefits and therapeutic insights. Life Sci. 2020, 240, 117104.
- Esteban, R.; Moran, J.F.; Becerril, J.M.; García-Plazaola, J.I. Versatility of carotenoids: An integrated view on diversity, evolution, functional roles and environmental interactions. Environ. Exp. Bot. 2015, 119, 63–75.
- 4. Sun, T.; Yuan, H.; Cao, H.; Yazdani, M.; Tadmor, Y.; Li, L. Carotenoid Metabolism in Plants: The Role of Plastids. Mol. P lant 2018, 11, 58–74.
- Jahns, P.; Holzwarth, A.R. The role of the xanthophyll cycle and of lutein in photoprotection of photosystem II. Biochim. Biophys. Acta 2012, 1817, 182–193.
- Jackson, H.; Braun, C.L.; Ernst, H. The Chemistry of Novel Xanthophyll Carotenoids. Am. J. Cardiol. 2008, 101, 50D–5 7D.
- 7. Nakano, T.; Wiegertjes, G. Properties of Carotenoids in Fish Fitness: A Review. Mar. Drugs 2020, 18, 568.
- 8. Pereira, A.G.; Otero, P.; Echave, J.; Casais, A.C.; Chamorro, F.; Collazo, N.; Jaboui, A.; Lopes, C.L.; Gandara, J.S.; Pri eto, M.A. Xanthophylls from the Sea: Algae as Source of Bioactive Carotenoids. Mar. Drugs 2021, 19, 188.
- 9. Bhosale, P.; Bernstein, P.S. Microbial xanthophylls. Appl. Microbiol. Biotechnol. 2005, 68, 445-455.
- 10. Claudia, S.-D. Engineering novel carotenoids in microorganisms. Curr. Opin. Biotechnol. 2000, 11, 255–261.
- Patel, A.K.; Tambat, V.S.; Chen, C.-W.; Chauhan, A.S.; Kumar, P.; Vadrale, A.P.; Huang, C.-Y.; Dong, C.-D.; Singhania, R.R. Recent advancements in astaxanthin production from microalgae: A review. Bioresour. Technol. 2022, 364, 12803
 0.
- Ma, R.; Tao, X.; Chua, E.T.; Ho, S.-H.; Shi, X.; Liu, L.; Xie, Y.; Chen, J. Enhancing astaxanthin production in Haematoco ccus pluvialis QLD by a pH steady NaHCO3-CO2-C/NH4CI-N culture system. Algal Res. 2022, 64, 102697.
- Yi, X.; Alper, H.S. Considering Strain Variation and Non-Type Strains for Yeast Metabolic Engineering Applications. Life 2022, 12, 510.
- 14. Ding, Q.; Diao, W.; Gao, C.; Chen, X.; Liu, L. Microbial cell engineering to improve cellular synthetic capacity. Biotechn ol. Adv. 2020, 45, 107649.
- 15. Nielsen, J.; Keasling, J.D. Engineering Cellular Metabolism. Cell 2016, 164, 1185–1197.
- 16. Pu, X.; Dong, X.; Li, Q.; Chen, Z.; Liu, L. An update on the function and regulation of methylerythritol phosphate and m evalonate pathways and their evolutionary dynamics. J. Integr. Plant Biol. 2021, 63, 1211–1226.
- 17. Niu, F.; Lu, Q.; Bu, Y.; Liu, J. Metabolic engineering for the microbial production of isoprenoids: Carotenoids and isopre noid-based biofuels. Synth. Syst. Biotechnol. 2017, 2, 167–175.
- 18. Wang, C.; Zhao, S.; Shao, X.; Park, J.-B.; Jeong, S.-H.; Park, H.-J.; Kwak, W.-J.; Wei, G.; Kim, S.-W. Challenges and t ackles in metabolic engineering for microbial production of carotenoids. Microb. Cell Fact. 2019, 18, 55.
- Jing, Y.; Wang, Y.; Zhou, D.; Wang, J.; Li, J.; Sun, J.; Feng, Y.; Xin, F.; Zhang, W. Advances in the synthesis of three typ ical tetraterpenoids including β-carotene, lycopene and astaxanthin. Biotechnol. Adv. 2022, 61, 108033.
- Jing, Y.; Guo, F.; Zhang, S.; Dong, W.; Zhou, J.; Xin, F.; Zhang, W.; Jiang, M. Recent advances on biological synthesis of lycopene by using industrial yeast. Ind. Eng. Chem. Res. 2021, 60, 3485–3494.
- Zafar, J.; Aqeel, A.; Shah, F.I.; Ehsan, N.; Gohar, U.F.; Moga, M.A.; Festila, D.; Ciurea, C.; Irimie, M.; Chicea, R. Bioche mical and Immunological implications of Lutein and Zeaxanthin. Int. J. Mol. Sci. 2021, 22, 10910.
- Gupta, I.; Adin, S.N.; Panda, B.P.; Mujeeb, M. β-Carotene-production methods, biosynthesis from Phaffia rhodozyma, f actors affecting its production during fermentation, pharmacological properties: A review. Biotechnol. Appl. Biochem. 20 22, 69, 2517–2529.
- Zhang, Y.; Liu, Z.; Sun, J.; Xue, C.; Mao, X. Biotechnological production of zeaxanthin by microorganisms. Trends Food Sci. Technol. 2018, 71, 225–234.

- Basiony, M.; Ouyang, L.; Wang, D.; Yu, J.; Zhou, L.; Zhu, M.; Wang, X.; Feng, J.; Dai, J.; Shen, Y.; et al. Optimization of microbial cell factories for astaxanthin production: Biosynthesis and regulations, engineering strategies and fermentatio n optimization strategies. Synth. Syst. Biotechnol. 2022, 7, 689–704.
- Wang, D.-N.; Feng, J.; Yu, C.-X.; Zhang, X.-K.; Chen, J.; Wei, L.-J.; Liu, Z.; Ouyang, L.; Zhang, L.; Hua, Q.; et al. Integr ated pathway engineering and transcriptome analysis for improved astaxanthin biosynthesis in Yarrowia lipolytica. Synt h. Syst. Biotechnol. 2022, 7, 1133–1141.
- 26. Rebelo, B.A.; Farrona, S.; Ventura, M.R.; Abranches, R. Canthaxanthin, a Red-Hot Carotenoid: Applications, Synthe-si s, and Biosynthetic Evolution. Plants 2020, 9, 1039.
- 27. Stachowiak, B.; Szulc, P. Astaxanthin for the food industry. Molecules 2021, 26, 2666.
- Takemura, M.; Sahara, T.; Misawa, N. Violaxanthin: Natural function and occurrence, biosynthesis, and heterologous pr oduction. Appl. Microbiol. Biotechnol. 2021, 105, 6133–6142.
- Bouvier, F.; Hugueney, P.; d'Harlingue, A.; Kuntz, M.; Camara, B. Xanthophyll biosynthesis in chromoplasts: Isolation a nd molecular cloning of an enzyme catalyzing the conversion of 5,6-epoxycarotenoid into ketocarotenoid. Plant J. 199 4, 6, 45–54.
- Jeknić, Z.; Morré, J.T.; Jeknić, S.; Jevremović, S.; Subotić, A.; Chen, T.H.H. Cloning and functional characterization of a gene for capsanthin-capsorubin synthase from tiger lily (Lilium lancifolium Thunb. 'Splendens'). Plant Cell Physiol. 201 2, 53, 1899–1912.
- 31. Bouvier, F.; D'Harlingue, A.; Backhaus, R.A.; Kumagai, M.H.; Camara, B. Identification of neoxanthin synthase as a car otenoid cyclase paralog. Eur. J. Biochem. 2000, 267, 6346–6352.
- 32. North, H.M.; Almeida, A.D.; Boutin, J.-P.; Frey, A.; To, A.; Botran, L.; Sotta, B.; Marion-Poll, A. The Arabidopsis ABA-deficient mutant aba4 demonstrates that the major route for stress-induced ABA accumulation is via neoxanthin isomers. Pl ant J. 2007, 50, 810–824.
- Jian, Y.; Zhang, C.; Wang, Y.; Li, Z.; Chen, J.; Zhou, W.; Huang, W.; Jiang, M.; Zheng, H.; Li, M.; et al. Characterization of the role of the neoxanthin synthase gene BoaNXS in carotenoid biosynthesis in Chinese Kale. Genes 2021, 12, 112 2.
- Ahmad, A.; Tiwari, A.; Srivastava, S. A Genome-Scale Metabolic Model of Thalassiosira pseudonana CCMP 1335 for a systems-level understanding of its metabolism and biotechnological potential. Microorganisms 2020, 8, 1396.
- 35. Dambek, M.; Eilers, U.; Breitenbach, J.; Steiger, S.; Büchel, C.; Sandmann, G. Biosynthesis of fucoxanthin and diadino xanthin and function of initial pathway genes in Phaeodactylum tricornutum. J. Exp. Bot. 2012, 63, 5607–5612.
- Chen, D.; Yuan, X.; Zheng, X.; Fang, J.; Lin, G.; Li, R.; Chen, J.; He, W.; Huang, Z.; Fan, W.; et al. Multi-omics analyse s provide insight into the biosynthesis pathways of fucoxanthin in Isochrysis galbana. Genom. Proteom. Bioinform. 202 2.
- Gómez-García, M.R.; Ochoa-Alejo, N. Biochemistry and molecular biology of carotenoid biosynthesis in chili peppers (Capsicum spp.). Int. J. Mol. Sci. 2013, 14, 19025–19053.
- Palozza, P.; Krinsky, N.I. Astaxanthin and canthaxanthin are potent antioxidants in a membrane model. Arch. Biochem. Biophys. 1992, 297, 291–295.
- Kobayashi, M. In vivo antioxidant role of astaxanthin under oxidative stress in the green alga Haematococcus pluvialis. Appl. Microbiol. Biotechnol. 2000, 54, 550–555.
- 40. Kennedy, L.E.; Abraham, A.; Kulkarni, G.; Shettigar, N.; Dave, T.; Kulkarni, M. Capsanthin, a plant-derived xanthophyll: A review of pharmacology and delivery strategies. AAPS PharmSciTech 2021, 22, 203.
- Pyne, M.E.; Narcross, L.; Martin, V.J.J. Engineering plant secondary metabolism in microbial systems. Plant Physiol. 2 019, 179, 844–861.
- Shen, H.-J.; Cheng, B.-Y.; Zhang, Y.-M.; Tang, L.; Li, Z.; Bu, Y.-F.; Li, X.-R.; Tian, G.-Q.; Liu, J.-Z. Dynamic control of the mevalonate pathway expression for improved zeaxanthin production in Escherichia coli and comparative proteome ana lysis. Metab. Eng. 2016, 38, 180–190.
- Nishizaki, T.; Tsuge, K.; Itaya, M.; Doi, N.; Yanagawa, H. Metabolic engineering of carotenoid biosynthesis in Escherichi a coli by ordered gene assembly in Bacillus subtilis. Appl. Environ. Microbiol. 2007, 73, 1355–1361.
- 44. Li, X.-R.; Tian, G.-Q.; Shen, H.-J.; Liu, J.-Z. Metabolic engineering of Escherichia coli to produce zeaxanthin. J. Ind. Mic robiol. Biotechnol. 2015, 42, 627–636.
- 45. Wu, Z.; Zhao, D.; Li, S.; Wang, J.; Bi, C.; Zhang, X. Combinatorial modulation of initial codons for improved zeaxanthin synthetic pathway efficiency in Escherichia coli. Microbiologyopen 2019, 8, e930.
- Xie, Y.; Chen, S.; Xiong, X. Metabolic Engineering of non-carotenoid-producing yeast Yarrowia lipolytica for the biosynt hesis of zeaxanthin. Front. Microbiol. 2021, 12, 699235.
- Sun, J.; Shao, Z.; Zhao, H.; Nair, N.; Wen, F.; Xu, J.-H.; Zhao, H. Cloning and characterization of a panel of constitutive promoters for applications in pathway engineering in Saccharomyces cerevisiae. Biotechnol. Bioeng. 2012, 109, 2082– 2092.

- 48. Liang, J.; Ning, J.C.; Zhao, H. Coordinated induction of multi-gene pathways in Saccharomyces cerevisiae. Nucleic Aci ds Res. 2013. 41. e54.
- Niu, G.; Guo, Q.; Wang, J.; Zhao, S.; He, Y.; Liu, L. Structural basis for plant lutein biosynthesis from α-carotene. Proc. Natl. Acad. Sci. USA 2020, 117, 14150–14157.
- Hannemann, F.; Bichet, A.; Ewen, K.M.; Bernhardt, R. Cytochrome P450 systems--biological variations of electron tran sport chains. Biochim. Biophys. Acta 2007, 1770, 330–344.
- Iyanagi, T.; Xia, C.; Kim, J.-J.P. NADPH-cytochrome P450 oxidoreductase: Prototypic member of the diflavin reductase family. Arch. Biochem. Biophys. 2012, 528, 72–89.
- 52. Takemura, M.; Kubo, A.; Watanabe, A.; Sakuno, H.; Minobe, Y.; Sahara, T.; Murata, M.; Araki, M.; Harada, H.; Terada, Y.; et al. Pathway engineering for high-yield production of lutein in Escherichia coli. Synth. Biol. 2021, 6, ysab012.
- 53. Bian, Q.; Zhou, P.; Yao, Z.; Li, M.; Yu, H.; Ye, L. Heterologous biosynthesis of lutein in S. cerevisiae enabled by tempor ospatial pathway control. Metab. Eng. 2021, 67, 19–28.
- 54. Bian, Q.; Jiao, X.; Chen, Y.; Yu, H.; Ye, L. Hierarchical dynamic regulation of Saccharomyces cerevisiae for enhanced I utein biosynthesis. Biotechnol. Bioeng. 2023, 120, 536–552.
- Takemura, M.; Maoka, T.; Misawa, N. Biosynthetic routes of hydroxylated carotenoids (xanthophylls) in Marchantia poly morpha, and production of novel and rare xanthophylls through pathway engineering in Escherichia coli. Planta 2015, 2 41, 699–710.
- 56. Zhu, H.Z.; Jiang, S.; Wu, J.J.; Zhou, X.R.; Liu, P.Y.; Huang, F.H.; Wan, X. Production of high levels of 3 S,3' S-astaxant hin in Yarrowia lipolytica via iterative metabolic engineering. J. Agric. Food Chem. 2022, 70, 2673–2683.
- Le-Feuvre, R.; Moraga-Suazo, P.; Gonzalez, J.; Martin, S.S.; Henríquez, V.; Donoso, A.; Muñoz, C.A. Biotechnology ap plied to Haematococcus pluvialis Fotow: Challenges and prospects for the enhancement of astaxanthin accumulation. J. Appl. Phycol. 2020, 32, 3831–3852.
- Zhu, X.; Meng, C.; Sun, F.; Wei, Z.; Chen, L.; Chen, W.; Tong, S.; Du, H.; Gao, J.; Ren, J.; et al. Sustainable production of astaxanthin in microorganisms: The past, present, and future. Crit. Rev. Food Sci. Nutr. 2022, 1–17.
- 59. Chang, J.-J.; Thia, C.; Lin, H.-Y.; Liu, H.-L.; Ho, F.-J.; Wu, J.-T.; Shih, M.-C.; Li, W.-H.; Huang, C.-C. Integrating an algal β-carotene hydroxylase gene into a designed carotenoid-biosynthesis pathway increases carotenoid production in yeas t. Bioresour. Technol. 2015, 184, 2–8.
- Lin, Y.-J.; Chang, J.-J.; Lin, H.-Y.; Thia, C.; Kao, Y.-Y.; Huang, C.-C.; Li, W.-H. Metabolic engineering a yeast to produce astaxanthin. Bioresour. Technol. 2017, 245, 899–905.
- 61. Araya-Garay, J.M.; Ageitos, J.M.; Vallejo, A.J.; Veiga-Crespo, P.; Sánchez-Pérez, A.; Villa, T.G. Construction of a novel Pichia pastoris strain for production of xanthophylls. AMB Express 2012, 2, 24.
- 62. Gao, J.; Xu, J.; Zuo, Y.; Ye, C.; Jiang, L.; Feng, L.; Huang, L.; Xu, Z.; Lian, J. Synthetic biology toolkit for marker-less int egration of multigene pathways into Pichia pastoris via CRISPR/Cas9. ACS Synth. Biol. 2022, 11, 623–633.
- 63. Henke, N.A.; Heider, A.E.S.; Peters-Wendisch, P.; Wendisch, V.F. Production of the marine carotenoid astaxanthin by metabolically engineered Corynebacterium glutamicum. Mar. Drugs. 2016, 14, 124.
- 64. Henke, N.A.; Wendisch, V.F. Improved astaxanthin production with Corynebacterium glutamicum by application of a me mbrane fusion protein. Mar. Drugs 2019, 17, 621.
- 65. Zhang, M.; Gong, Z.; Tang, J.; Lu, F.; Li, Q.; Zhang, X. Improving astaxanthin production in Escherichia coli by co-utilizi ng CrtZ enzymes with different substrate preference. Microb. Cell Fact. 2022, 21, 71.
- Nogueira, M.; Enfissi, M.A.E.; Welsch, R.; Beyer, P.; Zurbriggen, D.M.; Fraser, D.P. Construction of a fusion enzyme for astaxanthin formation and its characterisation in microbial and plant hosts: A new tool for engineering ketocarotenoids. Metab. Eng. 2019, 52, 243–252.
- 67. Gong, Z.; Wang, H.; Tang, J.; Bi, C.; Li, Q.; Zhang, X. Coordinated expression of astaxanthin biosynthesis genes for im proved astaxanthin production in Escherichia coli. J. Agric. Food Chem. 2020, 68, 14917–14927.
- 68. Ye, L.; Zhu, X.; Wu, T.; Wang, W.; Zhao, D.; Bi, C.; Zhang, X. Optimizing the localization of astaxanthin enzymes for im proved productivity. Biotechnol. Biofuels. 2018, 11, 278.
- 69. Lu, Q.; Bu, Y.-F.; Liu, J.-Z. Metabolic Engineering of Escherichia coli for Producing astaxanthin as the predominant caro tenoid. Mar. Drugs 2017, 15, 296.
- 70. Scaife, M.A.; Burja, A.M.; Wright, P.C. Characterization of cyanobacterial beta-carotene ketolase and hydroxylase gene s in Escherichia coli, and their application for astaxanthin biosynthesis. Biotechnol. Bioeng. 2009, 103, 944–955.
- Zhang, C.; Seow, V.Y.; Chen, X.; Too, H.-P. Multidimensional heuristic process for high-yield production of astaxanthin a nd fragrance molecules in Escherichia coli. Nat. Commun. 2018, 9, 1858.
- 72. Qi, D.-D.; Jin, J.; Liu, D.; Jia, B.; Yuan, Y.-J. In vitro and in vivo recombination of heterologous modules for improving bi osynthesis of astaxanthin in yeast. Microb. Cell Fact. 2020, 19, 103.
- 73. Jiang, G.; Yang, Z.; Wang, Y.; Yao, M.; Chen, Y.; Xiao, W.; Yuan, Y. Enhanced astaxanthin production in yeast via combi ned mutagenesis and evolution. Biochem. Eng. J. 2020, 156, 107519.

- 74. Jin, J.; Wang, Y.; Yao, M.; Gu, X.; Li, B.; Liu, H.; Ding, M.; Xiao, W.; Yuan, Y. Astaxanthin overproduction in yeast by stra in engineering and new gene target uncovering. Biotechnol. Biofuels. 2018, 11, 230.
- 75. Wang, R.; Gu, X.; Yao, M.; Pan, C.; Liu, H.; Xiao, W.; Wang, Y.; Yuan, Y. Engineering of β-carotene hydroxylase and ket olase for astaxanthin overproduction in Saccharomyces cerevisiae. Front. Chem. Sci. Eng. 2017, 11, 89–99.
- 76. Zhou, P.; Li, M.; Shen, B.; Yao, Z.; Bian, Q.; Ye, L.; Yu, H. Directed coevolution of β-carotene ketolase and hydroxylase and its application in temperature-regulated biosynthesis of astaxanthin. J. Agric. Food Chem. 2019, 67, 1072–1080.
- Zhou, P.; Xie, W.; Li, A.; Wang, F.; Yao, Z.; Bian, Q.; Zhu, Y.; Yu, H.; Ye, L. Alleviation of metabolic bottleneck by combin atorial engineering enhanced astaxanthin synthesis in Saccharomyces cerevisiae. Enzym. Microb. Technol. 2017, 100, 28–36.
- 78. Zhou, P.; Ye, L.; Xie, W.; Lv, X.; Yu, H. Highly efficient biosynthesis of astaxanthin in Saccharomyces cerevisiae by inte gration and tuning of algal crtZ and bkt. Appl. Microbiol. Biotechnol. 2015, 99, 8419–8428.
- Ukibe, K.; Hashida, K.; Yoshida, N.; Takagi, H. Metabolic engineering of Saccharomyces cerevisiae for astaxanthin pro duction and oxidative stress tolerance. Appl. Environ. Microbiol. 2009, 75, 7205–7211.
- 80. Ma, Y.; Li, J.; Huang, S.; Stephanopoulos, G. Targeting pathway expression to subcellular organelles improves astaxan thin synthesis in Yarrowia lipolytica. Metab. Eng. 2021, 68, 152–161.
- Tramontin, L.R.R.; Kildegaard, K.R.; Sudarsan, S.; Borodina, I. Enhancement of astaxanthin biosynthesis in oleaginous Yeast Yarrowia lipolytica via microalgal pathway. Microorganisms 2019, 7, 472.
- Kildegaard, K.R.; Pérez, B.A.; Belda, D.D.; Khangura, J.K.; Holkenbrink, C.; Borodina, I. Engineering of Yarrowia lipolyti ca for production of astaxanthin. Synth. Syst. Biotechnol. 2017, 2, 287–294.
- Scaife, M.A.; Prince, C.A.; Norman, A.; Armenta, R.E. Progress toward an Escherichia coli canthaxanthin bioprocess. P rocess Biochem. 2012, 47, 2500–2509.
- Chen, M.; Li, M.; Ye, L.; Yu, H. Construction of canthaxanthin-producing yeast by combining spatiotemporal regulation and pleiotropic drug resistance engineering. ACS Synth. Biol. 2022, 11, 325–333.
- 85. Marinn, E.; Nussaume, L.; Quesada, A.; Gonneau, M.; Sotta, B.; Hugueney, P.; Frey, A.; Poll, A.M. Molecular identificati on of zeaxanthin epoxidase of Nicotiana plumbaginifolia, a gene involved in abscisic acid biosynthesis and correspondi ng to the ABA locus of Arabidopsis thaliana. EMBO J. 1996, 15, 2331–2342.
- Bouvier, F.; d'Harlingue, A.; Hugueney, P.; Marini, E.; Polli, A.M.; Camara, B. Xanthophyll biosynthesis. Cloning, expres sion, functional reconstitution, and regulation of beta-cyclohexenyl carotenoid epoxidase from pepper (Capsicum annu um). J. Biol. Chem. 1996, 271, 28861–28867.
- Zhu, C.; Yamamura, S.; Nishihara, M.; Koiwa, H.; Sandmann, G. cDNAs for the synthesis of cyclic carotenoids in petals of Gentiana lutea and their regulation during flower development. Biochim. Biophys. Acta 2003, 1625, 305–308.
- 88. Eilers, U.; Dietzel, L.; Breitenbach, J.; Büchel, C.; Sandmann, G. Identification of genes coding for functional zeaxanthi n epoxidases in the diatom Phaeodactylum tricornutum. J. Plant Physiol. 2016, 192, 64–70.
- 89. Deng, Y.; Zhou, Q.; Wu, Y.; Chen, X.; Zhong, F. Properties and mechanisms of flavin-dependent monooxygenases and their applications in natural product synthesis. Int. J. Mol. Sci. 2022, 23, 2622.
- Iyanagi, T. Roles of Ferredoxin-NADP+ Oxidoreductase and Flavodoxin in NAD(P)H-dependent electron transfer syste ms. Antioxidants 2022, 11, 2143.
- Takemura, M.; Kubo, A.; Higuchi, Y.; Maoka, T.; Sahara, T.; Yaoi, K.; Ohdan, K.; Umeno, D.; Misawa, N. Pathway engin eering for efficient biosynthesis of violaxanthin in Escherichia coli. Appl. Microbiol. Biotechnol. 2019, 103, 9393–9399.
- Cataldo, V.F.; Arenas, N.; Salgado, V.; Camilo, C.; Ibáñez, F.; Agosin, E. Heterologous production of the epoxycarotenoi d violaxanthin in Saccharomyces cerevisiae. Metab. Eng. 2020, 59, 53–63.
- Babili, S.A.; Hugueney, P.; Schledz, M.; Welsch, R.; Frohnmeyer, H.; Laule, O.; Beyer, P. Identification of a novel gene coding for neoxanthin synthase from Solanum tuberosum. FEBS Lett. 2000, 485, 168–172.
- 94. Bakare, O.O.; Fadaka, A.O.; Akanbi, M.O.; Akinyede, K.A.; Klein, A.; Keyster, M. Evaluation of selected carotenoids of Lycopersicon esculentum variants as therapeutic targets for Alzheimer's disease: An in silico approach. BMC Mol. Cell Biol. 2021, 22, 49.
- Kumagai, M.H.; Keller, Y.; Bouvier, F.; Clary, D.; Camara, B. Functional integration of non-native carotenoids into chloro plasts by viral-derived expression of capsanthin-capsorubin synthase in Nicotiana benthamiana. Plant J. 1998, 14, 305 –315.
- 96. Piano, D.; Cocco, E.; Guadalupi, G.; Kalaji, H.M.; Kirkpatrick, J.; Farci, D. Characterization under quasi-native condition s of the capsanthin/capsorubin synthase from Capsicum annuum L. Plant Physiol. Biochem. 2019, 143, 165–175.
- Mialoundama, A.S.; Heintz, D.; Jadid, N.; Nkeng, P.; Rahier, A.; Deli, J.; Camara, B.; Bouvier, F. Characterization of pla nt carotenoid cyclases as members of the flavoprotein family functioning with no net redox change. Plant Physiol. 201 0, 153, 970–979.
- Hugueney, P.; Badillo, A.; Chen, H.C.; Klein, A.; Hirschberg, J.; Camara, B.; Kuntz, M. Metabolism of cyclic carotenoids: A model for the alteration of this biosynthetic pathway in Capsicum annuum chromoplasts. Plant J. 1995, 8, 417–424.

99. Furubayashi, M.; Kubo, A.; Takemura, M.; Otani, Y.; Maoka, T.; Terada, Y.; Yaoi, K.; Ohdan, K.; Misawa, N.; Mitani, Y. C apsanthin Production in Escherichia coli by Overexpression of Capsanthin/Capsorubin Synthase from Capsicum annuu m. J. Agric. Food Chem. 2021, 69, 5076–5085.

Retrieved from https://encyclopedia.pub/entry/history/show/101468