Antioxidant-Based Therapies in Male Infertility

Subjects: Others | Others Contributor: Giacomo Lazzarino

Under physiological conditions, reactive oxygen species (ROS) play pivotal roles in various processes of human spermatozoa. Indeed, semen requires the intervention of ROS to accomplish different stages of its maturation. However, ROS overproduction is a well-documented phenomenon occurring in the semen of infertile males, potentially causing permanent oxidative damages to a vast number of biological molecules (proteins, nucleic acids, polyunsaturated fatty acids of biological membrane lipids), negatively affecting the functionality and vitality of spermatozoa. ROS overproduction may concomitantly occur to the excess generation of reactive nitrogen species (RNS), leading to oxidative/nitrosative stress and frequently encountered in various human pathologies. Under different conditions of male infertility, very frequently accompanied by morpho-functional anomalies in the sperm analysis, several studies have provided evidence for clear biochemical signs of damages to biomolecules caused by oxidative/nitrosative stress. In the last decades, various studies aimed to verify whether antioxidant-based therapies may be beneficial to treat male infertility have been carried out.

Keywords: male infertility ; oxidative/nitrosative stress ; antioxidants ; seminal plasma ; spermatozoa

1. Introduction

Infertility is a far-reaching condition affecting 8–12% of reproductive-age couples worldwide ^[1]. It is defined as the inability to have conception after 12 months of unprotected sexual intercourse ^[2]. Historically, females have often been indicated as the sole ones responsible for pregnancy failures. During the last decades, thanks to the increasing number of epidemiological studies regarding infertility, it came forth into the light that the male factor is implicated in around 50% of cases of conceiving problems. Nowadays, among couples affected by this condition, the male factor is considered the unique cause in \approx 20% of all cases, together with a female factor co-contribution in the remaining 30% of cases ^{[3][4][5]}. The etiological causes of male infertility comprise a wide variety of factors, ranging from infections, anatomical and genetic abnormalities, neurological disease, environmental and lifestyle ^{[6][7][8]} and psychosocial factors, defining this condition as a multifactorial disease. In this context, it is worth recalling that even the COVID-19 pandemic has recently been included as a new external factor potentially negatively affecting couple fertility ^{[9][10]}. Indeed, since SARS-Cov-2 viral infection is responsible for systemic inflammatory processes, also leading to increased oxidative/nitrosative stress ^[11], it has been hypothesized that it may alter various fertility-related biochemical processes, thus resulting in decreased fertilization capacities of both males and females ^[12].

Male infertility is generally characterized by morpho-functional anomalies in the sperm analysis, including low sperm count, alteration of the sperm quality parameters (concentration, motility, morphology), or both. The main causes of male infertility are divided into: (1) extra-testicular (obstructive); (2) testicular (primary); (3) pretesticular (secondary); (4) idiopathic ^{[13][14]}. This last group, in whom the primary cause is not clearly manifest, accounts for 30–40% of infertile male patients ^[14].

2. Male infertility and Oxidative/Nitrosative Stress: A Rationale for Antioxidant-Based Therapies

One of the hallmarks of most of the aforementioned conditions affecting infertile males is the evidence of oxidative stress in their seminal fluid ^{[15][16][17][18]}. This condition is defined as the imbalance between the production of reactive oxygen species (ROS) and the protective action of the antioxidant systems responsible for their neutralization and removal.

Recent data have shown that disorders such as poor fertilization, pregnancy loss, birth defects and poor embryonic development are correlated with high susceptibility of spermatozoa to oxidative insult ^{[19][20]}. Since the excess production of ROS and RNS are able to induce irreversible modifications to biologically fundamental macromolecules of spermatozoa, such as lipids, proteins and nucleic acids, it has been found that oxidative/nitrosative stress provokes defects in sperm functions by decreasing their motility and fertilizing capacity. In particular, it has been demonstrated that

increased levels of ROS and RNS stimulate lipid peroxidation of polyunsaturated fatty acid-rich plasma membranes of spermatozoa ^{[21][22][23]} and DNA fragmentation ^{[24][25][26]}. Furthermore, as well as in other pathological conditions such as neurodegenerative diseases, ischemia and cancer ^{[27][28][29][30]}, the insurgence of oxidative/nitrosative stress in spermatozoa is associated with mitochondrial dysfunction. The damaging activities of ROS and RNS on proteins provoke permanent modifications of their structures through thiol oxidation, sulfonylation and tyrosine nitrosylation reactions. One of the targets of the harmful action of these toxic compounds is represented by altered expressions and catalytic activities of glycolytic enzymes, Krebs' cycle enzymes, protein complexes of the electron transfer chain (ETC) and oxidative phosphorylation (OXPHOS). Altogether, these changes of monosaccharide metabolism (glucose and fructose) of spermatozoa lead to reduced efficiency in ATP production, with a consequent energy crisis and mitochondrial dysfunction ^{[31][32]}. In this scenario, many studies, in recent years, focused on the correlation between oxidative/nitrosative stress and male infertility, identifying many promising biomarkers in the seminal fluid (derived by the damaging action of ROS and RNS overproduction on spermatozoa macromolecules) such as malondialdehyde (MDA), 4-hydroxynonenal (4-HNE), 8-hydroxy-2'-deoxyguanosine (8-OHdG), nitrites and nitrates (NO₂⁻ and NO₃⁻) ^{[33][34][35][36]}.

To counteract ROS and RNS overproduction and to prevent the insurgence of oxidative/nitrosative stress, cells of all oxygen-dependent organisms are provided with a pool of ROS and RNS scavenging enzymes (SOD, catalase, glutathione peroxidase, heme oxygenase, thioredoxin), representing the enzymatic category of the antioxidant defenses of the cells and capable to efficiently remove selected ROS ($O_2^{-\bullet}$, H_2O_2 , organic peroxides). To account for wider protection aimed to eliminate other dangerous species (OH $^{\bullet}$, ONOO⁻), cells contain various low-molecular-weight antioxidants having, as a common characteristic, remarkable reducing power ^[37]. Therefore, these compounds are capable of interacting nearly with any type of oxidant, including ROS and RNS. The gross division of low-molecular-weight antioxidants into hydrophilic (water-soluble) and hydrophobic (fat-soluble) confines them into water-rich (cytoplasm, mitochondrial matrix) and fat-rich (biological membranes) cell compartments, thus discriminating (and/or limiting) their ability to scavenge ROS and RNS. Figure 1 summarizes ROS and RNS damaging activities and the location of the main water- and fat-soluble low-molecular-weight antioxidants within cells.

Figure 1. Schematic representation of the main sources and damaging effects of reactive oxygen species (ROS) and reactive nitrogen species (RNS) overproduction. Mitochondria are both the main intracellular source and one of the main targets of ROS. Incorrect transport of electrons through the electron transport chain (ETC) exacerbates mitochondrial ROS generation, causes a decrease in oxidative phosphorylation (OXPHOS) efficiency with a consequent decrease in ATP production and energy crisis. Increased expression of the inducible form of nitric oxide synthase (iNOS) is responsible for increased nitric oxide production. Its decomposition and reaction with ROS generate further RNS. Sustained oxidative/nitrosative stress provokes a decrease in intracellular levels of ascorbic acid (AA) and reduced glutathione (GSH), as well as those of the membrane-bound α -tocopherol. The decrease in low-molecular-weight antioxidant defenses allows ROS and RNS to attack the most important biomolecules, causing: (i) the induction of the lipid peroxidation reaction chain by acting on the double bonds of the polyunsaturated fatty acids (PUFAs) of membrane phospholipids and leading to the formation of various decomposition products including malondialdehyde (MDA); (ii) the increase in protein misfolding and unfolding with loss of the protein biological activities, including the activities of glycolytic

enzymes that further imbalance mitochondrial functioning and energy metabolism; iii) the oxidative damage to DNA, causing its fragmentation and generation of DNA oxidative products such as 8-hydroxy-2'-deoxyguanosine (8-OHdG). Mitochondrial malfunctioning leads to loss of cytochrome c from the intermembrane space into the cytoplasm, acting as a signal for triggering apoptosis. Decreased morpho-functional parameters of spermatozoa, deeply affecting male fertility, are the ultimate consequences of ROS and RNS damaging activities.

2.1 ROS and RNS in Sperm Cells: the ROS Paradox and the Sperm Antioxidant Defenses

As for the other cell types of aerobic organisms, spermatozoa rely on oxidative phosphorylation (OXPHOS) as the major source of energy production in terms of ATP molecules. It has been demonstrated that OXPHOS mainly occurs in the sperm mid-piece, a mitochondrial rich district, differently from glycolysis that takes place in the head and in the fibrous sheath of the flagellum, where glycolytic enzymes are anchored ^[38]. Mitochondrial metabolism represents one of the major endogenous sources of ROS deriving from the activity of the electron transport chain (ETC) complexes. The electron flux through ETC and the concomitant transport of H⁺ ions in the intermembrane space of mitochondria performed by complexes I, III and IV are utilized to catalyze the ATP synthesis and the tetravalent reduction of O_2 to H_2O . Despite this process is extremely efficient and finely regulated, a considerable amount of superoxide and hydroxyl radicals are physiologically produced (representing about 1% of total oxygen consumed by the cell), in particular by complexes I and III ^{[39][40]}.

Another important endogenous source of ROS and RNS in semen is represented by immature spermatozoa deriving from defective stages of spermatogenesis. During the last phase of gamete differentiation (the spermiogenesis stage that occurs in the testis), spermatids undergo the maturation process, characterized by nucleus remodeling, chromatin condensation, acrosome and sperm tail formation and excess cytoplasm removal (the maturation stages occurring in the epididymis) ^[41]. Defects in this latter process give rise to immature spermatozoa, marked by retained cytoplasmic droplets in the mid-piece ^[42], containing glucose-6-phosphate dehydrogenase (G6PDH) enzyme. G6PDH is a cytoplasmic enzyme involved in the first reaction of the pentose phosphate pathway (PPP). It uses oxidized nicotinamide adenine dinucleotide phosphate (NADP⁺) to oxidize glucose-6-phosphate to 6-phosphogluconate, thus generating reduced nicotinamide adenine dinucleotide phosphate (NADP⁺) to oxidize glucose-6-phosphate to 6-phosphogluconate, thus generating reduced nicotinamide adenine dinucleotide phosphate (NADP⁺). Notwithstanding the maintenance of adequate levels of the reduced form of glutathione (GSH) needs high NADPH levels to ensure the correct activity of glutathione reductase, it has been demonstrated that high NADPH concentrations favor ROS production in immature spermatozoa through the activity of the Ca⁺⁺-dependent, membrane-bound isoform 5 of NADPH oxidase (NOX5) ^{[43][44][45]}. Additional endogenous ROS production is obtained, even if in modest contribution, from peroxisomal β -oxidation, the catabolic pathway of very long fatty acids and from the activity of the enzymes of the cytochrome P₄₅₀ family involved in xenobiotic detoxification ^{[40][46]}.

The above-depicted scenario highlights the evidence that a considerable amount of ROS and RNS are physiologically produced by the spermatozoa themselves, identifying these cells as "professional generators of ROS". Notwithstanding the aforementioned destructive effects that ROS and RNS can induce on sperm functions, many studies clearly demonstrated that controlled levels of ROS and RNS are necessary for spermatozoa to acquire their fertilization ability during the differentiation in the testis, the maturation in epididymis and the capacitation in the female reproductive tract. In particular, it has been shown that moderate H₂O₂ levels are required for correct spermatogenesis in the testis. Indeed, H₂O₂ regulates the stem cell population and the progenitor cells from which spermatozoa originate, the chromatin condensation in the nucleus and the mitochondria arrangement in the mid-piece of spermatozoa. This type of regulation is mainly performed by promoting the subsequent phosphorylation of different tyrosine and serine amino acidic residues of various target proteins [47][48], and it is also required for the maturation process occurring in the epididymis, in which spermatozoa undergo DNA compaction and acquisition of motility ^[49]. Furthermore, much evidence has demonstrated that moderate levels of ROS and RNS are necessary for the correct development of capacitation, again performed through the redox-modulated tyrosine phosphorylation of target proteins. In particular, H_2O_2 , $\bullet NO$, and $O_2^{-\bullet}$ levels regulate the phosphorylation of two tyrosine-rich proteins of 81 and 105 kDa, which are selectively overexpressed during capacitation. The post-translational modification of these proteins, occurring through the concomitant activation of tyrosine kinase and protein kinase C and inactivation of phosphatases, is fundamental to promote the subsequent oxidation of specific cysteine residues necessary for their ultimate activation ^[50]. Moreover, ROS, in conjunction with the intracellular Ca⁺⁺ entry via the cation channels of sperm CatSper, are able to stimulate the increase in intracellular cyclic AMP (cAMP) levels that leads to the activation of protein kinase A (PKA), indirectly increasing the tyrosine phosphorylation of proteins $^{(50)}$. This latter mechanism, again regulated by H₂O₂, •NO, and O₂^{-•}, is also extended to proteins located in the flagellar fibrous sheath involved in hyperactivation (i.e., the acquisition by spermatozoa of nonlinear motility) and to proteins involved in acrosome reaction and sperm-oocyte fusion [51][52]. In order to acquire the ability to perform the acrosome reaction and fusion with the oocyte, spermatozoa need to undergo modifications of their plasma membrane composition. In particular, it has been shown that ROS are necessary to stimulate cholesterol oxidation and its consequent removal

from the plasma membrane. This process is accompanied by the activation of phospholipase A2 leading to an increase in polyunsaturated fatty acids cleavage, ultimately causing an increase in plasma membrane fluidity ^{[53][54]}. Figure 2 summarizes ROS and RNS activities that are considered fundamental for the correct development and function of spermatozoa.



Figure 2. Schematic representation of the involvement of ROS and RNS to ensure the correct development and function

of spermatozoa. In functional mitochondria, the correct coupling between ETC and OXPHOS ensures an adequate ATP supply for cell energy demand, as well as a "physiologic" flux of ROS (mainly in the form of superoxide anion). At the same time, the endothelial (and possibly the neuronal) nitric oxide synthase isoform (eNOS) provides "physiologic" production of RNS (in the form of nitric oxide). These controlled amounts of ROS and RNS positively affects numerous intracellular processes of spermatozoa, including; (i) influx of calcium ions (Ca⁺⁺) triggering the cyclic AMP (cAMP) cascade (increase in adenylate cyclase activity) leading to activation of protein kinase A (PKA) which catalyzes the phosphorylation of numerous target proteins; (ii) the concomitant activation of protein kinase C (PKC), producing a further increase in protein phosphorylation; (iii) regulation of protein activities through nitrosylation reactions; (iiii) oxidation of cholesterol and removal from the plasma membrane. Some of these processes are deeply involved during spermatogenesis, maturation and capacitation, as well as in acrosome reaction and fusion with the oocyte. Due to these functions in their life cycle, spermatozoa may be defined as "professional generators" of ROS and RNS. During these ROS and RNS-mediated processes, no significant depletion in the antioxidant heritage of spermatozoa occurs.

In consideration of the abovementioned fundamental beneficial role of ROS and RNS in the differentiation and the acquisition of the fertilization ability by spermatozoa on one side, but on the other hand of their harmful role in the insurgence of oxidative/nitrosative stress, seminal plasma has evolved as one of the most powerful antioxidant-rich fluids in human beings, in order to finely control the amount of ROS and RNS. Despite spermatozoa are equipped with endogenous antioxidant defenses against oxidative/nitrosative stress, for obvious structural reasons, these cells are highly dependent on the powerful antioxidant properties of the fluids that surround them, i.e., the seminal plasma released from accessory glands, following spermatozoa release from the germinal epithelium. The antioxidant profile, characteristic of this body fluid, comprehends both enzymatic and nonenzymatic antioxidants that are able to detoxify ROS and RNS, avoiding the possible insurgence of oxidative/nitrosative stress. The enzymatic scavengers include catalase, superoxide dismutase (SOD), glutathione peroxidase (GPx) and glutathione-S-transferase (GST) [55][56][57]. The nonenzymatic antioxidant profile of seminal plasma is represented by a heterogeneous group of both endogenous and exogenous lowmolecular-weight scavengers divided into water-soluble (such as ascorbic acid, GSH, uric acid, polyphenols) and fatsoluble compounds (such as α-tocopherol, y-tocopherol, all-trans-retinoic acid, all-trans-retinol, carotenoids and coenzyme Q₁₀) [58][59][60]. Giving the importance of maintaining adequate levels of each component of this set of antioxidants to ensure the wellness of spermatozoa and, therefore, their intrinsic fertilizing capacity, it is a very common clinical practice to indiscriminately prescribe to infertile males repeated cycles of administration of various types of nutraceutical/drug-containing antioxidant to improve their fertilization capacity. This therapeutic approach is generally

adopted without previously screened patients to determine the concentration of antioxidants in seminal plasma, i.e., antioxidant treatment is very often administered without knowing its real need caused by a deficiency in one or more of the low-molecular-weight antioxidants.

With the aim to understand the actual effectiveness of antioxidant-based therapies in the treatment of male infertility, in this review, we analyzed data from the literature concerning the use of low-molecular-weight antioxidants to ameliorate the sperm quality and the spermatozoa fertilization ability of patients affected by different conditions of infertility. We limited the evaluation of the literature (using PubMed as the data bank for consultation, using all the different mix of the search words, antioxidants, male infertility, low molecular-weight antioxidants, humans, sperm, spermatozoa) by considering only clinical studies performed during the last ten years and focusing on those in which compounds having a direct and well-established antioxidant activity towards ROS and/or RNS were used.

3. Antioxidant-Based Therapies in the Treatment of Male Infertility

3.1. Ascorbic Acid (Vitamin C)

Ascorbic acid (AA), also known as vitamin C, is one of the most abundant water-soluble antioxidants within mammalian tissues, acting as a reducing cofactor in several enzymatic reactions [61]. AA is an essential cellular antioxidant also in conditions of abundance of GSH, protecting tissues, organs and cells from the harmful effects of ROS and RNS overproduction during oxidative/nitrosative stress [62]. The standard redox potential of the couple oxidized/reduced ascorbate (+0.08 V at pH 7.0) qualifies AA as a fairly good reducing agent, capable to quickly react with a wide range of ROS and RNS, including superoxide, peroxynitrite and hydroxyl radicals, through one or two electrons reactions generating, respectively, ascorbyl radical or dehydroascorbic acid (DHA). DHA can be reconverted back to the reduced AA form in a direct chemical reaction involving glutathione (GSH), even though NADPH-dependent enzymatic reactions involving glutaredoxin, thioredoxin and GSH reductases have also been demonstrated. The antioxidant protection performed by AA is also related to the recycling of oxidized α -tocopherol (vitamin E), in particular, contributing to the maintenance of the α-tocopherol redox state within biological membranes. Since AA is not synthesized by most mammals (including human beings), the human body depends on vitamin C-rich food ingestion to satisfy its AA need ^[63], with a recommended daily intake (RDI) of 65 to 90 mg/day. The inability to synthesize AA is due to the loss of function of the biosynthetic enzyme L-gulono-1,4-lactone oxidase [64]. Two distinct transporter families are involved in the cellular absorption of AA, depending on the oxidative state of the molecule. Indeed, facilitate DHA internalization is performed through the hexose transporters (GLUT1, GLUT3 and GLUT4). Differently, AA is transported by high-affinity, low-capacity active sodium-ascorbate co-transporters (SVCT1 and SVCT2) [65]. Notwithstanding that circulating levels of AA are relatively low (30 to 70 µmol/L in the serum of healthy humans) [58], concentrations of this important antioxidant in peripheral tissues are, interestingly, elevated. Particularly in the brain, AA is highly concentrated, usually reaching concentrations up to 3000 µmol/L brain water [66]. In the testis, AA levels hover around 3 mg/100 g wet tissue (about 140 µmol/L tissue water) [67]. Expression of SCVT2 and some isoforms of the GLUT transporters have been found in Sertoli cells that form the blood-testis barrier via tight and adherens junctions. These cells are able to regulate AA concentrations in the seminiferous epithelium, underlying the key role of AA in the spermatogenesis [68]. In seminal plasma, concentrations of AA have been found up to 10 times higher than those measured in plasma, with values ranging from 200 to 350 µmol/L. Explanations of the mechanisms allowing to reach such abundant vitamin C levels in this peculiar body fluid are still unclear [58][69]. In general, since AA concentrations are higher in peripheral tissues than in plasma, the cellular internalization of AA via SVCT1 and SVCT2 occurs against a concentration gradient; the energy to overcome the electrochemical barrier is provided by the favorable inward gradient for Na⁺, which is fostered by the regular extrusion of Na⁺ via the Na⁺/K⁺-ATPase pump. The electrogenic transport is exploited by SVCT1 only, since the SVCT2 transport mechanism also involves Ca^{2+} and Mg^{2+} . As mentioned above, AA is the most abundant water-soluble low-molecularweight antioxidants in the seminal plasma [33][58][69], playing significant protection against oxidative damage to DNA [70] and prevention of semen agglutination [71]. Furthermore, AA levels in seminal plasma positively correlate not only with normal semen parameters, such as sperm volume, sperm count and sperm motility ^[72] but also with GSH concentration and SOD activity, suggesting a close relationship between different antioxidant defensive mechanisms against ROSmediated damages [73][74]. Under pathological conditions leading to male infertility, in which the insurgence of oxidative stress is involved, it was observed an overall depletion of AA in the seminal fluid due to the sustained presence of ROS and RNS. In particular, in patients affected by varicocele of various grade, by genitourinary infections and by idiopathic infertility, a net decrease in AA concentration in the seminal fluid has been demonstrated [33][73][75][76].

Several studies focused on the association between AA dietary intake and semen quality. In a cross-sectional study conducted in 210 young, healthy subjects, ranging between 18 and 23 years of age, Mínguez-Alarcón L et al. ^[77] reported a positive correlation between the dietary intake of several antioxidant nutrients (β -cryptoxanthin, vitamin C, lycopene and

 β -carotene) and the total motile sperm count, underlying that an AA intake satisfying the RDI was positively associated with increased semen volume. In a case–control study conducted on 31 normozoospermic subjects and 30 infertile males affected by moderate or severe oligozoospermia and severe teratozoospermia, it has been shown that low intake of antioxidant nutrients seems to worsen semen quality while the regular ingestion of vitamin C-reach food improves the morpho-functional characteristics of the semen ^[78]. Furthermore, in a study realized on 107 asthenozoospermic patients and 235 age-matched controls, Eslamian et al. reported a significant negative relationship between a dietary pattern of antioxidant-rich foods (including vitamin E, vitamin D, vitamin C, zinc, folate, total fiber, selenium and PUFAs) and risks of asthenozoospermia in a cohort of Iranian men ^[79].

The majority of the clinical studies aimed to evaluate the effect of vitamin C supplementation on male infertility were carried out by applying the simultaneous administration of various antioxidants, including AA. The administration of AA only was conducted in three clinical trials, in which different semen parameters and fertilization capacity of infertile males were evaluated. When examining the effect of AA supplementation on parameters of the sperm analysis, results reported in the available studies are contradictory. In a controlled clinical trial, 115 infertile males affected by varicocele and subjected to varicocelectomy were divided into two groups receiving, twice a day, either 250 mg of AA (AA group) or placebo (control group) for the next three months after surgery. The AA group showed no increase in sperm count and significant amelioration of sperm motility and morphology, suggesting that AA administration positively affected only quality but not the number of spermatozoa [80]. Differently, Rafiee et al. [81] reported that the administration of a single dose of 1000 mg/day to 200 subjects with increasing body mass index and ages ranging from 20 to 60 year was able to improve the sperm concentration and motility, but not the semen volume and the percentage of normal spermatozoa morphology. In a cross-sectional study carried out in 120 male workers, occupationally exposed to lead and to show testicular dysfunction and high rate of DNA fragmentation, it was found that the administration of 1000 mg/day of AA for five consecutive days/week for up to 3 months, statistically increased sperm motility, sperm total count and significantly decreased the number of morphologically abnormal spermatozoa [82]. These results prompted the authors to suggest that AA treatment could offer protection against sperm damage caused by a lead-mediated increase of oxidative stress [82].

As previously said, in most studies, exogenous AA was administered in combination with other antioxidants (such as vitamin E, carotenoids and coenzyme Q₁₀), thus rendering it difficult to extrapolate the potential vitamin C beneficial effects on male patients affected by infertility. In a study conducted on 20 infertile males with known asthenoteratozoospermia, the combined daily supplementation with AA (60 mg), L-carnitine (1500 mg), coenzyme Q_{10} (20 mg), vitamin E (10 mg), vitamin B₉ (200 µg), vitamin B₁₂ (1 µg), zinc (10 mg) and selenium (50 µg) was able to induce, after 3 months of the administration, a statistical improvement of all semen parameters and DNA integrity. Although a remarkable variability in the inter-patient response to treatment was also shown, the authors reported a pregnancy rate of 5% [83]. Interestingly, the same administration protocol was applied to 20 males affected by grade I varicocele resulting in a significant reduction of DNA fragmentation and highly degraded sperm cells, a significant increase in total sperm cells, but no effects on other semen parameters [84]. In 210 infertile males affected by severe oligoasthenoteratozoospermia, Magdi et al. [85] tested the effects of the daily administration of AA (1 g), vitamin E (400 mg) and L-carnitine (2 g) as a possible management protocol of clinical utility. After six months of treatment, a significant increase in sperm concentration, total sperm count, percentage of spermatozoa with total motility and progressive motility, as well as a significant reduction of spermatozoa with abnormal morphology, was observed [85]. It should, however, be underlined that the real effects of this antioxidant protocol were confounded by the change in patient lifestyle adopted in the study protocol and aimed at minimizing adverse lifestyle factors [85]. Differently, the daily supplementation with 80 mg of AA, 120 mg of coenzyme Q_{10} and 40 mg of vitamin E was reported to be ineffective, after 3 and 6 months of treatment, in changing both atypical sperm cell number and semen volume, but capable of inducing significant improvements in sperm cell concentration and sperm motility in a cohort of 169 males affected by idiopathic oligoasthenoteratozoospermia [86]. Worthy of note is that after six months, a total of 48 pregnancies, comprising 16 spontaneous pregnancies (i.e., pregnancies occurring with no need of any type of ART) and 32 pregnancies requiring ART, were achieved by this group of patients using this treatment protocol [86]. In a study conducted on 15 infertile patients with oligozoospermia and/or asthenozoospermia, Terai et al. [87] reported that the combined administration of 1000 mg of AA, 750.1 mg of L-carnitine, 30 mg of zinc, 16.05 mg of astaxanthin, 90.26 mg of coenzyme Q_{10} , 60.1 μ g of vitamin B_{12} and 150 mg of vitamin E three times a day for 12 weeks, significantly improved only total motile sperm count, with no effects on the other semen parameters. In a single-blinded clinical trial, Sadaghiani et al. [88] reported that the daily administration for three consecutive months of 100 mg of AA, 30 mg of coenzyme Q10, 8 mg of zinc, 12 mg of vitamin E and 400 µg of folic acid, combined with that of 200 mg of selenium every other day, to 50 infertile oligozoospermic and asthenozoospermic male smokers, beneficially affected various morpho-functional semen parameters (concentration, pH, volume, total motility, morphology, count and progressive motility). A substantially different scenario was depicted by Steiner et al. [89], who conducted a randomized controlled trial on 174 infertile males (categorized into oligozoospermic, asthenozoospermic, teratozoospermic and DNA fragmented patients), analyzed for sperm analysis anomalies and clinical outcome measures

of fertility (pregnancy and live birth rates). In this study, infertile males were randomly assigned to a group receiving placebo and a group receiving the daily administration of an antioxidant formulation composed of 500 mg of AA, 400 mg of vitamin E, 0.20 mg of selenium, 1000 mg of L-carnitine, 20 mg of zinc, 1000 mg of folic acid, 10 mg of lycopene, and 2000 IU of vitamin D, for at least 3 months and up to 6 months. The authors reported that, compared to the placebo group, this antioxidant formulation and administration protocol did not result in any statistically significant improvement of the sperm analysis. Even more important, these results suggested that antioxidant treatment is not able to improve in vivo pregnancy or live-birth rates of couples with male factor infertility. A summary of the clinical studies concerning the effects of AA on male infertility is provided in Table 1.

Table 1. Summary of the main results obtained in the 10 papers focusing on the effect of AA administration for the treatment of male infertility.

Authors	Patients (Number and Type of Infertility)	Controls (Number)	Antioxidant(s) (Type(s), Doses and Administration Protocol)	Duration of Treatment (Months)	Improvement (Sperm Parameters, Pregnancy and Live Birth Rates)	No Improvement (Sperm Parameters, Pregnancy and Live Birth Rates)
Cyrus A. et al. ^[80]	46 = varicocele	69	AA 500 mg/day	3	Sperm motility and morphology	Sperm count
Rafiee B. et al. ^[81]	200 = increasing Body mass index	50	AA 1000 mg/day	6	Sperm concentration and motility	
Vani K. et al. ^[82]	120 = testicular dysfunction high rate of DNA fragmentation	No controls	AA 1000 mg/day for 5 day/week	3	Sperm motility, sperm total count and significantly decreased the number of morphologically abnormal spermatozoa	
Abad C. et al. ^[83]	20 = AT	No controls	AA (60 mg), L- carnitine (1500 mg), CoQ ₁₀ (20 mg), vitamin E (10 mg), vitamin B ₉ (200 μ g), vitamin B ₁₂ (1 μ g), zinc (10 mg) and selenium (50 μ g)/day	3	Sperm concentration, motility, vitality and morphology parameters, DNA integrity and 5% pregnancy rate	

Gual-Frau J. et al. ^[84]	20 = grade I varicocele	No controls	AA (60 mg), L- carnitine (1500 mg), CoQ ₁₀ (20 mg), vitamin E (10 mg), vitamin B ₉ (200 μ g), vitamin B ₁₂ (1 μ g), zinc (10 mg) and selenium (50 μ g)/day	3	Sperm count and DNA integrity	Concentration, progressive motility, vitality, and morphology
Magdi Y. et al. ^[85]	120 = severe OAT	No controls	AA (1 g), vitamin E (400 mg) and L-carnitine (2 g)/day	6	Sperm concentration, percentage of total motility and progressive motility	
Kobori Y. et al. ^[86]	169 = idiopathic OAT	No controls	AA (60 mg), CoQ ₁₀ (120 mg) and vitamin E (40 mg)/day	3 and 6	Sperm concentration and motility	Atypical sperm cell number and semen volume
Terai K. et al. ^[87]	15 = O and/or A	16 = administered with a Chinese herbal medicine	AA (1000 mg), L- carnitine (750.1 mg), zinc (30 mg), astaxanthin (16.05 mg), Co Q_{10} (90.26 mg), vitamin B ₁₂ (60.1 µg) and of vitamin E (150 mg) 3 times/day	3	Total motile sperm count	Semen volume, sperm concentration and sperm motility
Sadaghiani S. et al. ^[88]	50 = infertile O and A	No controls	AA (100 mg), CoQ ₁₀ (30 mg), zinc (8 mg), vitamin E (12 mg) and folic acid (400 μg)/day selenium (200 mg)/other day	3	Volume, morphology, motion, count, progressive motility and pH	

IT4 = O, A, T mg), selenium Sperm Steiner AZ. 174 = O, A, T carnitine (1000 motility, DNA Et al. [89] and DNA- fragmented patients 86 mg), zinc (20 3 and 6 integrity, no Mg), folic acid mg), folic acid different gregnancy active It000 mg), It000 mg), It000 mg), It000 mg),	Å and æs
lycopene (10 live-birth rat mg), and vitamin	es
D (2000 IU)/day	

AA = ascorbic acid; asthenozoospermia = A; asthenoteratozoospermia = AT; oligoasthenoteratozoospermia = OAT; oligoasthenozoospermia = OA; oligozoospermic = O; teratozoospermia = T.

3.2. N-Acetylcysteine (NAC)

GSH is the other most abundant hydrophilic low-molecular-weight antioxidant that, differently from AA, mammals are able to synthesize [90]. It is a tripeptide, y-L-glutamyl-L-cysteinyl glycine, present in concentrations ranging between 1 and 10 mM in animal tissues. GSH is synthesized via a two-step ATP-requiring enzymatic reactions; the first step is catalyzed by glutamate-cysteine ligase (GCL) that conjugates cysteine with glutamate, producing y-glutamylcysteine. The second step is catalyzed by GSH synthase, which adds glycine to y-glutamylcysteine producing the final product, GSH. This hydrophilic compound exists in the thiol-reduced (GSH) and disulfide-oxidized (GSSG) forms [91] GSH is the most abundant of the two species in eukaryotic cells (the GSH/GSSG ratio is about 100), mainly localized in cytoplasm and mitochondria, and having different concentrations in the various tissues. Notwithstanding that GSH is involved in the modulation of cell proliferation, apoptosis, inflammatory processes, redox signaling and detoxification of xenobiotics, probably the main function of this hydrophilic compound is to act as an antioxidant [90]. In particular, thanks to its sulfhydryl group (-SH) that confers active protection against ROS, GSH is involved in a redox cycle for the detoxification of hydrogen peroxide and lipid peroxides. These potentially dangerous compounds are reduced through the activity of the two selenium-dependent and selenium-independent forms of the enzyme GPx, in a reaction that oxidizes two GSH molecules into one GSSG molecule [92]. The regeneration of GSH is accomplished by glutathione reductase (GR), responsible for the maintenance of high values of the GSH/GSSG ratio, in the presence of reduced NADPH. Hence, GSH homeostasis is strictly connected to the PPP and the malic enzyme activity, i.e., the two main cellular sites for NADPH generation ^[93]. In addition to the involvement in the aforementioned redox cycle, GSH also acts as the cofactor of the enzyme family of GSTs. These protective enzymes are a large group of cytosolic and membrane-bound isoenzymes involved in the detoxification of xenobiotics (potentially toxic compounds often causing an increase in oxidative/nitrosative stress), obtained through the conjugation of GSH to xenobiotic substrates for their faster extrusion from the cell. In sperm cells, GSTs are involved not only in preventing lipid peroxidation in conditions of high levels of ROS and RNS but also in cell signaling regulation during spermatogenesis, sperm capacitation and fertilization [94].

Since the -SH group of GSH is supplied by cysteine, high levels of this amino acid are essential for GSH biosynthesis, therefore ensuring adequate GSH tissue levels. Cysteine normally derives from the diet, from protein breakdown and from the trans-sulfuration of methionine. Due to the fundamental role of GSH in detoxifying reactions, both to scavenge ROS and RNS and to eliminate xenobiotics, increased cysteine availability is needed under conditions of oxidative/nitrosative stress. As in the case of AA, GSH levels in human tissues are remarkably higher (2500-4000 µmol/L tissue water) than its circulating concentrations ranging between 15 and 20 µmol/L [58][95]. Seminal plasma values of GSH have been found equal or higher to those present in serum, indicating a free distribution of this molecule regulated on its relative concentrations in the two fluids [58][96]. Interestingly, in pathological conditions related to male infertility, characterized by sustained oxidative/nitrosative stress, data on variations of GSH levels are somehow controversial. Indeed, in viable spermatozoa of infertile patients, Moretti et al. [97] showed increased intracellular levels of total GSH (GSH + GSSG) with respect to values detected in spermatozoa of control fertile males. The possible explanation suggested by the authors is that the higher levels of ROS and RNS in spermatozoa of the infertile group increased the demand for antioxidants to counteract oxidative/nitrosative stress, thus leading to increasing values of GSH + GSSG. This hypothesized mechanism might have been accomplished by an augmented transport of GSH into the cell through membrane transporters located on the sperm surface ^[97]. These results were corroborated by those reported in a study by Kralikova et al. ^[98], who demonstrated significantly higher concentrations of homocysteine, cysteine, cysteinyl-glycine and GSH in spermatozoa of

patients with pathological sperm parameters compared to those found in normozoospermic fertile subjects. The authors reported the same aforementioned explanation for this phenomenon, i.e., the overproduction of ROS and RNS could cause an upregulation of thiol synthesis in order to protect spermatozoa against oxidative damages ^[98].

As said, diametrically opposite results reported an overall depletion of GSH in seminal plasma of infertile males, caused by increased oxidative/nitrosative stress. In a study conducted on 74 infertile patients affected by varicocele, genitourinary infections or idiopathic infertility, Micheli et al. ^[99] reported a significant concomitant decrease of GSH and AA only in seminal plasma of the varicocele group compared to the control group of fertile males. The non-significant reduction of GSH in the two other conditions of infertility was explained by the authors by the sparing effect induced by AA. Compared to a control group of 17 fertile males, Moretti et al. reported a net decrease of the GSH/GSSG ratio, in a cohort of 36 infertile patients affected by leuokocytospermia or varicocele, independently on the pathology causing infertility, even though the most striking differences were recorded in the varicocele group [100].

The aforementioned results suggest that drugs aimed to increase seminal plasma/spermatozoa levels of GSH may be successful for the treatment of male infertility. Since the oral administration of GSH is useless because the molecule is degraded before its absorption, pharmacological strategies are aimed to stimulate GSH biosynthesis to obtain a net increase of intracellular GSH concentrations. In this light, to improve the availability of cysteine, ultimately enhancing GSH synthesis, N-acetylcysteine (NAC) has widely been used as a potential treatment for male infertility (administered alone or in combination with other antioxidants), also thanks to its relevant properties to act as a direct ROS and RNS scavenger. In a randomized, blinded clinical trial performed by Jannatifar et al. [101], 50 infertile males with asthenoteratozoospermia received NAC orally for 3 months (600 mg/day), and their semen parameters and DNA fragmentation index were compared with those determined before the beginning of NAC treatment. The authors reported that the three months supplementation with NAC significantly improved sperm analysis parameters (count, motility and normal morphology) and significantly reduced DNA fragmentation, indicating that NAC supplementation is able to protect morpho-functional spermatozoa characteristics and to reduce ROS and RNS-mediated damages to DNA. The protective antioxidant effect of NAC against DNA damages was also reported by Barekat et al. [102], who conducted a study on 35 infertile men with varicocele. Following microsurgery, patients were randomly divided into control or treatment groups, with the former receiving no drug after varicocelectomy (n = 20) and the latter (n = 15) receiving the daily administration of NAC (200 mg/day) post-varicocelectomy for three months. In addition to the beneficial effects due to the microsurgery, the authors showed that, while control patients had a pregnancy rate of 10%, this value in patients receiving NAC supplementation increased significantly, up to 33.4%. According to these data, the authors suggested that administration of NAC following varicocelectomy may represent an added value not only for the improvement of sperm motility, a decrease of oxidative/nitrosative stress, recovery of GSH levels, protection of the DNA integrity, but also for the overall improvement of patient's fertilization capacity [102].

The potential benefit of NAC administration to infertile males was also reported when patients received NAC in combination with other compounds (mainly antioxidants). Dattilo et al. [103] conducted an uncontrolled study on 84 male partners of couples with at least 2 previous in vitro fertilization (IVF) failures, with no evidence of organic causes of infertility and with either measurable DNA fragmentation index (DFI) or nuclear decondensation index (SDI). Patients were administered with a single daily dose (for an average period of 130 days) of a tablet containing a proprietary extract of opuntia fig fruits (100 mg) delivering tailored amounts of quercetin (0.05 mg) and betalain (0.001 mg), plus a mix of vitamins B₂ (1.4 mg), B₃ (16 mg), B₆ (1.4 mg), B₉ (400 µg), B₁₂ (2.5 µg), zinc (12.5 mg), NAC (250 mg) and vitamin E (12 mg). The aim of the authors was to facilitate the endogenous production of GSH by increasing the rate of the so-called one-carbon cycle, a pathway that involves the trans-sulfuration of homocysteine favoring the de novo GSH biosynthesis. The authors reported a significant decrease of both DFI and SDI, but no modifications of the total sperm count, fast motility and normal morphology rates after the administration protocol. Notwithstanding, high clinical pregnancy rates, with 18 live births from 18 spontaneous pregnancy (i.e., pregnancy occurring with no need of any type of ART) and 15 live births from 22 initial clinical ART pregnancies (taking place at the second IVF attempt), were reported to occur in this cohort of patients [103]. Conversely, in a study conducted on 20 patients affected by β -thalassemia major, a condition characterized by hypogonadotropic hypogonadism mediated by iron deposition and generation of ROS with a consequent increase in damaged spermatozoa, a combined treatment with L-carnitine and NAC was ineffective in the improvement of fertility [104]. In this regard, the study showed that after six months of treatment (2 g/day of L-carnitine and 600 mg/day of NAC), patients had a significant increase in the SDI and teratozoospermia index. Apparently, the treatment accentuated sperm deformities, leading to the conclusion that the use of antioxidants in β-thalassemia major patients is not justified for the potential improvement of fertility [104]. A summary of the clinical studies concerning the effects of AA on male infertility is provided in Table 2.

No Improvement Antioxidant(s) Improvement Duration Patients (Sperm Controls of (Type(s), Doses (Sperm Parameters, Authors Treatment (Number and and Parameters, Pregnancy and (Number) Type of Infertility) Administration Pregnancy (Months) Live Birth Protocol) and Live Birth Rates) Rates) Sperm count, motility and Jannatifar normal NAC (600 No 50 = AT 3 R. et al. morphology, DNA controls mg)/day [<u>101</u>] integrity NAC (200 Sperm motility Barekat mg)/day postand 33.4% 15 = varicocele 20 3 F. et al. varicocelectomy pregnancy rate [<u>102</u>] NAC (250 mg), extract of opuntia fig fruits (100 mg), quercetin DNA integrity, (0.05 mg), spontaneous Total sperm betalain (0.001 pregnancy rate count, fast mg), vitamins B₂ (21%), Dattilo M. No motility and (1.4 mg), B₃ (16 84 = ART failures 4 pregnancy rate et al. [103] controls normal mg), B₆ (1.4 mg), (47.6%), and live morphology B₉ (400 µg), B₁₂ birth rate (39.3%) rates (2.5 µg), zinc after a new ART (12.5 mg), and attempt. vitamin E (12 mg)/day NAC (600 20 = BTM Increased Elsedfy mg)and L-No (hypogonadotropic 6 sperm H. et al. controls carnitine (2 [104] hypogonadism) deformities g)/day

Table 2. Summary of the main results obtained in the 4 papers focusing on the effect of *N*-acetylcysteine (NAC) administration for the treatment of male infertility.

NAC = *N*-acetylcysteine; asthenozoospermia = A; asthenoteratozoospermia = AT; oligoasthenoteratozoospermia = OAT; oligoasthenozoospermia = OA; oligozoospermic = O; teratozoospermia = T.

3.3. Tocopherols (Vitamin E)

The term vitamin E covers eight different substances, four tocopherols (α , β , γ , δ) and four tocotrienols (α , β , γ , δ). All these fat-soluble compounds contain a chroman ring with different numbers and positions of methyl groups and a side-chain linked to position 2 of the aforementioned ring. Tocopherols contain a saturated phytyl chain, while in tocotrienols, the tail contains three isolated double bonds ^[105]. Due to the presence of the hydrophobic side-chain, tocopherols are

mainly localized within the phospholipid bilayer of biological membranes, acting as the major lipid-soluble antioxidant that is able to block the lipid peroxidation reaction chain. This antioxidant action is exerted by the -OH group of the chromane ring that, under the condition of oxidative/nitrosative stress, donates hydrogen radical to a lipid radical or a lipid hydroperoxide radical, resulting in the termination of the lipid peroxidation reaction chain and in the formation of a relatively stable tocopheryl- or tocotrienyl-radical. These two partially oxidized forms of vitamin E will most likely depend on AA availability in the aqueous cellular compartments for their regenerations. In this oxidoreductive reaction, AA is oxidized to ascorbyl radical, subsequently undergoing either dismutation (two ascorbyl radicals), with the formation of one AA and one DHA, or to further one-electron oxidation with the generation of DHA. Notwithstanding α , β , γ and δ tocopherols have progressively decreasing ROS scavenging power [106], it has been demonstrated that y-tocopherol is the most effective tocopherol against RNS, with particular scavenging activity towards nitrogen dioxide with which it forms the stable nitro-adduct 5-nitro-y-tocopherol [107]. Among the eight congeners, α -tocopherol is the most abundant in the European diet and y-tocopherol in the American diet. In general, the RDI for vitamin E is around 7–15 mg/day ^[108], and in adults, severe vitamin E deficiency mainly occurs as a result of genetic defects in the α -tocopherol transfer protein (α -TTP), caused by an infrequent genetic disorder known as "isolated vitamin E deficiency" or 'ataxia with isolated vitamin E deficiency" (AVED) ^[109]. The most abundant tocopherol in human tissues and in blood plasma is α -tocopherol, the circulating levels of which range between 18 and 30 µmol/L [50]. The distribution to peripheral organs and tissues is achieved by the α -TTP that incorporates α -tocopherol into lipoproteins, and it is able to distinguish α -tocopherol from all incoming forms of vitamin $E^{[110]}$.

Due to their hydrophobicity and the lack of effective components (such as lipoproteins) that could facilitate their solubility, concentrations of tocopherols in seminal plasma, unlike AA, are about nine times lower (α -tocopherol ~3 µmol/L and γ -tocopherol ~0.06 µmol/L) than those detected in blood plasma ^[58], suggesting the lack of efficient carrier systems capable of accumulating these compounds in seminal plasma. Notwithstanding the low concentrations of seminal plasma tocopherols, the high levels of AA in this body fluid ensure the correct recycling of vitamin E and exploit its beneficial effect against ROS and RNS-induced lipid peroxidation. Because of their high content of polyunsaturated fatty acids, membranes of germ cells and spermatozoa are very sensitive to oxidation from high levels of ROS and RNS. Therefore, vitamin E represents for these cells one of the most effective fat-soluble antioxidants against oxidative/nitrosative stress-related damages ^[111].

In a case report of a patient affected by AVED, Rossato M. and Mariotti C. [109] showed normal spermatogenesis and semen parameters notwithstanding vitamin E deficiency, challenging the effective role of tocopherols in human spermatogenesis. On the other hand, several studies showed the correlation of decreased levels of tocopherols in human seminal plasma with augmented concentrations of ROS and RNS-mediated marker of lipid peroxidation (MDA) in infertile males [33][107][112][113]. In addition to the aforementioned study [71], the impact of dietary vitamin E intake was evaluated by Nadjarzadeh et al. [114] in a case control study conducted on 32 men with primary infertility due to idiopathic oligo and/or astheno- and/or teratozoospermia and 32 age-matched normal healthy donors. The authors highlighted the association between a lower intake of vitamin E, zinc and folate and a negative effect on sperm motility and morphology. Schmid et al. [115], in a cross-sectional study performed on 80 healthy fertile male volunteers (aged 22-80 years), showed that a higher dietary intake of vitamin E, AA, folate, and zinc positively correlated with lower damages to sperm DNA. Furthermore, the authors reported that older men who consumed higher levels of the aforementioned micronutrients had similar levels of sperm DNA damage than those found in young men, underlying the age-independent importance of the intake of adequate levels of antioxidants to counteract oxidative-mediated damages to sperm DNA [115]. Similarly, Salas-Huetos et al. [116] conducted a study in healthy individuals to evaluate the effect of chronic consumption of foods rich in monounsaturated fatty acids, polyunsaturated fatty acids (including α -linolenic acid, and total ω -3), magnesium and vitamin E on semen parameters related to male fertility. To this end, 50% of the volunteers fed for 14 weeks the usual Western-style diet enriched with 60 g of a mixture of nuts/day (30 g of walnuts, 15 g of almonds, and 15 g of hazelnuts) and the remaining 50% fed the usual Western-style diet avoiding nuts. The authors reported a significant improvement of the morpho-functional parameters of the sperm analysis (total sperm count, vitality, motility, and morphology) in subjects who fed a high nut diet, suggesting that these benefits may be due to a reduction in the DNA fragmentation mediated by the higher intake of the aforementioned fat-soluble antioxidants [116].

The effectiveness of the administration of vitamin E only in the treatment of male infertility is, unfortunately, particularly limited. In this regard, Ener et al. [117] conducted a study on 45 varicocele-affected patients who underwent sub-inguinal varicocelectomy. After surgery, patients were randomized into two groups, one of which (n = 22) received daily oral vitamin E capsules (2 × 300 mg) for 12 months, while the other (n = 23) served as the control group and did not receive any supplementation. Besides the positive effects caused by the operational procedure to remove varicocele (increase in both total sperm count and motile spermatozoa), compared to the control group, the authors failed to show any additional amelioration attributable to vitamin E administration, including the pregnancy rate [117]. In a study conducted on 134

subjects, no differences in seminal plasma levels of α -tocopherols in the different groups of participants, divided according to their sperm analysis alterations, were found. However, the authors reported significantly decreased concentrations of y-tocopherol only in the group of varicocele patients, underlying the importance of this isoform in a detoxifying varicocelerelated increase of RNS ^[107]. Additionally, no significant differences in any parameter of the sperm analysis after the supplementation of 800 IU of α -tocopherol for 3 months to ten infertile patients (selected from a larger cohort of 134 sperm donors, including infertile patients with both sperm analysis anomalies and varicocele) were observed. However, the authors showed an increase in blood α -tocopherol after 45 days (+88%), mirrored by a decrease in y-tocopherol (-33%) after 45 days of α -tocopherol supplementations. At the same time point, semen α -tocopherol increased by 38%, while no changes were found in y-tocopherol concentrations. The authors speculated that decrease in y-tocopherol levels after α -tocopherol supplementation could be due to the hepatic α -TTP, which preferentially incorporates α -tocopherol into the plasma, as well as increasing hepatic catabolism of y-tocopherol. Moreover, authors suggested that the decrease in y-tocopherol levels after α -tocopherol supplementation could be potentially deleterious and that oral supplements of vitamin E should contain a combination of both tocopherols, to avoid the replacement of y-tocopherol by α -tocopherol in lipid membranes, resulting in decreased blood plasma y-tocopherol levels ^[107].

Two independent studies evaluated the effect of the concomitant administration of vitamin E (as an antioxidant) and clomiphene citrate (as an antiestrogen) in patients with idiopathic oligoasthenozoospermia. The first (uncontrolled) study reported that after daily treatment for six months with vitamin E (400 mg) and clomiphene citrate (25 mg), idiopathic oligoasthenozoospermic patients (n = 30) had a significant improvement in the mean sperm concentration and in the mean total sperm motility ^[118]. The second (controlled) study was conducted on 60 infertile men with idiopathic oligoasthenozoospermia treated with the same administration protocol (400 mg/day of vitamin E and 25 mg day of clomiphene citrate for 6 months). A significant increase in sperm count and progressive sperm motility were again recorded. Interestingly, the improvement of semen parameters was mirrored by a significant increase in the pregnancy rate recorded at the end of the study (13.3% in the control and 36.7% in the group receiving drug treatment), corroborating the conclusion that the combination of an antioxidant and antiestrogen-based therapy is a valid option for the treatment of a selected group of men with unexplained isolated oligoasthenozoospermia ^[119].

As previously underlined, most of the studies focused their attention on the effectiveness of different vitamin E-containing antioxidant mixtures [83][84][85][86][87][88][89][103], thus rendering it difficult (if not impossible) to separate the effects of tocopherols from those of the other administered antioxidants. In this context, in a prospective, open-labeled, nonrandomized study conducted by Lipovac et al. [120], 299 subfertile males with at least one recent pathological semen analysis were treated for 3 months either with 500 mg L-carnitine/twice a day (n = 156) or with the combination of 440 mg L-carnitine + 250 mg L-arginine + 40 mg zinc +120 mg vitamin E + 80 mg GSH + 60 µg selenium + 15 mg coenzyme Q₁₀ + 800 µg folic acid/once a day (n = 143). The authors reported that both groups, after the administration period, experienced a significant improvement in all studied sperm parameters (volume, density, overall progressive motility); however, relative changes of sperm density and overall progressive motility (including fast motility) were found to be higher in the group receiving the combined micronutrient treatment. In a different study, Gvozdjáková et al. [121] enrolled 40 infertile men with oligoasthenozoospermia who were treated with a combination of 440 mg L-carnitine fumarate + 30 mg ubiquinol + 75 IU vitamin E + 12 mg vitamin C twice a day for the first 3 months and once a day for the remaining 3 months of treatment. The authors reported a significant increase in the concentrations of total coenzyme Q_{10} (ubiquinone + ubiguinol) and α -tocopherol in blood and seminal plasma of treated patients. Furthermore, a significant increase in sperm density, a decrease of the sperm anomalies and a high pregnancy rate (45%) were measured after treatment [121]. Encouraging results were also obtained by Moslemi et al. [122] in a study performed on 690 infertile males affected by idiopathic asthenoteratozoospermia, who received daily oral supplementation of 400 IU of synthetic α -tocopherol (as a protector of cellular membranes from oxidative/nitrosative stress) in combination with 200 µg of L-selenomethionine (with the aim to increase the levels of the ROS scavenging enzyme Se-dependent GPx) for 14 weeks. After treatment, significant amelioration of sperm analysis parameters (sperm motility and morphology) occurred in 52.6% (362 cases) of patients, while 253 of them (36.6%) were unresponsive to the therapy. However, the authors reported that in 10.8% of patients (75 cases), spontaneous pregnancy was recorded. On the basis of these results, it was concluded that the combined administration of vitamin E and Se in cases of asthenoteratozoospermia is a potentially useful therapy not only to improve the quality of semen parameters but, most importantly, also to increase the probabilities to achieve normal pregnancy [122]. A summary of the clinical studies concerning the effects of α -tocopherol on male infertility is provided in Table 3.

Table 3. Summary of the main results obtained in the 7 papers focusing on the effect of α -tocopherol administration for the treatment of male infertility.

Authors	Patients (Number and Type of Infertility)	Controls (Number)	Antioxidant(s) (Type(s), Doses and Administration Protocol)	Duration of Treatment (Months)	Improvement (Sperm Parameters, Pregnancy and Live Birth Rates)	No Improvement (Sperm Parameters, Pregnancy and Live Birth Rates)
Zerbinati C. et al. ^[107]	10 = sperm analysis anomalies and varicocele	No controls	α-Tocopherol (800 IU)/day	3		All parameters (data not shown)
Ener K. et al. [<u>117</u>]	22 = varicocele	23	Vitamin E (2×300 mg)/day post- varicocelectomy	12		Total sperm count, motile spermatozoa and no different pregnancy rates
El Sheikh MG. et al. [118]	60 = idiopathic OA	30	Vitamin E (400 mg) and clomiphene citrate (25 mg)/day	6	Means of sperm concentration and total sperm motility	
Ghanem H. et al. ^[119]	60 = idiopathic OA	30	Vitamin E (400 mg) and clomiphene citrate (25 mg)/day	6	Sperm count, progressive sperm motility and 36.7% pregnancy rate in treated patients vs. 13.3% in controls	Total sperm motility, percentage of abnormal forms and semen volume
Lipovac M. et al. ^[120]	143 = at least one recent (1 year) pathological semen analysis result	156	Vitamin E (120 mg), L-carnitine (440 mg) L-arginine (250 mg) zinc (40 mg), GSH (80 mg), selenium (4 µg), CoQ ₁₀ (15 mg) and folic acid (800 µg)/day), , 60 ₃	Volume, density, overall progressive motility (including slow and fast motility) and% of sperm with normal morphology	

Gvozdjáková et al. ^[121]	40 = OA	No controls	Vitamin E (75 IU), L- carnitine fumarate (440 mg), ubiquinol (30 mg) and AA (12 mg) twice a day for 3 months + once a day for 3 months	6	Sperm density, decrease of the sperm anomalies and 45% pregnancy rate	
Moslemi et al. ^[122]	690 = idiopathic AT	No controls	Vitamin E (400 IU) and Se (200 µg)/day	3.5	Sperm motility, morphology (52.6% cases) and 10.8% pregnancy rate	Sperm motility and morphology (36.6% cases)

 CoQ_{10} = oxidized coenzyme Q_{10} ; asthenozoospermia = A; asthenoteratozoospermia = AT; oligoasthenoteratozoospermia = OAT; oligoasthenozoospermia = OA; oligozoospermic = O; teratozoospermia = T.

3.4. Coenzyme Q₁₀

Coenzyme Q (ubiquinone) is a fat-soluble compound ubiquitously found as a component of the mitochondrial electron transport chain (ETC) in oxygen-dependent living organisms. It is characterized by a redox-active benzoquinone head group that is conjugated to a poly-isoprenoid side-chain of species-specific length. Indeed, depending on the organism considered, the hydrophobic chain made up of isoprene units can be repeated from a minimum of 3 to a maximum of 10 units (as it occurs in humans). This last form of coenzyme Q, named coenzyme Q_{10} (Co Q_{10}), is embedded in the mitochondrial inner membrane, actively participating in the ETC. Fully oxidized CoQ₁₀ (CoQ_{100x}) accepts one couple of electrons from complex I and complex II and passing them to complex III, where the so-called Q-cycle takes place. In this cycle, during which four protons are pumped into the intermembrane space, reduced ubiquinol (CoQ10H2) is oxidized to ubiquinone (CoQ_{100x}), losing its electrons via two separate one-electron reductions. One of these electrons will serve to reduce the oxidized form of cytochrome c (anchored to a specific site of complex III protruding in the intermembrane space). The other electron left on the partially reduced radical semiquinone form of ubiquinone (CoQ^{-•}) is internally transferred to cytochromes b_L and b_H and then used to reduce one molecule of CoQ_{100x} (bound to the so-called Q_0 site of complex III) to $CoQ^{-\bullet}$ [123]. The Q-cycle, during which from two $CoQ_{10}H_2$ molecules, one molecule of CoQ_{10ox} and one of CoQ10H2, are finally obtained, represents a critical process during the ETC since electrons may leak and react with oxygen, forming superoxide and rendering the Q-cycle as one of the major sites of mitochondrial ROS production. Due to its ability to participate in one-electron transfer reactions, relative long-life, the low reactivity of its semiquinone radical, and its high hydrophobicity (clustering CoQ_{10} within biological membranes), an additional role of $CoQ_{10}H_2$ is to act as a potential free radical scavenger antioxidant. Indeed, it has been demonstrated that CoQ10 is involved in the protection against lipid peroxidation of low-density lipoproteins (LDL) (where it is oxidized before vitamin E), of mitochondria (where it counteracts the inhibitory effects of ETC substrates) and of plasma membranes (where it acts as an intermediate electron carrier) [124]. CoQ10 is considered as an endogenous antioxidant compound since its biosynthesis occurs in almost all cell types, and its abundance is not dependent on dietary supply. Due to its main localization, the final steps of CoQ₁₀ biosynthesis are undertaken within the inner mitochondrial membrane and involve many regulatory components such as mitofusin 1 and 2, two proteins implicated in the mitochondrial fusion process [124][125]. The chemical features and the main biological role as a mobile electron transporter in the mitochondrial ETC render CoQ10 strictly clustered within mitochondria and poorly abundant in human body fluids, with concentrations ranging between 0.152 to 0.939 µmol/L in human serum [58][126] and between 0.0019 to 0.013 µmol/L in human seminal plasma [58][127]. However, since spermatozoa are particularly rich in mitochondria and require a relevant amount of ATP in order to correctly satisfy their energy demand, CoQ10 seems to play an important role in the physiology of these germinal cells. Furthermore, the wellestablished implications of oxidative/nitrosative stress in the insurgence of male infertility render CoQ10 potentially involved in the pathophysiology of this condition.

In patients affected by varicocele, oligozoospermia, asthenozoospermia or idiopathic infertility, decreased levels of CoQ_{10} in seminal plasma were measured when compared to control fertile subjects ^{[127][128][129]}. Notwithstanding the complexity of the process of absorption of CoQ_{10} , caused by its chemical features and by problems linked to the efficacy of the ubiquinol—ubiquinone formulations ^[130], the supplementation with exogenous CoQ_{10} , both as a single compound and in

combination with other substances, has widely been applied in the treatment of male infertility. The most effective results were obtained when administering CoQ₁₀H₂, not only for the improvement of morpho-functional semen parameters but also for decreasing oxidative/nitrosative stress. In this scenario, Safarinejad et al. [131] conducted a randomized, doubleblind, placebo-controlled trial on 228 men with idiopathic oligoasthenoteratozoospermia, in which 50% of the patients were daily treated for 26 weeks with 200 mg of $CoQ_{10}H_2$ and the remaining 50% with placebo. The authors reported a significant improvement in sperm density, sperm motility and sperm strict morphology in the group treated with $CoQ_{10}H_2$ compared to the placebo group. Interestingly, semen parameters gradually returned to baseline values at 12 weeks after the drug-off period, even though still significant differences in sperm density and motility were recorded [131]. The administration of CoQ10H2 has further been applied in three different studies for the treatment of idiopathic oligoasthenozoospermia. In the first prospective randomized clinical trial, the effect of the administration of 200 or 400 mg/die of oral CoQ10H2 on semen parameters of 65 infertile men with idiopathic oligoasthenoteratozoospermia was tested. After three months of treatment, results showed significant amelioration of the sperm analysis (sperm count and motility) and of semen antioxidant status, with higher improvements found in response to a dose of 400 mg/day [132]. In the second case–control study, the same number of patients affected by idiopathic oligoasthenozoospermia (n = 65) were treated for three months with 200 mg/day of an unspecified form of CoQ10. The authors reported a significant improvement of sperm analysis parameters (sperm concentration, progressive motility and total motility) compared to the baseline levels recorded before the beginning of treatment. Furthermore, CoQ10 supplementation increased the total antioxidant capacity and the glutathione peroxidase levels, with a consequent reduction in ROS levels and in the sperm DNA fragmentation index [133]. Similar results concerning the amelioration in sperm concentration, motility, and antioxidant status, were also obtained in the third study, in which 35 infertile males with idiopathic oligoasthenozoospermia were administered 200 mg/day of $CoQ_{10}H_2$ [134]. In two independent studies, the effectiveness of the 200 mg/die dosage of $CoQ_{10}H_2$ was evaluated by performing the administration of the drug in two daily doses of 100 mg each. In the first retrospective study, Cakiroglu et al. [135] reported significant changes in sperm morphology and motility in 62 patients with idiopathic asthenoteratozoospermia who received 100 mg of CoQ10H2 twice a day for six months. In the second randomized, placebo-controlled study, carried out on 47 infertile males affected by idiopathic oligoasthenoteratozoospermia, Nadjarzadeh et al. investigated the effects of the abovementioned CoQ₁₀H₂ administration protocol for 3 months on the expression of catalase and SOD and on the F2-isoprostanes levels in seminal plasma [136]. Besides showing a significant positive correlation between CoQ₁₀ levels and the percent of morphologically normal spermatozoa, results indicated increased activity of the aforementioned antioxidant enzymes, accompanied by a significant decrease of F2-isoprostanes levels in seminal plasma. The authors suggested that the supplementation of $CoQ_{10}H_2$ to infertile males with oligoasthenoteratozoospermia can attenuate oxidative stress in seminal plasma by increasing antioxidant enzyme activity, ultimately improving semen parameters [136]. Surprisingly, a significant improvement in sperm forward motility, sperm density and in total antioxidant capacity was also obtained by Festa et al. $\frac{[137]}{1}$, who administered a lower dose of 50 mg CoQ₁₀H₂ twice a day, for 12 consecutive weeks, to 38 infertile males affected by varicocele. However, it is worth underlining that no pregnancies were recorded in this cohort of patients [137]. Comparing the studies cited above, the beneficial effects of CoQ10 supplementation on sperm parameters, in particular on sperm density and motility, seem to be independent of the dosage and the etiology of infertility. Safarinejad [138] conducted a study on 287 infertile men with idiopathic oligoasthenoteratozoospermia treated with 300 mg CoQ₁₀ (unspecified form) twice a day for 12 months, with a follow-up of additional 12 months after CoQ₁₀ discontinuation. The Author reported a dramatic improvement in sperm density, motility, and morphology for treatment periods longer than 6 months. At the end of the 12 months of CoQ₁₀ supplementation, 70% to 100% increases in morpho-functional sperm parameters (density, motility, and morphology) were recorded. As a consequence of the improved semen quality induced by the administration protocol, high pregnancy rates (34.1%) were also observed when the CoQ₁₀ administration time was protracted beyond the 6th month, suggesting that oligoasthenoteratozoospermia may effectively be treated by the prolonged use of CoQ₁₀ [138]

In addition to the previously mentioned studies reporting the CoQ_{10} supplementation in combination with other compounds ^{[83][86][87][88][120][121]}, Tirabassi et al. ^[139] conducted an observational study on 20 patients affected by idiopathic asthenozoospermia and daily administered with 200 mg CoQ_{10} and 2660 mg of D-aspartate for 3 months. Notwithstanding no increment of CoQ_{10} in the seminal fluid of treated patients was recorded, the authors reported a significant improvement of sperm kinetic parameters and concomitantly decreased levels of • NO, peroxynitrites and nitrosative stress-related damages to DNA after 3 months of administration ^[139]. A summary of the clinical studies concerning the effects of CoQ_{10} on male infertility is provided in Table 4.

Table 4. Summary of the main results obtained in the 9 papers focusing on the effect of CoQ_{10} administration for the treatment of male infertility.

Authors	Patients (Number and Type of Infertility)	Controls (Number)	Antioxidant(s) (Type(s), Doses and Administration Protocol)	Duration of Treatment (Months)	Improvement (Sperm Parameters, Pregnancy and Live Birth Rates)	No Improvement (Sperm Parameters, Pregnancy and Live Birth Rates)
Safarinejad et al. ^[131]	114 = idiopathic OAT	114	CoQ ₁₀ H ₂ (200 mg)/day	6.5	Sperm density, sperm motility and sperm morphology	Semen volume
Alahmar AT. [<u>132]</u>	65 = idiopathic OAT	No controls	CoQ ₁₀ H ₂ (200 mg)/day (35 patients) CoQ ₁₀ H ₂ (400 mg)/day (30 patients)	3	Sperm concentration, progressive motility and total motility (greater improvement after 400 mg/day dose)	
Alahmar AT. at al. ^[133]	65 = idiopathic OAT	40	CoQ ₁₀ (unspecified form, 200 mg)/day	3	Sperm concentration, progressive motility, total motility and DNA integrity	
Alahmar AT. et al. ^[134]	35 = idiopathic OAT	35	CoQ ₁₀ H ₂ (200 mg)/day	3	Sperm concentration and progressive motility	Sperm morphology
Cakiroglu B. et al. ^{[<u>135]</u>}	62 = idiopathic AT	No controls	CoQ ₁₀ H ₂ (100 mg) twice/day	6	Sperm morphology and motility	Sperm concentration
Nadjarzadeh A. et al. ^[126]	47 = idiopathic OAT	30	CoQ ₁₀ H ₂ (100 mg) twice/day	3	Sperm morphology	Sperm concentration and motility

Festa et al. [<u>137]</u>	38 = varicocele	No controls	CoQ ₁₀ H ₂ (50 mg) twice/day	3	Sperm forward motility and sperm density	No pregnancies
Safarinejad MR. ^[<u>1</u>38]	287 = idiopathic OAT	No controls	CoQ ₁₀ (unspecified form, 300 mg) twice/day	12	Sperm concentration, motility percentage, total sperm count, morphology and 34.1% pregnancy rate	
Tirabassi et al. ^[139]	20 = idiopathic A	No controls	CoQ ₁₀ (unspecified form, 200 mg) and D-aspartate (2660 mg)/day	3	Total sperm motility	Sperm count and morphology

 $CoQ_{10}H_2$ = reduced coenzyme Q_{10} ; CoQ_{10} = oxidized coenzyme Q_{10} ; asthenozoospermia = A; asthenoteratozoospermia = AT; oligoasthenoteratozoospermia = OA; oligozoospermia = O; teratozoospermia = T.

4. Conclusions

A summary of the data from the literature of the last ten years, concerning the effects of the administration of lowmolecular-weight antioxidants (AA, NAC, α -tocopherol and CoQ₁₀) to treat male infertility and discussed in the present review, is provided in Tables 1–4. Surprisingly, only 30 clinical studies tested the effects of the administration of AA, NAC, α -tocopherol and CoQ₁₀ (each of them alone or in combination with other antioxidants and/or vitamins and/or micronutrients) to infertile males. It is even more surprising that only 33.3% of these studies (10/30) included, as an outcome measure to determine the effects of the therapy, the pregnancy and/or live birth rates. Of these 10 studies, only 4 were case–control studies, and only 2 of them found improvement of the pregnancy rate in the group of antioxidanttreated patients compared to untreated controls. Additionally, of the 30 studies considered in this review, only 43.3% (13/30) were case–control studies, 66.7% (20/30) enrolled a number of patients higher than 40 and 40% (16/40) carried out the administration of a single antioxidant (3 AA, 2 NAC, 2 α -tocopherol and 9 CoQ₁₀).

In addition to the aforementioned critical points, the difficulty emerging in determining the real effectiveness of antioxidantbased therapies to ameliorate male infertility may certainly be due to further factors, including (i) studies using the same antioxidant(s) enrolled infertile males affected by different pathological conditions, therefore rendering impossible the comparison of the results; (ii) when considering the same antioxidant, administration protocols were characterized, in the majority of cases, by different dosages and by different supplementation times. An additional, rather crucial limitation of the studies in which antioxidant-based therapies for the treatment of male infertility were tested is represented by the fact that patients enrolled in almost all of these studies were not selected on the basis of specific deficiency of one or more antioxidants in their seminal fluid. To our opinion, before starting any antioxidant administration, it should be highly recommended (if not mandatory) to determine the full antioxidant profile in the seminal fluid of infertile patients (using analytical methods allowing the quantification of the most important water- and fat-soluble antioxidants and avoiding applying the poorly informative measurements of the so-called "total antioxidant capacity"). Only after performing a precise quantification of the seminal fluid antioxidants of every single infertile male, a precise selection of the patients and of the antioxidant(s) to administer will be possible. With this approach, the probability of recording benefits in targeted antioxidant treatments aimed to replenish the seminal fluid with the deficient or decreased antioxidant(s) should sensibly increase. Furthermore, this personalized medicine approach should decrease any possible side effects that may arise when undiscriminating supplementations with casual antioxidant cocktails are performed. The last observations are that: (i) almost no data have been published concerning the administration of carotenoids and/or flavonoids in the treatment of infertile males, notwithstanding several compounds of these two classes are currently utilized as side-therapies for the treatments of other pathological states characterized by increased oxidative/nitrosative stress; (ii) since mitochondria play a pivotal role in oxidative/nitrosative stress, it is presumable that, to obtain better results, antioxidant administration should be accompanied by selected drugs favoring mitochondrial functions and energy metabolism.

In conclusion, due to the aforementioned deficiencies showed when evaluating results of the studies using low-molecularweight antioxidant administration to infertile males, it is not currently possible to establish, beyond any doubt, either whether the antioxidant-based therapies are effective in treating male infertility or what is the best candidate antioxidant to obtain positive results, or what is the group of infertile males more susceptible to have beneficial effects from this therapeutic approach. Further case–control and longitudinal studies, enrolling a larger number of patients, using similar administration protocols (including the type of antioxidant, dosage, mode and time of administration) and enrolling patients with the same clinical diagnosis of infertility, are needed to clearly define the usefulness of antioxidant administration to treat male infertility.

References

- 1. Inhorn, M.C.; Patrizio, P. Infertility around the globe: New thinking on gender, reproductive technologies and global mov ements in the 21st century. Hum. Reprod. Update 2015, 21, 411–426.
- Zegers-Hochschild, F.; Adamson, G.D.; de Mouzon, J.; Ishihara, O.; Mansour, R.; Nygren, K.; Sullivan, E.; Vanderpoel, S. International Committee for Monitoring Assisted Reproductive Technology; World Health Organization. International Committee for Monitoring Assisted Reproductive Technology (ICMART) and the World Health Organization (WHO) revi sed glossary of ART terminology, 2009. Fertil. Steril. 2009, 92, 1520–1524.
- 3. Winters, B.R.; Walsh, T.J. The epidemiology of male infertility. Urol. Clin. N. Am. 2014, 41, 195–204.
- Katz, D.J.; Teloken, P.; Shoshany, O. Male infertility–The other side of the equation. Aust. Fam. Physician. 2017, 46, 64 1–646.
- 5. Vander Borght, M.; Wyns, C. Fertility and infertility: Definition and epidemiology. Clin. Biochem. 2018, 62, 2–10.
- Anifandis, G.; Bounartzi, T.; Messini, C.I.; Dafopoulos, K.; Sotiriou, S.; Messinis, I.E. The impact of cigarette smoking a nd alcohol consumption on sperm parameters and sperm DNA fragmentation (SDF) measured by Halosperm(®). Arch. Gynecol. Obstet. 2014, 290, 777–782.
- 7. Anifandis, G.; Amiridis, G.; Dafopoulos, K.; Daponte, A.; Dovolou, E.; Gavriil, E.; Gorgogietas, V.; Kachpani, E.; Mamuri s, Z.; Messini, C.I.; et al. The In Vitro Impact of the Herbicide Roundup on Human Sperm Motility and Sperm Mitochond ria. Toxics 2017, 6, 2.
- 8. Anifandis, G.; Katsanaki, K.; Lagodonti, G.; Messini, C.; Simopoulou, M.; Dafopoulos, K.; Daponte, A. The Effect of Gly phosate on Human Sperm Motility and Sperm DNA Fragmentation. Int. J. Environ. Res Public Health 2018, 15, 1117.
- 9. Khalili, M.A.; Leisegang, K.; Majzoub, A.; Finelli, R.; Panner Selvam, M.K.; Henkel, R.; Mojgan, M.; Agarwal, A. Male Fe rtility and the COVID-19 Pandemic: Systematic Review of the Literature. World J. Mens Health. 2020, 38, 506–520.
- 10. Anifandis, G.; Messini, C.I.; Daponte, A.; Messinis, I.E. COVID-19 and fertility: A virtual reality. Reprod. Biomed. Online 2020, 41, 157–159.
- 11. Dutta, S.; Sengupta, P. SARS-CoV-2 and Male Infertility: Possible Multifaceted Pathology. Reprod. Sci. 2020, 10, 1-4.
- 12. Mahdian, S.; Shahhoseini, M.; Moini, A. COVID-19 Mediated by Basigin Can Affect Male and Female Fertility. Int. J. Fer til. Steril. 2020, 14, 262–263.
- Szkodziak, P.; Wozniak, S.; Czuczwar, P.; Wozniakowska, E.; Milart, P.; Mroczkowski, A.; Paszkowski, T. Infertility in the light of new scientific reports—focus on male factor. Ann. Agric. Environ. Med. 2016, 23, 227–230.
- 14. Ray, A.; Shah, A.; Gudi, A.; Homburg, R. Unexplained infertility: An update and review of practice. Reprod. Biomed. Onl ine 2012, 24, 591–602.
- 15. Emokpae, M.A.; Chima, H.N. Effect of senescence on some apoptosis and oxidative stress markers in infertile normoz ospermic and oligospermic men: A cross-sectional study. Int. J. Reprod. Biomed. 2018, 16, 435–442.
- 16. Bisht, S.; Faiq, M.; Tolahunase, M.; Dada, R. Oxidative stress and male infertility. Nat. Rev. Urol. 2017, 14, 470–485.
- 17. Tremellen, K. Oxidative stress and male infertility—A clinical perspective. Hum. Reprod. Update 2008, 14, 243–258.
- 18. Martin-Hidalgo, D.; Bragado, M.J.; Batista, A.R.; Oliveira, P.F.; Alves, M.G. Antioxidants and Male Fertility: From Molecu lar Studies to Clinical Evidence. Antioxidants 2019, 8, 89.

- 19. Fujii, J.; Imai, H. Redox reactions in mammalian spermatogenesis and the potential targets of reactive oxygen species under oxidative stress. Spermatogenesis 2014, 4, e979108.
- 20. Azantee, Y.A.W.; Lokman, M.I. The future of Azoospermic Patients: In vitro spermatogenesis. Andrology 2015, 4, 2.
- 21. Walters, J.L.H.; Gadella, B.M.; Sutherland, J.M.; Nixon, B.; Bromfield, E.G. Male Infertility: Shining a Light on Lipids an d Lipid-Modulating Enzymes in the Male Germline. J. Clin. Med. 2020, 9, 327.
- 22. Oborna, I.; Wojewodka, G.; De Sanctis, J.B.; Fingerova, H.; Svobodova, M.; Brezinova, J.; Hajduch, M.; Novotny, J.; R adova, L.; Radzioch, D. Increased lipid peroxidation and abnormal fatty acid profiles in seminal and blood plasma of no rmozoospermic males from infertile couples. Hum. Reprod. 2010, 25, 308–316.
- Aitken, R.J.; Gibb, Z.; Baker, M.A.; Drevet, J.; Gharagozloo, P. Causes and consequences of oxidative stress in sperma tozoa. Reprod. Fertil. Dev. 2016, 28, 1–10.
- 24. Drevet, J.R.; Aitken, R.J. Oxidative Damage to Sperm DNA: Attack and Defense. Adv. Exp Med. Biol. 2019, 1166, 107– 117.
- 25. Homa, S.T.; Vassiliou, A.M.; Stone, J.; Killeen, A.P.; Dawkins, A.; Xie, J.; Gould, F.; Ramsay, J.W.A. A Comparison Betw een Two Assays for Measuring Seminal Oxidative Stress and their Relationship with Sperm DNA Fragmentation and S emen Parameters. Genes 2019, 10, 236.
- 26. Bui, A.D.; Sharma, R.; Henkel, R.; Agarwal, A. Reactive oxygen species impact on sperm DNA and its role in male infer tility. Andrologia 2018, 50, e13012.
- 27. Bracko, O.; Di Pietro, V.; Lazzarino, G.; Amorini, A.M.; Tavazzi, B.; Artmann, J.; Wong, E.C.; Buxton, R.B.; Weller, M.; L uft, A.R.; et al. 3-Nitropropionic acid-induced ischemia tolerance in the rat brain is mediated by reduced metabolic activi ty and cerebral blood flow. J. Cereb. Blood Flow Metab. 2014, 34, 1522–1530.
- Lazzarino, G.; Amorini, A.M.; Petzold, A.; Gasperini, C.; Ruggieri, S.; Quartuccio, M.E.; Lazzarino, G.; Di Stasio, E.; Tav azzi, B. Serum Compounds of Energy Metabolism Impairment Are Related to Disability, Disease Course and Neuroima ging in Multiple Sclerosis. Mol. Neurobiol. 2017, 54, 7520–7533.
- 29. Cadenas, S. ROS and redox signaling in myocardial ischemia-reperfusion injury and cardioprotection. Free Radic. Biol. Med. 2018, 117, 76–89.
- Kudryavtseva, A.V.; Krasnov, G.S.; Dmitriev, A.A.; Alekseev, B.Y.; Kardymon, O.L.; Sadritdinova, A.F.; Fedorova, M.S.; Pokrovsky, A.V.; Melnikova, N.V.; Kaprin, A.D.; et al. Mitochondrial dysfunction and oxidative stress in aging and cance r. Oncotarget 2016, 7, 44879–44905.
- 31. Lazzarino, G.; Amorini, A.M.; Signoretti, S.; Musumeci, G.; Lazzarino, G.; Caruso, G.; Pastore, F.S.; Di Pietro, V.; Tavaz zi, B.; Belli, A. Pyruvate Dehydrogenase and Tricarboxylic Acid Cycle Enzymes Are Sensitive Targets of Traumatic Brai n Injury Induced Metabolic Derangement. Int. J. Mol. Sci. 2019, 20, 5774.
- 32. Tibullo, D.; Giallongo, C.; Romano, A.; Vicario, N.; Barbato, A.; Puglisi, F.; Parenti, R.; Amorini, A.M.; Wissam Saab, M.; Tavazzi, B.; et al. Mitochondrial Functions, Energy Metabolism and Protein Glycosylation are Interconnected Processes Mediating Resistance to Bortezomib in Multiple Myeloma Cells. Biomolecules 2020, 10, 696.
- 33. Lazzarino, G.; Listorti, I.; Muzii, L.; Amorini, A.M.; Longo, S.; Di Stasio, E.; Caruso, G.; D'Urso, S.; Puglia, I.; Pisani, G.; et al. Low-molecular weight compounds in human seminal plasma as potential biomarkers of male infertility. Hum. Repr od. 2018, 33, 1817–1828.
- 34. Cosci, I.; Moretti, E.; Collodel, G. Lipid peroxidation in human spermatozoa from men with genitourinary infections. Sys t. Biol. Reprod. Med. 2008, 54, 75–83.
- 35. Moazamian, R.; Polhemus, A.; Connaughton, H.; Fraser, B.; Whiting, S.; Gharagozloo, P.; Aitken, R.J. Oxidative stress and human spermatozoa: Diagnostic and functional significance of aldehydes generated as a result of lipid peroxidatio n. Mol. Hum. Reprod. 2015, 21, 502–515.
- Vorilhon, S.; Brugnon, F.; Kocer, A.; Dollet, S.; Bourgne, C.; Berger, M.; Janny, L.; Pereira, B.; Aitken, R.J.; Moazamian, A.; et al. Accuracy of human sperm DNA oxidation quantification and threshold determination using an 8-OHdG immun o-detection assay. Hum. Reprod. 2018, 33, 553–562.
- 37. Kuang, F.; Liu, J.; Tang, D.; Kang, R. Oxidative Damage and Antioxidant Defense in Ferroptosis. Front. Cell Dev. Biol. 2 020, 8, 586578.
- 38. Ferramosca, A.; Zara, V. Bioenergetics of mammalian sperm capacitation. Biomed. Res. Int. 2014.
- Ribas-Maynou, J.; Yeste, M. Oxidative Stress in Male Infertility: Causes, Effects in Assisted Reproductive Techniques, a nd Protective Support of Antioxidants. Biology 2020, 9, 77.
- 40. Baskaran, S.; Finelli, R.; Agarwal, A.; Henkel, R. Reactive oxygen species in male reproduction: A boon or a bane? And rologia 2020.

- 41. Sullivan, R.; Mieusset, R. The human epididymis: Its function in sperm maturation. Hum. Reprod. Update 2016, 22, 574 –587.
- 42. Ollero, M.; Gil-Guzman, E.; Lopez, M.C.; Sharma, R.K.; Agarwal, A.; Larson, K.; Evenson, D.; Thomas, A.J., Jr.; Alvare z, J.G. Characterization of subsets of human spermatozoa at different stages of maturation: Implications in the diagnosi s and treatment of male infertility. Hum. Reprod. 2001, 16, 1912–1921.
- Aitken, J.; Krausz, C.; Buckingham, D. Relationships between biochemical markers for residual sperm cytoplasm, react ive oxygen species generation, and the presence of leukocytes and precursor germ cells in human sperm suspensions. Mol. Reprod. Dev. 1994, 39, 268–279.
- 44. Said, T.M.; Agarwal, A.; Sharma, R.K.; Mascha, E.; Sikka, S.C.; Thomas, A.J., Jr. Human sperm superoxide anion gene ration and correlation with semen quality in patients with male infertility. Fertil. Steril. 2004, 82, 871–877.
- 45. Barati, E.; Nikzad, H.; Karimian, M. Oxidative stress and male infertility: Current knowledge of pathophysiology and role of antioxidant therapy in disease management. Cell Mol. Life Sci. 2020, 77, 93–113.
- 46. Hanukoglu, I. Antioxidant protective mechanisms against reactive oxygen species (ROS) generated by mitochondrial P 450 systems in steroidogenic cells. Drug Metab. Rev. 2006, 38, 171–196.
- 47. Conrad, M.; Ingold, I.; Buday, K.; Kobayashi, S.; Angeli, J.P. ROS, thiols and thiol-regulating systems in male gametoge nesis. Biochim. Biophys. Acta. 2015, 1850, 1566–1574.
- Cabrillana, M.E.; Monclus, M.L.Á.; Lancellotti, T.E.S.; Boarelli, P.V.; Vincenti, A.E.; Fornés, M.M.; Sanabria, E.A.; Forné s, M.W. Thiols of flagellar proteins are essential for progressive motility in human spermatozoa. Reprod. Fertil. Dev. 20 17, 29, 1435–1446.
- Andrade, M.B.R.; Bertolla, R.P.; Intasqui, P.; Antoniassi, M.P.; Tibaldi, D.S.; Belardin, L.B.; Spaine, D.M. Effect of orchie ctomy on sperm functional aspects and semen oxidative stress in men with testicular tumours. Andrologia 2019, 51, e1 3205.
- 50. Morielli, T.; O'Flaherty, C. Oxidative stress impairs function and increases redox protein modifications in human sperma tozoa. Reproduction 2015, 149, 113–123.
- De Lamirande, E.; O'Flaherty, C. Sperm activation: Role of reactive oxygen species and kinases. Biochim. Biophys. Act a. 2008, 1784, 106–115.
- 52. Ichikawa, T.; Oeda, T.; Ohmori, H.; Schill, W.B. Reactive oxygen species influence the acrosome reaction but not acrosi n activity in human spermatozoa. Int. J. Androl. 1999, 22, 37–42.
- Si Aitken, R.J.; Nixon, B. Sperm capacitation: A distant landscape glimpsed but unexplored. Mol. Hum. Reprod. 2013, 19, 785–793.
- 54. Flesch, F.M.; Gadella, B.M. Dynamics of the mammalian sperm plasma membrane in the process of fertilization. Biochi m. Biophys. Acta. 2000, 1469, 197–235.
- 55. Kampfer, C.; Spillner, S.; Spinnler, K.; Schwarzer, J.U.; Terradas, C.; Ponzio, R.; Puigdomenech, E.; Levalle, O.; Köhn, F.M.; Matzkin, M.E.; et al. Evidence for an adaptation in ROS scavenging systems in human testicular peritubular cells f rom infertility patients. Int. J. Androl. 2012, 35, 793–801.
- Alkan, I.; Simşek, F.; Haklar, G.; Kervancioğlu, E.; Ozveri, H.; Yalçin, S.; Akdaş, A. Reactive oxygen species production by the spermatozoa of patients with idiopathic infertility: Relationship to seminal plasma antioxidants. J. Urol. 1997, 15 7, 140–143.
- 57. Otasevic, V.; Kalezic, A.; Macanovic, B.; Jankovic, A.; Stancic, A.; Garalejic, E.; Korac, A.; Korac, B. Evaluation of the a ntioxidative enzymes in the seminal plasma of infertile men: Contribution to classic semen quality analysis. Syst. Biol. R eprod. Med. 2019, 65, 343–349.
- Lazzarino, G.; Listorti, I.; Bilotta, G.; Capozzolo, T.; Amorini, A.M.; Longo, S.; Caruso, G.; Lazzarino, G.; Tavazzi, B.; Bil otta, P. Water- and Fat-Soluble Antioxidants in Human Seminal Plasma and Serum of Fertile Males. Antioxidants 2019, 8, 96.
- 59. Rhemrev, J.P.; van Overveld, F.W.; Haenen, G.R.; Teerlink, T.; Bast, A.; Vermeiden, J.P. Quantification of the nonenzym atic fast and slow TRAP in a post-addition assay in human seminal plasma and the antioxidant contributions of various seminal compounds. J. Androl. 2000, 21, 913–920.
- 60. Smits, R.M.; Mackenzie-Proctor, R.; Yazdani, A.; Stankiewicz, M.T.; Jordan, V.; Showell, M.G. Antioxidants for male sub fertility. Cochrane Database Syst. Rev. 2019, 3, CD007411.
- 61. Lykkesfeldt, J.; Michels, A.J.; Frei, B. Vitamin C. Adv. Nutr. 2014, 5, 16–18.
- 62. Getoff, N. Vitamin C: Electron emission, free radicals and biological versatility. In Vivo 2013, 27, 565–570.
- 63. Granger, M.; Eck, P. Dietary Vitamin C in Human Health. Adv. Food Nutr. Res. 2018, 83, 281–310.

- 64. Nishikimi, M.; Yagi, K. Molecular basis for the deficiency in humans of gulonolactone oxidase, a key enzyme for ascorbi c acid biosynthesis. Am. J. Clin. Nutr. 1991, 54, 1203S–1208S.
- 65. Savini, I.; Rossi, A.; Pierro, C.; Avigliano, L.; Catani, M.V. SVCT1 and SVCT2: Key proteins for vitamin C uptake. Amino Acids. 2008, 34, 347–355.
- 66. Di Pietro, V.; Lazzarino, G.; Amorini, A.M.; Tavazzi, B.; D'Urso, S.; Longo, S.; Vagnozzi, R.; Signoretti, S.; Clementi, E.; Giardina, B.; et al. Neuroglobin expression and oxidant/antioxidant balance after graded traumatic brain injury in the ra t. Free Radic. Biol. Med. 2014, 69, 258–264.
- 67. Hornig, D. Distribution of Ascorbic Acid, Metabolites and Analogues in Man and Animals. Ann. N. Y. Acad. Sci. 1975, 25 8, 103–118.
- 68. Angulo, C.; Maldonado, R.; Pulgar, E.; Mancilla, H.; Córdova, A.; Villarroel, F.; Castro, M.A.; Concha, I.I. Vitamin C and oxidative stress in the seminiferous epithelium. Biol. Res. 2011, 44, 169–180.
- 69. Kanďár, R.; Drábková, P.; Hampl, R. The determination of ascorbic acid and uric acid in human seminal plasma using a n HPLC with UV detection. J. Chromatogr. B Analyt. Technol. Biomed. Life Sci. 2011, 879, 2834–2839.
- Guz, J.; Gackowski, D.; Foksinski, M.; Rozalski, R.; Zarakowska, E.; Siomek, A.; Szpila, A.; Kotzbach, M.; Kotzbach, R.; Olinski, R. Comparison of oxidative stress/DNA damage in semen and blood of fertile and infertile men. PLoS ONE 2013, 8, e68490.
- 71. Berger, G.K.; Smith-Harrison, L.I.; Sandlow, J.I. Sperm agglutination: Prevalence and contributory factors. Andrologia 2 019, 51, e13254.
- 72. Nsonwu-Anyanwu, C.A.; Ekong, R.E.; Offor, J.S.; Awusha, F.O.; Orji, C.O.; Umoh, I.E.; Owhorji, A.J.; Emetonjor, R.F.; Opara Usoro, A.C. Heavy metals, biomarkers of oxidative stress and changes in sperm function: A case-control study. I nt. J. Reprod. Biomed. 2019, 17, 163–174.
- 73. Dorostghoal, M.; Kazeminejad, S.R.; Shahbazian, N.; Pourmehdi, M.; Jabbari, A. Oxidative stress status and sperm DN A fragmentation in fertile and infertile men. Andrologia 2017, 49.
- 74. Yan, L.; Liu, J.; Wu, S.; Zhang, S.; Ji, G.; Gu, A. Seminal superoxide dismutase activity and its relationship with semen quality and SOD gene polymorphism. J. Assist. Reprod. Genet. 2014, 31, 549–554.
- 75. Abd-Elmoaty, M.A.; Saleh, R.; Sharma, R.; Agarwal, A. Increased levels of oxidants and reduced antioxidants in semen of infertile men with varicocele. Fertil. Steril. 2010, 94, 1531–1534.
- Tawadrous, G.A.; Aziz, A.A.; Mostafa, T. Seminal soluble fas relationship with oxidative stress in infertile men with varic ocele. Urology 2013, 82, 820–823.
- 77. Mínguez-Alarcón, L.; Mendiola, J.; López-Espín, J.J.; Sarabia-Cos, L.; Vivero-Salmerón, G.; Vioque, J.; Navarrete-Muñ oz, E.M.; Torres-Cantero, A.M. Dietary intake of antioxidant nutrients is associated with semen quality in young universi ty students. Hum. Reprod. 2012, 27, 2807–2814.
- Mendiola, J.; Torres-Cantero, A.M.; Vioque, J.; Moreno-Grau, J.M.; Ten, J.; Roca, M.; Moreno-Grau, S.; Bernabeu, R. A low intake of antioxidant nutrients is associated with poor semen quality in patients attending fertility clinics. Fertil. Steri I. 2010, 93, 1128–1133.
- 79. Eslamian, G.; Amirjannati, N.; Rashidkhani, B.; Sadeghi, M.R.; Hekmatdoost, A. Nutrient patterns and asthenozoosper mia: A case-control study. Andrologia 2017, 49.
- 80. Cyrus, A.; Kabir, A.; Goodarzi, D.; Moghimi, M. The effect of adjuvant vitamin C after varicocele surgery on sperm qualit y and quantity in infertile men: A double-blind placebo controlled clinical trial. Int. Braz J. Urol. 2015, 41, 230–238.
- 81. Rafiee, B.; Morowvat, M.H.; Rahimi-Ghalati, N. Comparing the Effectiveness of Dietary Vitamin C and Exercise Interve ntions on Fertility Parameters in Normal Obese Men. Urol. J. 2016, 13, 2635–2639.
- Vani, K.; Kurakula, M.; Syed, R.; Alharbi, K. Clinical relevance of vitamin C among lead-exposed infertile men. Genet Te st Mol. Biomark. 2012, 16, 1001–1006.
- Abad, C.; Amengual, M.J.; Gosálvez, J.; Coward, K.; Hannaoui, N.; Benet, J.; García-Peiró, A.; Prats, J. Effects of oral antioxidant treatment upon the dynamics of human sperm DNA fragmentation and subpopulations of sperm with highly degraded DNA. Andrologia 2013, 45, 211–216.
- 84. Gual-Frau, J.; Abad, C.; Amengual, M.J.; Hannaoui, N.; Checa, M.A.; Ribas-Maynou, J.; Lozano, I.; Nikolaou, A.; Benet, J.; García-Peiró, A.; et al. Oral antioxidant treatment partly improves integrity of human sperm DNA in infertile grade I v aricocele patients. Hum. Fertil. 2015, 18, 225–229.
- 85. Magdi, Y.; Darwish, E.; Elbashir, S.; Majzoub, A.; Agarwal, A. Effect of modifiable lifestyle factors and antioxidant treatm ent on semen parameters of men with severe oligoasthenoteratozoospermia. Andrologia 2017, 49.

- 86. Kobori, Y.; Ota, S.; Sato, R.; Yagi, H.; Soh, S.; Arai, G.; Okada, H. Antioxidant cosupplementation therapy with vitamin C, vitamin E, and coenzyme Q10 in patients with oligoasthenozoospermia. Arch. Ital. Urol. Androl. 2014, 86, 1–4.
- 87. Terai, K.; Horie, S.; Fukuhara, S.; Miyagawa, Y.; Kobayashi, K.; Tsujimura, A. Combination therapy with antioxidants im proves total motile sperm counts: A Preliminary Study. Reprod. Med. Biol. 2019, 19, 89–94.
- Sadaghiani, S.; Fallahi, S.; Heshmati, H.; Teshnizi, S.H.; Chaijan, H.A.; Ebrahimi, F.F.A.; Khorrami, F.; Poorrezaeian, M.; Alizadeh, F. Effect of antioxidant supplements on sperm parameters in infertile male smokers: A single-blinded clinic al trial. AIMS Public Health 2020, 7, 92–99.
- Steiner, A.Z.; Hansen, K.R.; Barnhart, K.T.; Cedars, M.I.; Legro, R.S.; Diamond, M.P.; Krawetz, S.A.; Usadi, R.; Baker, V.L.; Coward, R.M.; et al. Reproductive Medicine Network. The effect of antioxidants on male factor infertility: The Male s, Antioxidants, and Infertility (MOXI) randomized clinical trial. Fertil. Steril. 2020, 113, 552–560.
- 90. Lu, S.C. Glutathione synthesis. Biochim. Biophys. Acta. 2013, 1830, 3143-3153.
- 91. Flohé, L. The fairytale of the GSSG/GSH redox potential. Biochim. Biophys. Acta. 2013, 1830, 3139–3142.
- 92. Brigelius-Flohé, R.; Maiorino, M. Glutathione peroxidases. Biochim. Biophys. Acta. 2013, 1830, 3289–3303.
- Couto, N.; Wood, J.; Barber, J. The role of glutathione reductase and related enzymes on cellular redox homoeostasis network. Free Radic. Biol. Med. 2016, 95, 27–42.
- 94. Llavanera, M.; Mateo-Otero, Y.; Bonet, S.; Barranco, I.; Fernández-Fuertes, B.; Yeste, M. The triple role of glutathione S-transferases in mammalian male fertility. Cell Mol. Life Sci. 2020, 77, 2331–2342.
- 95. Look, M.P.; Rockstroh, J.K.; Rao, G.S.; Kreuzer, K.A.; Barton, S.; Lemoch, H.; Sudhop, T.; Hoch, J.; Stockinger, K.; Spe ngler, U.; et al. Serum selenium, plasma glutathione (GSH) and erythrocyte glutathione peroxidase (GSH-Px)-levels in asymptomatic versus symptomatic human immunodeficiency virus-1 (HIV-1)-infection. Eur. J. Clin. Nutr. 1997, 51, 266– 272.
- 96. Atig, F.; Raffa, M.; Habib, B.A.; Kerkeni, A.; Saad, A.; Ajina, M. Impact of seminal trace element and glutathione levels o n semen quality of Tunisian infertile men. BMC Urol. 2012, 12, 6.
- 97. Moretti, E.; Collodel, G.; Fiaschi, A.I.; Micheli, L.; Iacoponi, F.; Cerretani, D. Nitric oxide, malondialdheyde and non-enzy matic antioxidants assessed in viable spermatozoa from selected infertile men. Reprod. Biol. 2017, 17, 370–375.
- 98. Kralikova, M.; Crha, I.; Huser, M.; Melounova, J.; Zakova, J.; Matejovicova, M.; Ventruba, P. The intracellular concentrat ion of homocysteine and related thiols is negatively correlated to sperm quality after highly effective method of sperm ly sis. Andrologia 2017, 49.
- Micheli, L.; Cerretani, D.; Collodel, G.; Menchiari, A.; Moltoni, L.; Fiaschi, A.I.; Moretti, E. Evaluation of enzymatic and n on-enzymatic antioxidants in seminal plasma of men with genitourinary infections, varicocele and idiopathic infertility. A ndrology 2016, 4, 456–464.
- 100. Moretti, E.; Micheli, L.; Noto, D.; Fiaschi, A.I.; Menchiari, A.; Cerretani, D. Resistin in Human Seminal Plasma: Relations hip with Lipid Peroxidation, CAT Activity, GSH/GSSG Ratio, and Semen Parameters. Oxid. Med. Cell Longev. 2019.
- 101. Jannatifar, R.; Parivar, K.; Roodbari, N.H.; Nasr-Esfahani, M.H. Effects of N-acetyl-cysteine supplementation on sperm quality, chromatin integrity and level of oxidative stress in infertile men. Reprod. Biol. Endocrinol. 2019, 17, 24.
- 102. Barekat, F.; Tavalaee, M.; Deemeh, M.R.; Bahreinian, M.; Azadi, L.; Abbasi, H.; Rozbahani, S.; Nasr-Esfahani, M.H. A P reliminary Study: N-acetyl-L-cysteine Improves Semen Quality following Varicocelectomy. Int. J. Fertil. Steril. 2016, 10, 120–126.
- 103. Dattilo, M.; Cornet, D.; Amar, E.; Cohen, M.; Menezo, Y. The importance of the one carbon cycle nutritional support in h uman male fertility: A preliminary clinical report. Reprod. Biol. Endocrinol. 2014, 12, 71.
- 104. Elsedfy, H.; De Sanctis, V.; Ahmed, A.Y.; Mohamed, N.R.; Arafa, M.; Elalfy, M.S. A pilot study on sperm DNA damage in β-thalassemia major: Is there a role for antioxidants? Acta Biomed. 2018, 89, 47–54.
- 105. Comitato, R.; Ambra, R.; Virgili, F. Tocotrienols: A Family of Molecules with Specific Biological Activities. Antioxidants 20 17, 6, 93.
- 106. Niki, E.; Traber, M.G. A history of vitamin E. Ann. Nutr. Metab. 2012, 61, 207–212.
- 107. Zerbinati, C.; Caponecchia, L.; Fiori, C.; Sebastianelli, A.; Salacone, P.; Ciacciarelli, M.; Iuliano, L. Alpha- and gamma-t ocopherol levels in human semen and their potential functional implications. Andrologia 2020, 52, e13543.
- 108. Traber, M.G. Vitamin E inadequacy in humans: Causes and consequences. Adv. Nutr. 2014, 5, 503–514.
- 109. Rossato, M.; Mariotti, C. Normal spermatogenesis and sperm function in a subject affected by cerebellar ataxia due to congenital vitamin E deficiency. Andrologia 2014, 46, 322–324.

- 110. Hosomi, A.; Arita, M.; Sato, Y.; Kiyose, C.; Ueda, T.; Igarashi, O.; Arai, H.; Inoue, K. Affinity for alpha-tocopherol transfer protein as a determinant of the biological activities of vitamin E analogs. FEBS Lett. 1997, 409, 105–108.
- 111. Bolle, P.; Evandri, M.G.; Saso, L. The controversial efficacy of vitamin E for human male infertility. Contraception 2002, 65, 313–315.
- 112. Suleiman, S.A.; Ali, M.E.; Zaki, Z.M.; el-Malik, E.M.; Nasr, M.A. Lipid peroxidation and human sperm motility: Protective role of vitamin E. J. Androl. 1996, 17, 530–537.
- 113. Benedetti, S.; Tagliamonte, M.C.; Catalani, S.; Primiterra, M.; Canestrari, F.; De Stefani, S.; Palini, S.; Bulletti, C. Differe nces in blood and semen oxidative status in fertile and infertile men, and their relationship with sperm quality. Reprod. Biomed. Online 2012, 25, 300–306.
- 114. Nadjarzadeh, A.; Mehrsai, A.; Mostafavi, E.; Gohari, M.R.; Shidfar, F. The association between dietary antioxidant intak e and semen quality in infertile men. Med. J. Islam Repub. Iran 2013, 27, 204–209.
- 115. Schmid, T.E.; Eskenazi, B.; Marchetti, F.; Young, S.; Weldon, R.H.; Baumgartner, A.; Anderson, D.; Wyrobek, A.J. Micro nutrients intake is associated with improved sperm DNA quality in older men. Fertil. Steril. 2012, 98, 1130–1137.
- 116. Salas-Huetos, A.; Rosique-Esteban, N.; Becerra-Tomás, N.; Vizmanos, B.; Bulló, M.; Salas-Salvadó, J. The Effect of N utrients and Dietary Supplements on Sperm Quality Parameters: A Systematic Review and Meta-Analysis of Randomiz ed Clinical Trials. Adv. Nutr. 2018, 9, 833–848.
- 117. Ener, K.; Aldemir, M.; Işık, E.; Okulu, E.; Özcan, M.F.; Uğurlu, M.; Tangal, S.; Özayar, A. The impact of vitamin E supple mentation on semen parameters and pregnancy rates after varicocelectomy: A randomised controlled study. Andrologia 2016, 48, 829–834.
- 118. ElSheikh, M.G.; Hosny, M.B.; Elshenoufy, A.; Elghamrawi, H.; Fayad, A.; Abdelrahman, S. Combination of vitamin E an d clomiphene citrate in treating patients with idiopathic oligoasthenozoospermia: A prospective, randomized trial. Androl ogy 2015, 3, 864–867.
- 119. Ghanem, H.; Shaeer, O.; El-Segini, A. Combination clomiphene citrate and antioxidant therapy for idiopathic male infert ility: A randomized controlled trial. Fertil. 2010, 93, 2232–2235.
- 120. Lipovac, M.; Bodner, F.; Imhof, M.; Chedraui, P. Comparison of the effect of a combination of eight micronutrients versu s a standard mono preparation on sperm parameters. Reprod. Biol. Endocrinol. 2016, 14, 84.
- 121. Gvozdjáková, A.; Kucharská, J.; Dubravicky, J.; Mojto, V.; Singh, R.B. Coenzyme Q10, α-tocopherol, and oxidative stre ss could be important metabolic biomarkers of male infertility. Dis. Markers 2015.
- 122. Moslemi, M.K.; Tavanbakhsh, S. Selenium-vitamin E supplementation in infertile men: Effects on semen parameters an d pregnancy rate. Int. J. Gen. Med. 2011, 4, 99–104.
- 123. Liu, S.S. Mitochondrial Q cycle-derived superoxide and chemiosmotic bioenergetics. Ann. N. Y. Acad. Sci. 2010, 1201, 84–95.
- 124. Wang, Y.; Hekimi, S. Understanding Ubiquinone. Trends Cell Biol. 2016, 26, 367–378.
- 125. Di Pietro, V.; Lazzarino, G.; Amorini, A.M.; Signoretti, S.; Hill, L.J.; Porto, E.; Tavazzi, B.; Lazzarino, G.; Belli, A. Fusion or Fission: The Destiny of Mitochondria in Traumatic Brain Injury of Different Severities. Sci. Rep. 2017, 7, 9189.
- 126. Niklowitz, P.; Onur, S.; Fischer, A.; Laudes, M.; Palussen, M.; Menke, T.; Döring, F. Coenzyme Q10 serum concentratio n and redox status in European adults: Influence of age, sex, and lipoprotein concentration. J. Clin. Biochem. Nutr. 201 6, 58, 240–245.
- 127. Abdul-Rasheed, O.F.; Farid, Y.Y.; Al-Nasiri, U.S. Coenzyme Q10 and oxidative stress markers in seminal plasma of Iraq i patients with male infertility. Saudi Med. J. 2010, 31, 501–506.
- 128. Mancini, A.; Balercia, G. Coenzyme Q(10) in male infertility: Physiopathology and therapy. Biofactors 2011, 37, 374–38 0.
- 129. Eroglu, M.; Sahin, S.; Durukan, B.; Ozakpinar, O.B.; Erdinc, N.; Turkgeldi, L.; Sofuoglu, K.; Karateke, A. Blood serum a nd seminal plasma selenium, total antioxidant capacity and coenzyme Q10 levels in relation to semen parameters in m en with idiopathic infertility. Biol. Trace Elem. Res. 2014, 159, 46–51.
- 130. Mantle, D.; Dybring, A. Bioavailability of Coenzyme Q(10): An Overview of the Absorption Process and Subsequent Met abolism. Antioxidants 2020, 9, 386.
- 131. Safarinejad, M.R.; Safarinejad, S.; Shafiei, N.; Safarinejad, S. Effects of the reduced form of coenzyme Q10 (ubiquinol) on semen parameters in men with idiopathic infertility: A double-blind, placebo controlled, randomized study. J. Urol. 20 12, 188, 526–531.
- 132. Alahmar, A.T. The impact of two doses of coenzyme Q10 on semen parameters and antioxidant status in men with idio pathic oligoasthenoteratozoospermia. Clin. Exp. Reprod. Med. 2019, 46, 112–118.

- 133. Alahmar, A.T.; Calogero, A.E.; Sengupta, P.; Dutta, S. Coenzyme Q10 Improves Sperm Parameters, Oxidative Stress Markers and Sperm DNA Fragmentation in Infertile Patients with Idiopathic Oligoasthenozoospermia. World J. Mens H ealth 2020.
- 134. Alahmar, A.T.; Sengupta, P. Impact of Coenzyme Q10 and Selenium on Seminal Fluid Parameters and Antioxidant Stat us in Men with Idiopathic Infertility. Biol. Trace Elem. Res. 2020.
- 135. Cakiroglu, B.; Eyyupoglu, S.E.; Gozukucuk, R.; Uyanik, B.S. Ubiquinol effect on sperm parameters in subfertile men wh o have astheno-teratozoospermia with normal sperm concentration. Nephrourol. Mon. 2014, 6, e16870.
- 136. Nadjarzadeh, A.; Shidfar, F.; Amirjannati, N.; Vafa, M.R.; Motevalian, S.A.; Gohari, M.R.; Nazeri Kakhki, S.A.; Akhondi, M.M.; Sadeghi, M.R. Effect of Coenzyme Q10 supplementation on antioxidant enzymes activity and oxidative stress of seminal plasma: A double-blind randomised clinical trial. Andrologia 2014, 46, 177–183.
- 137. Festa, R.; Giacchi, E.; Raimondo, S.; Tiano, L.; Zuccarelli, P.; Silvestrini, A.; Meucci, E.; Littarru, G.P.; Mancini, A. Coen zyme Q10 supplementation in infertile men with low-grade varicocele: An open, uncontrolled pilot study. Andrologia 201 4, 46, 805–807.
- 138. Safarinejad, M.R. The effect of coenzyme Q10 supplementation on partner pregnancy rate in infertile men with idiopath ic oligoasthenoteratozoospermia: An open-label prospective study. Int. Urol. Nephrol. 2012, 44, 689–700.
- 139. Tirabassi, G.; Vignini, A.; Tiano, L.; Buldreghini, E.; Brugè, F.; Silvestri, S.; Orlando, P.; D'Aniello, A.; Mazzanti, L.; Lenz i, A.; et al. Protective effects of coenzyme Q10 and aspartic acid on oxidative stress and DNA damage in subjects affect ed by idiopathic asthenozoospermia. Endocrine 2015, 49, 549–552.

Retrieved from https://encyclopedia.pub/entry/history/show/18303