# GRIN2B-Related Neurodevelopmental Disorder

Subjects: Genetics & Heredity Contributor: Camila Xu

GRIN2B-related neurodevelopmental disorder is a condition that affects the nervous system.

genetic conditions

## 1. Introduction

Neurodevelopmental disorders result from impaired growth and development of the central nervous system, which includes the brain and spinal cord, and the nerves connecting them. These disorders often affect learning ability, memory, and behavior and can be associated with other neurological problems.

Individuals with *GRIN2B*-related neurodevelopmental disorder have mild to profound intellectual disability and delayed development of speech and motor skills, such as sitting and walking. Some affected individuals never develop speech or the ability to walk on their own. Many people with this condition have weak muscle tone (hypotonia), which can contribute to the problems developing motor skills and lead to difficulty eating. Some affected individuals have abnormal muscle stiffness (spasticity), which can also cause problems with movement.

Recurrent seizures (epilepsy) occur in about half of people with *GRIN2B*-related neurodevelopmental disorder. About one-quarter of affected individuals have features of autism spectrum disorder, which is characterized by impaired communication and social interaction. Other behavioral problems are also possible. These individuals may be hyperactive, impulsive, or easily distractible, and some are described as being overly friendly. Sleeping difficulties can also occur in this condition.

Less common features of *GRIN2B*-related neurodevelopmental disorder include structural brain abnormalities, an unusually small head size (microcephaly), impaired vision, and involuntary muscle movements.

## 2. Frequency

The prevalence of *GRIN2B*-related neurodevelopmental disorder is unknown. Fewer than 100 cases have been reported in the medical literature.

## 3. Causes

*GRIN2B*-related neurodevelopmental disorder is caused by mutations in a gene called *GRIN2B*. This gene provides instructions for making a protein called GluN2B, which is found in nerve cells (neurons) in the brain primarily during development before birth. This protein is a part of specialized protein structures called NMDA receptors, which are involved in normal brain development, changes in the brain in response to experience (synaptic plasticity), learning, and memory.

Some *GRIN2B* gene mutations lead to production of a nonfunctional GluN2B protein or prevent the production of any GluN2B protein from one copy of the gene in each cell. A shortage of this protein may reduce the number of functional NMDA receptors, which would decrease receptor activity in cells. Other mutations lead to production of abnormal GluN2B proteins that likely alter how the NMDA receptors function; some mutations reduce NMDA receptor signaling while others increase it. Researchers are unsure how abnormal activity of NMDA receptors prevents normal growth and development of the brain or why too much or too little activity lead to similar neurological problems in people with *GRIN2B*-related neurodevelopmental disorder.

#### 3.1. The gene associated with GRIN2B-related neurodevelopmental disorder

• GRIN2B

### 4. Inheritance

This condition is inherited in an autosomal dominant pattern, which means one copy of the altered gene in each cell is sufficient to cause the disorder.

Most cases of this condition result from new (de novo) mutations in the gene that occur during the formation of reproductive cells (eggs or sperm) in an affected individual's parent or in early embryonic development. These cases occur in people with no history of the disorder in their family.

## 5. Other Names for This Condition

- EIEE27
- epileptic encephalopathy, early infantile, 27
- GRIN2B encephalopathy
- GRIN2B related syndrome

#### References

- Bell S, Maussion G, Jefri M, Peng H, Theroux JF, Silveira H, Soubannier V, Wu H, Hu P, Galat E, Torres-Platas SG, Boudreau-Pinsonneault C, O'Leary LA, Galat V,Turecki G, Durcan TM, Fon EA, Mechawar N, Ernst C. Disruption of GRIN2B ImpairsDifferentiation in Human Neurons. Stem Cell Reports. 2018 Jul 10;11(1):183-196.doi: 10.1016/j.stemcr.2018.05.018.
- Fedele L, Newcombe J, Topf M, Gibb A, Harvey RJ, Smart TG. Disease-associated missense mutations in GluN2B subunit alter NMDA receptor ligand binding and ionchannel properties. Nat Commun. 2018 Mar 6;9(1):957. doi:10.1038/s41467-018-02927-4.
- Freunscht I, Popp B, Blank R, Endele S, Moog U, Petri H, Prott EC, Reis A, Rübo J, Zabel B, Zenker M, Hebebrand J, Wieczorek D. Behavioral phenotype in five individuals with de novo mutations within the GRIN2B gene. Behav Brain Funct.2013 May 29;9:20. doi: 10.1186/1744-9081-9-20.
- Hu C, Chen W, Myers SJ, Yuan H, Traynelis SF. Human GRIN2B variants inneurodevelopmental disorders. J Pharmacol Sci. 2016 Oct;132(2):115-121. doi:10.1016/j.jphs.2016.10.002.Sci. 2017 Apr;133(4):280.
- Platzer K, Lemke JR. GRIN2B-Related Neurodevelopmental Disorder. 2018 May 31. In: Adam MP, Ardinger HH, Pagon RA, Wallace SE, Bean LJH, Stephens K, Amemiya A, editors. GeneReviews® [Internet]. Seattle (WA): University of Washington,Seattle; 1993-2020. Available from http://www.ncbi.nlm.nih.gov/books/NBK501979/
- 6. Platzer K, Yuan H, Schütz H, Winschel A, Chen W, Hu C, Kusumoto H, Heyne HO,Helbig KL, Tang S, Willing MC, Tinkle BT, Adams DJ, Depienne C, Keren B, MignotC, Frengen E, Strømme P, Biskup S, Döcker D, Strom TM, Mefford HC, Myers CT, MuirAM, LaCroix A, Sadleir L, Scheffer IE, Brilstra E, van Haelst MM, van der SmagtJJ, Bok LA, Møller RS, Jensen UB, Millichap JJ, Berg AT, Goldberg EM, De Bie I,Fox S, Major P, Jones JR, Zackai EH, Abou Jamra R, Rolfs A, Leventer RJ, LawsonJA, Roscioli T, Jansen FE, Ranza E, Korff CM, Lehesjoki AE, Courage C, LinnankiviT, Smith DR, Stanley C, Mintz M, McKnight D, Decker A, Tan WH, Tarnopolsky MA,Brady LI, Wolff M, Dondit L, Pedro HF, Parisotto SE, Jones KL, Patel AD, FranzDN, Vanzo R, Marco E, Ranells JD, Di Donato N, Dobyns WB, Laube B, Traynelis SF, Lemke JR. GRIN2B encephalopathy: novel findings on phenotype, variant clustering,functional consequences and treatment aspects. J Med Genet. 2017Jul;54(7):460-470. doi: 10.1136/jmedgenet-2016-104509.
- Swanger SA, Chen W, Wells G, Burger PB, Tankovic A, Bhattacharya S, Strong KL, Hu C, Kusumoto H, Zhang J, Adams DR, Millichap JJ, Petrovski S, Traynelis SF, Yuan H. Mechanistic Insight into NMDA Receptor Dysregulation by Rare Variants in the GluN2A and GluN2B Agonist Binding Domains. Am J Hum Genet. 2016 Dec1;99(6):1261-1280. doi: 10.1016/j.ajhg.2016.10.002.

Retrieved from https://encyclopedia.pub/entry/history/show/11472